A cullin-RING ubiquitin ligase promotes thermotolerance as part of
the Intracellular Pathogen Response in *C. elegans.*

Johan Panek, Spencer S. Gang, Kirthi C. Reddy, Robert J. Luallen, Amitkumar Fulzele, Eric J.
Bennett, Emily R. Troemel.

Division of Biological Sciences, Section of Cell and Developmental Biology, University of
California, San Diego, La Jolla, CA 92093 USA.

Corresponding author:

Emily R. Troemel

Address: UC San Diego, 9500 Gilman Dr, #0349, 4205 Bonner Hall, La Jolla, CA 92093-0349

Office phone: 858-246-0708

Email: etroemel@ucsd.edu

https://orcid.org/0000-0002-6026-9875

https://orcid.org/0000-0003-2422-0473

Classification

Major: Biological Sciences

Minor: Immunology and Inflammation.

Keywords

Proteostasis, Intracellular Pathogen Response, heat shock, *C. elegans*, cullin-RING ubiquitin
ligase complex.

Author Contributions

JP SG KR RL AF EB ET

Designed research JP, SG, EB, ET; Performed research JP, SG, AF; Contributed new reagents
or analytic tools SG, RL, KR, ET; Analyzed data JP, AF, SG; Wrote the paper JP, SG, AF, ET

This PDF file includes:

Main Text
Figures 1 to 6
31

Abstract

Intracellular pathogen infection leads to proteotoxic stress in host organisms. Previously we described a physiological program in the nematode *C. elegans* called the Intracellular Pathogen Response (IPR), which promotes resistance to proteotoxic stress and appears to be distinct from canonical proteostasis pathways. The IPR is controlled by PALS-22 and PALS-25, proteins of unknown biochemical function, which regulate expression of genes induced by natural intracellular pathogens. We previously showed that PALS-22 and PALS-25 regulate the mRNA expression of the predicted ubiquitin ligase component cullin *cul-6*, which promotes thermotolerance in *pals-22* mutants. However, it was unclear whether CUL-6 acted alone, or together with other ubiquitin ligase components. Here we use co-immunoprecipitation studies paired with genetic analysis to define the cullin-RING ligase components that act together with CUL-6 to promote thermotolerance. First, we identify a previously uncharacterized RING domain protein in the TRIM family we named RCS-1, which acts as a core component with CUL-6 to promote thermotolerance. Next, we show that the Skp-related proteins SKR-3, SKR-4 and SKR-5 act redundantly to promote thermotolerance with CUL-6. Finally, we screened F-box proteins that co-immunoprecipitate with CUL-6 and find that FBXA-158 promotes thermotolerance. In summary, we have defined the three core components and an F-box adaptor of a cullin-RING ligase complex that promotes thermotolerance as part of the IPR in *C. elegans*, which adds to our understanding of how organisms cope with proteotoxic stress.

Significance Statement

Intracellular pathogen infection in the nematode *Caenorhabditis elegans* induces a robust transcriptional response as the host copes with infection. This response program includes several ubiquitin ligase components that are predicted to function in protein quality control. In this study, we show that these infection-induced ubiquitin ligase components form a protein complex that promotes increased tolerance of acute heat stress, an indicator of improved protein homeostasis capacity. These findings show that maintaining protein homeostasis may be a critical component of a multifaceted approach allowing the host to deal with stress caused by intracellular infection.

Main Text

Introduction

Maintaining protein homeostasis (proteostasis) after exposure to environmental stressors is critical for organismal survival (1). Several signaling pathways have been identified that help organisms cope with stressors that perturb proteostasis. For example, elevated temperature triggers the conserved Heat Shock Response (HSR) pathway, which helps organisms survive the toxic effects of heat (2). The HSR upregulates expression of chaperones that help with refolding of misfolded proteins, to prevent the formation of protein aggregates and restore proteostasis (1). Disruptions of proteostasis and the formation of protein aggregates in humans are associated with severe neurodegenerative and age-related diseases, such as Alzheimer's and Huntington's diseases (1, 3–5).

Pathogen infection can perturb proteostasis and several studies in the nematode *Caenorhabditis elegans* have demonstrated intriguing connections between immune responses to extracellular pathogens and canonical proteostasis pathways (6–11). More recently, examining the *C. elegans* host response to intracellular pathogens has uncovered a novel stress response pathway that promotes proteostasis (12, 13). Microsporidia are intracellular, fungal-like pathogens that are the most common cause of infection of *C. elegans* in the wild, with *Nematocida parisii* being the most commonly found microsporidian species in *C. elegans* (14). *N. parisii* replicates inside the *C. elegans* intestine, and the infection is associated with hallmarks of perturbed proteostasis in the

host, such as the formation of large ubiquitin aggregates in the intestine (12). Interestingly, the
host transcriptional response to this infection is very similar to the host transcriptional response to
another natural intracellular pathogen of the *C. elegans* intestine, the Orsay virus (12, 15, 16).
These molecularly distinct pathogens induce a common mRNA expression pattern in *C. elegans*
that we termed the "Intracellular Pathogen Response" or IPR (13).

Functional insights into the IPR came from analysis of mutants that constitutively express IPR
genes. Forward genetic screens identified two genes encoding proteins of unknown biochemical
function called *pals-22* and *pals-25* that comprise an ON/OFF switch for the IPR, with wild-type
pals-25 acting as an activator of the IPR, which is repressed by wild-type *pals-22* (13, 17).
Constitutive upregulation of IPR gene expression in *pals-22* loss-of-function mutants is
accompanied by a rewiring of *C. elegans* physiology, including increased resistance against
natural pathogens like *N. parisii* and the Orsay virus, as well as slowed growth and shortened
lifespan. *pals-22* mutants also have increased proteostasis capacity characterized by improved
thermotolerance and lowered levels of aggregated proteins (13). All of the *pals-22* mutant
phenotypes are reversed in *pals-22 pals-25* loss-of-function double mutants (17). Interestingly,
these phenotypes appear to be independent of canonical proteostasis factors (13), such as the
transcription factors HSF-1 and DAF-16, which mediate the HSR (18), and the SKN-1/Nrf2
transcription factor, which mediates the proteasomal bounceback response(17, 19). Instead, the
proteostasis phenotypes of the IPR require a cullin gene called *cul-6*, which is transcriptionally
regulated by *pals-22/pals-25* and by infection (13, 17). Cullins are components of multi-subunit E3
ubiquitin ligases, which are enzymes that catalyze transfer of ubiquitin onto substrate proteins in
order to alter their fate (20). Based on these findings we hypothesized that CUL-6-mediated
ubiquitylation of target proteins may act as a protein quality control mechanism in the IPR to
respond to proteotoxic stress.

CUL-6 belongs to the cullin-RING Ligase (CRL) superfamily, which is found throughout
eukaryotes (21). CRLs are multi-subunit enzyme complexes, a subset of which are Skp, cullin, F-
box (SCF) complexes consisting of four subunits: a RING-box/RBX protein, a Skp, a cullin and an
F-box protein, which serves as a substrate adaptor. Interestingly, the SCF class of ubiquitin
ligases appears to have undergone a significant expansion in the evolutionary lineage that gave
rise to the nematode *C. elegans* (22, 23). For example, there are around 520 F-box proteins in
the *C. elegans* genome (22), in comparison to around 68 in humans (24), 22 in *Drosophila* and 11
in *Saccharomyces cerevisiae* (25). In addition, the number of core SCF components has
increased in nematodes, with *C. elegans* having 22 Skp-related proteins in comparison to six in
Drosophila, just one in *S. cerevisiae*, and one in humans (26). The SCF components upregulated
as part of the IPR include not only *cul-6* as mentioned above, but also the Skp-related proteins
skr-3, skr-4 and skr-5, and several F-box proteins (12). If *cul-6* were functioning as part of a
ubiquitin ligase complex to promote proteostasis as part of the IPR, it should be acting with *skrs*
and other CRL components. Here we use a combination of biochemistry and genetics to describe
how CUL-6 acts together with a RING domain protein, three Skp-related proteins and an F-box
protein to promote proteostasis in *C. elegans*.

Results
122
The cullin CUL-6 acts in the intestine and pharynx to promote thermotolerance

In comparison to wild-type animals, *pals-22* loss-of-function mutants have increased
thermotolerance, which is reduced to wild-type levels in *pals-22; cul-6* double mutants (13). Our
previous results indicated that *pals-22* regulates thermotolerance in the intestine, where it also
regulates *cul-6* mRNA expression (13). Because *cul-6* is expressed in the both the intestine and
the pharynx (Fig. 1*A*), we investigated where *cul-6* acts to promote thermotolerance. We
designed tissue-specific rescue constructs with *cul-6* cDNA using the Mos1-mediated Single
Copy Insertion (MosSCI) system (27), to drive expression of GFP-tagged versions of *cul-6* using
intestinal (*vha-6p*) or pharyngeal promoters (*myo-2p*). Here we found that both the intestinal and

pharyngeal strains expressed GFP::CUL-6 with the expected tissue distribution pattern (Fig. 1*A*).
We then crossed these tissue-specific CUL-6 MosSCI transgenes into *pals-22; cul-6* double
mutants to test for rescue of thermotolerance. Here we found that expression of *cul-6* in either the
intestine or the pharynx was sufficient to increase the thermotolerance of *pals-22; cul-6* double
mutants (Fig. 1*B*).

Previous studies had only found a functional role for CUL-6 in a *pals-22* mutant background. Here
we found that overexpression of CUL-6 from a multi-copy array (CUL-6 tagged at the C-terminus
with GFP and 3xFLAG, surrounded by ~20kb endogenous regulatory region (28)) increased
thermotolerance in a wild-type animal background (Fig. 1*A* and *B*). Furthermore, we found that
either pharyngeal or intestinal expression of *cul-6* cDNA promoted thermotolerance in a wild type
background (Fig. 1*B*). Importantly, transgenic strains with *vha-6* and *myo-2* promoters driving
genes other than wild-type *cul-6* did not have increased thermotolerance (see text below, and Fig.
1*B*). Thus, increased expression of *cul-6* in a wild-type background under its own promoter, or
only in the intestine or only in the pharynx leads to increased thermotolerance.

We also generated a *myo-3p::cul-6* construct to determine whether expression in body wall
muscle could promote thermotolerance. Here, we failed to recover *myo-3p::cul-6* transgenic
animals after several rounds of germline injections, suggesting expression of CUL-6 in muscles
may be lethal. To quantify this effect, we injected either red fluorescent protein markers together
with *myo-3p::cul-6*, or red fluorescent protein markers alone (see Materials and Methods). Here
we found none of the eggs expressing red fluorescent protein markers hatched when co-injected
with *myo-3p::cul-6* (0/87), while more than half of the eggs hatched when injected with the red
fluorescent protein markers alone (48/84). These results suggest that ectopic expression of CUL-
6 in muscles is toxic.

The activity of cullin-RING ubiquitin ligases can be increased by neddylation, which is the process
of conjugating the ubiquitin-like protein Nedd8 onto a cullin protein at a conserved lysine residue
(29). To determine whether CUL-6 might be regulated by neddylation, we mutated the lysine
residue that would likely be targeted for neddylation into an arginine, which would be predicted to
disrupt neddylation (Fig. S1) (30). We used the MosSCI technique to generate a strain that
contains this *vha-6p::cul-6(K673R)* transgene (expression visualized in Fig. 1*A*) and found that it
could not rescue the thermotolerance of *pals-22*; *cul-6* mutants (Fig. 1*B*). Furthermore, unlike
vha-6p::cul-6(wild-type), the *vha-6p::cul-6(K673R)* transgene did not promote thermotolerance in
a wild-type background. These results suggest that CUL-6 requires neddylation to promote
thermotolerance, which is consistent with CUL-6 acting as part of a ubiquitin ligase complex.

Co-immunoprecipitation/mass spectrometry analysis identifies CUL-6 binding partners

Next we performed co-immunoprecipitation mass spectrometry (co-IP/MS) analysis to identify
binding partners of CUL-6 (Dataset S1). Here we used the *C. elegans* strain with GFP::3xFLAG-
tagged CUL-6, which is functional for thermotolerance (Fig. 1*A*). We also used similar
GFP::3xFLAG-tagged strains for PALS-22 and PALS-25 (13, 17). Through analysis of binding
partners for PALS-22 and PALS-25, we sought to obtain insight into their biochemical function,
which is currently unknown. Two proteins, GFP::3xFLAG alone and an unrelated protein
F42A10.5::GFP::3xFLAG, were added as controls for the co-IPs. For each strain used for co-
IP/MS analysis we confirmed transgene expression using immunoblotting and microscopy. When
we treated animals with the proteasome inhibitor Bortezomib to induce proteotoxic stress and IPR
gene expression, we saw an increase in CUL-6 expression by both Western and microscopy
analysis, as expected from previous studies (12, 17)(Fig. S2*A* and *B*). CUL-6 expression was
seen most strongly in the pharynx and the anterior-most intestinal cells. PALS-22 and PALS-25
were broadly expressed throughout animal and, consistent with previous studies, their expression
was not affected by IPR activation (Fig. S2*A* and *B*) (17).

Co-IP/MS of PALS-22 identified 23 binding partners, including PALS-25 as one of the most highly
enriched binding partners, as compared to co-IP/MS of control proteins (Fig. S3*A*). PALS-22 also
physically associated with PALS-23, which is a PALS protein of unknown function, as well as with
F26F2.1, which is a protein of unknown function previously shown to be induced by intracellular
infection (12). Co-IP/MS of PALS-25 identified 7 binding partners, with PALS-22 being the most
highly enriched hit when compared with co-IP/MS of either control protein (Fig. S3*B*). These
reciprocal co-IP results suggest that PALS-22 and PALS-25 are in a physical complex together.

Co-IP/MS of CUL-6 identified 26 significant binding partners. These proteins included predicted
SCF ubiquitin ligase components, such as the Skp-related protein SKR-3 and the F-box protein
FBXA-158 (Fig. 2*A* and *B*). Additionally, 6 subunits of the 26S proteasome were identified (RPT-
3, 4 and RPN-5, 6.1, 8, 9). An SCF ubiquitin ligase complex canonically contains an RBX RING
box protein, which interacts with a cullin. *C. elegans* has two RBX proteins, RBX-1 and RBX-2,
but neither of these proteins were identified as significant binding partners for CUL-6 in the co-
IP/MS. Instead, we identified a single RING domain protein, C28G1.5. Because of results
described below, we renamed C28G1.5 as <u>R</u>ING protein acting with <u>c</u>ullin and <u>S</u>kr proteins (RCS-
1). Interestingly, *rcs-1* mRNA expression, like *cul-6* mRNA expression, is higher in *pals-22*
mutants when compared to either wild-type animals (log2 FC=2.81, adjusted p-value= 4.83E-08),
or compared to *pals-22 pals-25* mutants (log2 FC=2.10, adjusted p-value=3.898E-07) (17).

RING domain protein RCS-1 acts with CUL-6 to promote thermotolerance in pals-22 mutants

The *rcs-1* gene had no previously described role in *C. elegans* and has two isoforms: *rcs-1b* has
a RING domain and a B-box domain, and *rcs-1a* has the RING domain only. The closest potential
homolog of *rcs-1* in humans is the tripartite motif-containing protein 23 (TRIM23). The TRIM
family is named for having three motifs (RING finger, B box domain, and coiled coil domain), and
many TRIM proteins have E3 ubiquitin ligase activity (31). Phylogenetic analysis of the full-length
RCS-1 (RCS-1B) indicated that it is part of the *C. elegans* TRIM protein family (Fig. 3*A*) (32), but
several *C. elegans* proteins like the ADP-Ribosylation Factor related proteins (ARF-3 and ARF-6)
are more closely related to TRIM23 than RCS-1. To determine the expression pattern of RCS-1
we generated transgenic *C. elegans* containing a 3xFLAG and GFP tagged version of RCS-1 as
a multi copy array under the control of its endogenous regulatory region (28). The resulting strain
expressed GFP throughout the intestine of the worms, with particularly strong expression in the
anterior-most intestinal cells (Fig. 3*B*), where CUL-6 is also expressed.

Next we used CRISPR-Cas9 to generate two independent deletion alleles of *rcs-1* (*jy84* and
jy105), which we crossed into *pals-22* mutants (Fig. 3*C*). We found that, for both *rcs-1* alleles,
pals-22; rcs-1 double mutants had thermotolerance similar to wild-type animals, indicating that
rcs-1 is required for the increased thermotolerance of *pals-22* mutants (Fig. 3*D* and *E*). Similar to
cul-6, *rcs-1* mutations had no effect in a wild-type background (Fig. 3*D* and *E*). If RCS-1 were
acting in a SCF together with CUL-6, then loss of *rcs-1* would not further lower thermotolerance in
a *pals-22; cul-6* mutants. Indeed, we found that *pals-22; cul-6; rcs-1* triple mutants had a similar
level of thermotolerance to *pals-22; cul-6* mutants, as well as to *pals-22; rcs-1* mutants and wild-
type animals. These findings, together with CUL-6 co-IP results and RCS-1::GFP expression
pattern, are consistent with RCS-1 being the RING domain protein that acts with CUL-6 in a
ubiquitin ligase complex.

SKP-related proteins SKR-3, SKR-4 and SKR-5 act redundantly to promote thermotolerance in pals-22 mutants

In addition to a cullin and a RING protein, SCF ubiquitin ligase complexes contain a Skp protein.
Expression of three Skp-related genes (*skr-3, skr-4 and skr-5*) is upregulated by both intracellular
infection and mutation of *pals-22*, similar to *cul-6* expression (12, 13). We previously found that
mutation of either *skr-3, skr-4* or *skr-5* alone in a *pals-22* mutant background had no effect on

thermotolerance (Fig. 4*A*) (13). Therefore, we made all of the possible combinations of *skr-3*, *skr-*
4 and *skr-5* as double mutants and then crossed them into a *pals-22* mutant background to
determine whether they may act redundantly. Here we found that *pals-22; skr-3 skr-5* and *pals-*
22; skr-5 skr-4 triple mutants had a significant reduction in thermotolerance compared to *pals-22*
mutants, with levels similar to wild-type animals. In contrast, *pals-22; skr-3 skr-4* mutants had
thermotolerance similar to *pals-22* mutants (Fig. 4*B*). These results indicate that either SKR-3,
SKR-4, or SKR-5 can act together with CUL-6 in a SCF to promote thermotolerance, with SKR-5
being the most important. Consistent with the idea that SKR-5 acts together with CUL-6 and
RCS-1, we found that SKR-5::GFP::3xFLAG under control of endogenous regulatory regions is
strongly expressed in the anterior-most cells of the intestine (Fig. 4*C*).

Our results indicated than CUL-6 overexpression in the intestine was sufficient to increase
thermotolerance in a wild-type background (Fig. 1*B*). To investigate whether CUL-6 acts together
with SKR proteins in this context, we crossed the *skr-3 skr-5* double mutant into the CUL-6
overexpressing strain (*vha-6p::cul-6*). As predicted, the resulting strain had a thermotolerance
phenotype comparable to wild-type animals, consistent with CUL-6 acting together with SKR
proteins (Fig. 4*D*).

Next we sought to use RNA interference (RNAi) to further validate the redundancy of SKR
proteins acting with CUL-6. However, we found that *pals-22* mutants do not have increased
thermotolerance compared to wild-type animals when fed on the standard *E. coli* bacteria used
for feeding RNAi (HT115) (Fig. S4*A*). This effect may be due to dietary differences between
HT115 and the OP50 strain used for thermotolerance experiments described above (33).
Therefore, we tested thermotolerance with an OP50 strain (R)OP50 that was modified to enable
feeding RNAi studies (34). Here we found that *pals-22* mutant animals fed on these RNAi-
competent OP50 bacteria (transformed with empty vector L4440) have increased
thermotolerance compared to wild-type animals, which is restored back to wild-type levels in *pals-*
22; cul-6 mutants (Figure S4*A*). Using this system, we successfully knocked down expression of
cul-6 with RNAi as assessed by lowered CUL-6::GFP::3xFLAG transgene expression (Figure
S4*B* and *C*). Here we found that *cul-6* RNAi suppressed the enhanced thermotolerance of *pals-22*
mutants (Figure S4*D*). With this system, we then confirmed that *skr-3* and *skr-5* act redundantly
to promote thermotolerance, as RNAi against *skr-3* suppressed thermotolerance of *pals-22; skr-5*
double mutants but not in *skr-5* mutants (Fig. 4*E*).

Analysis of CUL-6, SKR-3,4,5 and RCS-1 in other phenotypes mediated by pals-22

Previous analysis of IPR genes indicated that RNAi knock-down of *cul-6*, *skr-3* or *skr-5* increased
susceptibility to intracellular infection in a sterile but otherwise wild-type background *C. elegans*
strain (12). Here we investigated whether *cul-6* mutants, *skr-3 skr-5* or *skr-4 skr-5* double
mutants, or *rcs-1* mutants could suppress the increased pathogen resistance of *pals-22* mutants
to microsporidia. In contrast to the complete suppression we found for increased thermotolerance
of *pals-22* mutants, we found only a minor suppression of pathogen resistance by these mutants
in a *pals-22* mutant background (Fig. S5). We also found that mutations in *rcs-1*, *skr-3 skr-4*, *skr-*
3 skr-5, or *skr-5 skr-4* did not suppress the slowed developmental rate in *pals-22* mutants, similar
to *cul-6* (Fig. S6). Therefore, *cul-6, rcs-1,* and *skr-3,4,5* appear to be important for executing the
thermotolerance phenotype of *pals-22* mutants, but not other phenotypes.

FBXA-158 acts with CUL-6 to promote thermotolerance in pals-22 mutants

Next we sought to identify the F-Box protein that functions as a substrate adaptor with CUL-6,
RCS-1 and SKR-3,4,5. Here we used RNAi to screen through the F-box proteins that were found
to physically associate with CUL-6 from co-IP/MS. In addition to FBXA-158 and FBXA-75 as
significant hits, there were other F-box proteins that we identified below the significance
threshold, including FBXA-54, FBXA-188 and FBXA-11. Because there were RNAi clones
available for *fbxa-158*, *fbxa-54*, *fbxa-188* and *fbxa-11* we used the (R)OP50 RNAi system to

screen these genes in a *pals-22* mutant background. When treated with RNAi, only *fbxa-158*
showed significant suppression of the increased thermotolerance phenotype (Fig. 5*A*). To further
validate the results from our initial screen we also tested *fbxa-158* RNAi in the CUL-6
overexpression strain *vha-6p::cul-6*. Here, *fbxa-158* RNAi treatment similarly suppressed the *vha-*
6p::cul-6 increased thermotolerance phenotype (Fig. 5*B*). Moreover, when treated with *fbxa-158*
RNAi, wild-type animals did not show further decreased thermotolerance, thus indicating that
FBXA-158 is not acting independently from CUL-6 (Fig. 5*B*).

To confirm our RNAi results we used CRISPR-Cas9 to generate two independent deletion alleles
of *fbxa-158* (*jy145* and *jy146*) and crossed them with *pals-22* mutants (Fig. 5*C*). When tested for
thermotolerance, both *fbxa-158* deletion alleles suppressed the increased thermotolerance of
pals-22 mutants while in wild-type animals the survival rate was unchanged (Fig. 5*D*). Similar to
cul-6, rcs-1 and *skr-3,4,5*, *fbxa-158* mRNA expression is higher in *pals-22* mutants when
compared to wild-type animals (log2 FC=2.81, adjusted p-value= 4.83E-08), and in *pals-22*
mutants compared to *pals-22 pals-25* mutants (log2 FC=6.95, adjusted p-value=1.2349E-06)
(17). *fbxa-158* expression is also enriched in the intestine (35). Together, these results indicate
that FBXA-158 acts as an F-box adaptor protein in a CUL-6/RCS-1/SKR-3,4,5 ubiquitin ligase
complex that promotes thermotolerance in *C. elegans*.

Discussion

Thermal stress is one of many types of proteotoxic stress that can impair organismal health and
survival. Here we used a combination of genetics and biochemistry to broaden our understanding
of a recently identified proteostasis pathway called the IPR, which enables animals to survive
exposure to thermal stress in a manner distinct from the canonical heat shock response (HSR).
Specifically, we demonstrate that overexpression of CUL-6/cullin alone promotes
thermotolerance, and it can act in either the intestine or the pharynx of *C. elegans*. Importantly,
we found that CUL-6 acts together with other ubiquitin ligase components, including a previously
uncharacterized RING protein we named RCS-1, as well as the Skp-related proteins SKR-3,
SKR-4, and SKR-5 and the F-box protein FBXA-158 (Fig. 6). We propose that this RCS-1/CUL-
6/SKR-3,4,5/FBXA-158 ubiquitin ligase complex is able to target proteins for ubiquitin-mediated
proteasomal degradation and that this activity is a critical part of the IPR program. Consistent with
this model, our co-IP/MS identified several proteasomal subunits that interact with CUL-6 (Fig. 2).

We also investigated protein-protein interactions of the PALS-22 and PALS-25 proteins, which
comprise an ON/OFF switch in the IPR that regulates mRNA expression of *cul-6, skr-3, skr-4,*
skr-5, rcs-1 and *fbxa-158*. Previous studies indicated that *pals-22* and *pals-25* are in the same
operon and interact genetically, and our co-IP/MS studies indicate they also interact
biochemically. Notably, the *pals* gene family has expanded in the *C. elegans* genome, with 39
genes in *C. elegans*, in comparison to only 1 *pals* gene each in mouse and human (36). Although
the divergent '*pals*' protein signature that defines PALS proteins is of unknown function, PALS-22
does have weak homology with F-box proteins (36), leading to the speculative idea that PALS
proteins function as adaptor proteins in ubiquitin ligase complexes.

Like the *pals* gene family, *C. elegans* has a greatly expanded repertoire of SCF ligase
components. This expansion has been suggested to reflect the results of an arms race against
intracellular pathogens, as SCF components are among the most rapidly diversifying genes in the
C. elegans genome (23). The most dramatically expanded class of SCF components includes
~520 F-box adaptor proteins in *C. elegans* (22). *C. elegans* also has 22 SKR proteins, in
comparison to only 1 Skp in humans, indicating there has been an expansion of core SCF
components as well. Here we identified SKR-3 as a binding partner for CUL-6, which is consistent
with previous 2-hybrid results (26). Interestingly, we found redundancy in the role of SKRs at the
phenotypic level: either SKR-3, SKR-4 or SKR-5 appear capable of acting with CUL-6 to promote
thermotolerance, with SKR-5 being the most important. We also found that FBXA-158, one of the
two F-box proteins identified as binding partners of CUL-6 by co-IP/MS acts to promote

thermotolerance. Together, with the Skp-related proteins, the F-box protein are responsible for
substrate specificity of the SCF (37). Hence the identification of FBXA-158 and SKR-3,4,5 as
member of the complex will help in future studies to identify what specific proteins are
ubiquitylated by a CUL-6 SCF.

Canonical CRLs contain an RBX protein as the RING domain protein, which interacts with both a
cullin and an E2 ubiquitin ligase (21). However, our co-IP/MS studies with CUL-6 did not identify
RBX-1 or RBX-2 as interacting partners for CUL-6, but rather identified RCS-1. Given our genetic
and biochemical results, we propose that RCS-1 plays the same role as an RBX protein would in
a canonical SCF complex (Fig. 6). RCS-1 does not have a clear human ortholog, but the closest
human protein is TRIM23 (38). Interestingly, the TRIM family contains 68 genes in humans, many
of which encode single subunit E3 ubiquitin ligases, including those that restrict viral infection,
and regulate inflammatory signaling (31). *C. elegans* has 18 TRIM proteins, and they appear to
have a simpler structure than human TRIM proteins, given that absence of additional motifs in the
C-terminal domains normally found in the majority of mammalian TRIM proteins (32). If these
other *C. elegans* TRIM proteins can act in SCF ligases like RCS-1, it suggests there may also be
an expansion of the RING core SCF components in *C. elegans*, in addition to the previously
described expansion of SKRs and adaptor proteins.

The role of CUL-6 in promoting thermotolerance was first demonstrated in *pals-22* mutants,
where there is an upregulation of *cul-6* mRNA as well as several other SCF components,
including *skr-3, skr-4, skr-5, rcs-1* and *fbxa-158* (13, 17). However, here we found that animals
overexpressing only CUL-6, without overexpression of the other components of the SCF, have
increased thermotolerance. One explanation for this result is that CUL-6 is the limiting factor in a
SCF that promotes thermotolerance in the IPR. Consistent with this idea, other components of
the complex like SKR-3,4,5 are functionally redundant for thermotolerance, so even basal
expression level might be sufficient to build a functional SCF, once CUL-6 expression increases
past a certain threshold level.

What substrate(s) is targeted by the RCS-1/CUL-6/SKR-3,4,5/FBXA-158 ubiquitin ligase
complex? It is possible that the effects of this complex are mediated through targeting a single
regulatory protein for ubiquitylation and degradation. For example, ubiquitylation of the DAF-2
insulin receptor by the ubiquitin ligase CHIP can alter DAF-2 trafficking, and it appears that
ubiquitylation of just this one target has significant effects on proteostasis in *C. elegans* (39). In
contrast, CRL2 and CRL4 complexes in humans have recently been shown to target the C-
termini of a large number of proteins for degradation as part of a newly identified protein quality
control system (40, 41), Given the modular nature of the SCF ligase family, the large number of
F-box substrate adaptor proteins in *C. elegans*, and the redundancy we found in the SKR-3,4,5
proteins, it seems possible that there are multiple substrates and adaptors used in the IPR. An
exciting possibility is that the IPR involves distinct CRLs that ubiquitylate several different targets
to improve proteostasis and tolerance against environmental stressors, including infections.
Identifying these factors will be the subject of future studies.
367
368   **Materials and Methods**
369
Cloning and generation of cul-6 tissue-specific rescue strains

A full list of strains used in this study is in Table S1. A full list of constructs used in this study is in
Table S2. To generate the *vha-6p::CUL-6* transgene (pET499), *vha-6p::SBP::3XFLAG*, *cul-6*
cDNA, and the *unc-54* 3' UTR were assembled in pCFJ150 using Gateway LR. To generate *myo-*
2p::CUL-6 (pET686) and *myo-3p::CUL-6* (pET687) constructs, the promoters *myo-2p*, *myo-3p*
and the pET499 linearized backbone without the *vha-6p* promoter were amplified by PCR from
pCFJ90, pCFJ104 and pET499, respectively and assembled by Gibson recombination (42). To
generate the *cul-6(K673R)* construct (pET688) a 86 bp single strand oligonucleotide and a
linearized pET499 backbone that had been amplified by PCR were assembled by Gibson

recombination. To generate the *spp-5p::3XFLAG::GFP* (pET555) transgene, *spp-*
5p::3XFLAG::GFP and *let-858* 3' UTR were assembled in pCFJ150 using Gateway LR. Mos1-
mediated Single Copy Insertion (MosSCI) was performed as described previously (27). Briefly,
the plasmid of interest (25 ng/µl) was injected with pCFJ601 (50 ng/µl), pMA122 (10 ng/µl), pGH8
(10 ng/µl), pCFJ90 (2.5 ng/µl), and pCFJ104 (5 ng/µl) into the EG6699 strain. Injected animals
were incubated at 25°C for 7 days and then subjected to heat shock for 2h at 34°C. After 5 h non-
Unc animals were selected and the presence of the insertion was verified by PCR and
sequencing.

Transgenic strains with TransgeneOme fosmids (Table S2) were generated as extrachromosomal
arrays (28) by injecting into *ttTi5605; unc-119(ed3)* worms (strain EG6699) and then selecting
non-Unc worms.

Lethality scoring of myo-3p::cul-6 tissue-specific rescue strains

391 To score the lethality of *myo-3p::cul-6* expression, worms were injected with a complete MosSCI
392 mix (see above) containing either *myo-3p::cul-6* as the plasmid of interest, or water. Injected
393 animals were incubated at 25°C and after 1 day, eggs expressing red fluorescence were
394 transferred onto new plate at 25°C for 24 h. The hatching ratio of transferred eggs was then
395 scored for both conditions. The assay was repeated two independent times.

Thermotolerance assays

Animals were grown on standard NGM plates at 20°C. L4 stage animals were transferred onto
fresh NGM plates seeded with OP50-1 and then subjected to heat shock for 2 h at 37.5°C. The
plates were then placed in a single layer on a benchtop at room temperature for 30 min, and then
transferred to a 20°C incubator. Then, 24 h later the survival was scored in a blinded manner.
Worms not responding to touch were scored as dead, and 30 worms were scored per plate.
Three replicate plates were scored for each strain per experiment, and each experiment was
performed at least three independent times. Statistical significance was tested using one-way
ANOVA and Tukey's HSD for *post-hoc* multiple comparisons.

Co-immunoprecipitation

Each sample for co-IP/MS was prepared in 3 independent experimental replicates. For each
sample, 200,000 synchronized L1 animals were transferred onto NGM plates and grown for 48 h
at 20°C. Bortezomib was added to reach a final concentration of 22 µM or the equivalent volume
of DMSO for the control plates. After 6 h at 20°C worms were washed off of the plates with M9,
washed twice with M9, resuspended in 500 µl of ice-cold lysis buffer (50mM HEPES, pH7.4, 1mM
EGTA, 1mM MgCl2, 100mM KCl, 1% glycerol, 0.05% NP40, 0.5mM DTT, 1x protease inhibitor
tablet) and immediately frozen dropwise in liquid N_2. Frozen pellets were ground into powder with
a pre-chilled mortar and pestle. Protein extracts were spun for 15 min 21,000g at 4°C and
supernatants were filtered on 0.45 µm filters (Whatman). Protein concentration was determined
using Pierce 660nm protein assay and adjusted to 1 µg/µl with fresh lysis buffer. 1 mg of each
sample was mixed with 25 µl of ANTI-FLAG M2 Affinity Gel (Sigma) and incubated at 4°C with
rotation (12 rpm) for 1h. The resin was washed twice with 1 ml lysis buffer, twice with 1 ml lysis
buffer for 5min, twice with 1 ml wash buffer (50mM HEPES, pH7.4, 1mM MgCl2, 100mM KCl)
and 20min with 1 ml wash buffer with rotation. The liquid was removed and the beads were then
stored at -80°C.

Trypsin digestion

The immunoprecipitated proteins bound to the beads were digested overnight in 400 ng trypsin
(Sigma, V511A) in 25 mM ammonium bicarbonate (Sigma) at 37°C. Samples were then reduced
with 1mM final concentration of Dithiothreitol (DTT, Acros Organics) for 30min and alkylated with

5mM final concentration of Iodoacetamide (IAA, MP Biomedicals, LLC) for 30min in dark. The
peptides were extracted from the beads by adding 50 µL of 5% Formic acid (Sigma). The
extraction was repeated one more time and the eluted peptides were combined. Digested
peptides were desalted using Stage-Tip, C18 peptide cleanup method. The eluates were vacuum
dried, and peptides were reconstituted in 15 µL of 5% Formic acid, 5% Acetonitrile solution for
LC-MS-MS analysis.

431 *LC-MS-MS Analysis*

Samples were analyzed in triplicate by LC-MS-MS using an EASY-nLC 1000 HPLC (Thermo
Scientific) and Q-Exactive mass spectrometer (Thermo Scientific, San Jose, CA) as described
previously (43) with the following modifications. The peptides were eluted using a 60min
acetonitrile gradient (45 min 2%-30% ACN gradient, followed by 5 min 30-60% ACN gradient, a
2min 60-95% ACN gradient, and a final 8min isocratic column equilibration step at 0% ACN) at
250nL/minute flow rate. All the gradient mobile phases contained 0.1% formic acid. The data
dependent analysis (DDA) was done using top 10 method with a positive polarity, scan range
400-1800 m/z, 70,000 resolution, and an AGC target of 3e6. A dynamic exclusion time of 20 s
was implemented and unassigned, singly charged and charge states above 6 were excluded for
the data dependent MS/MS scans. The MS2 scans were triggered with a minimum AGC target
threshold of 1e5 and with maximum injection time of 60 ms. The peptides were fragmented using
the normalized collision energy (NCE) setting of 25. Apex trigger and peptide match settings were
disabled.

445 *Peptide and Protein Identification and Quantification*

The RAW files obtained from the MS instrument were converted into mzXML format. The
SEQUEST search algorithm was used to search MS/MS spectra against the concatenated target
decoy database comprised of forward and reverse, reviewed *C. elegans* FASTA sequences from
Uniprot (downloaded on 6/8/2015) along with GFP and *E. coli* sequences appended in the same
file. The search parameters used were as follows: 20 ppm peptide mass tolerance; 0.01 Da
fragment ion tolerance; Trypsin (1 1 KR P) was set as the enzyme; maximum 2 missed cleavages
were allowed; Oxidation on methionine (15.99491 Da) and n-term acetylation (42.01056 Da) were
set as differential modifications; static modification (57.02146 Da) was set on cysteine for alkyl
modification. Peptide matches were filtered to a peptide false discovery rate (FDR) of 2% using
the linear discrimination analysis. The protein level matches were filtered at 2% FDR using the
protein sieve analysis. The spectral counts from the triplicates were then summed and used for
the data analysis.

458 *Analysis of mass spectrometry data*

Peptide spectral counts were used to calculate fold change ratio, p-value and adjusted p-value
between sample IPs and control IPs (GFP and F42A10.5) using the DEP package in R (44).
Briefly the data were filtered to keep only the peptides present in at least two replicates in one
condition. Filtered data were normalized using Variance Stabilizing Normalization. Missing values
were imputed using the MiniProb method from the DEP package by randomly selecting values
from a Gaussian distribution centered on a minimal value of the dataset. Fold change ratio and
adjusted p-values were calculated. Proteins with adjusted $P < 0.05$ and log2 fold change > 1 in
comparison with at least one of the controls were considered as significant. Any protein with a
negative log2 fold change with one control or the other (*i.e.* more affinity to the control protein
than the tested bait) was considered as non-significant.

469 *Phylogenic analysis of RCS-1*

Amino acid sequences of 15 proteins were aligned using MUSCLE (version 3.7) and trimmed with
trimAI (version 1.3) (45) using phylemon2 online platform (46). Bayesian Markov chain Monte

Carlo inference (LG + I + G + F) was performed using BEAST (version 1.10.4) (47). Analysis was
run using a Yule model tree prior, an uncorrelated relaxed clock (lognormal distribution) and a
chain length of 10 million states sampled every 1,000 iterations. Results were assessed with
Tracer (version 1.7.1), maximum clade credibility tree was built after a 25% burn-in. Posterior
probability values greater than 0.5 are marked on branch labels.

CRISPR deletions of rcs-1 and fbxa-158

To generate deletions of *rcs-1* and *fbxa-158,* a co-CRISPR strategy was used, adapted from the
IDT proposed method for *C. elegans*. To generate the ~2.18 kb deletion alleles of *rcs-1* we
designed two crRNA encompassing the whole *rcs-1* locus (crRNA1: 5'-
GTTTGTTGAAGGAAATGCACAGG-3', crRNA2: 5'-GGTTTCCTATAGCTGTGACACGG-3').
These two oligonucleotides were synthetized by IDT and used with a crRNA targeting the *dpy-10*
gene (*dpy-10* crRNA3: 5'-GCTACCATAGGCACCACGAG-3') and assembled with commercial
tracrRNA (50 μM crRNA1 and crRNA2, 25 μM *dpy-10* crRNA, and 40 μM tracrRNA). After an
annealing step for 5 min at 95°C, the resulting guide RNA mixture was added to CAS9-NLS
protein (27 μM final – ordered from QB3 Berkeley) and microinjected into N2 worms. F1 Dpy
progeny were screened by PCR, confirmed by sequencing for a deletion and positive lines were
backcrossed 4 times to the N2 background before testing. The same co-CRISPR strategy was
used to generate the ~1.65 kb deletion alleles of *fbxa-158*. Two crRNAs encompassing the whole
fbxa-158 locus (crRNA1: 5'-ATAGTCGGGTACAAAACAAATGG-3', crRNA2:
CTACTCCATCTTTAAGAACACGG-3') were used with the same *dpy-10* crRNA and assembled
as described above. Deletion positive lines were backcrossed 2 times to the N2 background
before testing.

RNA interference assays

RNA interference assays were performed using the feeding method. Overnight cultures of HT115
or OP50 strain (R)OP50 modified to enable RNAi (34) (gift from Meng Wang lab, Baylor College
of Medicine) were plated on RNAi plates (NGM plates supplemented with 5 mM IPTG, 1 mM
carbenicillin) and incubated at 20°C for 3 days. Gravid adults were transferred to these plates,
and their F1 progeny (L4 stage) were transferred onto new RNAi plates before being tested for
thermotolerance as previously described.

Western blot analysis

For each strain, 1500 synchronized L1 worms were placed on NGM plates seeded with OP50
bacteria and incubated at 20°C for 48 h. These animals were then treated with Bortezomib (22
μM final) or control DMSO for 6 h and washed off the plate with M9. Proteins were extracted in
lysis buffer (50 mM HEPES, pH7.4, 1 mM EGTA, 1 mM MgCl2, 100 mM KCl, 1% glycerol, 0.05%
NP40, 0.5 mM DTT, 1x protease inhibitor tablet) as previously described (13). Protein levels were
determined using the Pierce 660nm assay. Equal amount of proteins (5 μg) were boiled in protein
loading buffer, separated on a 5-20% SDS-PAGE precast gel (Bio-Rad) and transferred onto
PVDF membrane. Nonspecific binding was blocked using 5% nonfat dry milk in PBS-Tween
(0.1%) for 1 hour at room temperature. The membranes were incubated with primary antibodies
overnight at 4°C (mouse anti-FLAG diluted 1:1,000 and mouse anti-Tubulin diluted 1:7,500),
washed 5 times in PBS-Tween and blotted in horseradish peroxidase-conjugated secondary
antibodies at room temperature for 2 h (Goat anti-mouse diluted 1:10,000). Membranes were
then washed 5 times in PBS-Tween, treated with ECL reagent (Amersham), and imaged using a
Chemidoc XRS+ with Image Lab software (Bio-Rad).

Fluorescence microscopy

517 For *cul-6, rcs-1* and *skr-5* tissue-specific expression lines shown in Fig. 1, Fig. 3 and Fig. 4,
518 respectively, images were taken using a Zeiss LSM700 confocal microscope with 10X and 40X

objectives. For the expression analysis of GFP::3xFLAG-tagged proteins PALS-22, PALS-25, CUL-6 and F42A10.5 in Fig S2, image were taken with a Zeiss LSM700 confocal microscope with 10X objective. For RNAi knock-down in Fig. S4, L4 stage F1 progeny of ERT422 worms fed with OP50 expressing *cul-6* RNAi were anesthetized using 10 μM levamisole in M9 buffer and mounted on 2% agarose pads for imaging with a Zeiss LSM700 confocal. Using ImageJ software (version 1.52e), GFP signal in the pharynx and the first intestinal cells ring was measured, as well as three adjacent background regions. The total corrected fluorescence (TCF) was calculated as TCF = integrated density – (area of selected cell × mean fluorescence of background readings). For each condition, 30 worms were imaged. Significance was assessed with a Student's t-test.

Measurement of developmental rate

40 – 50 gravid adults were transferred onto standard 10 cm NGM plate and incubated at 20°C to lay eggs for 2h before being removed. These plates were incubated at 20°C and the proportion of eggs that hatched and developed into L4's was scored at 48 h, 64 h, and 72 h by scoring 100 animals each replicate.

Acknowledgments

We thank Vladimir Lazetic, Ivana Sfarcic, Jessica Sowa, and Eillen Tecle for comments on the manuscript. This work was supported by NIH under R01 AG052622 to ERT and EJB, and a Burroughs Wellcome Fund Investigators in the Pathogenesis of Infectious Diseases fellowship to ERT.

References

1. J. Labbadia, R. I. Morimoto, The biology of proteostasis in aging and disease. *Annu. Rev. Biochem.* **84**, 435–64 (2015).

2. R. Gomez-Pastor, E. T. Burchfiel, D. J. Thiele, Regulation of heat shock transcription factors and their roles in physiology and disease. *Nat. Rev. Mol. Cell Biol.* **19**, 4–19 (2017).

3. M. S. Hipp, S.-H. H. Park, F. U. Hartl, Proteostasis impairment in protein-misfolding and -aggregation diseases. *Trends Cell Biol.* **24**, 1–9 (2014).

4. R. C. Taylor, A. Dillin, Aging as an Event of Proteostasis Collapse. *Cold Spring Harb. Perspect. Biol.* **3**, a004440–a004440 (2011).

5. N. Kourtis, N. Tavernarakis, Cellular stress response pathways and ageing: intricate molecular relationships. *EMBO J.* **30**, 2520–2531 (2011).

6. E. J. Tillman, *et al.*, Endoplasmic Reticulum Homeostasis Is Modulated by the Forkhead Transcription Factor FKH-9 During Infection of *Caenorhabditis elegans*. *Genetics* **210**, 1329–1337 (2018).

7. V. Singh, A. Aballay, Heat-shock transcription factor (HSF)-1 pathway required for Caenorhabditis elegans immunity. **103** (2006).

8. M. W. Pellegrino, *et al.*, Mitochondrial UPR-regulated innate immunity provides resistance to pathogen infection. *Nature* **516**, 414–7 (2014).

9. A. Mohri-Shiomi, D. A. Garsin, Insulin signaling and the heat shock response modulate protein homeostasis in the Caenorhabditis elegans intestine during infection. *J. Biol. Chem.* **283**, 194–201 (2008).

10. J. Miles, R. Scherz-Shouval, P. van Oosten-Hawle, Expanding the Organismal
Proteostasis Network: Linking Systemic Stress Signaling with the Innate Immune
Response. *Trends Biochem. Sci.* (2019) https:/doi.org/10.1016/j.tibs.2019.06.009.

11. D. O'Brien, *et al.*, A PQM-1-Mediated Response Triggers Transcellular Chaperone
Signaling and Regulates Organismal Proteostasis. *Cell Rep.* **23**, 3905–3919 (2018).

12. M. Bakowski, *et al.*, Ubiquitin-Mediated Response to Microsporidia and Virus Infection in
C. elegans. *PLoS Pathog.* **10** (2014).

13. K. C. Reddy, *et al.*, An Intracellular Pathogen Response Pathway Promotes Proteostasis
In C. elegans. *Curr. Biol.* (2017) https:/doi.org/10.1016/j.cub.2017.10.009 (August 28,
2017).

14. G. Zhang, *et al.*, A Large Collection of Novel Nematode-Infecting Microsporidia and Their
Diverse Interactions with Caenorhabditis elegans and Other Related Nematodes. *PLOS*
Pathog. **12**, e1006093 (2016).

15. P. Sarkies, A. Ashe, J. Le Pen, M. A. McKie, E. A. Miska, Competition between virus-
derived and endogenous small RNAs regulates gene expression in Caenorhabditis
elegans. *Genome Res.* **23**, 1258–1270 (2013).

16. K. Chen, C. J. Franz, H. Jiang, Y. Jiang, D. Wang, An evolutionarily conserved
transcriptional response to viral infection in Caenorhabditis nematodes. *BMC Genomics*
18, 303 (2017).

17. K. C. Reddy, *et al.*, Antagonistic paralogs control a switch between growth and pathogen
resistance in C. elegans. *PLOS Pathog.* **15**, e1007528 (2019).

18. A.-L. Hsu, C. T. Murphy, C. Kenyon, Regulation of Aging and Age-Related Disease by
DAF-16 and Heat-Shock Factor. *Science (80-.).* **300**, 1142–1145 (2003).

19. X. Li, *et al.*, Specific SKN-1/NrF stress responses to perturbations in translation elongation
and proteasome activity. *PLoS Genet.* **7**, 9–11 (2011).

20. N. Zheng, N. Shabek, Ubiquitin Ligases: Structure, Function, and Regulation. *Annu. Rev.*
Biochem. **86**, 129–157 (2017).

21. M. D. Petroski, R. J. Deshaies, Function and regulation of cullin–RING ubiquitin ligases.
Nat. Rev. Mol. Cell Biol. **6**, 9–20 (2005).

22. J. H. Thomas, Adaptive evolution in two large families of ubiquitin-ligase adapters in
nematodes and plants. *Genome Res.* **16**, 1017–30 (2006).

23. E. Kipreos, Ubiquitin-mediated pathways in C. elegans. *WormBook*, 1–24 (2005).

24. J. Jin, *et al.*, Systematic analysis and nomenclature of mammalian F-box proteins. *Genes*
Dev. **18**, 2573–2580 (2004).

25. E. T. Kipreos, M. Pagano, The F-box protein family. *Genome Biol.* **1**, REVIEWS3002
(2000).

26. S. Nayak, *et al.*, The *Caenorhabditis elegans* Skp1-related gene family: diverse functions
in cell proliferation, morphogenesis, and meiosis. *Curr. Biol.* **12**, 277–87 (2002).

27. C. Frøkjær-Jensen, M. W. Davis, M. Ailion, E. M. Jorgensen, Improved Mos1-mediated

transgenesis in C. elegans. *Nat. Methods* **9**, 117–118 (2012).

28.   M. Sarov, *et al.*, A genome scale resource for in vivo tag-based protein function
exploration in C. elegans. *Cell* **150**, 855–866 (2012).

29.   D. M. Duda, *et al.*, Structural Insights into NEDD8 Activation of Cullin-RING Ligases:
Conformational Control of Conjugation. *Cell* **134**, 995–1006 (2008).

30.   K. Wu, A. Chen, Z.-Q. Pan, Conjugation of Nedd8 to CUL1 Enhances the Ability of the
ROC1-CUL1 Complex to Promote Ubiquitin Polymerization. *J. Biol. Chem.* **275**, 32317–
32324 (2000).

31.   S. Hatakeyama, TRIM Family Proteins: Roles in Autophagy, Immunity, and
Carcinogenesis. *Trends Biochem. Sci.* **42**, 297–311 (2017).

32.   M. Sardiello, S. Cairo, B. Fontanella, A. Ballabio, G. Meroni, Genomic analysis of the
TRIM family reveals two groups of genes with distinct evolutionary properties. *BMC Evol.*
Biol. **8**, 225 (2008).

33.   S. Pang, S. P. Curran, Adaptive Capacity to Bacterial Diet Modulates Aging in C. elegans.
Cell Metab. **19**, 221–231 (2014).

34.   D. A. Lynn, *et al.*, Omega-3 and -6 fatty acids allocate somatic and germline lipids to
ensure fitness during nutrient and oxidative stress in *Caenorhabditis elegans*. *Proc. Natl.*
Acad. Sci. **112**, 15378–15383 (2015).

35.   S. M. Lingala, M. G. M. Mhs. Ghany, Chromosomal clustering and GATA transcriptional
regulation of intestine-expressed genes in *C. elegans*. **25**, 289–313 (2016).

36.   E. Leyva-Díaz, *et al.*, Silencing of Repetitive DNA Is Controlled by a Member of an
Unusual *Caenorhabditis elegans* Gene Family. *Genetics* **207**, genetics.300134.2017
(2017).

37.   J. R. Skaar, J. K. Pagan, M. Pagano, Mechanisms and function of substrate recruitment
by F-box proteins. *Nat. Rev. Mol. Cell Biol.* **14**, 369–81 (2013).

38.   D. M. Dawidziak, J. G. Sanchez, J. M. Wagner, B. K. Ganser-Pornillos, O. Pornillos,
Structure and catalytic activation of the TRIM23 RING E3 ubiquitin ligase. *Proteins* **85**,
1957–1961 (2017).

39.   R. Tawo, *et al.*, The Ubiquitin Ligase CHIP Integrates Proteostasis and Aging by
Regulation of Insulin Receptor Turnover. *Cell* **169**, 470-482.e13 (2017).

40.   I. Koren, *et al.*, The Eukaryotic Proteome Is Shaped by E3 Ubiquitin Ligases Targeting C-
Terminal Degrons. *Cell* **173**, 1622-1635.e14 (2018).

41.   H.-C. Lin, *et al.*, C-Terminal End-Directed Protein Elimination by CRL2 Ubiquitin Ligases.
Mol. Cell **70**, 602-613.e3 (2018).

42.   D. G. Gibson, *et al.*, Enzymatic assembly of DNA molecules up to several hundred
kilobases. *Nat. Methods* **6**, 343–345 (2009).

43.   J. M. Gendron, *et al.*, Using the Ubiquitin-modified Proteome to Monitor Distinct and
Spatially Restricted Protein Homeostasis Dysfunction. *Mol. Cell. Proteomics* **15**, 2576–93
(2016).

44. X. Zhang, *et al.*, Proteome-wide identification of ubiquitin interactions using UbIA-MS. *Nat.*
Protoc. **13**, 530–550 (2018).

45. S. Capella-Gutierrez, J. M. Silla-Martinez, T. Gabaldon, trimAl: a tool for automated
alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**, 1972–1973
(2009).

46. R. Sanchez, *et al.*, Phylemon 2.0: a suite of web-tools for molecular evolution,
phylogenetics, phylogenomics and hypotheses testing. *Nucleic Acids Res.* **39**, W470–
W474 (2011).

47. M. A. Suchard, *et al.*, Bayesian phylogenetic and phylodynamic data integration using
BEAST 1.10. *Virus Evol.* **4**, vey016 (2018).

652
653
654 **Figures and Tables**
655
Figure 1. CUL-6 expression in the intestine or in the pharynx promotes thermotolerance. *(A)*
Confocal fluorescence images of L4 or adult animals with *cul-6::GFP* transgenes driven by either
the *vha-6* or *myo-2* promoter and integrated with MosSCI, or, in the case of *cul-6p::cul-6::GFP*,
driven by the endogenous promoter and expressed from a multi-copy array (28). *(B)* Survival of
animals after 2 h of 37.5°C heat shock treatment, followed by 24 h at 20°C. Strains were tested in
triplicate experiments, with three plates per experiment, 30 animals per plate. The genotypes
myo-2p::APX::GFP and *jyIs8[pals-5p::GFP; myo-2p::mCherry]* were tested as controls for *myo-2p*
driven expression. Each dot represents a plate and different shapes represent the experimental
replicates done on different days. Mean fraction alive of the nine replicates is indicated by black
bar with errors bars as SD. *** $P < 0.001$, one-way ANOVA with Tukey's *post-hoc* multiple
comparisons test.

Figure 2. Co-immunoprecipitation mass spectrometry analysis identifies binding partners for
CUL-6. Volcano plot of proteins significantly enriched in CUL-6 IP compared to F42A10.5 IP *(A)*
or to GFP IP *(B)*. Proteins significantly more abundant compared to either of the control IP's (GFP
alone control or F42A10.5 control, at adjusted $P < 0.05$ and log2 fold change > 1) were
considered interacting proteins (Dataset S1). Gray dots indicate non-significant proteins, red dots
indicate significant proteins, green dots indicate significant SCF proteins and blue dots indicate
significant proteasome subunits.

Figure 3. RING domain protein RCS-1 (C28G1.5) promotes thermotolerance in *pals-22* mutants.
(A) Phylogenetic relationships of RCS-1 protein with TRIM23 homologs proteins (in red),
canonical RBX proteins (in black) and known TRIM proteins (in blue – all are *C. elegans* except
noted human gene). The tree was built from a protein alignment using the Bayesian MCMC
method. Posterior probabilities are indicated on the branches. *(B)* Confocal fluorescence images

of L4 animals with *rcs-1::GFP* driven by the endogenous promoter and expressed from a multi-
copy array (28). *(C) rcs-1* isoforms and exon/intron structures. Protein domains are colored in
green and yellow. *jy84* and *jy105* are deletion alleles. *(D and E)* Survival of animals after 2 h of
37.5°C heat shock treatment, followed by 24 h at 20°C. Strains were tested in triplicate
experiments, with three plates per experiment, 30 animals per plate. Each dot represents a plate,
and different shapes represent the experimental replicates done on different days. Mean fraction
alive of the nine replicates is indicated by black bar with errors bars as SD. *** *P* < 0.001, one-
way ANOVA with Tukey's *post-hoc* multiple comparisons test.

Figure 4. SKR-3, SKR-4 and SKR-5 act redundantly to promote thermotolerance in *pals-22*
mutants. *(A) skr-3, skr-4 and skr-5* gene exon/intron structure. *ok365* and *ok3068* are deletion
alleles, *gk759439* is a premature stop mutation. *(B)* Survival of animals after 2 h of 37.5°C heat
shock treatment, followed by 24 h at 20°C. *** *P* < 0.001, one-way ANOVA with Tukey's *post-hoc*
multiple comparisons test. n = 9 replicates per strain. *(C)* Confocal fluorescence images of L4
animals with *skr-5::GFP* driven by the endogenous promoter and expressed from a multi-copy
array (28). *(D)* Survival of animals after 2 h of 37.5°C heat shock treatment, followed by 24 h at
20°C. *** *P* < 0.001, ** *P* < 0.01, one-way ANOVA with Tukey's *post-hoc* multiple comparisons
test. n = 7 replicates per strain. *(E)* Survival of animals after 2 h of 37.5°C heat shock treatment,
followed by 24 h at 20°C. *skr-5* and *pals-22*; *skr-5* mutants were fed on (R)OP50 expressing
either L4440 (control vector) or *skr-3* RNAi. *** *P* < 0.001, two-way ANOVA with Sidak's multiple
comparisons test. n = 9 replicates per condition. For *(B, D* and *E)* strains were tested in triplicate
experiments, 30 animals per plate. Mean fraction alive of the replicates is indicated by black bar
with errors bars as SD. Each dot represents a plate, and different shapes represent the
experimental replicates done on different days.

Figure 5. FBXA-158 promotes thermotolerance in *pals-22* mutants. *(A)* Survival of animals after 2
h of 37.5°C heat shock treatment, followed by 24 h at 20°C. *pals-22* mutants were fed on
(R)OP50 expressing either L4440 (control vector) or RNAi for the indicated genes. *pals-22*
mutants were tested in duplicate experiments, with two plates per experiment, 30 animals per
plate. *** *P* < 0.001, ** *P* < 0.01, one-way ANOVA with Tukey's *post-hoc* multiple comparisons
test. *(B)* Survival of animals after 2 h of 37.5°C heat shock treatment, followed by 24 h at 20°C.
Wild type and *vha-6p::cul-6* animals were fed on (R)OP50 expressing either L4440 (control
vector), *cul-6* or *fbxa-158* RNAi. Strains were tested in quadruplicate experiments, with three
plates per experiment, 30 animals per plate. *** *P* < 0.001, * *P* < 0.05, two-way ANOVA with
Sidak's multiple comparisons test. *(C) fbxa-158* isoforms and exon/intron structures. *jy145* and
jy146 are deletion alleles. *(D)* Survival of animals after 2 h of 37.5°C heat shock treatment,
followed by 24 h at 20°C. *** *P* < 0.001, one-way ANOVA with Tukey's *post-hoc* multiple

comparisons test. *(A, B* and *D)* each dot represents a plate, and different shapes represent the

experimental replicates done on different days. Mean fraction alive of the replicates is indicated

by black bar with errors bars as SD.

Figure 6. Model for a RCS-1/CUL-6/SKR/FBXA-158 ubiquitin ligase that promotes proteostasis.

Figure 2

Figure 3

Figure 4

Supplementary Information for:

A cullin-RING ubiquitin ligase promotes thermotolerance as part of the Intracellular Pathogen Response in *C. elegans*.

Johan Panek, Spencer S. Gang, Kirthi C. Reddy, Robert J. Luallen, Amitkumar Fulzele, Eric J. Bennett, Emily R. Troemel.

Emily R. Troemel
Email: etroemel@ucsd.edu

This PDF file includes:

> Figures S1 to S6
> Table S1 to S2
> Legend for Dataset S1
> SI References

Other supplementary materials for this manuscript include the following:

> Dataset S1

Neddylation site

Consensus - - - - - - - - - - - - - - - - A - I V R I M K - - - - - - H
CUL-6 (*C. elegans*) T D A V Q N T V E S D R K Y E I K A C I V R I M K T R K S L T H
CUL-1 (*C. elegans*) T E N V Q K N V E E D R K S V I S A C I V R I M K T R K R V Q H
CUL-1 (*Homo sapiens*) Q E T T H K N I E E D R K L L I Q A A I V R I M K M R K V L K H
CDC53 (*Saccharomyces cerevisiae*) T A S S V D T Y D N E I V M E L S A I I V R I M K T E G K L S H

Fig. S1. Alignment of the C-terminal region of *cul-6* with other cullin genes. The conserved lysine residue targeted by neddylation is indicated by a red circle.

Fig. S2. Expression analysis of GFP::3xFLAG-tagged proteins used for Co-IP/MS studies. *(A)* Western blot analysis of total protein lysate from transgenic adult animals containing fosmid transgenes expressing GFP::3xFLAG tagged protein and treated with Bortezomib or DMSO as diluent control. Proteins were detected with anti-FLAG, and anti-tubulin antibody was used as a loading control. Expected sizes; CUL-6::GFP::3xFLAG (116 kD), PALS-22::GFP::3xFLAG (64.8 kD), PALS-25 GFP::3xFLAG (66.9 kD), F42A10.5::GFP::3xFLAG (61.3 kD), GFP-3::FLAG (34 kD). *(B)* Confocal fluorescence images of L4 animals with fosmid transgenes expressing GFP tagged proteins from endogenous promoters, after exposure to DMSO or Bortezomib (diluted in DMSO). Images are overlays of green and phase contrast channels. Scale bar is 100 µm.

Fig. S3. Co-immunoprecipitation mass spectrometry analysis identifies binding partners for PALS-22 and PALS-25. Volcano plot of proteins significantly enriched in PALS-22 *(A)* and PALS-25 *(B)* IP's compared to F42A10.5 IP or GFP IP. Proteins significantly more abundant compared to either of the control IP's (GFP alone control or F42A10.5 control, at adjusted *P* < 0.05 and log2 fold change > 1) were considered interacting proteins (Dataset S1). Gray dots indicate non-significant proteins, red dots indicate significant proteins and blue dots indicate significant proteasome subunits.

Fig. S4. Development of an RNAi system for analyzing thermotolerance in *pals-22* mutants. *(A)* Survival of animals after 2 h of 37.5°C heat shock treatment, followed by 24 h at 20°C either fed on OP50 strain (R)OP50 or HT115 *E. coli*. Each dot represents a plate, and different shapes represent the experimental replicates done on different days. Mean fraction alive of the nine replicates is indicated by black bar with errors bars as SD. *** *P* < 0.001 with Student's t-test. *(B)* Quantification of GFP signal in L4 animal expressing CUL-6::GFP grown on (R)OP50 expressing either L4440 (control vector) or *cul-6* RNAi. GFP Signal was measured with ImageJ in the pharynx and the first intestinal cells ring together with 3 adjacent background area and the Total Corrected Fluorescence (TCF) was calculated. Error bars are SD. *** *P* < 0.001 with Student's t-test. *(C)* Western blot analysis on total protein lysate from adult animals with fosmid transgenes expressing CUL-6::GFP::3XFLAG treated with control or *cul-6* RNAi (OP50). CUL-6::GFP::3XFLAG protein was detected with anti-FLAG, and anti-tubulin antibody was used as a loading control. *(D)* Survival of *pals-22* mutants after 2 h of 37.5°C heat shock treatment, followed by 24 h at 20°C. Animals were fed on (R)OP50 expressing either L4440 (control vector) or *cul-6* RNAi. Each dot represents a plate, and different shapes represent the experimental replicates done on different days. Mean fraction alive of the nine replicates is indicated by black bar with errors bars as SD. *** *P* < 0.001 with Student's t-test.

Fig. S5. Analysis of pathogen resistance for *pals-22, cul-6, rcs-1* and *skr* double mutants in a *pals-22* mutant background. *N. parisii*-specific FISH probe (red) was quantified using a COPAS Biosort machine as mean red fluorescence normalized by length of individual animals. Strains were tested in triplicate infection experiments, three plates per experiment, approximately 1200 animals per plate. Each dot represents an individual animal and different shapes represent infection experiments conducted on different days. The number of animals analyzed across three separate infections is indicated for each strain. *** *P* < 0.001, Student's t-test as compared to *pals-22* mutants. Error bars indicate mean and SD.

Fig. S6. Analysis of developmental timing for *pals-22, rcs-1* and *skr* mutants. *skr* double and triple mutants in *pals-22* background do not suppress the developmental delay of *pals-22* mutants. Percentage of animals reaching the L4 larval stage at time points after eggs were laid is indicated. Results shown are the average of 3 independent biological replicates, with 100 animals assayed in each replicate.

Table S1. List of *C. elegans* strains used in this publication.

Strain Name	Genotype (transgene or mutant allele details)	Source
N2	Wild type	Caenorhabditis Genetics Center
EG6699	*ttTi5605 II; unc-119(ed3) III*	(Frøkjær-Jensen *et al.*, 2012)
RB2266	*skr-5(ok3068) V*	Caenorhabditis Genetics Center
ERT54	*jyIs8[pals-5p::GFP; myo-2::mCherry] X*	(Bakowski *et al.*, 2014)
ERT356	*pals-22(jy1) III*	(Reddy *et al.*, 2017)
ERT365	*unc-119(ed3) III; jyEx193[pals-22::EGFP::3xFLAG, unc-119(+)]*	(Reddy *et al.*, 2017)
ERT413	*jySi21[pET555(spp-5p::strepII3xFLAG_GFP::let858 3'UTR; unc-119(+))] II; unc-119(ed3) III*	This paper
ERT422	*unc-119(ed3) III; jyEx224[cul-6::EGFP::3xFLAG, unc-119(+)]*	(Reddy *et al.*, 2017)
ERT441	*pals-22(jy1) III; cul-6(ok1614) IV*	(Reddy *et al.*, 2017)
ERT443	*jySi22[pET592(myo-2p::GFP::APX_NLS::unc-54; unc-119(+))] II; unc-119(ed3) III*	(Reddy *et al.*, 2017)
ERT465	*unc-119(ed3) III; jyEx237[pals-25::EGFP::3xFLAG, unc-119(+)]*	(Reddy *et al.*, 2019)
ERT479	*pals-22(jy3) III; skr-4(gk759439) V*	(Reddy et al., 2017)
ERT488	*unc-119(ed3) III; jyEx253[F42A10.5::EGFP::3xFLAG, unc-119(+)]*	(Reddy *et al.*, 2017)
ERT554	*pals-22(jy1) III; skr-3(ok365) V*	(Reddy *et al.*, 2017)
ERT555	*pals-22(jy1) III; skr-5(ok3068) V*	(Reddy *et al.*, 2017)
ERT571	*jySi42[pET499(vha-6p::GFP::cul-6::unc-54 3' UTR, unc-119(+))] II; unc-119(ed3) III*	This paper
ERT638	*jySi42 II; unc-119(ed3) pals-22(jy1) III; cul-6(ok1614) IV*	This paper
ERT691	*rcs-1(jy84) X*	This paper
ERT692	*pals-22(jy1) III; rcs-1(jy84) X*	This paper
ERT694	*skr-3(ok365) skr-5(ok3068) V*	This paper
ERT717	*skr-3(ok365) skr-4(gk759439) V*	This paper
ERT718	*skr-5(ok3068) skr-4(gk759439) V*	This paper
ERT720	*pals-22(jy1) III; skr-3(ok365) skr-4(gk759439) V*	This paper
ERT721	*pals-22(jy1) III; skr-3(ok365) skr-5(ok3068) V*	This paper
ERT722	*pals-22(jy1) III; cul-6(ok1614) IV; rcs-1(jy84) X*	This paper
ERT723	*rcs-1(jy105) X*	This paper
ERT724	*pals-22(jy1) III; rcs-1(jy105) X*	This paper
ERT727	*pals-22(jy1) III; skr-5(ok3068) skr-4(gk759439) V*	This paper
ERT739	*jySi45[pET686(myo-2p::GFP::cul-6::unc-54 3' UTR, unc-119(+))] II; unc-119(ed3) III*	This paper
ERT740	*jySi46[pET688(vha-6p::GFP::cul-6(K673R)::unc-54 3' UTR, unc-119(+))] II; unc-119(ed3) III*	This paper
ERT741	*jySi45 II; pals-22(jy1) unc-119(ed3) III; cul-6(ok1614) IV*	This paper
ERT742	*pals-22(jy1) III; cul-6(ok1614) IV; rcs-1(jy105) X*	This paper
ERT752	*jySi46 II; pals-22(jy1) unc-119(ed3) III; cul-6(ok1614) IV*	This paper
ERT817	*jySi42 II; unc-119(ed3) III; skr-3 (ok365) skr-5 (ok3068) V*	This paper
ERT819	*unc-119(ed3) III; jyEx286[pET711(rcs-1::EGFP::3xFLAG, unc-119(+))]*	This paper
ERT820	*unc-119(ed3) III; jyEx287[pET561(skr-5::EGFP::3xFLAG, unc-119(+))]*	This paper
ERT852	*fbxa-158(jy145) II*	This paper
ERT853	*fbxa-158(jy146) II*	This paper
ERT854	*fbxa-158(jy145) II; pals-22(jy1) III*	This paper
ERT855	*fbxa-158(jy146) II; pals-22(jy1) III*	This paper

Table S2. List of DNA constructs used in this publication.

Construct Name	Genotype (transgene or mutant allele details)	Source
pET499	*vha-6p::SBP_3xFLAG_GFP_cul-6::unc-54 3'UTR* in pCFJ150	This paper
pET555	*spp-5p::strepII3xFLAG_GFP::let-858-3UTR* in *pCFJ150*	This paper
pET686	*myo-2p::SBP_3xFLAG_GFP_cul-6::unc-54 3'UTR* in pCFJ150	This paper
pET687	*myo-3p::SBP_3xFLAG_GFP_cul-6::unc-54 3'UTR* in pCFJ150	This paper
pET688	*vha-6p::SBP_3xFLAG_GFP_cul-6(K673R)::unc-54 3'UTR* in pCFJ150	This paper
pCFJ150 - Addgene Plasmid #19329	*pDESTttTi5605[R4-R3]*	(Frøkjær-Jensen *et al.*, 2012)
pCFJ601 - Addgene Plasmid #34874	*eft-3p::Mos1* transposase	(Frøkjær-Jensen *et al.*, 2012)
pMA122 - Addgene Plasmid #34873	*peel-1* negative selection	(Frøkjær-Jensen *et al.*, 2012)
pGH8 - Addgene Plasmid #19359	*rab-3p::mCherry*	(Frøkjær-Jensen *et al.*, 2012)
pCFJ90 - Addgene Plasmid #19327	*myo-2p::mCherry*	(Frøkjær-Jensen *et al.*, 2012)
pCFJ104 - Addgene Plasmid #19328	*myo-3p::mCherry*	(Frøkjær-Jensen *et al.*, 2012)
873721959883807 G09	*cul-6::EGFP::3xFLAG* Fosmid; *unc-119* selection	(Sarov *et al.*, 2012)
9830596596427236 B12	*pals-22::EGFP::3xFLAG* Fosmid; *unc-119* selection	(Sarov *et al.*, 2012)
18995122782808704 G01	*pals-25::EGFP::3xFLAG* Fosmid; *unc-119* selection	(Sarov *et al.*, 2012)
8218370932910004 G03	*F42A10.5::EGFP::3xFLAG* Fosmid; *unc-119* selection	(Sarov *et al.*, 2012)
2491680425634929 B04	*rcs-1::EGFP::3xFLAG* Fosmid; *unc-119* selection	(Sarov *et al.*, 2012)
5202602939198744 E05	*skr-5::EGFP::3xFLAG* Fosmid; *unc-119* selection	(Sarov *et al.*, 2012)

Dataset S1 (separate file). Statistical analysis of co-IP experiments. Each tab shows the results for one experimental IP, CUL-6, PALS-22 or PALS-25. Each column indicates the fold change or adjusted p-value of the experimental IP relative to either F42A10.5 or GFP control IPs. Proteins indicated as "TRUE" were significantly more abundant in the experimental IP compared to control IPs (adjusted $P < 0.05$ and log2 fold change > 1).

References:

1. C. Frøkjær-Jensen, M. W. Davis, M. Ailion, E. M. Jorgensen, Improved Mos1-mediated transgenesis in C. elegans. Nat. Methods **9**, 117–118 (2012).

2. M. Bakowski, et al., Ubiquitin-Mediated Response to Microsporidia and Virus Infection in C. elegans. PLoS Pathog. **10** (2014).

3. K. C. Reddy, et al., An Intracellular Pathogen Response Pathway Promotes Proteostasis In C. elegans. Curr. Biol. (2017) https:/doi.org/10.1016/j.cub.2017.10.009 (August 28, 2017).

4. K. C. Reddy, et al., Antagonistic paralogs control a switch between growth and pathogen resistance in C. elegans. PLOS Pathog. **15**, e1007528 (2019).

5. M. Sarov, et al., A genome scale resource for in vivo tag-based protein function exploration in C. elegans. Cell **150**, 855–866 (2012).