Estimating and accounting for genotyping errors in RAD-seq experiments

Luisa Bresadola[1,*], Vivian Link[1,2,*], C. Alex Buerkle[3], Christian Lexer[4], Daniel Wegmann[1,2]

[1] Department of Biology, University of Fribourg, Fribourg, Switzerland

[2] Swiss Institute of Bioinformatics, Fribourg, Switzerland

[3] Department of Botany, University of Wyoming, Laramie WY, USA

[4] Department of Botany and Biodiversity Research, University of Vienna, Vienna, Austria

[*] These authors contributed equally

Abstract

In non-model organisms, evolutionary questions are frequently addressed using reduced representation sequencing techniques due to their low cost, ease of use, and because they do not require genomic resources such as a reference genome. However, evidence is accumulating that such techniques may be affected by specific biases, questioning the accuracy of obtained genotypes, and as a consequence, their usefulness in evolutionary studies. Here we introduce three strategies to estimate genotyping error rates from such data: through the comparison to high quality genotypes obtained with a different technique, from individual replicates, or from a population sample when assuming Hardy-Weinberg equilibrium. Applying these strategies to data obtained with Restriction site Associated DNA sequencing (RAD-seq), arguably the most popular reduced representation sequencing technique, revealed per-allele genotyping error rates that were much higher than sequencing error rates, particularly at heterozygous sites that were wrongly inferred as homozygous. As we exemplify through the inference of genome-wide and local ancestry of well characterized hybrids of two Eurasian poplar (*Populus*) species, such high error rates may lead to wrong biological conclusions. By properly accounting for these error rates in downstream analyses, either by incorporating genotyping errors directly or by recalibrating genotype likelihoods, we were nevertheless able to use the RAD-seq data to support biologically meaningful and robust inferences of ancestry among *Populus* hybrids. Based on these findings, we strongly recommend carefully assessing genotyping error rates in reduced representation sequencing experiments, and to properly account for these in downstream analyses, for instance using the tools presented here.

Introduction

Despite the impressive advancements in sequencing techniques and the decrease of related costs, whole genome sequencing (WGS) remains prohibitively expensive when working with a large number of samples or species with large genomes. Since many applications do not require information on the whole genome, reduced representation sequencing techniques are valuable alternatives and have become widely used for genome-wide SNP discovery and genotyping, especially in species with poor genomic resources (Andrews, Good, Miller, Luikart, & Hohenlohe, 2016; Narum, Buerkle, Davey, Miller, & Hohenlohe, 2013).

A commonly used reduced representation sequencing technique is Restriction site Associated DNA sequencing (Baird et al., 2008; Miller, Dunham, Amores, Cresko, & Johnson, 2007), which allows the

sequencing of massively multiplexed samples at minimal costs by focusing on the sequences adjacent
to restriction sites. Since restriction sites are often shared between individuals within a species and
often also between closely related species (Cariou, Duret, & Charlat, 2013), focusing on adjacent
sequences guarantees that sequenced loci are mostly overlapping across samples.

Briefly, the first step in the original RAD-seq protocol is the digestion of genomic DNA with a
restriction enzyme. The resulting fragments are ligated to an adaptor and a unique barcode for each
sample, and multiple individuals are pooled. The fragments are then sheared using a sonicator and
those showing the proper size are selected and amplified through polymerase chain reaction (PCR). At
this point the library is suitable for sequencing. By focusing the sequencing effort on tagged
restriction sites, rather than on all randomly sheared genomic fragments (Arnold, Corbett-Detig, Hartl,
& Bomblies, 2013; Rowe, Renaut, & Guggisberg, 2011), the number of markers can be customized
through the choice of restriction enzymes. This choice will also influence which features of the
genome are sampled, since certain enzymes preferentially cut in exonic regions, while others target
intergenic and intronic regions (Arnold et al., 2013; Pootakham et al., 2016).

Several alternative RAD-seq protocols allow for ample customization of this methodology. These
include the elimination of the sonication step (Andrews et al., 2016), ddRAD (Peterson, Weber, Kay,
Fisher, & Hoekstra, 2012), which uses two restriction enzymes rather than one, 2bRAD (Wang,
Meyer, McKay, & Matz, 2012), which uses IIb-type restriction enzymes and produces fragments of
36 bp, and ezRAD (Toonen et al., 2013), in which DNA is digested with isoschizomers (a pair of
restriction enzymes recognizing the same sequence). However, each method has different advantages
and pitfalls, and a specific protocol may be more suitable for some applications than for others (Puritz
et al., 2014). Due to this versatility, RAD-seq has been used in diverse applications, including the
study of the genomics of adaptation (Andrews et al., 2016), hybridization and speciation (Marques et
al., 2016), inbreeding depression (Hoffman et al., 2014), genetic associations (Nadeau et al., 2014),
genetic mapping (Chutimanitsakun et al., 2011) and phylogeographic and phylogenomic analyses
(Emerson et al., 2010; Leaché et al., 2015).

Despite this widespread use, genotypes called from RAD-seq data have been associated with several
biases, many of which are specific to RAD-seq. Several major biases potentially affect alleles
differently, which may lead to their unequal representation in sequencing data, and hence to
genotyping errors at heterozygous sites (Davey et al., 2013). For instance, polymorphisms occurring
in the restriction site may result in one allele not being cut and therefore not sequenced, potentially
causing linked sites to be erroneously called homozygous ("allele dropout"). Polymorphisms at
neighbouring restriction sites may also result in genotyping biases, for example, if the length of the
fragment of one allele falls short of the selected size range (Andrews et al., 2016). Yet size
differences among longer fragments were also found to result in unequal sequencing depth at linked
sites because sonicators shear shorter fragments less efficiently than longer fragments (Sambrook &
Russell, 2006). Finally, the PCR step present in most RAD-seq protocols may contribute to
genotyping errors through unequal amplification of the two alleles (Casbon, Osborne, Brenner, &
Lichtenstein, 2011) or through so-called PCR duplicates, the sequencing of misleading clonal copies
of the same initial molecule (Andrews & Luikart, 2014, Euclid 2019). Since many protocols produce
single-end libraries or libraries where both ends are defined by restriction sites (e.g. ddRAD), PCR
duplicates cannot be reliably identified bioinformatically unless very many different adapter
sequences are used (Schweyen, Rozenberg, & Leese, 2014). This is particularly problematic in the
case of PCR errors that might be sequenced in many copies, resulting in wrongly called heterozygous
genotypes (Andrews et al., 2016).

95
Some consequences of these biases in downstream analyses are well documented. Gautier et al.,
(2013), for instance, found that under certain circumstances allele dropout leads to incorrect estimates
of genetic diversity. Arnold et al. (2013) demonstrated through both simulations and real data that
estimates of summary statistics commonly used to infer diversity and past demography from RAD
data are severely affected by missing haplotypes and may show strong deviations from true values.
Cariou, Duret, & Charlat (2016), finally, illustrated that allele dropout can lead to underestimation of
diversity, especially in highly polymorphic species.
103
In light of these potentially common issues, our aims here were to develop strategies (1) to estimate
genotyping error rates in RAD data, and (2) to properly incorporate the resulting genotyping
uncertainty in downstream analyses to mitigate the consequences of errors. For this we present
methods to estimate genotyping errors in three different ways: First, by taking advantage of available
genotyping data based on a different, more reliable method (e.g. using a chip or high-depth
sequencing). Second, by using independent RAD-seq replicates of individuals. And third, by
assuming Hardy-Weinberg proportions among population samples. Using simulations we show that
all these methods are powerful in inferring error rates even if limited samples are available. As a case
study, we then applied these methods to RAD-seq data of the two widespread, genetically and
ecologically divergent tree species *Populus alba* (White poplar) and *P. tremula* (European aspen) and
inferred high genotyping error rates of multiple percent. By properly accounting for genotyping
uncertainty, however, we obtain biologically meaningful estimates of genome-wide and local
ancestry.
117

Materials and Methods

118
119
120 *Estimation of genotyping error rates*
121
Let us denote by g_{il} the observed genotype of individual $i = 1, ..., I$ at locus $l = 1, ..., L$, where
$g_{il} = 0,1,2$ reflects the number of copies of the alternative allele at a bi-allelic locus. Given per-allele
genotyping error rates ε_0 and ε_1 at homozygous and heterozygous sites, respectively, the probabilities
$P(g_{il}|\gamma, \varepsilon_0, \varepsilon_1)$ of observing genotype g_{il} at a locus with true genotype $\gamma = 0,1,2$ are given in Table
1. We next describe three strategies to estimate the genotyping error rates ε_0 and ε_1 from called
genotypic data. While we present these methods in the context of Restriction site Associated DNA
sequencing (RAD-seq), we stress that they are applicable to genotypic data obtained from any data
source.
130
131 **From a truth set.** Consider accurate genotypes γ_{il} obtained independently for some individuals and
132 loci. Assuming all γ_{il} to be correct and genotyping errors to be independent between sites and
133 individuals, the likelihood of the observed genotypes $\boldsymbol{g} = \{g_{11}, ..., g_{I1}, ... g_{IL}\}$ is then given by

$$P(\boldsymbol{g}|\boldsymbol{\gamma}, \varepsilon_0, \varepsilon_1) = \prod_{l=1}^{L} \prod_{i=1}^{I} P(g_{il}|\gamma_{il}, \varepsilon_0, \varepsilon_1),$$

134 where $\boldsymbol{\gamma} = \{\gamma_{11}, ..., \gamma_{I1}, ..., \gamma_{IL}\}$ and $P(g_{il}|\gamma_{il}, \varepsilon_0, \varepsilon_1)$ is given in Table 1. We obtain maximum
135 likelihood estimates $\hat{\varepsilon}_0$ and $\hat{\varepsilon}_1$ through numerical maximization (see Supporting Information).
136

137 **From individual replicates.** Consider individuals for which multiple independent sequencing
138 experiments were conducted. Let us denote by $g_{il}^{(j)}$ the inferred genotype of individual i at locus l in
139 replicate $j = 1, \dots, r_1$. The likelihood of the full data is then given by

$$P(\boldsymbol{g}|\varepsilon_0, \varepsilon_1, \boldsymbol{f}) = \prod_{l=1}^{L} \prod_{i=1}^{I} \sum_{\gamma=0}^{2} \left[P(\gamma|f_{i\gamma}) \prod_{j=1}^{r_i} P(g_{il}^{(j)}|\gamma, \varepsilon_0, \varepsilon_1) \right],$$

140 where $\boldsymbol{g}$ denotes the vector of all observed genotypes, γ the unobserved true genotype, $P(\gamma|f_{i\gamma}) = f_{i\gamma}$
141 the frequency of genotype γ among all loci of individual i, $\boldsymbol{f} = \{f_{10}, \dots, f_{I0}, \dots, f_{I2}\}$ and
142 $P(g_{il}^{(j)}|\gamma, \varepsilon_0, \varepsilon_1)$ is given in Table 1. We obtain maximum likelihood estimates $\hat{\varepsilon}_0$, $\hat{\varepsilon}_1$ and $\hat{\boldsymbol{f}}$ with an
143 Expectation-Maximization (EM) algorithm as detailed in the Supporting Information.
144
145 **From population samples.** Consider individuals sampled from a random mating population such that
146 the distribution of the true genotypes is well described by Hardy-Weinberg proportions. While the
147 locus-specific allele frequencies f_l are unknown, let us assume that they follow a Beta distribution
148 with parameters α, β such that $f_l \sim Beta(\alpha, \beta)$ as is expected under neutrality (Wright, 1931). The
149 likelihood of the full data is then given by

$$P(\boldsymbol{g}|\varepsilon_0, \varepsilon_1, \alpha, \beta) = \prod_{l=1}^{L} \int_{0}^{1} P(f_l|\alpha, \beta) \prod_{i=1}^{I} \sum_{\gamma=0}^{2} [P(g_{il}|\gamma, \varepsilon_0, \varepsilon_1) P(\gamma|f_l)] \, df_l,$$

where the sum runs over the unknown true genotype γ, $P(\gamma|f_{i\gamma})$ are the Hardy-Weinberg proportions
given allele frequencies $f_{i\gamma}$ and $P(g_{il}|\gamma, \varepsilon_0, \varepsilon_1)$ is given in Table 1. In contrast to the other two
models, the computationally efficient maximum likelihood scheme resulted in biased estimates when
working with population samples. We therefore resort to an MCMC approach under a Bayesian
scheme (see Supporting Information) with exponential priors $\varepsilon_0, \varepsilon_1 \sim \text{Exp}(\lambda)$ truncated at 0.5 and
normal priors $\log(\alpha), \log(\beta) \sim N(\mu, \sigma^2)$. We used $\lambda = 5$, $\mu = \log(0.5)$ and $\sigma^2 = 0.25$ throughout.
We also extend the model to consider data from multiple populations jointly, given that multiple
samples are available for each population (see Supporting Information).
158
Set specific error rates. All methods introduced above are readily extended to jointly infer error rates
for multiple sets of genotype calls that differ in their error rates, such as genotypes obtained with
different sequencing depth or from different batches, for example different sequencing libraries (see
Supporting Information). Inferring the error rates of all sets jointly is beneficial in the case of
individual replicates or population samples, as information about hierarchical parameters such as
individual genotype frequencies $f_i = \{f_{i0}, f_{i1}, f_{i2}\}$ or the parameters α, β of the Beta distribution are
shared across them.
166
167 **Recalibrating genotype likelihoods.** We recalibrate genotype likelihoods by determining the
168 likelihoods $P(g_{il}|\gamma, \varepsilon_0, \varepsilon_1)$ of the called genotype g_{il} for all $\gamma = 0,1,2$ according to Table 1 and using
169 parameter estimates ε_0 and ε_1 obtained for the relevant set. If a truth set is available, we also calculate
170 the empirical likelihoods $P(g_{il}|\gamma_{il} = g)$ across all genotypes $\gamma_{il} = g$ of a particular set.
171
Implementation: We implemented all algorithms developed here in the open-source C++ program
Tiger (Tools to Incorporate Genotyping ERrors), available through the git repository at
https://bitbucket.org/wegmannlab/tiger. Tiger works directly off VCF files in format 4.2, which
contains genotype calls encoded by the mandatory GT field. Optionally, the depth encoded by the
field DP is used to bin genotype calls into different sets, which may further be stratified on a per
sample basis. All methods readily handle missing data.

Simulations. We used simulations to assess the power of the methods introduced above to infer
genotyping error rates. All simulations were generated directly under the assumed models using
routines we implemented in *Tiger*. Under the truth set or replicate model, each simulated data set
consisted of 50% heterozygous genotypes. Under the population samples model we drew true allele
frequencies from a Beta distribution with parameters $\alpha = \beta = 0.7$, implying about 29% of
heterozygous genotypes.

To assess the bias in inferring error rates from a truth set obtained with shotgun sequencing data, we
simulated sequencing data for 10^4 truly heterozygous, $5 \cdot 10^3$ truly homozygous reference and
$5 \cdot 10^3$ truly homozygous alternative loci at different mean depth μ. Specifically, we drew locus
specific depths $d_l \sim \mathrm{Pois}(\mu)$ from a Poisson distribution with mean μ and then simulated the number
of observed reference alleles $r_l \sim \mathrm{Bin}(d_l, p_l)$ from a binomial distribution with probability $p_l = 1 -$
$e, 0.5, e$ for homozygous reference, heterozygous and homozygous alternative loci, respectively,
assuming a sequencing error rate $e = 0.01$. We next determined the maximum likelihood genotypes
$\hat{\gamma}_l$ under the same model and used those to infer error rates $\hat{\varepsilon}_0$ and $\hat{\varepsilon}_l$ from genotypes g_l we simulated
according to Table 1 with $\varepsilon_0 = \varepsilon_1 = 0.01$ or 0.1. We then generated 10^6 replicates for each mean
depth $\mu = 1, 2, \ldots, 30$ and quantified the biases $b_0 = (\hat{\varepsilon}_0 - \varepsilon_0)/\varepsilon_0$ and $b_1 = (\hat{\varepsilon}_1 - \varepsilon_1)/\varepsilon_1$.

Application to Populus *species*

Study system and plant material. We generated RAD-seq data of 139 individuals of the two
widespread tree species *Populus alba* and *P. tremula*, and their hybrids (*P. x canescens*) in two sets
ideally suited to test the inference of error rates from a truth set and individual replicates, respectively.
The first set consisted of 136 individuals (Supplementary Table S1) grown with minimal interference
in a common garden established at the University of Fribourg (Switzerland) and previously genotyped
with a genotyping-by-sequencing (GBS) protocol (Lindtke, Gompert, Lexer, & Buerkle, 2014). All
these individuals grew from seeds collected from 15 mother trees in a natural hybrid zone in the Parco
Lombardo della Valle del Ticino in Northern Italy where individuals of the two species and their
hybrids grow side by side (Christe et al., 2016; Lindtke et al., 2012).

The second set consisted of four individuals for which we generated multiple replicates: a hybrid
individual (F039_05) also included among the samples of the first set, a second hybrid individual
(I373_A) also from the Ticino hybrid zone but grown in a common garden in Salerno, Italy, a pure *P.*
alba individual (J1) from the Jalón river in the Ebro watershed (Northeast of the Iberian Peninsula),
an assumed F1-hybrid tree (BET) from a population in the Tajo river headwaters (Central Iberian
Peninsula). The two Iberian individuals were previously genotyped using microsatellites (Macaya-
Sanz et al., 2011).

DNA extraction and RAD sequencing. For all samples, DNA was extracted from 15-20 mg of
silica-dried leaf material with the Qiagen DNeasy Plant Mini Kit (Valencia, CA). The concentration
of DNA was measured with a Qubit 2.0 Fluorometer using the dsDNA HS assay kit (Invitrogen), and
its integrity verified with electrophoresis on 1.5% agarose gels (1X TBE). Concentrations were
standardized to 20 ng/µl and individual samples were submitted for RAD library preparation and
sequencing to Floragenex (Eugene, OR). There, all extractions of the individuals of the first set, as
well as the replicate extractions of F039_05 and I373_A, were processed (together with additional
samples prepared in the same way), in five libraries of 95 individuals each. These libraries were

prepared according to Floragenex' standard commercial protocol: genomic DNA was digested with
the restriction endonuclease *PstI* (Christe et al., 2016; chosen according to previous studies on these
species - Stölting et al., 2013) and RAD libraries were prepared with a method similar to the one
described in Baird et al. (2008). This protocol included 18 PCR cycles, after which DNA fragments
ranging from 300 to 500 bp were retained. All five libraries were sequenced in a single run on an
Illumina HiSeq 2500 instrument, but on individual lanes. This data is available in the Sequence Read
Archive through bioproject PRJNA528699.

Following the same protocol, an additional library was generated and sequenced by Floragenex in a
separate experiment, consisting of two and three replicate extractions of J1 and BET, respectively, as
well as extractions from offspring of a controlled cross between them. This data is available in the
Sequence Read Archive through bioproject PRJNA528706.

Bioinformatic data processing. We assigned reads to individuals or replicates with *fastq-multx* (ea-
utils; Aronesty, 2011), allowing one mismatch in the 15 bp including barcode and restriction site.
Read quality was checked with *FastQC 0.10.1* (S. Andrews, 2010) and low quality bases and reads
were removed with *condetri* v.2.3 (Smeds & Künstner, 2011) using default parameters, except for the
options *-hq* (high quality threshold) and *-lfrac* (maximum acceptable fraction of bases after quality
trimming with quality scores lower than the threshold *-lq*), for which a value of 15 and 0.1 were
chosen, respectively.

High-quality reads were aligned against the *P. tremula* mitochondrial reference sequence (Kersten et
al., 2016) and against the nuclear reference genome of *P. trichocarpa* (Ptrichocarpa_210_v3.0;
Tuskan et al., 2006) using *Bowtie2 2.3.0* (Langmead & Salzberg, 2012) with "end-to-end" and "very
sensitive" settings. Reads with mapping quality lower than 20 were discarded using *samtools 1.3* (Li
et al., 2009) and read group information was added with *picard tools 1.139*
(http://broadinstitute.github.io/picard). We then used the tools *TargetCreator* and *IndelRealigner* of
GATK 3.8 (DePristo et al., 2011) to realign around indels, and recalibrated base quality scores for
each individual using the method by Kousathanas et al. (2017) implemented in *ATLAS* (Link et al.,
2017) on mitochondrial sequences. This method does not require *a priori* information on genotypes
and instead learns base qualities from haploid regions while integrating over genotype uncertainty.
Finally, we called genotypes with *UnifiedGenotyper* in *GATK 3.8* (DePristo et al., 2011).

To then only retain reliable sites for comparison, we filtered resulting variants using *vcftools*
(Danecek et al., 2011) and custom R scripts: first, we removed sites with an average depth across
individuals ≥24 (the 98.7% quantile) to exclude potentially paralogous loci. Second, we only kept
Single Nucleotide Variants (SNVs) with at most two segregating alleles. Third, we removed all
variant sites within 5 bp of an indel to avoid SNVs due to misalignments.

Truth Set. Genotypes of the 136 individuals grown in the common gardens were previously obtained
(Lindtke et al., 2014) with a genotyping-by-sequencing (GBS) protocol very similar to the ddRAD
protocol (Peterson et al., 2012) and using the restriction enzymes *EcoRI* and *MseI*. Importantly,
(Lindtke et al., 2014) generated sequencing data also for the 15 mother trees and used sibships in a
Bayesian approach to infer genotypes while accounting for familial relationships.

To compare these high quality genotypes to those obtained from our own RAD-seq experiment, loci
covered in both studies had to be identified first. Since Lindtke et al. (2014) used an older *P. tremula*
reference, we extracted from this reference windows of 201 bp around each locus in the GBS data set

(100 bp on either side). We then mapped these extracted sequences against the *P. trichocarpa*
reference with *Bowtie2 2.3.0* (Langmead & Salzberg, 2012) with "end-to-end" and "very sensitive"
settings and retained only those sequences that mapped uniquely with quality of 20 or more. We then
kept all loci overlapping between the two data sets, but removed four loci for which different
alternative alleles were called. To ensure high accuracy of the GBS data, we restricted all comparisons
to genotypes called with a posterior probability ≥99% by Lindtke et al. (2014).
279
280 **Estimating genotyping error rates.** We estimated genotyping error rates from our RAD-seq data
281 from both individual replicates as well as through the comparison to the truth data described above.
282 Since our data does not reflect random samples of a population, we could not benefit from the
283 population sample method introduced above.
284
Estimating of genome-wide and interspecific ancestry. For each of the 136 common garden
samples we estimated the genome-wide proportion of *P. alba* ancestry (genome-wide ancestry q) and
the proportion of the genome heterozygous for the two ancestries (interspecific ancestry Q_{12}) using
entropy (Gompert et al., 2014), a program that implements a model similar to the admixture model in
structure (Pritchard, Stephens, & Donnelly, 2000). In contrast to *structure*, however, *entropy* can also
make use of uncertain genotypes from low depth sequence data by working directly with genotype
likelihoods, rather than genotype calls. Here we ran *entropy* on the raw genotype likelihoods provided
by UnifiedGenotyper, as well as on genotype likelihoods recalibrated using empirical likelihoods
estimated from the comparison to a truth set, excluding sites with >50% missing data in both cases.
To stratify the estimates and have sufficient observations to estimate these probabilities reliably, we
considered five RAD-seq depth classes: 1-3, 4-7, 8-15, 16-31 and ≥32.
296
Inference of locus-specific ancestry. To infer locus-specific ancestry, we ran *RASPberry* (Wegmann
et al., 2011), which implements a Hidden Markov Model (HMM) to explain haplotypes of admixed
individuals as a mosaic of provided reference haplotypes for each species. We obtained suitable
reference haplotypes by phasing previously characterized pure *P. alba* and pure *P. tremula* individuals
(51 each) from the Italian, Austrian and Hungarian hybrid zones (Christe et al., 2016) using
FastPhase (Scheet & Stephens, 2006), building input files with *fcGENE* (Roshyara & Scholz, 2014).
For use in *RASPberry*, individuals in the reference panels were not allowed to have missing data. We
thus restricted the comparison to only the SNVs covered in all parental individuals.
305
To compute HMM transition probabilities in *RASPberry*, we used a default recombination rate of 5
cM/Mb as estimated by Tuskan et al. (2006) in *P. trichocarpa* and the estimates of the genome-wide
ancestry q for each sample obtained with *entropy* from the error corrected data. For most other
parameters we used previous estimates for *P. alba* and *P. tremula* hybrid zones (Christe et al., 2016),
but scaled these as proposed by Wegmann et al. (2011) to reflect the size of the reference panel. These
include the ancestral population recombination rates (315 and 900 for *P. alba* and *P. tremula*,
respectively), mutation rates (0.00185 and 0.00349, respectively), the miscopying rate (0.01) and the
miscopy mutation rate (0.01). However, we set the time since admixture to five (rather than one) to
reflect the different sampling strategy.
315
To account for genotyping errors, we determined a per-allele genotyping error ε as the weighted
average across depth-specific error rates obtained under the truth set model, with the constraint
$\varepsilon_0 = \varepsilon_1$ and weights reflecting the distribution of depth in the sites included in the analysis. We then
added this error rate estimate to the population specific and miscopy mutation rates. Under the
RASPberry copying model, these parameters control the rate at which the sample genotypes differ

from the reference haplotype from which the sample is copying. That rate thus depends on the
reference panel size, but also on genotyping errors.
323
324   We called ancestry segments as any stretch on a chromosome within which the posterior probabilities
325   for a particular ancestry (homozygous *P. alba*, heterozygous ancestry or homozygous *P. tremula*) was
326   >0.5 at all SNVs, and measured its length from the first to the last SNV.
327

Results

329
330   *Power to infer genotyping error rates*
331
Simulations suggest that a few thousand loci are sufficient to accurately estimate genotyping error
rates even from just a few samples (Figure 1). Due to the extra information provided, the smallest
estimation errors were obtained when using a truth set: >90% of all estimates fell within a range from
half to two-fold the true value (Q2, e.g., within [0.005, 0.02] for a true error rate of 0.01) if estimated
genotypes were compared at 200 and 400 sites for ε_0 and ε_1, respectively. Similar accuracy was
achieved under the population samples model as soon as 5,000 sites were used.
338
The lowest accuracy was observed under the replicates model, especially if error rates were low.
Using 5,000 sites, for instance, all estimates were within Q2 for a true value of $\varepsilon_0 = \varepsilon_1 = 0.1$, but
only slightly above 80% of all estimates for a true value of $\varepsilon_0 = \varepsilon_1 = 0.01$. This is readily explained
by the fact that only very limited information about the true genotype is available: if the two replicates
differ in their genotype, it is not clear which one is correct. Consequently, the accuracy of inference is
much increased if more than two replicates are available per individual (Supplementary Figure S1).
Nonetheless, even small error rates can be estimated relatively accurately, as >90% of all estimates for
true value of $\varepsilon_0 = \varepsilon_1 = 0.01$ fell within Q2 as soon as 10^4 or more comparisons were used.
347
Importantly, the accuracy is a function of the fraction of truly heterozygous genotypes in the data set,
with the accuracy of ε_1 increasing with more heterozygous calls, and that of ε_0 and ε_0 decreasing.
Here we simulated 50% heterozygous genotypes for the truth set and replicate models and about 29%
heterozygous genotypes for the population sample model. If the number of heterozygous genotypes
were lower, proportionally more sites would be needed to achieve the accuracy presented here. If 20%
of all genotypes were heterozygous, for instance, around 10^5 sites are required under the replicate
model with two replicates per sample.
355
356   *Using Whole-Genome Sequencing to Establish a Truth Set*
357
As shown above, genotyping error rates are inferred particularly accurately when comparing calls to a
truth set. One possibility to establish such a truth set is to shot-gun sequence a subset of individuals
and to infer genotypes from this data. This is also feasible in case no reference genome is available as
the shot-gun reads can be mapped directly against the RAD-loci. But since genotyping errors in the
truth set results in an overestimation of genotyping error rates, sufficiently high depth must be used.
Here we used simulations to assess this bias for the case of shot-gun sequencing one individual at 10^4
homozygous and 10^4 heterozygous RAD loci.
365
As shown in Figure 2 for $\varepsilon_0 = \varepsilon_1 = 0.01$, genotyping error rates may be overestimated considerably
unless the depth used to establish truth set was relatively high. At a mean depth of 10x, for instance,

the median relative bias was 1.92 and 0.50 for ε_0 and ε_1, respectively, and hence on the same order as
the inferred error rate. At a mean depth of 30x, the median overestimation dropped below 0.01 in both
cases. Importantly, larger quantiles drop much slower, suggesting that a minimal bias is hard to
overcome even with very high depths. Noteworthy, we observed a stronger bias for ε_0 than ε_1 at all
depths, a result of more often calling truly heterozygous sites homozygous than vice versa from next-
generation sequencing data. Results were virtually identical for the case of $\varepsilon_0 = \varepsilon_1 = 0.1$, but scaled
due to the larger true error (Supporting Figure S2).

High genotyping error rates in RAD-seq

We next used our inference methods to quantify genotyping errors from our RAD-seq experiment of
137 individuals of the two widespread tree species *Populus alba* and *P. tremula*, and their hybrids (*P.*
x canescens). On average, our experiment resulted in 831,161 (sd 153,434) reads per sample that
passed quality trimming and mapped against the reference genome of *P. trichocarpa* with mapping
quality ≥20. From those, we called 529,305 Single Nucleotide Variants (SNVs), after removing multi-
allelic sites, those with excessive depth, indels and variant sites around indels. We estimated per-allele
genotyping error rates from these SNVs through a comparison with previously published, high-quality
genotypes (truth set), and from multiple replicate libraries sequenced for a subset of our samples
(replicates).

Truth set. We estimated per-allele genotyping errors by comparing genotype calls from our RAD-seq
experiment to those of a previously published GBS dataset (Lindtke et al., 2014) for 136 individuals
present in both studies. In total, 7,426 SNVs overlapped between experiments, at which we could use
a total of 16,610 genotype comparisons. Of those, only 69.9% matched, with matching rates
increasing with RAD-seq depth (Figure 3A). Strikingly, RAD-seq genotypes were much less often
heterozygous than GBS genotypes (Figure 3B), especially at low depth. In line with these
observations, we inferred per-allele genotyping error rates >10% whenever RAD-seq depth was ≤ 35
and when using a model assuming a single error rate ($\varepsilon_0 = \varepsilon_1$), driven by an exceptionally high error
rate at truly heterozygous sites ε_1 (Figure 2C).

These are surprisingly high error rates, particularly when considering that sequencing error rates of
Illumina machines are estimated at <1% (Nielsen, Paul, Albrechtsen, & Song, 2011). Importantly, the
bias towards homozygous genotype calls is not simply explained by low depth. Indeed, RAD-seq still
resulted in less then half as many heterozygous calls at depths ≥40x, which are usually considered
more than sufficient for accurate genotype calling (Nielsen et al., 2011). Instead, our results suggest
an inherent bias in the RAD-seq data analyzed here.

We confirmed our suspicion by analyzing allelic imbalance at sites thought to be heterozygous based
on the GBS genotypes, limiting the analyses to genotypes covered at up to 50x to avoid spurious
alignments (Supplementary Figure S3). Only a single allele was for instance observed at 69% of all
heterozygous genotypes with depth ≥ 5x, a fraction that dropped to 60% and 52% for genotypes with
depth ≥ 20x and depth ≥ 30x, respectively, indicative of allelic dropout at a large scale.

However, our estimates rely on the assumption that the GBS data reflect true genotypes. This is based
on good evidence: First, Lindtke et al. (2014) additionally sequenced the mother trees of all
individuals considered here and estimated posterior genotypes using a hierarchical ancestry model
that incorporated familial relationships with mothers and among siblings. These updated estimates

correlated with the raw maximum likelihood genotype estimates ignoring familial data at 0.985 and
differed from those in <0.02% of all calls. Second, we restricted this comparison to GBS genotypes
with a posterior probability ≥99%. Third, the fraction of concordant genotype calls between the GBS
and RAD-seq data increased with RAD-seq depth. If the mismatches were driven by errors in the
GBS data, no such dependence should be observed.

420
Replicates. We next benefitted from two sets of replicate libraries to estimate per-allele genotyping
error rates. The first set consisted of two replicate libraries of each of two individuals (F039_05 and
I373_A) sequenced along all other samples. Error rates estimated from that data corroborated the
conclusion obtained from the comparison with GBS data (Figure 3D): error rates at truly homozygous
sites (ε_0) were on the order of 1% or less, and those at truly heterozygous sites (ε_1) were equal or
close to 50%, which is the largest value possible under our model.

427
In contrast to the estimates obtained in comparison to GBS genotypes, the error rates at truly
heterozygous sites (ε_1) dropped to about 20% at high depth (≥20x). This difference might in part be
driven by errors in the GBS data slightly inflating error rate estimates. However, given the high
quality of the GBS data, it appears more likely that the error rates from replicates are underestimated.
Polymorphisms in restriction cut sites or unequal PCR amplification rates of alleles, for instance,
affect replicates systematically, while the statistical inference must assume independence of errors
between replicates.

435
To verify that high error rates are not specific to the RAD-seq run performed on these hybrids, we
carried out a second RAD-seq experiment including two and three replicates of a pure *P. alba*
individual (J1) and a putative F1 *P. alba* x *P. tremula* hybrid (BET), respectively. This experiment
resulted in 694,031 (sd 187,957) reads per sample that passed quality trimming and mapped against
the reference genome of *P. trichocarpa* with quality ≥20. Per-allele error rates estimated from this
data were indeed lower, with error rates at truly homozygous sites (ε_0) on the order of 0.2% and those
at truly heterozygous sites (ε_1) starting out at 50% and dropping to about 3% at a depth of 25x (Figure
3D). However, these lower errors still point to a particular issue in calling heterozygous genotypes: of
all truly heterozygous sites with depth ≥25x in our data, >5% are expected to be called homozygous.
(The lower output of this sequencing experiment does not allow us to make reliable statements for
higher depths).

447
448 *Estimation of genome-wide and interspecific ancestry*

449
450 We investigated the impact of genotyping errors in our RAD-seq data on the inference of genome-
451 wide ancestry *q* as well as interspecific ancestry Q_{12}, which reflects the proportion of loci of
452 heterospecific ancestry. We estimated these ancestry components with *entropy* from 230,805 SNVs,
453 and compared them to estimates from GBS obtained by Lindtke *et al.* (2014).

454
The model implemented in *entropy* accounts for the genotyping uncertainty reflected in the genotype
likelihoods. However, the raw genotype likelihoods obtained from our RAD-seq data are misleading:
for sites with considerable depth, the RAD-seq genotype likelihoods often suggest almost certainty for
wrong genotypes. Of all genotypes wrongly called as homozygous at depth ≥30x (judged by the
comparison to GBS data), 90% had a variant quality of 77 or more. (A variant quality of 77 implies
that it is more than 5 billion times less likely to observe the obtained data from a heterozygous than
homozygous site).

As a result, ancestry estimates differed considerably between the GBS and RAD-seq data sets (Figure
4). Interestingly, the estimates of the genome-wide ancestry q were much less affected by genotyping
errors than the estimates of the interspecific ancestry Q_{12}. This is readily explained, however, by the
directionality of the most common error, which is to wrongly infer homozygous genotypes at
heterozygous sites. We found these errors do result more frequently in a homozygous reference than
homozygous alternative call (65.0%), particularly at low depth (84.5% at depth ≤5x, 49.9% at depth
≥20x). But this did not introduce a bias in q towards one of the species since we used the sequence of
the outgroup *P. trichocarpa* as reference. The estimates of Q_{12}, however, were very sensitive to an
underestimation of heterozygosity.

To improve these estimates, we propose to directly account for the elevated genotyping error rates in
RAD-seq data by adjusting the genotype likelihoods according to the observed genotyping
uncertainty. By treating the genotype calls as data, we can determine the probabilities $P(g|\gamma)$ of
observing a RAD-seq genotype call g given the true genotype γ either by using estimates of the per-
allele error rates $\varepsilon_0, \varepsilon_1$, or empirically from the comparison to a truth set. Using the latter approach on
our data (individually for each depth) resulted in estimates of q and Q_{12} that were much closer to those
obtained by Lindtke *et al.* (2014; Figure 4).

Some differences between the point estimates of Q_{12} remain, likely due to 1) differences in the models
and their information-sharing among individuals, as Lindtke *et al.* (2014) used information about
maternal plants and sibships as part of the model, and 2) the extent to which information in uncertain
genotypes was outweighed by hierarchical prior probabilities related to ancestry. Nonetheless, these
results demonstrate the importance of accounting for uncertainty in genotyping data since the
estimates of interspecific ancestry with and without correction lead to a very different biological
interpretation: With correction, the hybrid individuals appear to be mostly early generation hybrids,
suggesting meaningful reproductive isolation between the species. Without correction, the large
number of individuals with low Q_{12} but intermediate values of q suggests considerable gene flow
between the species, an interpretation at odds with recent work (Christe et al., 2016, 2017; Macaya-
Sanz et al., 2011).

Inference of locus-specific ancestry

We next evaluated the impact of genotyping errors on local ancestry inference. Hybrid zones between
P. alba and *P. tremula* are dominated by pure parental individuals and early hybrids (mostly F1), with
only few adult recombinant hybrids (Christe et al., 2016; Lindtke et al., 2014). We chose ten
individuals among our samples spanning that spectrum according to q and Q_{12} values from Lindtke *et*
al. (2014): a putatively pure *P. alba* individual (F039_01), a putatively pure *P. tremula* individual
(F030_01), two putative backcrosses to *P. alba* (F020_04 and F032_08), two putative backcrosses to
P. tremula (I345_02 and I345_03), two putative F1 hybrids (I373_03 and F030_05) and two putative
hybrids of later generations (F022_03 and F026_05). We then used *RASPberry* (Wegmann et al.,
2011) to infer local ancestry along chromosomes 1 through 5, restricting our inference to the 6,445
SNVs that did not have missing data in the parental reference haplotypes we took from Christe *et al.*
(2016).

In line with these expected simple ancestry make-ups, we inferred many large ancestry blocks often
spanning almost entire chromosomes (Figure 5). Surprisingly, however, we inferred most of these
blocks to be of homospecific ancestry, and also inferred many short segments, which are difficult to
reconcile with the putative ancestries of our samples (Figure 5). As an example, consider the

individual I373_03 in Figure 5 that was classified as an F1 hybrid by Lindtke et al. (2014), but for
which we inferred homospecific ancestry blocks for both parental species. Such artifacts could arise
from the reference panels being too small to properly reflect the haplotypes found in our hybrid
individuals, large gaps between neighboring SNVs limiting the power of the HMM implemented in
RASPberry, but most likely by genotyping errors towards homozygous genotypes.
516
While *RASPberry* does not account for genotyping uncertainty via genotype likelihoods, the
implemented copying-model allows for "mutations", or differences between the observed genotype of
an admixed individual and the reference haplotypes from which it is copying. We thus repeated the
inference by adding an estimate of the average per-allele genotyping error rate obtained under the
truth set model (13.7%) to the mutation rate parameters to account for the high genotyping error in
our RAD-seq data (Figure 5). Accounting for genotyping errors indeed improved our estimates. For
the putative backcrosses, for instance, we called fewer segments homozygous for the "wrong"
ancestry (11 versus 34) and these covered a smaller fraction of the first five chromosomes (6.1%
versus 11.1% of the parts at which ancestry could be called). Similarly, we called a higher fraction of
the first five chromosomes to be of heterozygous ancestry (45.0% versus 37.2% of the parts at which
ancestry could be called). While these results corroborate the importance of accounting for the true
uncertainty in genotypes in downstream analysis, they also illustrate that a method accounting for a
uniform error fails to fully mitigate the bias against heterozygous genotypes present in our RAD-seq
data sets.
531

Discussion

533
534 *High Genotyping Error Rates in RAD-seq*
535
Here, we report high genotyping error rates in two independent RAD-seq data sets. We obtained these
estimates by comparing RAD-seq calls of two independent experiments, either to published genotype
calls for the same individuals (Lindtke et al., 2014) obtained with a different sequencing method
(GBS), or to calls obtained from independent replicates of the same individuals. Both approaches
provide evidence for high per-allele genotyping errors of several percent and show that the RAD-seq
data analyzed here has a strong bias towards calling homozygous genotypes at heterozygous sites that
is not overcome with higher sequencing depth.
543
While these results do not directly generalize to other RAD-seq studies or other RAD-seq protocols,
the few estimates of genotyping errors reported to date agree with the high estimates we obtained
here. Luca et al. (2011), for instance, compared genotypes of human samples obtained with a
technique similar to RAD to those available in a public database, and estimated that between 6.3 and
9.7% of heterozygous sites were called as homozygous. Similarly, Mastretta-Yanes et al. (2015)
found that, depending on the parameter settings chosen for the *de-novo* assembly, between 5.9 and
8.8% of alleles were not concordantly called between replicates.
551
Several factors could explain the high genotyping error and the lack of heterozygous genotypes
frequently reported in RAD-seq data. For example, one allele might not have been sequenced or
sequenced only at very low depth because of differences in fragment length that can lead to
amplification bias, less efficient shearing, or loss in size selection. This is a likely explanation, since
Davey and colleagues (2013) showed that there is a high correlation between read depth and fragment
length and the protocol used here included 18 cycles of PCR. Similarly, differential efficiency of PCR

among alleles could have masked one allele (i.e. PCR duplicates), causing it to be represented in a
very low number of reads (Schweyen et al., 2014). However, PCR duplicates were recently reported
to be of only minor concern in RAD-seq studies, at least if only up to 12 PCR cycles were used
(Euclid 2019). Finally, the well-known issue of allele dropout due to polymorphisms in the restriction
site and the "loss" of one allele at heterozygous sites may have contributed to the inaccurate, low
observed heterozygosity (Andrews et al., 2016; Davey et al., 2013; Puritz et al., 2014). However, this
problem is a less likely explanation as it should not affect RAD-seq at a higher rate than the double-
digest GBS protocol we used for the error estimation.
566
567 *Accounting for Genotyping Error Rates*
568
Several bioinformatic solutions have been suggested to mitigate the apparent biases in RAD-seq.
Both Arnold et al. (2013) and Gautier et al. (2013) recommend the comparison of read depth across
sites, to identify loci likely exhibiting allele dropout. In our case, however, depth varied substantially
across sites, because of PCR duplicates or stochastic events, rendering such an approach difficult.
Davey et al. (2013) also noted that alleles present in two copies at homozygous sites have higher
depth compared to alleles present in single copy at heterozygous loci, but read depths for the two sets
of alleles overlap, inhibiting the accurate detection of loci with allele dropout by using depth alone.
To improve upon this, Cooke et al. (2016) developed a method to infer the likelihood of observing
allele dropout at a site on the basis of the sequencing depth of each sample, and suggested to ignore
sites where this likelihood is high. Finally, it was also suggested to discard any locus with a missing
genotype, since this might indicate a polymorphism in the restriction site. However, in many studies
with moderate depth, including ours, the amount of missing data prevents the adoption of such drastic
solutions. In summary, all these filtering suggestions result in a massive reduction in usable loci, and
hence further accentuate the already limited genome-wide coverage of reduced library techniques
such as RAD.
584
As a model-based alternative, we propose here to properly account for the high genotyping errors in
downstream analysis. A first such attempt was proposed by Cariou et al. (2016), who developed an
Approximate Bayesian Computation (ABC) method to estimate genetic diversity while accounting for
allele dropout, but found this method not to be accurate under elevated levels of diversity. A more
general solution, we believe, is to make use of the large number of recently developed tools that do
not require genotype calls but rather work directly from genotype likelihoods to account for
uncertainty in the data (Fumagalli, Vieira, Linderoth, & Nielsen, 2014; Jørsboe, Hanghøj, &
Albrechtsen, 2017; Korneliussen, Albrechtsen, & Nielsen, 2014; Kousathanas et al., 2017). Using
such tools minimizes the necessity to filter data stringently and is readily applied to low-depth data
(Nielsen et al., 2011).
595
For such methods to work properly, however, the genotype likelihoods need to accurately reflect the
uncertainty in genotypes. While all modern genotype callers also calculate genotype likelihoods, these
do not reflect biases specific to individual sequencing protocols such as RAD-seq, as we illustrated
here, and must thus be recalibrated. Here we propose three recalibration strategies: If accurate
genotype calls are available for a subset of the individuals and markers, empirical genotype
likelihoods can be obtained by comparing those to calls from a reduced representation sequencing
experiment. Alternatively, individual replicates or population samples may be used to infer per-allele
genotyping error rates, from which recalibrated genotype likelihoods are readily calculated. Tools for
these strategies are available through the software *Tiger*, which also accounts for sequencing depth as
an additional covariate. While we found sequencing depth to be a particularly important predictor, the

model is also readily extended to additional covariates such as the raw genotype likelihood or genotype call, which might provide additional information about genotyping error rates.

However, all these strategies might be biased. Genotyping errors in a set of genotype calls considered to be accurate (the truth set), for instance, will result in an overestimation of genotyping error rates. On the other hand, consistent biases in genotype calls affecting replicates similarly, such as polymorphisms in restriction cut sites or unequal PCR amplification rates of alleles, might be difficult to infer and result in an underestimation of genotyping error rates. Finally, the assumption of strict random mating may often be violated, leading to an overestimation when inferring error rates from population samples. But despite these caveats, recalibrated genotype likelihoods are likely to reflect genotype uncertainty much more accurately, particularly for protocols with error rates as high as those reported here for RAD-seq. Indeed, we found that genotype recalibration was essential to avoid drawing inaccurate conclusions and instead recovered biologically meaningful results about the ancestry of *Populus* hybrids.

We also emphasize that if no tools accepting genotype likelihoods as input information are available for specific applications, this should not discourage users from incorporating genotyping uncertainty in the analyses. We have shown here that local ancestries were more reliably estimated by *RASPberry* (Wegmann et al., 2011), a tool requiring genotype calls, when adding the estimated genotyping error rate to the parameters of the model. But we note that given the particular lack of heterozygous genotypes in the RAD-seq data analyzed here, a model using a single per-genotype error rate as implemented in *RASPberry* was not sufficient to overcome all biases.

Recommendations and Conclusions

Based on these results and in line with others (Cariou et al., 2016; Cooke et al., 2016; Mastretta-Yanes et al., 2015), we strongly suggest to carefully assess genotyping error rates in reduced representation sequencing experiments, and to properly account for these in downstream analyses. Here we present *Tiger*, a suite of tools to estimate genotyping error rates directly from hard genotype calls as produced by all commonly used RAD-seq analysis pipelines. Knowledge of these error rates then allows for fine-tuning the bioinformatic analysis to reduce these errors, as well as for recalibrating genotype likelihoods and hence properly accounting for genotyping errors in downstream analyses, rather than losing a large amount of information due to stringent filtering.

Among the different methods to estimate genotyping error rates, the use of population samples appears as a particularly cost-effective strategy, since no extra individuals have to be sequenced. However, not all applications include sufficiently many samples from randomly mating populations, and our simulations suggested that 20 samples are at the lower end, unless very many polymorphic sites are available. While *Tiger* can handle also multiple population samples with fewer individuals each, we expect much lower accuracy if fewer than, say, ten individuals were used per population. Also, we caution that the violation of random mating such as population sub-structure or inbreeding that causes a deficit of heterozygous genotypes directly translates into an overestimation of the heterozygous genotypes (Supplementary Figure S4).

For many studies, the preferred approach might thus be to prepare replicate libraries for a few individuals, not least because we found that just a few replicates (around ten) resulted in rather accurate estimates of genotyping errors. However, as already mentioned above, systematic biases affecting all replicates similarly may result in a slight underestimation of error rates.

Based on our simulations, the most accurate method to estimate genotyping errors is based on a truth
set against which genotype calls are compared. However, this is likely also the most expensive
approach. Since errors in the truth set will be wrongly interpreted as errors in the RAD-seq data, the
truth set must consist of highly accurate calls. As examined here using simulations for the case of
shot-gun sequencing, such a high accuracy requires considerable sequencing depth (>25x), which is
likely an underestimate given the simplicity of the model used here. More concerningly, genotyping
errors will likely prevail at low rates even in truth sets generated at very high depth, as was also
reported previously (Wall et al., 2014). But given the high error rates we observed for RAD-seq data,
the resulting slight overestimation of RAD-seq genotyping errors may be tolerable.

Nonetheless, and thanks to the ever dropping costs for both sequencing and library preparation, low-
depth whole genome sequencing may soon become a valuable alternative to reduced representation
sequencing for many applications. Since low-depth shotgun sequencing is much less prone to biases,
it not only allows to interrogate a much larger fraction of the genome, but also yields accurate and
precise estimates of population genetics parameters (e.g. Buerkle & Gompert, 2013; Kousathanas et
al., 2017; Rustagi et al., 2017). The problem of high error rates in RAD-seq and other reduced
representation libraries is thus likely transient and we expect that the field will quickly adopt new
sequencing technologies that circumvent it entirely.

Acknowledgements
674

We thank Kai N. Stölting, Camille Christe and Margot Paris for helpful insights and discussions,
Thelma Barbará for help in the laboratory, David Frey for collecting seeds in the hybrid zone, and
Santiago González Martínez for providing the Spanish samples. This work was supported by grant
31003A_149306 of the Swiss National Foundation to CL. All calculations were done on Vital-IT and
the Bioinformatic Core Facilities of the Universities of Bern and Fribourg, and computers at the
University of Wyoming.

Data Accessibility Statement
683

685 The code of Tiger is available through a git repository at https://bitbucket.org/wegmannlab/tiger. The
686 RAD-seq data are available in the Sequence Read Archive through bioprojects PRJNA528699 and
687 PRJNA528706. The called genotypes used to estimate genotyping errors are available at Zenodo
688 (DOI 10.5281/zenodo.2604109 and 10.5281/zenodo.2604124).

Author Contributions
690

LB and DW conceived the study; CL and DW provided funding; LB collected genetic data; LB, CAB,
VL and DW performed the analyses; CAB, CL and DW supervised the study; LB and DW wrote the
manuscript with input and revisions from all co-authors.

References

Andrews, K. R., Good, J. M., Miller, M. R., Luikart, G., & Hohenlohe, P. A. (2016). Harnessing the

power of RADseq for ecological and evolutionary genomics. *Nature Reviews. Genetics*, *17*(2),

81–92.

Andrews, K. R., & Luikart, G. (2014). Recent novel approaches for population genomics data

analysis. *Molecular Ecology*, *23*, 1661–1667.

Andrews, S. (2010). *FastQC: a quality control tool for high throughput sequence data*. Retrieved

from http://www.bioinformatics.babraham.ac.uk/projects/fastqc

Arnold, B., Corbett-Detig, R. B., Hartl, D., & Bomblies, K. (2013). RADseq underestimates diversity

and introduces genealogical biases due to nonrandom haplotype sampling. *Molecular Ecology*,

22(11), 3179–3190.

Aronesty, E. (2011). *ea-utils: Command-line tools for processing biological sequencing data*.

Retrieved from https://github.com/ExpressionAnalysis/ea-utils

Baird, N. A., Etter, P. D., Atwood, T. S., Currey, M. C., Shiver, A. L., Lewis, Z. A., … Johnson, E. A.

(2008). Rapid SNP discovery and genetic mapping using sequenced RAD markers. *PloS One*,

3(10), e3376.

Buerkle, A. C., & Gompert, Z. (2013). Population genomics based on low coverage sequencing: how

low should we go? *Molecular Ecology*, *22*(11), 3028–3035.

Cariou, M., Duret, L., & Charlat, S. (2013). Is RAD-seq suitable for phylogenetic inference? An in

silico assessment and optimization. *Ecology and Evolution*, *3*(4), 846–852.

Cariou, M., Duret, L., & Charlat, S. (2016). How and how much does RAD-seq bias genetic diversity

estimates? *BMC Evolutionary Biology*, *16*(1), 240.

Casbon, J. A., Osborne, R. J., Brenner, S., & Lichtenstein, C. P. (2011). A method for counting PCR

template molecules with application to next-generation sequencing. *Nucleic Acids Research*,

39(12), e81.

Christe, C., Stölting, K. N., Bresadola, L., Fussi, B., Heinze, B., Wegmann, D., & Lexer, C. (2016).

Selection against recombinant hybrids maintains reproductive isolation in hybridizing Populus

species despite F1 fertility and recurrent gene flow. *Molecular Ecology*, *25*(11), 2482–2498.

Christe, C., Stölting, K. N., Paris, M., Fraïsse, C., Bierne, N., & Lexer, C. (2017). Adaptive evolution and segregating load contribute to the genomic landscape of divergence in two tree species connected by episodic gene flow. *Molecular Ecology*, *26*(1), 59–76.

Chutimanitsakun, Y., Nipper, R. W., Cuesta-Marcos, A., Cistué, L., Corey, A., Filichkina, T., … Hayes, P. M. (2011). Construction and application for QTL analysis of a Restriction Site Associated DNA (RAD) linkage map in barley. *BMC Genomics*, *12*, 4.

Cooke, T. F., Yee, M.-C., Muzzio, M., Sockell, A., Bell, R., Cornejo, O. E., … Kenny, E. E. (2016). GBStools: A Statistical Method for Estimating Allelic Dropout in Reduced Representation Sequencing Data. *PLoS Genetics*, *12*(2), e1005631.

Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., … 1000 Genomes Project Analysis Group. (2011). The variant call format and VCFtools. *Bioinformatics* , *27*(15), 2156–2158.

Davey, J. W., Cezard, T., Fuentes-Utrilla, P., Eland, C., Gharbi, K., & Blaxter, M. L. (2013). Special features of RAD Sequencing data: implications for genotyping. *Molecular Ecology*, *22*(11), 3151–3164.

DePristo, M. A., Banks, E., Poplin, R., Garimella, K. V., Maguire, J. R., Hartl, C., … Daly, M. J. (2011). A framework for variation discovery and genotyping using next-generation DNA sequencing data. *Nature Genetics*, *43*(5), 491–498.

Emerson, K. J., Merz, C. R., Catchen, J. M., Hohenlohe, P. A., Cresko, W. A., Bradshaw, W. E., & Holzapfel, C. M. (2010). Resolving postglacial phylogeography using high-throughput sequencing. *Proceedings of the National Academy of Sciences of the United States of America*, *107*(37), 16196–16200.

Euclid, P. T., McKinney, G. J., Bootsma, M., Tarsa, C., Meek, M. H., Larson, W. A. (2019) Attack of the PCR clones: Rates of clonality have little effect on RAD-seq genotype calls. *Molecular Ecology Resources (in press).*

Fumagalli, M., Vieira, F. G., Linderoth, T., & Nielsen, R. (2014). ngsTools: methods for population genetics analyses from next-generation sequencing data. *Bioinformatics* , *30*(10), 1486–1487.

Gautier, M., Gharbi, K., Cezard, T., Foucaud, J., Kerdelhué, C., Pudlo, P., … Estoup, A. (2013). The

effect of RAD allele dropout on the estimation of genetic variation within and between

populations. *Molecular Ecology*, *22*(11), 3165–3178.

Gompert, Z., Lucas, L. K., Alex Buerkle, C., Forister, M. L., Fordyce, J. A., & Nice, C. C. (2014).

Admixture and the organization of genetic diversity in a butterfly species complex revealed

through common and rare genetic variants. *Molecular Ecology*, *23*(18), 4555–4573.

Hoffman, J. I., Simpson, F., David, P., Rijks, J. M., Kuiken, T., Thorne, M. A. S., … Dasmahapatra,

K. K. (2014). High-throughput sequencing reveals inbreeding depression in a natural population.

Proceedings of the National Academy of Sciences, *111*(10), 3775–3780.

Jørsboe, E., Hanghøj, K., & Albrechtsen, A. (2017). fastNGSadmix: admixture proportions and

principal component analysis of a single NGS sample. *Bioinformatics* , *33*(19), 3148–3150.

Kersten, B., Faivre Rampant, P., Mader, M., Le Paslier, M.-C., Bounon, R., Berard, A., … Fladung,

M. (2016). Genome Sequences of Populus tremula Chloroplast and Mitochondrion: Implications

for Holistic Poplar Breeding. *PloS One*, *11*(1), e0147209.

Korneliussen, T. S., Albrechtsen, A., & Nielsen, R. (2014). ANGSD: Analysis of Next Generation

Sequencing Data. *BMC Bioinformatics*, *15*, 356.

Kousathanas, A., Leuenberger, C., Link, V., Sell, C., Burger, J., & Wegmann, D. (2017). Inferring

Heterozygosity from Ancient and Low Coverage Genomes. *Genetics*, *205*(1), 317–332.

Langmead, B., & Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nature Methods*,

9(4), 357–359.

Leaché, A. D., Chavez, A. S., Jones, L. N., Grummer, J. A., Gottscho, A. D., & Linkem, C. W.

(2015). Phylogenomics of phrynosomatid lizards: conflicting signals from sequence capture

versus restriction site associated DNA sequencing. *Genome Biology and Evolution*, *7*(3), 706–

719.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., … 1000 Genome Project Data

Processing Subgroup. (2009). The Sequence Alignment/Map format and SAMtools.

Bioinformatics , *25*(16), 2078–2079.

Lindtke, D., Buerkle, C. A., Barbará, T., Heinze, B., Castiglione, S., Bartha, D., & Lexer, C. (2012).

Recombinant hybrids retain heterozygosity at many loci: new insights into the genomics of

reproductive isolation in Populus. *Molecular Ecology*, *21*(20), 5042–5058.

Lindtke, D., Gompert, Z., Lexer, C., & Buerkle, C. A. (2014). Unexpected ancestry of Populus

seedlings from a hybrid zone implies a large role for postzygotic selection in the maintenance of

species. *Molecular Ecology*, *23*(17), 4316–4330.

Link, V., Kousathanas, A., Veeramah, K., Sell, C., Scheu, A., & Wegmann, D. (2017). *ATLAS:*

Analysis Tools for Low-depth and Ancient Samples. doi: 10.1101/105346

Luca, F., Hudson, R. R., Witonsky, D. B., & Di Rienzo, A. (2011). A reduced representation approach

to population genetic analyses and applications to human evolution. *Genome Research*, *21*(7),

1087–1098.

Macaya-Sanz, D., Suter, L., Joseph, J., Barbará, T., Alba, N., González-Martínez, S. C., … Lexer, C.

(2011). Genetic analysis of post-mating reproductive barriers in hybridizing European Populus

species. *Heredity*, *107*(5), 478–486.

Marques, D. A., Lucek, K., Meier, J. I., Mwaiko, S., Wagner, C. E., Excoffier, L., & Seehausen, O.

(2016). Genomics of Rapid Incipient Speciation in Sympatric Threespine Stickleback. *PLoS*

Genetics, *12*(2), e1005887.

Mastretta-Yanes, A., Arrigo, N., Alvarez, N., Jorgensen, T. H., Piñero, D., & Emerson, B. C. (2015).

Restriction site-associated DNA sequencing, genotyping error estimation and de novo assembly

optimization for population genetic inference. *Molecular Ecology Resources*, *15*(1), 28–41.

Miller, M. R., Dunham, J. P., Amores, A., Cresko, W. A., & Johnson, E. A. (2007). Rapid and cost-

effective polymorphism identification and genotyping using restriction site associated DNA

(RAD) markers. *Genome Research*, *17*(2), 240–248.

Nadeau, N. J., Ruiz, M., Salazar, P., Counterman, B., Medina, J. A., Ortiz-Zuazaga, H., … Papa, R.

(2014). Population genomics of parallel hybrid zones in the mimetic butterflies, H. melpomene

and H. erato. *Genome Research*, *24*(8), 1316–1333.

Narum, S. R., Buerkle, C. A., Davey, J. W., Miller, M. R., & Hohenlohe, P. A. (2013). Genotyping-

by-sequencing in ecological and conservation genomics. *Molecular Ecology*, *22*(11), 2841–2847.

Nielsen, R., Paul, J. S., Albrechtsen, A., & Song, Y. S. (2011). Genotype and SNP calling from next-

generation sequencing data. *Nature Reviews. Genetics*, *12*(6), 443–451.

Peterson, B. K., Weber, J. N., Kay, E. H., Fisher, H. S., & Hoekstra, H. E. (2012). Double digest

RADseq: an inexpensive method for de novo SNP discovery and genotyping in model and non-

model species. *PloS One*, *7*(5), e37135.

Pootakham, W., Sonthirod, C., Naktang, C., Jomchai, N., Sangsrakru, D., & Tangphatsornruang, S.

(2016). Effects of methylation-sensitive enzymes on the enrichment of genic SNPs and the

degree of genome complexity reduction in a two-enzyme genotyping-by-sequencing (GBS)

approach: a case study in oil palm (Elaeis guineensis). *Molecular Breeding: New Strategies in*

Plant Improvement, *36*(11), 154.

Pritchard, J. K., Stephens, M., & Donnelly, P. (2000). Inference of population structure using

multilocus genotype data. *Genetics*, *155*(2), 945–959.

Puritz, J. B., Matz, M. V., Toonen, R. J., Weber, J. N., Bolnick, D. I., & Bird, C. E. (2014).

Demystifying the RAD fad. *Molecular Ecology*, *23*(24), 5937–5942.

Roshyara, N. R., & Scholz, M. (2014). fcGENE: a versatile tool for processing and transforming SNP

datasets. *PloS One*, *9*(7), e97589.

Rowe, H. C., Renaut, S., & Guggisberg, A. (2011). RAD in the realm of next-generation sequencing

technologies. *Molecular Ecology*, *20*(17), 3499–3502.

Rustagi, N., Zhou, A., Watkins, W. S., Gedvilaite, E., Wang, S., Ramesh, N., … Xing, J. (2017).

Extremely low-coverage whole genome sequencing in South Asians captures population

genomics information. *BMC Genomics*, *18*(1), 396.

Sambrook, J., & Russell, D. W. (2006). Fragmentation of DNA by Sonication. *Cold Spring Harbor*

Protocols, *2006*(23), db.prot4538 – pdb.prot4538.

Scheet, P., & Stephens, M. (2006). A fast and flexible statistical model for large-scale population

genotype data: applications to inferring missing genotypes and haplotypic phase. *American*

Journal of Human Genetics, *78*(4), 629–644.

Schweyen, H., Rozenberg, A., & Leese, F. (2014). Detection and removal of PCR duplicates in

population genomic ddRAD studies by addition of a degenerate base region (DBR) in

sequencing adapters. *The Biological Bulletin*, *227*(2), 146–160.

Smeds, L., & Künstner, A. (2011). ConDeTri - A Content Dependent Read Trimmer for Illumina

Data. *PloS One*, *6*(10), e26314.

Stölting, K. N., Nipper, R., Lindtke, D., Caseys, C., Waeber, S., Castiglione, S., & Lexer, C. (2013).

Genomic scan for single nucleotide polymorphisms reveals patterns of divergence and gene flow

between ecologically divergent species. *Molecular Ecology*, *22*(3), 842–855.

Toonen, R. J., Puritz, J. B., Forsman, Z. H., Whitney, J. L., Fernandez-Silva, I., Andrews, K. R., &

Bird, C. E. (2013). ezRAD: a simplified method for genomic genotyping in non-model

organisms. *PeerJ*, *1*, e203.

Tuskan, G. A., Difazio, S., Jansson, S., Bohlmann, J., Grigoriev, I., Hellsten, U., …, Rokhsar, D.

(2006). The genome of black cottonwood, Populus trichocarpa (Torr. & Gray). *Science*,

313(5793), 1596–1604.

Wall, J. D., Tang, L. F., Zerbe, B., Kvale, M. N., Kwok, P.-Y., Schaefer, C., & Risch, N. (2014).

Estimating genotype error rates from high-coverage next-generation sequence data. *Genome*

Research, *24*(11), 1734–1739.

Wang, S., Meyer, E., McKay, J. K., & Matz, M. V. (2012). 2b-RAD: a simple and flexible method for

genome-wide genotyping. *Nature Methods*, *9*(8), 808–810.

Wegmann, D., Kessner, D. E., Veeramah, K. R., Mathias, R. A., Nicolae, D. L., Yanek, L. R., …

Novembre, J. (2011). Recombination rates in admixed individuals identified by ancestry-based

inference. *Nature Genetics*, *43*(9), 847–853.

Wright, S. (1931). Evolution in Mendelian Populations. *Genetics*, *16*(2), 97–159.

[dataset]Bresadola L, Link V, Buerkle CA, Lexer D, Wegmann D; 2019; Common garden seedlings

of Populus hybrids; Sequence Read Archive; Bioproject ID: PRJNA528699.

[dataset]Bresadola L, Link V, Buerkle CA, Lexer D, Wegmann D; 2019; Controlled cross of Populus

hybrids; Sequence Read Archive; Bioproject ID: PRJNA528706.

[dataset]Bresadola L, Link V, Buerkle CA, Lexer D, Wegmann D; 2019; VCF File containing

genotype calls for 136 Populus alba x Populus tremula hybrids obtained through both RAD-seq

and GBS; Zenodo; DOI: 10.5281/zenodo.2604109.

[dataset]Bresadola L, Link V, Buerkle CA, Lexer D, Wegmann D; 2019; VCF File containing

genotype calls for 136 Populus alba x Populus tremula hybrids obtained through both RAD-seq

and GBS; Zenodo; DOI: 10.5281/zenodo.2604109.

Tables

870 **_Table 1_**. _Per-allele genotyping error model. Shown are the probabilities of observing a RAD-seq_
871 _genotype given the true genotype and the per-allele genotyping error rate ε._

	Observed genotype		
True genotype	0	1	2
0	$(1-\varepsilon_0)^2$	$2\varepsilon_0 * (1-\varepsilon_0)$	ε_0^2
1	$\varepsilon_1 * (1-\varepsilon_1)$	$(1-\varepsilon_1)^2 + \varepsilon_1^2$	$\varepsilon_1 * (1-\varepsilon_1)$
2	ε_0^2	$2\varepsilon_0 * (1-\varepsilon_0)$	$(1-\varepsilon_0)^2$

Figure Legends

Fig. 1: *Accuracy of error rate estimates. Shown are the estimated error rates relative to the true error rates (red line) of 100 replicates for different samples sizes n (shown on top), the two error rates 0.1 and 0.01 (first and last two columns, respectively) and different numbers of unlinked loci. The horizontal dashed lines indicate Q2, the interval within which an estimate is less than a factor of two away from the true value (i.e. within half and two times the true value). Simulations were generated for the truth set (top row), the individual replicate (two replicates per sample, middle row) and population samples models (bottom row).*

Fig. 2: *Distribution of the relative bias when inferring error rates from a truth set generated with next-generation sequencing data with a specific mean depth. Each line indicates a quantile m such that a smaller bias was observed in a fraction m of 10^6 simulations of 10^4 homozygous and 10^4 heterozygous loci with $\varepsilon_0 = \varepsilon_1 = 0.01$.*

Fig. 3: *Estimates of genotyping errors in RAD-seq data. **A**. Maximum likelihood estimate (MLE) of the frequency of genotypes estimated from GBS and RAD-seq data that match as a function of RAD-seq sequencing depth. The gray region indicates the range of frequencies within two log-likelihood units of the MLE. **B**. MLE estimate of the frequency of heterozygous calls with RAD-seq among all calls of a particular depth overlapping with the GBS data. Gray region as in A. The heterozygosity observed among all overlapping GBS genotypes is given as dashed line. **C**. Per-allele RAD-seq genotyping error rates for homozygous (ε_0, black open symbols) and heterozygous (ε_1, blue filled symbols) genotypes estimated using a GBS truth set, limited to depth class with at least 100 comparisons. Estimates for a model assuming $\varepsilon_0 = \varepsilon_1$ are shown with yellow closed circles. **D**. Per-allele RAD-seq genotyping error rates obtained from two replicates of F039_05 and I373_A each (circles) and two and three replicates from J1 and Bet, respectively (triangles), limited to depth class with at least 1000 sites. Colors as in C.*

Fig. 4: *Comparison of genome-wide (q) and interspecific (Q_{12}) ancestry estimates of 136 individuals obtained from GBS data (Lindtke et al., 2014) and estimates from RAD-seq data (black circles) using either the raw (left panel) or corrected (right panel) genotype likelihoods. Grey lines connect values obtained for the same individual.*

Fig. 5: *Comparison of local ancestry patterns on chromosomes 1-5 for (top to bottom) a putatively pure* P. tremula *individual (P. trem, F030_01), two putative backcrosses to* P. tremula *(BC trem, I345_02 and I345_03), two putative hybrids of later generation (Fn, F022_03 and F026_05), two putative F1 hybrids (F1, I373_03 and F030_05), two putative backcrosses to* P. alba *(BC alba, F020_04 and F032_08) and a putatively pure* P. alba *individual (P. alba, F039_01). For each individual,* RASPberry *results with (+) and without (-) correction are shown. Blue represents* P. alba *ancestry, orange* P. tremula *ancestry and grey heterospecific ancestry, with darker shades showing higher confidence in the ancestry estimates. To facilitate visualization, only sites with data are shown.*