1
2
A phenotypically plastic magic trait promoting reproductive isolation in sticklebacks?

Monica V. Garduno Paz[1,2], Felicity A. Huntingford[1] Sean Garrett[1] and Colin E. Adams[1,3]

[1]*Scottish Centre for Ecology and the Natural Environment, IBAHCM, University of Glasgow,*
Rowardennan, Glasgow G63 0AW, UK

[2]*Current academic address: Facultad de Ciencias, Universidad Autónoma del Estado de México,*
Instituto Literario no. 100 Colonia Centro, Toluca, CP50000, México

[3] Corresponding author Email: colin.adams@glasgow.ac.uk

Adams - Orcid identifier number: 0000-0003-2470-9754

Running title: A plastic magic trait in sticklebacks

KEY WORDS: Assortative mating, epigenetics, reproductive isolation, trophic polymorphism.

Wordcount: 3474

Figures: 1

Tables:1

Abstract

This study identifies one possible mechanism whereby gene flow is interrupted in populations

undergoing evolutionary divergence in sympatry; this is an important issue in evolutionary biology

that remains poorly understood. Variation in trophic morphology was induced in three-spined

stickleback by exposing them from an early age either to large benthic or to small pelagic prey. At

sexual maturity, females given a choice between two breeding males, showed positive assortative

mate choice for males raised on the same diet as themselves. The data indicate that this was

mediated through a preference for males with trophic morphology similar to that of fish with which

the females were familiar (from their pre-testing holding tanks). In trials where the female did not

choose the most familiar male, the evidence suggests that either she had difficulty discriminating

between two similar males or was positively choosing males with more extreme morphologies (more

benthic-like or pelagic -like). This study has shown for the first time that expression of a plastic trait

induced at an early age, not only results in specialisation for local foraging regimes but can also play

a significant role in mate choice. This is equivalent to an environmentally induced, plastic version of

the "magic traits" that promote ecologically-driven divergence in sympatry, hence the proposed

descriptor "plastic magic trait".

Introduction

The process whereby gene flow is interrupted in populations undergoing evolutionary divergence when in sympatry is an important issue in evolutionary biology that remains poorly understood. One potential mechanism is where an ecologically important trait which is under divergent selection also contributes to reproductive isolation and is thus a so-called "magic trait" (Gavrilets 2004; Servedio et al. 2011). Although evidence of such traits in nature is sparse (Servedio et al. 2011), magic traits are usually envisaged as inherited and linked to mate choice through pleiotropy. However, much research interest has focussed on the possible role of phenotypic plasticity in the initiation of evolutionary change through the development of discrete alternative phenotypes (West-Eberhard 1989, 2003; Fitzpatrick 2012; Skulason et al. 2019). If expressed alternative phenotypic traits induced by the environment through plasticity, also form part of the mate choice system of the diverging organism, then assortative mating resulting from mate choice based on such traits have the potential to generate reproductive barriers between individuals expressing different phenotypes (Fitzpatrick 2012). Such traits may thus act as a magic trait without the requirement of pleiotropy.

Discrete alternative phenotypes associated with foraging, or trophic polymorphisms (*sensu* Skúlason & Smith (Skulason and wund 1995), have been strongly implicated in sympatric speciation events (Dieckmann and Doebeli 1999). Divergent morphological traits are often the result of foraging conditions experienced during development (Day and McPhail 1996; Adams et al. 2003), so can only result in evolutionary change if mechanisms exist that result in gene pool segregation (West-Eberhard 1989; Smith and Skulason 1996; Schluter 2003). Here we explore one possible mechanism, namely morph-specific mate choice by breeding females. Using the three-spined stickleback (*Gasterosteus aculeatus*) as a model system, we present an example of a developmentally-plastic, trophic specialisation acting as a magic trait generating reproductive isolation and suggest a mechanism through which this comes about.

Trophic polymorphism is particularly common among freshwater fishes including sticklebacks; it often takes the form of co-existing but discrete phenotypes with morphological and behavioural specialisations for feeding on benthic invertebrates in the littoral zone or zooplankton in the pelagic zone (Skulason and Smith 1996; Adams and Huntingford 2002a; Proulx and Magnan 2004). Typically, the benthic form is robust, with a large mouth, small eyes and few short gill rakers, while the pelagic form is lightly built, with a relatively small mouth, large eyes and longer and more numerous gill rakers (Adams et al. 1998; Adams and Huntingford 2002a). Although in some cases the two sympatric forms are fully reproductively isolated, more commonly reproductive isolation is partial, weak or non-existent (Schluter and McPhail 1992; Hendry et al. 2009). Sticklebacks also exhibit scope for the

expression of characteristics under phenotypic plasticity (Wund et al. 2008, 2012; Garduño-Paz et al.

2010; Baker et al. 2013) including plasticity in morphological characteristics that define sympatric

forms described from the wild (Day and McPhail 1996; Garduño-Paz et al. 2010).

The main aim of this study was, having induced variable trophic morphology in three-spined

sticklebacks from a single population by manipulating early feeding regimes, to determine whether

these plastic, diet-induced differences in trophic morphology were associated with different patterns

of mate choice. A second aim was to seek possible behavioural mechanisms that might explain the

observed patterns of mating.

Methods

Diet Treatments

240 juvenile three-spined sticklebacks fry (5-9 mm length) were collected by dip nets from a small

freshwater pond in Scotland (56°3'N; 004°21'W) and transported to rearing facilities at the Scottish

Centre for Ecology and the Natural Environment (SCENE), Glasgow University, Loch Lomond. Fish were

assigned randomly in groups of 40 to 6 rearing aquaria (21L) and raised in the laboratory for 11

months, during which time they were fed twice daily to satiation on one of two diet treatments known

to induce differences in trophic morphology (Day and McPhail 1996). Half of the groups were fed on

frozen *Daphnia* spp in a bag hanging at the water surface, simulating pelagic prey; the rest were fed

on frozen chironomid prey placed on the bottom of the tank, simulating benthic prey.

87
Analysis of Induced Morphological Differences

After 10 months, the sticklebacks were anaesthetised with benzocaine and photographed on their

left side with a Canon EOS digital 350D camera (8.0 megapixels). Female fish that were gravid and

male fish which were coloured were identified as sexually mature. All fish used in mate choice

experiments were re-photographed at 11 months immediately following the mate choice

experiments. Body shape was quantified on the basis of 20 landmarks (Figure S1), placed using the

program "tpsUtil" ad digitised using the program "tpsDIg2" (Rohlf, 2006). Generalized least squares

procrustes superimposition was used to translate, scale and rotate raw landmark coordinates to

minimize the summed, squared, inter-landmark distances among fish; this procedure removes the

effect of fish size (Rohlf & Slice, 1990). Relative Warp Analysis was conducted on the partial warp

scores to reduce the number of informative shape variables (Rohlf, 2006). The second relative warp

(analogous to a Principal Component), which explained 13% of the total shape variation, separated

traits typical of pelagic and benthic feeders (Day and McPhail 1996; Figure S1).

101
Mate Choice Trials

Twenty eight females (21 from the chironomid diet and 7 from the *Daphnia.* diet) and 36 males (21

from the chironomid diet and 15 from the *Daphnia* diet) were used in trials of mate choice. Female

mate choice was examined using a well-tested methodology, widely used in previous studies (Milinski

and Bakker 1990; Kraak and Bakker 1998; Boulcott et al. 2005; Rick et al. 2006; Rick and Bakker 2008;

Heuschele et al. 2009) in which a single gravid female was placed alone in an aquarium (35 x 25 x 20

cms, screened on 3 sides), allowed to settle for 12h and was then presented simultaneously with two

breeding males in equally-sized sections (25 x 35 x 20 cms) of an adjacent aquarium and thus not

subject to olfactory cues. During trials, females could see both males, but the females had no olfactory

contact with males and the two males were separated by an opaque partition and so did not have

visual contact with each other. In any trial, the female was presented with one *Daphnia*-fed and one

chironomid-fed male. To avoid effects of size and familiarity with specific males, the two males in any

given trial were size-matched as far as possible and importantly taken from a different rearing tank

from the female. To avoid females potentially making choices based on nest construction, males were

not provided with nesting material.

Each trial lasted for 5 minutes, during which, the time the female spent on the side of the tank adjacent

to each male was recorded. Three replicates of each pairing trial were conducted, swapping the male

position each time. A female was deemed to have chosen a male if she spent at least 60% of the total

time of the trial near that male. Association time has been shown to be a strong predictor of eventual

mate choice in this species in a number of other studies (McLellan and McPhail; Rowland et al. 1995;

Milinski et al. 2005; Rick and Bakker 2008). Males and females were used maximally in four trials on

different days; males were re-used in fresh combinations so that the female was never exposed to the

same pair of males. Although male pairs were matched in size (by fork length) as nearly as possible,

small discrepancies between pairs remained. Retrospective analysis detected no significant difference

in body size (fork length) between chosen and rejected males (mean $\pm$ SE size differences between

accepted and rejected males =0.04 cm $\pm$ 0.02 paired t test: t = 1.68, p = 0.10).

Body shape, as defined by the second relative warp varied markedly both between and within diets

(Fig.1). Effects of sex (ANOVA: $F_{1,60}$=3.11; p=0.08) and of sex by diet (ANOVA: $F_{1,60}$=2.86; p= 0.09) were

not significant. However, there was a highly significant difference in morphology between the

chironomid fed (mean $\pm$ SE = 6.753 $\pm$ 0.23) and *Daphnia* fed sticklebacks (mean $\pm$ SE = 4.82 + 0.34.

ANOVA: $F_{1,60}$= 22.2; p< 0.0001). The higher scores of fish fed on the benthic diet reflected shorter

heads, shorter maxillary bones, smaller eyes and deeper bodies. This score was transformed to create

only positive values (by adding 6 and multiplying by 100) and hereafter this dimension of shape

variation is referred to as the pelagic-benthic (PB) shape score. Lower PB scores indicate shapes

tending towards a more typical of a pelagic foraging fish; higher scores tending towards a more

benthic foraging fish shape (Fig.1). To enable testing of the body shape of chosen and rejected males,

the difference in PB shape score between the chosen male and the rejected male for each pair was

determined as: male PB score difference =PB score chosen male – PB score rejected male. Thus a large

positive number in this metric indicates the choice of a male from the pair with a higher PB score; thus

tending towards a shape more typical of a benthic foraging fish. Conversely a large negative number

in the difference between chosen and rejected male PB scores indicates a choice of male with a shape

tending towards that typical of a more pelagic forging fish.

In nature, female sticklebacks review a number of males before selecting a nest in which to lay her

eggs and the females in this study were always tested with different pairs of unfamiliar males.

However female identity was included as a random factor in mixed effects models examining female

mate choice. A total of 96 trials of mate choice by females was analysed using "R" (R Development

Core Team 2017).

Results

Mate Choice in Females Exposed to Benthic and Pelagic Diets

To test for the effect of the previous diet experience of the female on female mate choice, the PB

score difference between male pairs in each trial was categorised as either a negative value, indicating

the choice of a male with a more pelagic-like body shape, or positive, indicating a choice of the male

from the pair with a more benthic-like body shape. The previous diet experience of the female (benthic

(chironomid) or pelagic (*Daphnia*) diet exposure) was used as a fixed factor and female I.D. as random

factor, in a mixed effects model with a binomial (positive or negative male PB score difference)

probability distribution. Female diet exposure significantly predicted the choice of male body shape

in pairwise tests ($p<0.01$) indicating non-random mating on the basis of previous diet exposure.

However the total explained variance was low (6.8%) and this effect was mostly driven by females

with previous exposure of a pelagic prey diet choosing the male with the more pelagic body shape (a

negative male PB score difference) on 66% of occasions (Table 1). In contrast, females exposed to the

benthic prey diet chose the male from the pair with the more benthic body shape (a positive male PB

score difference) on only 54% of occasions.

167
168
Behavioural Mechanisms Of Mate Choice

To explore possible behavioural mechanisms for the observed female preference by diet, mate choice

data were analysed in more detail. To test the possibility that females are making a choice of male

based on their own body shape, the male PB score difference in each trial, categorised as either a

negative value, (indicating the choice of a male of a more pelagic-like body shape), or a positive value

(indicating a choice of the male with a more benthic-like body shape) was modelled using female PB

score as a fixed factor and female I.D. as random factor, in a mixed effects model with a binomial

probability distribution. Female body shape did not predict the choice of male ($p<0.45$). Thus the

assortative mating effect predicted by female previous diet exposure does not appear to be driven by

the body shape of the female.

Another possible behavioural mechanism by which rearing diet might influence a female stickleback's

mate choice is through previous experience of the fish with which she was reared, whose shape will,

on average, reflect their common rearing diet. To test this possibility, we took advantage of the

variability in PB scores between rearing tanks on a given diet. The male PB score difference in each

trial, categorised as either a negative value, or a positive value was modelled using the mean PB score

of fish in the tank from which the female was drawn, as a fixed covariate and female I.D. as random

factor, in a mixed effects model with a binomial probability distribution.

The mean PB score of fish from the tank from which the female originated significantly predicted the

female's choice of male ($p<0.02$; $r^2= 6.9\%$). Indicating that females are choosing of males on the basis

of body shapes with which they are familiar. Despite this a considerable amount of the variation in the

choice of male was not explained by the body shape with which they were familiar.

One possible explanation is that in trails where females chose males that were not closer in body

shape to those with which she was familiar (from the same rearing tank) and thus she may have had

difficulty discriminating between body shapes. If this was the case, then one expectation would be

that in such trials the two males are likely to be closer in body shape (similar PB scores) to each other

than in trials where the female chose the male closest to that with which she is familiar. This was

tested; the PB score difference between males in trials where the female chose the most familiar

shape (2.22 ± 1.36; mean $\pm$ S.D.) was significantly greater (t test: $p=0.041$) than the PB score difference

between males from trails where she chose the less familiar male (1.85 ± 0.94). This indicates that in

at least some trails females may have had difficulty distinguishing between males of similar body

shape.

Another possible explanation for the outcome of those trails where the female did not chose the male

with a body shape closest to the mean of fish that she was familiar with, is that she may avoid choosing

males with extreme body shapes even if they are closer to the body shape with which she is familiar.

To test this the Extreme Shape Index was calculated as the deviation in body shape for chosen and

rejected male from the average of all fish combined (that is the PB score of each fish - the mean of all

fish combined irrespective of sign. As these data deviated significantly from normality, the Extreme

Shape Index for chosen and rejected males from trials where the female chose the male with a shape

that she was less familiar with, were compared in a paired Wilcoxon test. In these trials the chosen

male was much more likely to have a more extreme body shape (i.e a higher Extreme Shape Index)

$(2.09 \pm 1.08;$ mean $\pm$ S.D.) than the rejected male $(1.01 \pm 0.86).$

Discussion

Our results confirm the findings of previous studies demonstrating a plastic response of

morphological traits to rearing diet in three-spined sticklebacks (Day and McPhail 1996). More

significantly, they have demonstrated for the first time that exposure to different diets during the

juvenile phase can influence the mating preferences shown by breeding females. Thus, females

reared on the pelagic diet tended to prefer the male with a more pelagic-like morphology; females

reared on a benthic diet however mated randomly with respect to trophic morphology. Thus there is

partial assortative mating by diet-induced phenotype. Unlike the case of assortative mating on the

basis of diet specialisation in the mustard leaf beetle, which appears to use olfactory cues to identify

mates (Geiselhardt et al. 2012) the sticklebacks in this experiment only had visual cues are available

to them. However, it is quite possible that olfactory cues might also have affected mate choice had

they been available. In addition, the effect reported here did not result from female familiarity with

specific individual males, as females were never tested with males from the same rearing tank.

Additionally we show that mate choice was not dependent directly of the female's own

trophic morphology. Arguably, this is not surprising, since it is difficult to see how a female

stickleback could know what her own morphology is like. Instead the differences in mate choice

must be a consequence (direct or indirect) of the experience of being raised on a pelagic or a benthic

diet. Making use of the significant variation in morphology between and within rearing tanks

exposed to different and the same diets, we show that the expressed morphology of other fish with

which the female is familiar (from the same rearing tank) is a good predictor of mate choice. It is

highly likely that in the wild also sticklebacks grow up with fish exploiting a similar diet to themselves

and thus with similar diet-induced morphology, as individuals exploiting the same foraging resources

are more likely to come into contact with each other, than those that do not share a common diet

(Garduño-Paz and Adams 2010).

Despite a clear tendency for assortative mating by trophic morphology, females quite often made

the opposite choice. This was most often the case when the difference between the two males was

relatively small, but also occurred when if the morphology of the predicted choice male was of an

extreme benthic or pelagic-type morphology. One can envisage at least two plausible mechanistic

explanations for this, which are not mutually exclusive. It may be that, rather than responding to

familiarity *per se*, females have learned about the foraging efficacy of fish with the range of

morphologies that she has experienced during development. If this were the case, then females

might actively choose males of a more extreme morphology, which may well be more efficient at

foraging on the two alternative diets presented, even if this morphology is less common in her

previous experience familiar to her. We are not able to test directly this possibility using the data

from this study.

Although coexisting trophic morphs are thought to be an important step in evolutionary divergence

in sympatry (Skulason et al. 1999), speciation however is unlikely to be completed without some

mechanism for morph-specific assortative mating (Skulason et al. 1999). Several routes though

which this might occur have been suggested. For example specialist morphs might occupy different

habitats. Olafsdottir and co-workers (Olafsdóttir et al. 2006) for example, showed that sticklebacks

specialising in living in habitats with little vegetation had reduced nest building behaviour and as a

result weed-living specialists from the same lake mated assortatively with other weed-living

specialists when using nest quality as a mate choice criterion. Disruptive sexual selection is also

known to play a significant role in the divergence of recently evolved African cichlid species (Stelkens

et al. 2008). Here uniquely we demonstrate assortative mating on the basis of morphological traits

that frequently express as discrete forms in the wild, have strong functional significance for

resource acquisition (Adams and Huntingford 2002b) are thought to be under strong selection

pressure and the expression of which is significantly modulated by plasticity effects. This result

indicates that trophic morphology is both a plastic and a magic trait for sticklebacks, thus that

pleiotropy may not always be required for traits to operate as magic traits.

Acknowledgements

We thank Rona Brennan for technical support. M.V.G-P. was supported by a Mexican Council for
Science and Technology (CONACYT) scholarship. The authors declare no conflicts of interest.

Compliance with ethical standards

This study was conducted in accordance with UK legislation under Home Office Licence Number: PPL
70/8794.

References

Adams C, Huntingford F (2002a) Inherited differences in head allometry in polymorphic charr from
Loch Rannoch, Scotland. J Fish Biol 60:515–520. doi: 10.1006/jfbi.2002.1867

Adams C, Huntingford F (2002b) The functional significance of inherited differences in feeding
morphology in a sympatric polymorphic population of Arctic charr. Evol Ecol 16:15–25

Adams CE, Fraser D, Huntingford FA, et al (1998) Trophic polymorphism amongst Arctic charr from
Loch Rannoch, Scotland. J Fish Biol 52:1259–1271

Adams CE, Woltering C, Alexander G (2003) Epigenetic regulation of trophic morphology through
feeding behaviour in Arctic charr, *Salvelinus alpinus*. Biol J Linn Soc 78:43–49. doi:
10.1046/j.1095-8312.2003.00126.x

Baker JA, Räsänen K, Moore JS, Hendry AP (2013) Genetic and plastic contributions to trait
divergence between parapatric habitats: Female life-history traits in three-spine stickleback
within the Misty Lake system. Evol Ecol Res 15:473–487

Boulcott PD, Walton K, Braithwaite VA (2005) The role of ultraviolet wavelengths in the mate-choice
decisions of female three- spined sticklebacks. J Exper Biol 208:1453–1458. doi:
10.1242/jeb.01569

Day T, McPhail J (1996) The effect of behavioural and morphological plasticity on foraging efficiency
in the threespine stickleback (*Gasterosteus* sp.). Oecologia 108:380–388

Dieckmann U, Doebeli M (1999) On the origin of species by sympatric speciation. Nature 400:354–
357

Fitzpatrick B (2012) Underappreciated consequences of phenotypic plasticity for ecological
speciation. Int J Ecol 2012:Article ID 256017

Garduño-Paz M V, Adams CE (2010) Discrete prey availability promotes foraging segregation and
early divergence in Arctic charr, *Salvelinus alpinus*. Hydrobiologia 650:15–26. doi:
10.1007/s10750-009-0055-8

Garduño-Paz M V, Couderc S, Adams CE (2010) Habitat complexity modulates phenotype expression
through developmental plasticity in the threespine stickleback. Biol J Linn Soc 100:407–413.
doi: 10.1111/j.1095-8312.2010.01423.x

Gavrilets S (2004) Fitness landscapes and the origin of species. Princton University Press, Princton

Geiselhardt S, Otte T, Hilker M, Letters E (2012) Looking for a similar partner: host plants shape
mating preferences of herbivorous insects by altering their contact pheromones. Ecol Lett
15:971–7. doi: 10.1111/j.1461-0248.2012.01816.x

Hendry AP, Huber SK, De León LF, et al (2009) Disruptive selection in a bimodal population of
Darwin's finches. Proc Biol Sci 276:753–759. doi: 10.1098/rspb.2008.1321

Heuschele J, Mannerla M, Gienapp P, Candolin U (2009) Environment-dependent use of mate choice
cues in sticklebacks. Behav Ecol 20:1223–1227. doi: 10.1093/beheco/arp123

Kraak S, Bakker T (1998) Mutual mate choice in sticklebacks: attractive males choose big females,
which lay big eggs. Anim Behav 56:859–866

McLellan D, McPhail J (1990) Experimental investigations of the evolutionary significance of sexually
dimorphic nuptial colouration in Gasterosteus aculeatus (L.): the relationship between male
colour and female behaviour. Can J Zool 68:482–492

Milinski M, Bakker TCM (1990) Female sticklebacks use male coloration in mate choice and hence
avoid parasitized males. Nature 344:330–333. doi: 10.1038/344330a0

Milinski M, Griffiths S, Wegner K, et al (2005) Mate choice decisions of stickleback females
predictably modified by MHC peptide ligands. Proc Natl Acad Sci 102:4414–4418

Olafsdóttir GA, Ritchie MG, Snorrason SS (2006) Positive assortative mating between recently
described sympatric morphs of Icelandic sticklebacks. Biol Lett 2:250–252. doi:
10.1098/rsbl.2006.0456

Proulx R, Magnan P (2004) Contribution of phenotypic plasticity and heredity to the trophic
polymorphism of lacustrine brook charr (*Salvelinus fontinalis*). Evol Ecol Res 6:503–522

R Development Core Team (2017) R: A language and environment for statistical computing. R
Foundation for Statistical Computing. Vienna, Austria

Rick IP, Bakker TCM (2008) Color signaling in conspicuous red sticklebacks: Do ultraviolet signals
surpass others? BMC Evol Biol 8:1–9. doi: 10.1186/1471-2148-8-189

Rick IP, Modarressie R, Bakker TCM (2006) UV wavelengths affect female mate choice in three-
spined sticklebacks. Anim Behav 71:307–313. doi: 10.1016/j.anbehav.2005.03.039

Rohlf F (2000) Statistical power comparisions among alternative morphometric methods. Am J Phys
Anthropol 111:463–478

Rohlf F, Slice D (1990) Extensions of the Procrustes method for the optimal superimposition of
landmarks. Syst Zool 39:40–59

Rowland W, Bolyard K, Halpern A (1995) The dual effect of stickleback nuptial coloration on rivals:

manipulation of a graded signal using video playback. Anim Behav 50:267–272

Schluter D (2003) Frequency Dependent Natural Selection during character displacement in

sticklebacks. Evolution 57:1142–1150

Schluter D, McPhail JD (1992) Ecological character displacement and speciation in sticklebacks. Am

Nat 140:85–108

Servedio MR, Van Doorn GS, Kopp M, et al (2011) Magic traits in speciation: "magic" but not rare?

Trends Ecol Evol 26:389–97. doi: 10.1016/j.tree.2011.04.005

Skulason S, Parsons KJ, Svanback R, et al (2019) A way forward with eco evo devo : an extended

theory of resource polymorphism with postglacial fishes as model systems. Biol Rev 94:1786-

1808. doi: 10.1111/brv.12534

Skulason S, Smith TB (1995) Resource polymorphisms in vertebrates. Trends Ecol Evol 10:366–70

Skulason S, Smith TB (1996) The ecology of resource polymorphism in vertebrates - Reply. Trends

Ecol Evol 11:26. doi: 10.1016/0169-5347(96)81063-9

Skulason S, Snorrason S, Jonsson B (1999) Sympatric morphs populations and speciation in

freshwater fish with emphasis on arctic charr. In: Magurran A and May R (eds) Evolution of

Biological Diversity. Oxford University Press, Oxford, pp 70–92

Smith TB, Skulason S (1996) Evolution significance of resource polymorphisms in fishes, amphibians,

and birds. In: Fautin DG (ed) Annual Review of Ecology and Systematics. pp 111–133

Stelkens RB, Pierotti MER, Joyce DA, et al (2008) Disruptive sexual selection on male nuptial

coloration in an experimental hybrid population of cichlid fish. Philos Trans R Soc London B

363:2861–2870. doi: 10.1098/rstb.2008.0049

West-Eberhard MJ (1989) Phenotypic plasticity and the origins of diversity. Annu Rev Ecol Syst

20:249–278

West-Eberhard MJ (2003) Developmental plasticity and evolution. Oxford University Press, Oxford

Wund M, Baker J, Clancy B, et al (2008) A test of the "flexible stem" model of evolution: ancestral

plasticity, genetic accommodation, and morphological divergence in the threespine stickleback

radiation. Am Nat 172:449–462

Wund MA, Valena S, Wood S, Baker JA (2012) Ancestral plasticity and allometry in threespine

stickleback reveal phenotypes associated with derived, freshwater ecotypes. Biol J Linn Soc

Fig 1

Table 1. The frequency with which the male with the higher pelagic-benthic score (more benthic-
like) was chosen and rejected by female sticklebacks exposed to alternative diets (a pelagic like diet
(*Daphnia*) and a benthic-like diet (chironomid larvae).

	Male with more positive P-B score		
Female diet	Chosen	Rejected	χ^2 DF P
Benthic	34	29	0.4; 1; >0.53
Pelagic	9	24	6.8; 1; <0.01

376
377
378

380   Figure Legends
381
382   Figure 1. The landmark configurations used in the morphometric analysis of trophic morphology
383   in sticklebacks. The landmarks are connected to aid visualisation of fish shape. Arrows represent
384   vectors describing deformations that change the mean shape of sticklebacks fed on benthic prey
385   compared with the mean shape of those fed on pelagic prey.