Title: Seascape genomics as a new tool to empower coral reef conservation strategies: an

example on north-western Pacific *Acropora digitifera*.

Running Title: Seascape genomics of north-western Pacific *Acropora digitifera*

List of Authors: Oliver Selmoni[1], Estelle Rochat[1], Gael Lecellier[2,3], Veronique Berteaux-

Lecellier[2], Stéphane Joost[1].

Institutional Affiliations:

[1] Laboratory of Geographic Information Systems (LASIG), Ecole Polytechnique Fédérale de

Lausanne (EPFL), 1015 Lausanne, Switzerland

[2] UMR250/9220 ENTROPIE IRD-CNRS-UR, Labex CORAIL, Promenade Roger-Laroque,

Noumea cedex, New Caledonia

[3] UVSQ, Université de Paris-Saclay, 45 avenue des Etats-Unis, Versailles Cedex, France

Contact Information:

Oliver Selmoni: oliver.selmoni@epfl.ch , +41216935787

Abstract

Coral reefs are suffering a major decline due to the environmental constraints imposed by

climate change. Over the last 20 years, three major coral bleaching events occurred in

concomitance of anomalous heat waves, provoking a severe loss of coral world-wide. Recent

works, however, reported of reefs at which recurrent bleaching events resulted in increased

resistance to thermal stress, suggesting that adaptation might have occurred. The conservation

strategies for preserving the reefs, as they are conceived now, cannot cope with global

climatic shifts. In this regard, researchers advocated the set-up of a preservation framework to

reinforce coral adaptive potential. The main obstacle to this implementation is that studies on

coral adaption are usually hard to generalize at the scale of a reef system. In this work, we

combined a seascape genomics and connectivity analysis to characterize the adaptive

potential of a flagship coral species of the Ryukyu Archipelago (Japan). By associating

genotype frequencies with descriptors of historical environmental conditions, we discovered

six genomic regions hosting polymorphisms that might promote resistance against thermal

stress. Remarkably, annotations of genes in these regions are consistent with molecular roles

in heat responses. Furthermore, we confronted genetic distances between reefs with their

spatial separation according to sea currents to predict connectivity patterns across the region.

The cross-talk between the results of these analyses portrayed the adaptive potential of this

population: we were able to identify reefs carrying potential adaptive traits and to understand

how they disperse to neighbouring reefs. This information was summarized by objective,

quantifiable and mappable indexes covering the whole region that can be extremely useful for

prioritization in conservation plans. This framework is transferable to any coral species and

on any reef system and therefore represents a valuable tool for empowering preservation

efforts dedicated to the protection of coral reef in warming oceans.

Keywords (6-10)

Coral reef – Local adaptation – climate change – seascape genomics – coral bleaching –

conservation genomics – Acropora digitifera – Ryukyu Archipelago

Introduction

Coral reefs are suffering a severe decline due to the effects of climate change (Hughes et al., 2017). Loss of reef is already showing catastrophic consequences for marine wildlife depending on these structures (Pratchett, Thompson, Hoey, Cowman, & Wilson, 2018) and disastrous aftermaths are expected for human economies as well (Moberg & Folke, 1999). One of the major threat to the persistence of this ecosystem is coral bleaching (Bellwood, Hughes, Folke, & Nyström, 2004): a physiological response induced by environmental stress that provokes hard-skeleton coral, the cornerstone of the reef, to separate from the symbiotic microbial algae living within their cells (Mydlarz, McGinty, & Harvell, 2010). These algae, called Symbiodiniacea (LaJeunesse et al., 2018), are crucial for the host sustainment and therefore a persistent bleached state leads coral to death (Mydlarz et al., 2010). Three mass bleaching events occurred over the last 20 years in concomitance with anomalously high sea water temperatures, reason why thermal stress is now considered the main driver of this phenomenon (Hughes et al., 2017). These events struck world-wide and lead to local losses of coral cover up to 50% (Hughes et al., 2018, 2017). Besides, coral bleaching was found to be associated with other stressors linked to human activity, such as ocean acidification, water eutrophication, sedimentation and overfishing (Anthony, Kline, Diaz-Pulido, Dove, & Hoegh-Guldberg, 2008; Ateweberhan et al., 2013; Maina, Venus, McClanahan, & Ateweberhan, 2008).

Conservation efforts facing the plague of coral bleaching aim to restore reefs that underwent severe losses and to limit the impact of future damages (Baums, 2008; Bellwood et al., 2004; Young, Schopmeyer, & Lirman, 2012). For these purposes, two tools are nowadays usually used: the establishment of marine protected areas (MPAs) and the set-up of coral nurseries (Baums, 2008; Bellwood et al., 2004; Young et al., 2012). MPAs are designed zones in which human access and activities are severely restricted in order to alleviate the effects of local

anthropogenic stressors (Lester et al., 2009). Coral nurseries are underwater gardens of transplanted colonies that can then be employed to restore damaged reefs (Baums, 2008; Young et al., 2012). In both cases, researchers advocated the use of design systems accounting for demographic connectivity so that the position of a conservation measure can promote resistance and resilience even on neighbouring sites (Baums, 2008; Krueck et al., 2017; Lukoschek, Riginos, & van Oppen, 2016; Palumbi, 2003; Shanks, Grantham, & Carr, 2003). Even though beneficial effects of this kind of conservation policies have been observed worldwide (Cinner et al., 2016; Rodgers et al., 2017; Selig & Bruno, 2010), these solutions do not confer resistance against the temperature oscillations associated with the last mass bleaching events (Baums, 2008; Hughes et al., 2017). Coral reefs that experienced past thermal stress were found to be more resistant to subsequent heat waves (Hughes et al., 2019; Krueger et al., 2017; Penin, Vidal-Dupiol, & Adjeroud, 2013; Thompson & van Woesik, 2009) but to date this information is neglected in conservation actions (Baums, 2008; Maina, McClanahan, Venus, Ateweberhan, & Madin, 2011). There is an urgent need to understand whether these observations are due to evolutionary processes and, if so, to figure out how the underlying adaptive potential could be included in predictions of climate change responses and in conservation plans (Baums, 2008; Logan, Dunne, Eakin, & Donner, 2014; Maina et al., 2011; van Oppen, Oliver, Putnam, & Gates, 2015).

To this end, seascape genomics tools are likely to play an important role. Seascape genomics is the marine counterpart of Landscape genomics, a branch of population genomics that investigates adaptive potential in field-based experiments (S. Joost et al., 2007; Balkenhol et al., 2017; Rellstab, Gugerli, Eckert, Hancock, & Holderegger, 2015). Samples are collected across a landscape and then genotyped using next-generation-sequencing techniques, providing thousands of genetic variants (Rellstab et al., 2015). Simultaneously, the environmental characterization of the study area is carried out mainly based on remote-

sensing data describing specific local climatic conditions (Rellstab et al., 2015). Genomics

and environmental information are then combined in order to detect genetic polymorphisms

associated with peculiar conditions (*i.e.,* potentially adaptive traits; Rellstab et al., 2015).

This approach has already been applied to many land species and is increasingly used to

analyse marine animals in what is referred to as *seascape genomics* (exhaustively reviewed in

Riginos, Crandall, Liggins, Bongaerts, & Treml, 2016). As far as we know, no seascape

genomics experiment was applied to reef corals yet. In fact, adaptation of these species has

been mostly studied by performing transplantation assays followed by conditioning in

aquarium, a time and resource-demanding approach that often focuses on a couple of

contrasting reef (Howells, Berkelmans, van Oppen, Willis, & Bay, 2013; Krueger et al.,

2017; Palumbi, Barshis, Traylor-Knowles, & Bay, 2014; Sampayo et al., 2016; Ziegler,

Seneca, Yum, Palumbi, & Voolstra, 2017). Genotype-environment associations studies have

also been conducted on corals but they focused on a limited number of markers (Lundgren,

Vera, Peplow, Manel, & van Oppen, 2013) or on a restricted number of locations (Bay &

Palumbi, 2014; Thomas, Kennington, Evans, Kendrick, & Stat, 2017). Unlike these works, a

seascape genomics approach should cover ecologically meaningful spatial scales, allowing to

distinguish pressures of different climatic conditions and to account for confounding effects

of demographic processes (Balkenhol et al., 2017). Consequently, the main (adaptive) and

collateral (connectivity) information generated by this method provides a valuable

contribution for preservation strategies (Baums, 2008; Krueck et al., 2017; Palumbi, 2003).

In this work, we applied a seascape genomics framework to detect coral reefs carrying

potential adaptive traits to show how conservation policies could implement such findings.

Our study focuses on *Acropora digitifera* of the Ryukyu Archipelago in Japan (Fig. 1), an

emblematic species of the Indo-Pacific and flagship organism for studies on corals genomics

(Shinzato et al., 2011). We first analysed the convergence between genomic and

environmental information i) to detect loci potentially conferring a selective advantage and ii) to develop a model describing connectivity patterns. Next, we took advantage of these findings to indicate which reefs were more likely to be carrying adaptive traits and to evaluate how well such sites are interconnected with the rest of the reef system. Finally, we propose an approach to implement the results obtained in conservation planning. In general, our work provides tools at the interface of conservation genomics and environmental sciences likely to empower preservation strategies for coral reefs.

Figure 1. The Study area. The Ryukyu Archipelago extends for more than 1,000 km in the north-western Pacific Ocean. The red circles display the sites where samples were collected for the seascape genomics analysis (adapted from Shinzato, Mungpakdee, Arakaki, & Satoh, 2015).

Materials and methods

Materials and methods

The input genomics data used in this paper comes from a published work from Shinzato *et*

al. (2015) describing the population structure of *A. digitifera* of the Ryuyku Archipelago. Our

framework is structured in two axes of analysis and prediction, one focusing on the presence

of adaptive traits (seascape genomics) and the other on connectivity (Fig. 2). The seascape

genomics analysis (Fig. 2A) crosses genomics data with environmental information at

sampling sites to discover potentially adaptive genotypes. The model describing this

relationship is then used to predict, at the scale of the whole study area, the probability of

presence of adaptive genotypes (Fig. 2B). In the connectivity study (Fig. 2C), we built a

model describing how sea-distances between sites (computed out of remote sensing data)

correspond to genetic-distances. This model is then used to predict connectivity at scale of

the whole study area (Fig. 2D). Finally, the predictions on where the adaptive traits are more

likely to appear and on how the reef system is interconnected allow to assess the adaptive

potential across the whole study area (Fig. 2E).

Figure 2. Workflow of the methods. The starting point for the analysis is the generation of genetic data describing the genotypes observed at different sampling locations (in this example, 4 sampling sites). In the seascape genomics analysis (A) this data is crossed with environmental information to discover genotypes whose frequencies associate with specific climatic conditions (*ENV*). Such genotypes are defined a potentially adaptive. The model describing this link is then applied to environmental data at the scale of the whole reef system (B), to predict the probability of presence of the adaptive genotype (green: high probability, red: low probability). The genetic data is also crossed with sea-current information to build a connectivity model (C) describing how sea-distances between sampling sites correspond to genetic-distances. This model is fit with sea-distances between all the reefs of the study area to predict (D) patterns of connectivity from (*outbound*) and to (*inbound*) each reef (green: high connectivity, red: low connectivity). Finally, predictions on the presence of adaptive genotypes and connectivity patterns are combined to assess the adaptive potential across the study area (E): reefs that are connected with sites likely to carry the adaptive traits will be have a high adaptive potential (green), and inversely for those isolated (red).

Genomic dataset

The genomic data used come from a publicly available dataset consisting of 155 geo-

referenced colonies of *A. digitifera* from 12 sampling locations (13±5 colonies per site) of the

Ryukyu Archipelago in Japan (Fig. 1; Bioproject Accession PRJDB4188). These samples

were sequenced using a Whole-Genome Sequencing approach in the+ scope of a population

genomics study. Details on how samples were collected and processed for genomic analysis

can be found in Shinzato et al., (2015).

Genomics data were processed using the Genome Analysis Toolkit (GATK; McKenna et al.,

2010) framework following the recommended pipeline with modifications necessary to cope

with the absence of reliable databases of known variants for this species (Van der Auwera et

al., 2013). In short, the *A. digitifera* reference genome (v. 1.1, GenBank accession:

GCA_000222465.2; Shinzato et al., 2011) was indexed using bwa (v. 0.7.5a, Li & Durbin,

2009), samtools (v. 1.9, Heng Li et al., 2009) and picardtools (v. 1.95,

http://broadinstitute.github.io/picard) and raw sequencing reads were aligned using the bwa

mem algorithm. The resulting alignments were sorted, marked for duplicate reads, modified

for read-group headers and indexed using picard-tools. Next, each alignment underwent an

independent variant discovery using the GATK HaplotypeCaller tool (using the ERC mode

and setting the --minPruning flag to 10) and genotypes were then jointly called by the GATK

GenotypeGVCFs tool in random batches of 18 samples to match our computational power

(18 CPUs). The variant-calling matrices of the different batches were then joined and filtered

in order to keep only bi-allelic Single Nucleotide Polymorphisms (SNPs) using the GATK

CombineVariants and SelectVariants tools, respectively. This resulted in a raw genotype

matrix counting ~1.2 M of SNPs. Subsequently, we used the GATK VariantAnnotator tool to

annotate variants for Quality-by-depth and filtered for this value (<2), read coverage (>5 and

<100 within a sample), minor allele frequency (>0.05), major genotype frequency (<0.95)

and missing rate of both individuals and SNPs (<0.1) using the GATK VariantFiltrationTool

and custom scripts in the R environment (v. 3.5.1, R Core Team, 2016). This pipeline

produced the filtered genotype matrix consisting of 136 individuals and 15,516 SNPs.

Environmental data

Eleven georeferenced datasets describing atmospheric and seawater conditions were retrieved

from publicly available resources (EU Copernicus Marine Service, 2017; NASA, 2016;

National Oceanic and Atmospheric Administration, 2017; Tab S1). These remote sensing

derived datasets provided monthly or daily variables measured across the whole study area,

with a spatial resolution ranging from 25 km to 4 km (Tab S1). We processed these variables

in the R environment using the *raster* package (v. 2.8, Hijmans, 2016) to compute monthly

averages and standard deviations for the whole studied period. Furthermore, Sea Surface

Temperature (SST) measurements were used to compute a Degree Heat Week (DHW)

frequency index, representing the % of days during which the bi-weekly accumulated heat

stress exceeded the 4 °C (Liu, Strong, & Skirving, 2003; Logan et al., 2014). SST and sea

surface salinity records were combined to produce estimates of seawater pH (Covington &

Whitfield, 1988), dissolved inorganic carbon (Loukos, Vivier, Murphy, Harrison, & Le

Quéré, 2000) and alkalinity (Lee et al., 2006). Bathymetry data (Ryan et al., 2009) were used

to retrieve the depth at sampling locations. Finally, population density data (CIESIN

Columbia University, 2010) were averaged in a 50 km buffer area to produce a surrogate-

variable for anthropogenic pressure (Welle, Small, Doney, & Azevedo, 2017).

We used the geographic coordinates associated with each sample to characterize the

environmental conditions using the QGIS point sampling tool (v. 2.18.25, QGIS

development team, 2009). For predictive step of our study demanding to work at the scale of

the whole reef system (Fig. 2C), we retrieved the shapes of the reefs of the region (UNEP-

WCMC, WorldFish-Center, WRI, & TNC, 2010) and reported them into a regular grid (cell

size of 5x5 km) using QGIS. In this case, we also included reefs from the neighbouring

regions (Taiwan and Philippines, Fig. S7) to avoid border-effects in computations.

Environmental conditions were assigned to each reef-cell using average function of the QGIS

zonal statistics tool.

Seascape genomics

We performed the genotype-environment association analysis using the logistic regression

method implemented within the SamBada software (v. 0.7; Stucki et al., 2017), customized to

fasten multivariate models computation in the Python environment (v. 2.7; Python Software

Foundation, 2018) using the *pandas* (v. 0.23.4; McKinney, 2010) and *statsmodels* (v. 0.9;

Seabold & Perktold, 2010) libraries. The SamBada approach allows to include proxy

variables for genetic structure in the analysis in order to avoid a possible confounding effect

(pattern of neutral genetic variation mimicking signals of adaptation to the local

environment). Here we performed a discriminant analysis of principal components (DAPC)

on the SNPs genotype matrix using the R package *adegenet* (v. 2.1.1; Jombart, 2008). This

procedure highlighted a main separation between two groups of samples along the first

discriminant function (Fig. S1). The latter was therefore used as co-variable in association

models. We then ran the pipeline for the discovery of adaptive signals described hereafter and

summarized in a schematic view in Fig. S2. In order to speed up calculations, the 315

environmental variables were grouped into 29 clusters of highly correlated ($|R|{>}0.7$)

descriptors in the R environment (Fig. S2A). For each of these groups, one variable was

randomly selected to build logistic models against the three genotypes of each SNP (Fig.

S2B). In total, the SamBada instance evaluated 1,349,892 association-models (29

environmental groups vs. 3 x 15,516 SNPs; Fig. S2C, S2D) that were subsequently analysed

in the R environment. For each association-model related to the same environmental variable,

p-values of G-scores (G) and Wald-scores (W) were corrected for multiple testing using the

R *q-value* package (v. 2.14, Storey, 2003). Association-models scoring q<0.001 for both

statistics were deemed significant (Fig. S2E). If a SNP was found in more than one

significant association, only the best model (according to the value of G) was kept. For each

significant association retained, we then calculated all the models built with the involved

genotype vs. the other environmental variables from the same cluster of correlated descriptors

and we looked for the best association model (according to G; Fig. S2F). Finally, we

visualized the logistic regression of each significant association-model using the R *popbio*

library (v. 2.4.4; Stubben & Milligan, 2007) and manually checked the annotations around

significant SNPs on the NCBI genome browser (NCBI, 1988). We set the size of the search

window to ± 250 kbs since the gene(s) potentially linked to a mutation can lay up to

hundreds of kbs away (Brodie, Azaria, & Ofran, 2016; Visel, Rubin, & Pennacchio, 2009).

This window size was chosen because it corresponds approximately to the scaffold N50

statistics of the reference genome (*i.e.* half of the genome is contained within scaffolds of this

size or longer).

Probability of Presence of Adaptive Genotypes

The results of the seascape genomics analysis were then used to predict the probability of

presence of adaptive traits at the scale of the whole Ryukyu Archipelago (Fig. 2B). Indeed,

for each significant association between a genotype and an environmental variable, the

SamBada approach provides parameters of a logistic regression which links the probability of

occurrence of the genotype with the value of the environmental variable (Stucki et al., 2017).

Those logistic models can therefore be used to estimate the probability of presence of the

genetic variant for any value of the environmental variable (Stéphane Joost, 2006; Rochat, E.,

Leempoel, K., Vajana, E., Colli, L., Ajmone-Marsan, P., Joost, 2016). We assumed that

potentially adaptive traits could reach any reef of the region since several studies reported

high gene flow between distant islands of the Archipelago (Nakajima, Nishikawa, Iguchi, &

Sakai, 2010; Nishikawa, 2008; Shinzato et al., 2015). Finally, these probabilities where

combined in an average probability (the arithmetical mean of the five probabilities) of

carrying adaptive traits (P_a) at each reef of the Ryukyu Archipelago.

Sea current data

Daily records of sea surface current were retrieved from publicly available databases (zonal

and meridional surface velocities from the *global-reanalysis-phy-001-030* product; EU

Copernicus Marine Service, 2017) and used to compute the direction and speed of currents in

the R environment using the *raster* library. These day-by-day calculations were then piled-up

to retrieve, for each pixel of the study area (~0.083°), the cumulated speed (the conductance)

toward each of the eight neighbouring pixels. This information was used to calculate

dispersal costs (the inverse of conductance) and was summarized in a transition matrix in the

format of the R *gdistance* package (v. 1.2, van Etten, 2018). For the connectivity analysis

(Fig. 2C), the transition matrix was used to calculated sea-distances (*i.e.* the least-cost path)

between sampling sites of the genotyped colonies. For the connectivity predictions (Fig. 2D),

we calculated the sea-distances between all the reefs of the study area, whose coordinates

were the centroids of the reef-cells computed as described in the environmental variables

section. Importantly, for every couple of reefs (for instance $reef_1$ and $reef_2$) two sea-distances

were computed, one per direction (*i.e.* from $reef_1$ to $reef_2$ and from $reef_2$ to $reef_1$).

Connectivity analysis

Genomics data was processed by the means of a principal component analysis (PCA; Fig. S3)

on the filtered genotype matrix. Prior to PCA, in order to cope with missing genotypes in the

SNP matrix, we applied the *random* imputation method of the LEA R package (v.2.4.0;

Frichot & François, 2015). The values of the principal components (PCs) were used to

calculate the Euclidean distances between each couple of samples that were subsequently

averaged by site in order to obtain Genetic Distances (GDs) between reefs. Next, we built a

linear model (hereafter referred to as the connectivity model) to estimate GDs out of log-

transformed sea-distances and ran a t-test to check for the statistical significance of the

relationship. Since we wanted to consider the genetic distances within reef (*i.e.* sea-distance =

0) we added +0.005 to all the sea-distances prior to the log-transformation to avoid obtaining

non-definite values (*i.e.* the logarithm of zero). As a comparison, we also built a connectivity

model using log-transformed aerial-distances (*i.e.* Euclidean distance of coordinates) as

independent variable while maintaining GD as response variable. We then confronted models

based on sea- and aerial-distance by comparing the coefficients of determination (R^2) and the

Akaike information criterion (AIC; Bozdogan, 1987).

Connectivity Predictions

We then used the connectivity model to predict GDs between any couple of reefs of Ryukyu

Archipelago. This was done by translating sea-distances in GDs using the connectivity model

trained during the connectivity analysis. Importantly, since there were two sea-distances for

every couple of reefs (*i.e.* from $reef_1$ to $reef_2$ and *vice versa*), two GDs were computed as

well.

Based on these predictions, we were able to calculate, for each reef-cell, two indices (Fig.

S4):

- *outbound connectivity* (OC; Fig. S4B): the % of cells reachable from the focal cell

within a GD-threshold (tGD).

- *inbound connectivity* (IC; Fig. S4C): % of cells that can reach the focal cell within a tGD.

Since these indices were conceived to display contrasts in connectivity among reefs of the

Ryukyu Archipelago, we looked for the tGD value maximizing their variances (Fig. S5). The

values corresponding to the 25%-quantile and the 60%-quantile of the GD distribution were

those boosting the differences in both connectivity indices. To avoid the artificial bias of

border effects, we preferred working at a more local scale and set tGD at the 25%-quantile of

the GD distribution (*i.e.* tGD=79.861).

Evaluation of the adaptive potential

The adaptive potential was evaluated by combining the predictions on the presence of

adaptive traits and on connectivity patterns across the reefs of Ryuyku Archipelago (Fig. 2E).

The previously computed average probability of carrying adaptive traits (P_a) is used to

identify the putatively adaptive reefs. These are the reefs approaching the maximal values of

P_a ($P_a > 0.6$). We then used the GDs calculated during the connectivity analysis to retrieve, for

each reef-cell, the GD from the closest putatively adapted reef. This results in an index

describing the adaptive potential of every reef of the study area: the distance from the closest

adaptive reef.

Data Access and Code Availability

All the data and codes used in this article are publicly available on Zenodo

(DOI: 10.5281/zenodo.2643454).

Results

Seascape Genomics

We detected 15 significant genotype-environment association-models (A1-A15; q_G and

$q_W<0.001$; Tab.1), referring to genetic markers from 6 distinct scaffolds of the *A. digitifera*

reference genome. Among them, one association was related to SST variation in April, one to

changes in sea current velocity in May and 13 to DHW frequency (or to a highly correlated

variable such as variation in cloud fraction in August).

The SNP in the significant association-model A1 is related to SST standard deviation in April

and lies within a genomic window (± 250 kb) counting 18 predicted genes, 7 of which are

annotated with a known function (Box S1). Association-models A2 to A5 include four SNP

lying at two opposite sides of a long scaffold (~1.3 Mb). The SNPs located on one side

(involved in association-models A2 and A3) are associated with DHW frequency and are

surrounded by 17 predicted genes (of which 12 have a putative encoded protein with known

function; Box S2) while the SNPs located on the other side (association-models A4 and A5)

are related to Sea current velocity variance in May and are surrounded by twelve putative

genes (Box S4). All the remaining association-models (A6 to A15) involved only DHW.

Association-model A6 concerns a SNP surrounded by 14 putative genes, but only one is

annotated, the related predicted protein being *JNK1/MAPK8-associated membrane protein-*

like (NCBI accession: LOC107334815; Box S5). In association-model A7, the SNP of

interest lies within in a window showing eleven predicted genes (only two with a known

match by similarity; Box S6). Association-models A8 to A14 refer to seven SNPs present in

the same scaffold (~195 kb long). This region contains five putative genes of which 2

annotated by similarity (Box S6). In association-model A15, finally, the SNP associated with

DHW frequency is located on a ~75 kb scaffold that contains only one putative gene

predicted as *ATP-dependent DNA helicase RecQ-like* (LOC107328703; Box S7).

Table 1. Significant genotype-environment associations. The seascape genomics analysis using the SamBada method detected 15 significant association-models (q_G and q_W<0.001). This table shows, for each association-model, the genomic position (scaffold of the reference genome and relative situation), the genotype involved (0 and 2=homozygous, 1=heterozygous), the concerned environmental variables (in brackets is displayed a variable from the same group of correlated descriptors that resulted in a model with a higher G-score, if any), the q-values associated with the G-score (q_G) and the Wald-score (q_W), the number of predicted genes in a ± 250 kb window around the SNP (with the number of SNPs annotated by similarity with a known gene, in brackets) and the annotations from the reference genome that are in the neighbourhood (± 250 kb) of the SNP and that corroborate an adaptive role against the respective environmental variable.

Association ID	Scaffold ID	Position on Scaffold	GT	Environmental Variable	q_G	q_W	Number of Predicted Genes	Relevant Genome Annotations
A1	NW_015441190.1	582812	1	SST variance in April	1.5E-08	2.6E-04	18 (7)	*Helicase RecQ at 60 kbs*
A2	NW_015441080.1	208387	0	DHW frequency (Cloud Fraction variance in August)	5.8E-08	2.0E-04	17 (12)	*Cytochrome p450 at 150 kbs*
A3	NW_015441080.1	208400	0	DHW frequency	2.3E-09	6.4E-05	17 (12)	*Cytochrome p450 at 150 kbs*
A4	NW_015441080.1	963844	0	DHW frequency (Cloud Fraction variance in August)	4.0E-09	5.8E-04	12 (7)	
A5	NW_015441080.1	963851	0	Sea current velocity variance in May	1.5E-07	9.3E-04	12 (7)	
A6	NW_015441121.1	665651	0	DHW frequency	5.9E-08	1.6E-04	14 (1)	*MAPK8/JNK1 at 200 kbs*
A7	NW_015441282.1	27616	0	DHW frequency	5.5E-06	5.8E-04	7 (2)	*Helicase Q at 250 kbs*
A8	NW_015441785.1	16151	1	DHW frequency	6.1E-06	8.6E-04	5 (2)	*Helicase RecQ at 150kbs*
A9	NW_015441785.1	68415	1	DHW frequency	5.8E-08	1.1E-04	5 (2)	*Helicase RecQ at 90kbs*
A10	NW_015441785.1	68439	1	DHW frequency	3.1E-07	3.0E-04	5 (2)	*Helicase RecQ at 90kbs*
A11	NW_015441785.1	68444	1	DHW frequency	5.2E-06	8.6E-04	5 (2)	*Helicase RecQ at 90kbs*
A12	NW_015441785.1	68515	1	DHW frequency (Cloud Fraction variance in August)	5.9E-08	1.6E-04	5 (2)	*Helicase RecQ at 90kbs*
A13	NW_015441785.1	68521	1	DHW frequency	5.8E-08	2.0E-04	5 (2)	*Helicase RecQ at 90kbs*
A14	NW_015441785.1	68528	1	DHW frequency	6.9E-08	6.7E-04	5 (2)	*Helicase RecQ at 90kbs*
A15	NW_015442197.1	32233	0	DHW frequency	4.4E-06	8.6E-04	1 (1)	*Helicase RecQ at 20kbs*

Probability of Presence of Adaptive Genotypes

The significant association-models of the seascape genomics analysis were then used as

starting point for predicting the probability of presence of adaptive genotypes at the scale of

the whole Ryukyu Archipelago (Fig. 2B). These genotypes can be theoretically associated to

any environmental descriptor but, since most of the putative adaptive genotypes were found

to be associated with DHW frequency (Tab. 1), we designated this variable as the

environmental pressure of interest (association models A3, A6, A7, A9 and A15, Tab. 1). The

average of these probabilities (P_a) ranges from 0.40 to 0.62 (Fig. S7). Potentially adapted

reefs (*i.e.* reefs with P_a>0.6) are located in the islands of Okinawa and Miyako, in the remote

Japanese islands east of the Ryukyu Archipelago and in the Northern part of Philippines (Fig.

S7).

Connectivity modelling

The two dispersal models respectively based on sea- and aerial-distances resulted in a

significant association between genetic-distance (GD) and spatial separation ($p<10^{-9}$, Fig.

S6). We found that the model based on sea-distances better explained the variation in GD

(R2=0.57) than the aerial-distances model (R2=0.45). The AIC is lower for the model based

on sea-distances (AIC=315) than for the aerial-distances model (AIC=343). The sea-

distances dispersal model was therefore used for the following inbound and outbound

connectivity predictions.

The reefs around the islands of Tokara were those with the highest values of Inbound

Connectivity index (IC=0.49±0.01; Fig. 3a). The reefs in the central islands of Ryukyu

Archipelago (Okinawa and Amami) showed higher values of the IC index (IC=0.41±0.03;

Fig. 3a) as compared to those located at the edges of the reef system (Yaeyama in South,

Osumi in the North) which showed the lowest values (IC=0.27±0.03). As regards the

Outbound Connectivity index, reefs located in the southern part (Yaeyama and Miyako) are

those predicted with the highest values (OC=0.42±0.03; Fig. 3b) while those in the northern

part (Osumi and Tokara) display the lowest ones (OC=0.20±0.13).

Figure 3. Connectivity indices. The two maps show how dispersal to and from specific reefs is expected to vary across the Ryukyu Archipelago. Using the first quartile of the GD distribution as a threshold value (tGD=80.32), inbound connectivity (IC, a) was calculated as the % of cells that can reach the focal cell within tGD. Conversely, outbound connectivity (OC, b) represents the % of cells reachable from the focal cell within tGD.

a) Inbound Connectivity   b) Outbound Connectivity

Evaluation of the adaptive potential

380 Reefs in the northern Islands of the Archipelago (Osumi, Tokara and Amami) and in the

381 southern Islands of Yaeyama were those predicted as the genetically farthest (GD=79.4±0.3

382 and GD=79.0±0.1, respectively) from reefs potentially adapted against thermal stress (P_a>0.6;

383 Fig. 4). Conversely, the southern islands of Miyako and those around Okinawa resulted as the

384 closest ones, despite a considerable variance (GD=73.6±3.1 and GD=76.4±2.8, respectively).

Figure 4. Distance from Potentially Adaptive Reefs. The map displays the predicted genetic-distance (GD) from the closest reef expected to carry adaptive genotypes against thermal stress. Adaptive sites are designed as those having a cumulative probability of carrying adaptive genotypes (P_a, Fig. S7) higher than 0.6. The GDs are computed out of sea-distances using the dispersal models developed in the connectivity analysis.

Discussion

Adaptation to thermal stress

Thermal stress is expected to be one of the major threat to coral reef survival and the quest for adaptive traits is becoming of paramount importance (Baums, 2008; Logan et al., 2014; Maina et al., 2011). In this work, the seascape genomics analysis of *A. digitifera* of the Ryukyu Archipelago revealed the presence of 6 genomic regions hosting genetic variants that might confer a selective advantage against thermal stress (Tab. 1). None of these SNPs directly lay within a coding sequence of a putative gene, but this is rarely the case for causative-mutations (Brodie et al., 2016). In fact, genetic variants in intergenic regions that play a modulatory action on the expression of neighbouring genes are more frequent and can

influence loci at 1-2 Mb of distance (Brodie et al., 2016). The fragmentation of the reference

genome forced us to limit our search window to ±250 Kb around each SNP and still we found

several annotations corroborating a response to heat stress.

The SNP in association-model A6 is related to "JNK1/MAPK8-associated membrane

protein", the closest gene with a predicted function (Box S4). Mitogen-activated-protein

kinases (MAPKs) are proteins known to be involved in cellular responses to stress in a wide

range of taxa (Neupane, Nepal, Benson, Macarthur, & Piya, 2013) and the c-Jun-N-terminal

kinase (JNK) was experimentally proved to be activated under thermal stress in coral

Stylopohora pistillata (Courtial et al., 2017). Of the 26,060 predicted genes in the *A.*

digitifera genome (Shinzato et al., 2011)*, 35 are annotated as "mitogen-activated protein

kinase" and only one refers to the "JNK1/MAPK8-associated membrane protein". Another

annotation supports these findings and concerns gene-models predicted as "ATP-dependent

DNA helicase Q" and "RecQ" (A1, A7-15; Box S1, Box S5-7). These are proteins involved

in the DNA repairing mechanism to UV-light damage in prokaryotes (Courcelle & Hanawalt,

1999) and for which light-stress driven effects were observed in eukaryotic cells as well

(Sharma, Doherty, Brosh, & Jr, 2006). Annotations related to these proteins are not

uncommon (43 in the reference genome) but it is remarkable to find them around significant

SNPs from four distinct genomic regions. A third situation occurs with genes assigned to

cytochrome P450 (A2, A3; Box S2), a large family of proteins involved in cellular responses

to various types of stresses including heat shocks (Rosic, Pernice, Dunn, Dove, & Hoegh-

Guldberg, 2010; Sang, Ma, Qiu, Zhu, & Lei, 2012; Yampolsky et al., 2014) and for which

differential patterns of gene expression were observed in *Acropora hyacinthus* under distinct

thermal regimes (Palumbi et al., 2014). We detected 4 annotations related to these genes in

the proximity of one region with significant SNPs (models A2 and A3) but the relatively high

frequency of this gene-family (62 occurrences across the genome) invites to a cautious

interpretation.

Seascape/Landscape genomics studies are susceptible to high false discovery rates, especially

when neutral genetic structure is not accounted for (Rellstab et al., 2015). To take this

element into account, we processed data following a conservative pipeline and used models

explicitly integrating demographic processes. We also set a restrictive threshold to filter

significant association-models (*i.e.* $q<0.001$ in two distinct tests). Most of the significant

association-models found were linked to DHW, most probably because this variable

represents one of the main constraint to coral survival (Hughes et al., 2017). It is also possible

that the initial application of a sampling scheme adapted to seascape genomics (unlike the

one used by Shinzato et al. 2015 who did not consider environmental variability) would have

increased sensitivity to other types of adaptive signals (Riginos et al., 2016).

All the genotypes we found associated to thermal stress also displayed annotations suggesting

a functional role in cellular heat responses. Nonetheless, annotations analysis is only a first

step in the validation of the association-models detected. Ideally, further assays as reciprocal

transplantations (Palumbi et al., 2014), experimental conditioning (Krueger et al., 2017) and

molecular analysis (Courtial et al., 2017) could ascertain and help quantifying the link

between each genotype and the putative selective advantage it might confer. Nevertheless,

the high frequency of SNPs associated to helicase RecQ genes family together with their

location on 4 distinct scaffolds is strongly unlikely. This suggests that the maintenance of

ability to open the double strand of DNA for transcription, replication and repair is a key

point for *A. Digitifera* in Ryukyu archipelago to resist on the pressure applied by DHW.

Patterns of connectivity

Understanding connectivity patterns across a reef system is essential in order to plan effective

management strategies. This is usually carried out through the quantification of the dispersal

trajectories of larvae (Storlazzi, van Ormondt, Chen, & Elias, 2017) or through the study of

the population structure using genetic data (*e.g.* Lukoschek et al., 2016). The first approach

has the main drawback of being difficult and expensive because it requires an *a priori*

knowledge of the dispersal of the studied species (McCook et al., 2009). In population

genetics data, on the other hand, sampling often occurs at a few locations and the results are

therefore hard to generalize to a whole seascape extent (Beger et al., 2014). The method we

applied combines these two approaches. Firstly, genetic data are used to calculate genetic

distances between the sampled locations. The latter were then used to train dispersal models,

which could in turn predict genetic distances at the scale of the whole reef system.

In model training we found that sea-distances clearly represented a better indicator of spatial

separation, compared with aerial-distances (Fig. S6). This happens probably because coral

dispersal is driven by the water flow (Paris-Limouzy, 2011), which is strongly asymmetrical

in this region (north-east oriented) because of the Kuroshio current (Nishikawa, 2008). By

fitting this dispersal model with sea-distances among any reef of the region we were able to

evaluate the patterns of connectivity. Reefs in the extreme North (Osumi) face both *inbound*

and *outbound* isolation (Fig. 3), a situation that might lead to inbreeding depression (Keller &

Waller, 2002). Reefs in the extreme south-west of the system (Yaeyama) appear also

endangered: despite their proximity to Miyako, the strong north-eastward directionality of

sea-currents contrasts the arrival of recruits from the rest of the Archipelago.

The dispersal framework we propose here is potentially applicable to any coral species for

which geo-referenced genetic data are available and the indices associated can be extremely

valuable in management planning (Beger et al., 2014). However, a substantial part of GD

variation remains unexplained by our models and connectivity models developed at a finer

geographical and temporal scales might better describe the situation (Storlazzi et al., 2017).

Conservation and Adaptive Potential

Conservation strategies to preserve coral reefs need to account for and even to reinforce the

importance of the adaptive potential to resist to thermal stress (Baums, 2008; Logan et al.,

2014; Maina et al., 2008; van Oppen et al., 2015). In the perspective of its integration in

prioritization programs, adaptive potential needs to be objective and quantifiable (OECD,

2017). Previous works suggested to use climatic records (Logan et al., 2014; Maina et al.,

2011), but this approach neglects how organisms modulate a response in front of an

environmental constraint. The use of genetic data has also been advocated, but findings from

population genetics analyses are often hard to be extrapolated at large scales and therefore

includes in conservation plans (Beger et al., 2014). Furthermore, the established guidelines

(e.g. Beger et al., 2014) discuss how to reinforce conservation planning with genetic data (*i.e.*

a few loci), which is less informative for adaptive processes, compared to genomic data. Here

we showed how a combination seascape genomics and connectivity analysis makes it

possible to assess the adaptive potential at the scale of a reef system.

In the specific case of the Ryukyu Archipelago, we found that reef from Miyako and

Okinawa islands are those more likely to carry adaptive traits against thermal stress (Fig. 4).

Previous works reported indeed that adapted colonies from Okinawa might have resisted

during the 1998 and 2001 bleaching events (Van Woesik, Irikawa, & Loya, 2004). In

contrast, reefs in the northern part of the Archipelago (Amami, Tokara and Osumi) did not

experience the same thermal stress before and therefore are not predicted to show the same

adaptive capacities (Fig. 4).

The indicators on adaptive potential and connectivity produced in this work provide valuable

descriptors of the state of the reef. Furthermore, they can be used for predicting the expected

impact of distinct preservation measures and support therefore spatial planning in

conservation (Box. 1). It is unknown whether the connectivity patterns and the adaptive

signals observed in *A. digitifera* are representative for those of other taxa. In this concern, we

advocate that future research should focus on a multi-species implementation of the methods

proposed here.

Conclusions

This study highlights the value of the seascape genomics approach to support the

conservation of corals. We applied this approach to a flagship coral species of Ryukyu

Archipelago and identified genetic variants that might underpin adaptation to thermal stress.

Coupling this information with a genetic analysis of connectivity enabled the evaluation of

the adaptive potential at the scale of the whole study area. The outputs of this analysis are

quantitative indices likely to support objective prioritization of reefs in conservation plans.

This framework is transferable to any coral species on any seascape and therefore constitutes

a useful conservation tool to take into account and evaluate the genomic adaptive potential of

coral reefs worldwide.

Box. 1 Adaptive potential in conservation scenarios.

In this box we show how information on adaptive potential can be used to compare different conservation scenarios. We focus on two typical preservation initiatives, marine protected areas (MPAs) and coral nurseries (CNs). For each of them, we compare how the set-up of the measure at two different locations differentially drives the adaptive potential of the reef system.

The starting point for the predictions is the map showing the distribution of the probability of presence of adaptive genotypes (Fig. S7). We modify the values in this map to imitate the effects of preservation strategies. In the case of MPAs, we maintain potentially adaptive reefs ($P_a > 0.6$) only at locations where the protected area is established. The underlying idea is to emulate the way MPAs reduces coral loss due to local stressors (Selig & Bruno, 2010). In the CNs scenarios, potentially adaptive reefs ($P_a > 0.6$) are added where absent. Here we follow the good practices for this kind of measure that suggest to transplant adapted colonies at sites lacking them (Baums, 2008). After modifying the values of P_a, we re-calculate the genetic distance (GD) from adaptive sites (Fig. 4). This is the variable that we use to describe adaptive potential. We then compare differences in this value (ΔGD) between different conservation scenarios to investigate beneficial and negative effects.

Marine Protected Areas

We compare two MPAs established in Miyako-Yaeyama area (MPA1) or in Okinawa (MPA2), both covering the same surface ($\sim$15'000 km^2). The overall difference in GD (ΔGD=+0.17) indicates that MPA2 globally reduces GDs from potentially adapted reefs more than MPA1. This is probably due to the fact that MPA2 includes more reefs that are putatively adaptive (Fig. S7) and is better connected to the reef-dense area around Amami. MPA1, on the other hand, is far from this reef-dense area and shows beneficial effects only in Yaeyama.

Coral Nurseries

Two CNs emplacements were compared: in Amami (CN1) or in Tokara (CN2). The overall difference in GD (ΔGD=-0.28) suggests a more beneficial effect on adaptive potential under CN2, rather than CN1. In fact, the existence of a coral nursery in Amami (CN1) could provide adaptive genetic material to a larger number of reefs, compared with a set-up established in Tokara (CN2), because of a higher outbound-connectivity potential (Fig. 3b).

Acknowledgements

517 The authors received no specific financial support for the research, authorship, and/or

518 publication of this article. The authors have no conflict of interest to declare.

References

Anthony, K. R. N., Kline, D. I., Diaz-Pulido, G., Dove, S., & Hoegh-Guldberg, O. (2008).

Ocean acidification causes bleaching and productivity loss in coral reef builders.

Proceedings of the National Academy of Sciences of the United States of America,

105(45), 17442–17446. https://doi.org/10.1073/pnas.0804478105

Ateweberhan, M., Feary, D. A., Keshavmurthy, S., Chen, A., Schleyer, M. H., & Sheppard,

C. R. C. (2013). Climate change impacts on coral reefs: Synergies with local effects,

possibilities for acclimation, and management implications. *Marine Pollution Bulletin*,

74(2), 526–539. https://doi.org/10.1016/j.marpolbul.2013.06.011

Balkenhol, N., Dudaniec, R. Y., Krutovsky, K. V., Johnson, J. S., Cairns, D. M., Segelbacher,

G., … Joost, S. (2017). Landscape Genomics: Understanding Relationships Between

Environmental Heterogeneity and Genomic Characteristics of Populations (pp. 1–62).

Springer, Cham. https://doi.org/10.1007/13836_2017_2

Baums, I. B. (2008). A restoration genetics guide for coral reef conservation. *Molecular*

Ecology, *17*(12), 2796–2811. https://doi.org/10.1111/j.1365-294X.2008.03787.x

Bay, R. A., & Palumbi, S. R. (2014). Multilocus Adaptation Associated with Heat Resistance

in Reef-Building Corals. *Current Biology*, *24*(24), 2952–2956.

https://doi.org/10.1016/j.cub.2014.10.044

Beger, M., Selkoe, K. A., Treml, E., Barber, P. H., von der Heyden, S., Crandall, E. D., …

Riginos, C. (2014). Evolving coral reef conservation with genetic information. *Bulletin*

of Marine Science, *90*(1), 159–185. https://doi.org/10.5343/bms.2012.1106

Bellwood, D. R., Hughes, T. P., Folke, C., & Nyström, M. (2004). Confronting the coral reef

crisis. *Nature*, *429*(6994), 827–833. https://doi.org/10.1038/nature02691

Bozdogan, H. (1987). Model selection and Akaike's Information Criterion (AIC): The

general theory and its analytical extensions. *Psychometrika*, *52*(3), 345–370.

https://doi.org/10.1007/BF02294361

Brodie, A., Azaria, J. R., & Ofran, Y. (2016). How far from the SNP may the causative genes

be? *Nucleic Acids Research*, *44*(13), 6046–6054. https://doi.org/10.1093/nar/gkw500

CIESIN Columbia University. (2010). Gridded Population of the World (GPW), v3.

Retrieved January 18, 2019, from http://sedac.ciesin.columbia.edu/data/set/gpw-v3-

population-count

Cinner, J. E., Huchery, C., MacNeil, M. A., Graham, N. A. J., McClanahan, T. R., Maina, J.,

… Mouillot, D. (2016). Bright spots among the world's coral reefs. *Nature*, *535*(7612),

416–419. https://doi.org/10.1038/nature18607

Courcelle, J., & Hanawalt, P. C. (1999). RecQ and RecJ process blocked replication forks

prior to the resumption of replication in UV-irradiated Escherichia coli. *Molecular &*

General Genetics : MGG, *262*(3), 543–551. Retrieved from

http://www.ncbi.nlm.nih.gov/pubmed/10589843

Courtial, L., Picco, V., Grover, R., Cormerais, Y., Rottier, C., Labbe, A., … Ferrier-Pagès, C.

(2017). The c-Jun N-terminal kinase prevents oxidative stress induced by UV and

thermal stresses in corals and human cells. *Scientific Reports*, *7*(1), 45713.

https://doi.org/10.1038/srep45713

Covington, A. K., & Whitfield, M. (1988). Recommendations for the determination of pH in

sea water and estuarine waters. *Pure and Applied Chemistry*, *60*(6), 865–870.

https://doi.org/10.1351/pac198860060865

EU Copernicus Marine Service. (2017). Global Ocean - In-Situ-Near-Real-Time

Observations. Retrieved February 2, 2017, from http://marine.copernicus.eu

Frichot, E., & François, O. (2015). LEA : An R package for landscape and ecological

association studies. *Methods in Ecology and Evolution*, *6*(8), 925–929.

https://doi.org/10.1111/2041-210X.12382

Hijmans, R. J. (2016). raster: Geographic Data Analysis and Modeling. Retrieved from

https://cran.r-project.org/package=raster

Howells, E. J., Berkelmans, R., van Oppen, M. J. H., Willis, B. L., & Bay, L. K. (2013).

Historical thermal regimes define limits to coral acclimatization. *Ecology*, *94*(5), 1078–

1088. https://doi.org/10.1890/12-1257.1

Hughes, T. P., Anderson, K. D., Connolly, S. R., Heron, S. F., Kerry, J. T., Lough, J. M., …

Wilson, S. K. (2018). Spatial and temporal patterns of mass bleaching of corals in the

Anthropocene. *Science*, *359*(6371), 80–83. https://doi.org/10.1126/science.aan8048

Hughes, T. P., Kerry, J. T., Álvarez-Noriega, M., Álvarez-Romero, J. G., Anderson, K. D.,

Baird, A. H., … Wilson, S. K. (2017). Global warming and recurrent mass bleaching of

corals. *Nature*, *543*(7645), 373–377. https://doi.org/10.1038/nature21707

Hughes, T. P., Kerry, J. T., Connolly, S. R., Baird, A. H., Eakin, C. M., Heron, S. F., …

Torda, G. (2019). Ecological memory modifies the cumulative impact of recurrent

climate extremes. *Nature Climate Change*, *9*(1), 40–43. https://doi.org/10.1038/s41558-

018-0351-2

Jombart, T. (2008). adegenet: a R package for the multivariate analysis of genetic markers.

Bioinformatics, *24*(11), 1403–1405. https://doi.org/10.1093/bioinformatics/btn129

Joost, S. (2006). *The geographical dimension of genetic diversity: a GIScience contribution*

for the conservation of animal genetic resources. EPFL. https://doi.org/10.5075/EPFL-

THESIS-3454

Joost, S., Bonin, A., Bruford, M. W., Després, L., Conord, C., Erhardt, G., & Taberlet, P.

(2007). A spatial analysis method (SAM) to detect candidate loci for selection: Towards

a landscape genomics approach to adaptation. *Molecular Ecology*, *16*(18), 3955–3969.

https://doi.org/10.1111/j.1365-294X.2007.03442.x

Keller, L., & Waller, D. M. (2002). Inbreeding effects in wild populations. *Trends in Ecology*

& Evolution, *17*(5), 230–241. https://doi.org/10.1016/S0169-5347(02)02489-8

Krueck, N. C., Ahmadia, G. N., Green, A., Jones, G. P., Possingham, H. P., Riginos, C., …

Mumby, P. J. (2017). Incorporating larval dispersal into MPA design for both

conservation and fisheries. *Ecological Applications*, *27*(3), 925–941.

https://doi.org/10.1002/eap.1495

Krueger, T., Horwitz, N., Bodin, J., Giovani, M.-E., Escrig, S., Meibom, A., & Fine, M.

(2017). Common reef-building coral in the Northern Red Sea resistant to elevated

temperature and acidification. *Royal Society Open Science*, *4*(5), 170038.

https://doi.org/10.1098/rsos.170038

LaJeunesse, T. C., Parkinson, J. E., Gabrielson, P. W., Jeong, H. J., Reimer, J. D., Voolstra,

C. R., & Santos, S. R. (2018). Systematic Revision of Symbiodiniaceae Highlights the

Antiquity and Diversity of Coral Endosymbionts. *Current Biology*, *28*(16), 2570–

2580.e6. https://doi.org/10.1016/j.cub.2018.07.008

Lee, K., Tong, L. T., Millero, F. J., Sabine, C. L., Dickson, A. G., Goyet, C., … Key, R. M.

(2006). Global relationships of total alkalinity with salinity and temperature in surface

waters of the world's oceans. *Geophysical Research Letters*, *33*(19), L19605.

https://doi.org/10.1029/2006GL027207

Lester, S., Halpern, B., Grorud-Colvert, K., Lubchenco, J., Ruttenberg, B., Gaines, S., …

Warner, R. (2009). Biological effects within no-take marine reserves: a global synthesis.

Marine Ecology Progress Series, *384*, 33–46. https://doi.org/10.3354/meps08029

Li, H., & Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler

transform. *Bioinformatics*, *25*(14), 1754–1760.

https://doi.org/10.1093/bioinformatics/btp324

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., … 1000 Genome

Project Data Processing Subgroup, 1000 Genome Project Data Processing. (2009). The

Sequence Alignment/Map format and SAMtools. *Bioinformatics (Oxford, England)*,

25(16), 2078–2079. https://doi.org/10.1093/bioinformatics/btp352

Liu, G., Strong, A. E., & Skirving, W. (2003). Remote sensing of sea surface temperatures

during 2002 Barrier Reef coral bleaching. *Eos, Transactions American Geophysical*

Union, *84*(15), 137–141. https://doi.org/10.1029/2003EO150001

Logan, C. A., Dunne, J. P., Eakin, C. M., & Donner, S. D. (2014). Incorporating adaptive

responses into future projections of coral bleaching. *Global Change Biology*, *20*(1),

125–139. https://doi.org/10.1111/gcb.12390

Loukos, H., Vivier, F., Murphy, P. P., Harrison, D. E., & Le Quéré, C. (2000). Interannual

variability of equatorial Pacific CO_2 fluxes estimated from temperature and salinity

data. *Geophysical Research Letters*, *27*(12), 1735–1738.

https://doi.org/10.1029/1999GL011013

Lukoschek, V., Riginos, C., & van Oppen, M. J. H. (2016). Congruent patterns of

connectivity can inform management for broadcast spawning corals on the Great Barrier

Reef. *Molecular Ecology, 25*(13), 3065–3080. https://doi.org/10.1111/mec.13649

Lundgren, P., Vera, J. C., Peplow, L., Manel, S., & van Oppen, M. J. H. (2013). Genotype -

environment correlations in corals from the Great Barrier Reef. *BMC Genetics*, *14*(1), 9.

https://doi.org/10.1186/1471-2156-14-9

Maina, J., McClanahan, T. R., Venus, V., Ateweberhan, M., & Madin, J. (2011). Global

gradients of coral exposure to environmental stresses and implications for local

management. *PLoS ONE*, *6*(8), e23064. https://doi.org/10.1371/journal.pone.0023064

Maina, J., Venus, V., McClanahan, T. R., & Ateweberhan, M. (2008). Modelling

susceptibility of coral reefs to environmental stress using remote sensing data and GIS

models. *Ecological Modelling*, *212*(3), 180–199.

https://doi.org/10.1016/j.ecolmodel.2007.10.033

McCook, L. J., Almany, G. R., Berumen, M. L., Day, J. C., Green, A. L., Jones, G. P., …

Thorrold, S. R. (2009). Management under uncertainty: guide-lines for incorporating

connectivity into the protection of coral reefs. *Coral Reefs*, *28*(2), 353–366.

https://doi.org/10.1007/s00338-008-0463-7

McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., …

DePristo, M. A. (2010). The Genome Analysis Toolkit: a MapReduce framework for

analyzing next-generation DNA sequencing data. *Genome Research*, *20*(9), 1297–1303.

https://doi.org/10.1101/gr.107524.110

McKinney, W. (2010). Data Structures for Statistical Computing in Python. Retrieved from

http://conference.scipy.org/proceedings/scipy2010/mckinney.html

Moberg, F., & Folke, C. (1999). Ecological goods and services of coral reef ecosystems.

Ecological Economics, *29*(2), 215–233. https://doi.org/10.1016/S0921-8009(99)00009-9

Mydlarz, L. D., McGinty, E. S., & Harvell, C. D. (2010). What are the physiological and

immunological responses of coral to climate warming and disease? *The Journal of*

Experimental Biology, *213*(6), 934–945. https://doi.org/10.1242/jeb.037580

Nakajima, Y., Nishikawa, A., Iguchi, A., & Sakai, K. (2010). Gene Flow and Genetic

Diversity of a Broadcast-Spawning Coral in Northern Peripheral Populations. *PLoS*

ONE, *5*(6), e11149. https://doi.org/10.1371/journal.pone.0011149

NASA. (2016). NASA Earth Observations (NEO). Retrieved May 10, 2017, from

http://neo.sci.gsfc.nasa.gov/

National Oceanic and Atmospheric Administration. (2017). NOAA DATA CATALOG.

Retrieved February 2, 2017, from https://data.noaa.gov/dataset/

NCBI. (1988). National Center for Biotechnology Information. Retrieved January 18, 2019,

from https://www.ncbi.nlm.nih.gov/

Neupane, A., Nepal, M. P., Benson, B. V, Macarthur, K. J., & Piya, S. (2013). Evolutionary

history of mitogen-activated protein kinase (MAPK) genes in Lotus, Medicago, and

Phaseolus. *Plant Signaling & Behavior*, *8*(11), e27189.

https://doi.org/10.4161/psb.27189

Nishikawa, A. (2008). Degree and Pattern of Gene Flow in Several Scleractinian Corals in

the Ryukyu Archipelago, Southern Japan. *Https://Doi.Org/10.2984/1534-*

6188(2008)62[413:DAPOGF]2.0.CO;2, *62*(3), 413–422. https://doi.org/10.2984/1534-

6188(2008)62[413:DAPOGF]2.0.CO;2

OECD. (2017). *Marine protected areas : economics, management and effective policy mixes.*

Paris: OECD Publishing.

Palumbi, S. R. (2003). Population genetics, demographic connectivity, and the design of

marine reserves. *Ecological Applications*, *13*(1 SUPPL.), 146–158.

https://doi.org/10.1890/1051-0761(2003)013[0146:PGDCAT]2.0.CO;2

Palumbi, S. R., Barshis, D. J., Traylor-Knowles, N., & Bay, R. A. (2014). Mechanisms of

reef coral resistance to future climate change. *Science*, *344*(6186). Retrieved from

http://science.sciencemag.org/content/344/6186/895

Paris-Limouzy, C. B. (2011). Reef Interconnectivity/Larval Dispersal (pp. 881–889).

Springer, Dordrecht. https://doi.org/10.1007/978-90-481-2639-2_138

Penin, L., Vidal-Dupiol, J., & Adjeroud, M. (2013). Response of coral assemblages to

thermal stress: are bleaching intensity and spatial patterns consistent between events?

Environmental Monitoring and Assessment, *185*(6), 5031–5042.

https://doi.org/10.1007/s10661-012-2923-3

Pratchett, M. S., Thompson, C. A., Hoey, A. S., Cowman, P. F., & Wilson, S. K. (2018).

Effects of Coral Bleaching and Coral Loss on the Structure and Function of Reef Fish

Assemblages (pp. 265–293). Springer, Cham. https://doi.org/10.1007/978-3-319-75393-

5_11

Python Software Foundation. (2018). Python Language Refrence, version 3.5. Retrieved from

www.python.org

QGIS development team. (2009). QGIS Geographic Information System. Open Source

Geospatial Foundation Project. Retrieved from http://www.qgis.org/

R Core Team. (2016). R: A Language and Environment for Statistical Computing. Retrieved

from https://www.r-project.org/

Rellstab, C., Gugerli, F., Eckert, A. J., Hancock, A. M., & Holderegger, R. (2015). A

practical guide to environmental association analysis in landscape genomics. *Molecular*

Ecology, *24*(17), 4348–4370. https://doi.org/10.1111/mec.13322

Riginos, C., Crandall, E. D., Liggins, L., Bongaerts, P., & Treml, E. A. (2016). Navigating

the currents of seascape genomics: how spatial analyses can augment population

genomic studies. *Current Zoology*, *62*, doi: 10.1093/cz/zow067.

https://doi.org/10.1093/cz/zow067

Rochat, E., Leempoel, K., Vajana, E., Colli, L., Ajmone-Marsan, P., Joost, S. and the N. C.

(2016). Map of genotype frequency change in autochtonous Moroccan sheep breeds due

to global warming. In *Conference on Conservation genomics, May 3-6, 2016, Vairao,*

Portugal.

Rodgers, K. S., Lorance, K., Richards Donà, A., Stender, Y., Lager, C., & Jokiel, P. L.

(2017). Effectiveness of coral relocation as a mitigation strategy in Kāne'ohe Bay,

Hawai'i. *PeerJ*, *5*, e3346. https://doi.org/10.7717/peerj.3346

Rosic, N. N., Pernice, M., Dunn, S., Dove, S., & Hoegh-Guldberg, O. (2010). Differential

regulation by heat stress of novel cytochrome P450 genes from the dinoflagellate

symbionts of reef-building corals. *Applied and Environmental Microbiology*, *76*(9),

2823–2829. https://doi.org/10.1128/AEM.02984-09

Ryan, W. B. F., Carbotte, S. M., Coplan, J. O., O'Hara, S., Melkonian, A., Arko, R., …

Zemsky, R. (2009). Global Multi-Resolution Topography synthesis. *Geochemistry,*

Geophysics, Geosystems, *10*(3), n/a-n/a. https://doi.org/10.1029/2008GC002332

Sampayo, E. M., Franceschinis, L., Roff, G., Ridgway, T., Dove, S., & Hoegh-Guldberg, O.

(2016). Coral symbioses under prolonged environmental change: living near tolerance

range limits. *Scientific Reports*, *6*(1), 36271. https://doi.org/10.1038/srep36271

Sang, W., Ma, W.-H., Qiu, L., Zhu, Z.-H., & Lei, C.-L. (2012). The involvement of heat

shock protein and cytochrome P450 genes in response to UV-A exposure in the beetle

Tribolium castaneum. *Journal of Insect Physiology*, *58*(6), 830–836.

https://doi.org/10.1016/j.jinsphys.2012.03.007

Seabold, S., & Perktold, J. (2010). Statsmodels: econometric and statistical modeling with

Python. In *9th Python in Science Conference* (pp. 57–61). Retrieved from

http://statsmodels.sourceforge.net/

Selig, E. R., & Bruno, J. F. (2010). A global analysis of the effectiveness of marine protected

areas in preventing coral loss. *PloS One*, *5*(2), e9278.

https://doi.org/10.1371/journal.pone.0009278

Shanks, A. L., Grantham, B. A., & Carr, M. H. (2003). Propagule Dispersal Distance and the

Size and Spacing of Marine Reserves. *Ecological Applications*, *13*(1).

https://doi.org/https://doi.org/10.1890/1051-0761(2003)013[0159:PDDATS]2.0.CO;2

Sharma, S., Doherty, K. M., Brosh, R. M., & Jr. (2006). Mechanisms of RecQ helicases in

pathways of DNA metabolism and maintenance of genomic stability. *The Biochemical*

Journal, *398*(3), 319–337. https://doi.org/10.1042/BJ20060450

Shinzato, C., Mungpakdee, S., Arakaki, N., & Satoh, N. (2015). Genome-wide SNP analysis

explains coral diversity and recovery in the Ryukyu Archipelago. *Scientific Reports*,

5(1), 18211. https://doi.org/10.1038/srep18211

Shinzato, C., Shoguchi, E., Kawashima, T., Hamada, M., Hisata, K., Tanaka, M., … Satoh,

N. (2011). Using the Acropora digitifera genome to understand coral responses to

environmental change. *Nature*, *476*(7360), 320–323.

https://doi.org/10.1038/nature10249

Storey, J. D. (2003). The Positive False Discovery Rate: A Bayesian Interpretation and the q-

Value. *Annals of Statistics*, *31*(6), 2013–2035.

Storlazzi, C. D., van Ormondt, M., Chen, Y.-L., & Elias, E. P. L. (2017). Modeling Fine-

Scale Coral Larval Dispersal and Interisland Connectivity to Help Designate Mutually-

Supporting Coral Reef Marine Protected Areas: Insights from Maui Nui, Hawaii.

Frontiers in Marine Science, *4*, 381. https://doi.org/10.3389/fmars.2017.00381

Stubben, C., & Milligan, B. (2007). Estimating and Analyzing Demographic Models Using

the **popbio** Package in *R*. *Journal of Statistical Software*, *22*(11), 1–23.

https://doi.org/10.18637/jss.v022.i11

Stucki, S., Orozco-terWengel, P., Bruford, M. W., Colli, L., Masembe, C., Negrini, R., …

Consortium, the N. (2017). High performance computation of landscape genomic

models integrating local indices of spatial association. *Molecular Ecology Resources*,

17(5), 1072–1089. https://doi.org/10.1111/1755-0998.12629

Thomas, L., Kennington, W. J., Evans, R. D., Kendrick, G. A., & Stat, M. (2017). Restricted

gene flow and local adaptation highlight the vulnerability of high-latitude reefs to rapid

environmental change. *Global Change Biology*, *23*(6), 2197–2205.

https://doi.org/10.1111/gcb.13639

Thompson, D. M., & van Woesik, R. (2009). Corals escape bleaching in regions that recently

and historically experienced frequent thermal stress. *Proceedings. Biological Sciences /*

The Royal Society, *276*(1669), 2893–2901. https://doi.org/10.1098/rspb.2009.0591

UNEP-WCMC, WorldFish-Center, WRI, & TNC. (2010). Global distribution of warm-water

coral reefs, compiled from multiple sources including the Millennium Coral Reef

Mapping Project. Version 1.3. Retrieved May 9, 2017, from http://data.unep-

wcmc.org/datasets/1

Van der Auwera, G. A., Carneiro, M. O., Hartl, C., Poplin, R., del Angel, G., Levy-

Moonshine, A., … DePristo, M. A. (2013). From FastQ Data to High-Confidence

Variant Calls: The Genome Analysis Toolkit Best Practices Pipeline. In *Current*

Protocols in Bioinformatics (Vol. 43, p. 11.10.1-11.10.33). Hoboken, NJ, USA: John

Wiley & Sons, Inc. https://doi.org/10.1002/0471250953.bi1110s43

van Etten, J. (2018). gdistance: Distances and Routes on Geographical Grids. Retrieved from

https://cran.r-project.org/package=gdistance

van Oppen, M. J. H., Oliver, J. K., Putnam, H. M., & Gates, R. D. (2015). Building coral reef

resilience through assisted evolution. *Proceedings of the National Academy of Sciences*

of the United States of America, *112*(8), 2307–2313.

https://doi.org/10.1073/pnas.1422301112

Van Woesik, R., Irikawa, A., & Loya, Y. (2004). Coral Bleaching: Signs of Change in

Southern Japan. In *Coral Health and Disease* (pp. 119–141). Berlin, Heidelberg:

Springer Berlin Heidelberg. https://doi.org/10.1007/978-3-662-06414-6_5

Visel, A., Rubin, E. M., & Pennacchio, L. A. (2009). Genomic views of distant-acting

enhancers. *Nature*, *461*(7261), 199–205. https://doi.org/10.1038/nature08451

Welle, P. D., Small, M. J., Doney, S. C., & Azevedo, I. L. (2017). Estimating the effect of

multiple environmental stressors on coral bleaching and mortality. *PloS One*, *12*(5),

e0175018. https://doi.org/10.1371/journal.pone.0175018

Yampolsky, L. Y., Zeng, E., Lopez, J., Williams, P. J., Dick, K. B., Colbourne, J. K., &

Pfrender, M. E. (2014). Functional genomics of acclimation and adaptation in response

to thermal stress in Daphnia. *BMC Genomics*, *15*(1), 859. https://doi.org/10.1186/1471-

2164-15-859

Young, C. N., Schopmeyer, S. A., & Lirman, D. (2012). A review of reef restoration and

Coral propagation using the threatened genus Acropora in the Caribbean and western

Atlantic. *Bulletin of Marine Science*, *88*(4), 1075–1098.

https://doi.org/10.5343/bms.2011.1143

Ziegler, M., Seneca, F. O., Yum, L. K., Palumbi, S. R., & Voolstra, C. R. (2017). Bacterial

community dynamics are linked to patterns of coral heat tolerance. *Nature*

Communications, *8*, 14213. https://doi.org/10.1038/ncomms14213