De novo transcriptome assembly of the African bullfrog *Pyxicephalus*

adspersus for molecular analysis of aestivation

Naoki Yoshida[*], Chikara Kaito

Graduate School of Medicine, Dentistry, and Pharmaceutical Sciences, Okayama

University

*Corresponding author

E-mail: kamerou2005@yahoo.co.jp

Abstract

The molecular mechanisms of aestivation, a state of dormancy that occurs under dry conditions at ordinary temperature, are not yet clarified. Here, we report the first *de novo* transcriptome assembly of the African bullfrog *Pyxicephalus adspersus*, which aestivates for 6-10 months during the hot and dry season. Polyadenylated RNA from tissues was sequenced to 75,320,390 paired-end reads, and the *de novo* assembly generated 101,682 transcripts. Of these, 100,093 transcripts had open reading frames encoding more than 25 amino acids. BLASTx analysis against the Uniprot *Xenopus tropicalis* protein database revealed 64,963 transcripts having little homology with an E value higher than 1E-5 and 8,147 transcripts having no homology, indicating that the African bullfrog has many novel genes that are absent in *X. tropicalis*. The other 28,570 transcripts had homology with an E value lower than 1E-5 for which molecular functions were estimated by gene ontology (GO) analysis and found to contain the aestivation-related genes conserved among other aestivating organisms, including the African lungfish. This study is the first to identify a comprehensive set of genes expressed in the African bullfrog, thus providing basic information for molecular level analysis of aestivation.

Introduction

Dormancy is a phenomenon by which organisms temporarily decrease their physiologic activity under conditions such as low temperature, drying, and depletion of energy sources, and is observed in many organisms from microorganisms to mammals. In vertebrates, hibernation and aestivation are two common types of dormancies performed under either a low temperature condition or a drying and ordinary temperature condition, respectively. Molecular biologic and physiologic analyses of vertebrate hibernation have been performed for the Siberian chipmunk and Northern leopard frog. In Siberian chipmunks, hibernation is initiated by two cues: a seasonal change in the photoperiod depending on the melatonin level, and a hibernation-specific protein complex induced by cold temperature [1,2]. Energy-related genes are regulated to decrease body temperature and to suppress the contractile activity of cardiac myocytes. In the Northern leopard frog, glucose and glycerol concentrations in the body are increased to confer freeze tolerance, and the mammalian target of rapamycin (mTOR) is activated to prevent muscle atrophy during hibernation [3,4,5]. These molecular and physiologic aspects of hibernation have attracted attention as the potential basis for novel therapeutics against malignant tumors or tissue necrosis.

Aestivation in vertebrates is observed in many amphibian and fish species. Aestivation is similar to hibernation in terms of the reduced metabolic activity, but it is clearly distinguished from hibernation in terms of the environmental conditions; aestivation is performed under dry conditions at ordinary temperatures, whereas hibernation is performed under aqueous conditions at low temperatures [6]. Therefore, the mechanisms of aestivation are assumed to differ from those of hibernation, such as

suppression of energy metabolism at an ordinary temperature or tolerance against

dehydration. Molecular level analyses of aestivation are important to uncover

mechanisms of dormancy that have not been clarified in studies of hibernation, but such

studies are scarce at present.

So far, three species that undergo aestivation – the Green striped burrowing frog

(*Cyclorana alboguttata),* Couch's spadefoot toad (*Scaphiopus couchii*), and African

lungfish (*Protopterus annectens*) – have been analyzed at the transcriptional level to

identify genes whose expression is induced or suppressed during aestivation [7,8,9].

Very little is known about the molecular mechanisms of aestivation, however, including

how energy metabolism is suppressed or how tissue damage is minimized in the

aestivating organisms.

In the present study, we aimed to identify genes expressed in the African bullfrog,

P. adspersus using *de novo* transcriptome assembly. The African bullfrog, a species

belonging to the family of *Pyxicephalidae*, is a large frog – body size of males is larger

than 20 cm and that of females is ~14 cm. This frog inhabits a savanna area of east

Africa, south Africa, and the southern part of central Africa, where the air temperature

ranges from 20-30˚C throughout the year, and there are pronounced dry and rainy

seasons. During the dry season, the frogs burrow underground and form a tough cocoon

to reduce evaporative water loss and decrease the respiration rate to less than 10% that

during the active stage [10,11]. The aestivation of the frog continues for 6-10 months.

Because aestivation of the frog can be artificially induced and breeding methods are

established for breeding the frogs as pets, the African bullfrog is a suitable research

model animal to investigate the molecular mechanisms of aestivation. Furthermore, the

African bullfrog is phylogenetically and geographically distant from the other two

aestivating frogs: the Green striped burrowing frog and Couch's spadefoot toad, which

inhabit Australia and North America, respectively. Therefore, studies of the aestivation

of the African bullfrog will help to elucidate the molecular mechanisms of aestivation

that are conserved among different aestivating frog species. In this study, we identified

and characterized the genes of the African bullfrog to provide basic information for

investigating the aestivation mechanism.

Materials and Methods

Ethics statement

The Animal Care and Use Committee of Okayama University approved this work

(Approval number, OKU-2019300). This study was performed in strict accordance with

the recommendation of the Fundamental Guidelines for Proper Conduct of Animal

Experiment and Related Activities in Academic Research Institutions under the

jurisdiction of the Ministry of Education, Culture, Sports, Science, and Technology in

Japan, 2006.

Experimental animal

A captive bred African bullfrog was purchased from a specialty reptile and

amphibian store (Hachurui Club, Nakano, Japan). The purchased frog was maintained

in a plastic container with coarse sand (5-7 mm diameter) and water. The frog was fed

every day with house crickets (*Acheta domestica*) or wax worms (the larvae of the

Galleria mellonella) for the first 3 weeks, and then fed every day with artificial diets

(Samuraijapan, Ibaraki, Japan) for 2 months.

Isolation of genomic DNA and total RNA

The young adult frog (15 g body mass) was kept without feeding for 36 h. The frog

was anesthetized by placing it into crushed ice for 10 min and then dissected on ice. The

intestines were separated, the intestinal contents removed in phosphate buffered saline

(pH7.4), and the intestines frozen in liquid nitrogen. The other tissues, including inner

organs, muscle, and skin, were quickly excised to 3 - 5 mm^3 (0.1 - 0.3 g) and frozen in

liquid nitrogen. The frozen samples were maintained at -80˚C.

Genomic DNA was isolated by a standard phenol/chloroform extraction method

[12]. Frozen muscle tissues (5 mm^3, 0.2 g) were submerged in a mixture of 5 ml of

phenol/chloroform/isoamyl alcohol (25:24:1) and 5 ml of TE Buffer (10 mM Tris-HCl

[pH 8.0], 1mM EDTA). Then, the tissues were homogenized at 10,000 rpm three times

for 30 s each with a Polytron benchtop homogenizer (Kinematica, Switzerland). The

sample was centrifuged at room temperature at 2300 g for 10 min, and the upper

aqueous phase (1.0 ml) was transferred to a new tube. The sample was centrifuged at

room temperature at 20,400 g for 5 min to completely remove the organic solvents. The

upper aqueous phase (0.7 ml) was transferred to a new tube, vortexed with an

equivalent volume of chloroform, and centrifuged at room temperature for 5 min at

20,400 g. The upper aqueous phase (0.5 ml) was transferred to a new tube, mixed with a

2.5-fold amount of ethanol and a 0.1-fold amount of 3M sodium acetate, and

centrifuged at 20,400 g for 15 min at room temperature. The precipitate was washed

with 70% ethanol and resolved in 50 µl of TE buffer.

Total RNAs were extracted from the inner organs, intestines, muscles, and skin

using the chaotropic extraction protocol for mouse pancreatic RNA, described by

Robert [13]. Frozen tissues (3 - 5 mm^3, 0.1 - 0.3 g) were submerged in 10 ml of

TRIZOL Reagent (Life Technologies, Carlsbad, CA, USA), an amount three times

larger than that recommended by the supplier. The tissues were then homogenized at

14,000 rpm three times for 30 s each using the Polytron homogenizer. After incubation

for 5 min at room temperature, 2 ml of chloroform was added and the mixture was

vortexed for 15 s. The mixture was incubated for 3 min at room temperature and

centrifuged at 12,000 g for 10 min at 4˚C. The upper aqueous phase (4 ml) was

transferred to a fresh 50-ml tube and mixed with an equivalent amount of isopropyl

alcohol. The sample was centrifuged at 12,000 g for 10 min and an RNA pellet was

obtained. The RNA pellet was vortexed with 10 ml of 75% ethanol and centrifuged at

7500 g for 5 min at 4˚C. The RNA pellet was air-dried and dissolved in 200 µl of

RNase-free water by incubating for 10 min at 55˚C. Further purification to remove

contaminated genomic DNA was performed using a Monarch Total RNA Miniprep Kit

(New England Biolabs, MA, USA), according to the supplier's protocol. The RNA was

eluted from a column by 100 µl of RNase-free water and kept at -80˚C. The

concentration and purity of the isolated RNA was determined using a spectrophotometer

and denaturing agarose gel electrophoresis. The concentrations of RNA isolated from 11

tissue segments of inner organs, intestines, muscle, skin, and head ranged from

0.52-3.34 µg/µl or 0.38-1.60 µg/mg tissue.

Determination of mitochondrial 16S/12S rRNA sequence

Primers (16SA-L-frog: 5'-CGCCTGTTTACCAAAAACAT-3', 16SB-H-frog:

5'-CCGGTCTGAACTCAGATCACGT-3',   12SB-F:

5'-AAACTGGGATTAGATACCCCACTAT-3',   12SB-H:

5'-GAGGGTGACGGGCGGTGTGT-3') used to amplify mitochondrial 16S or 12S

rRNA partial regions were identical to those used in phylogenetic classification of

Anura, reported by Hirota [14]. Polymerase chain reaction (PCR) was performed in 25

µl of 1x Q5 reaction buffer (0.5 units of Q5 High-Fidelity DNA Polymerase (New

England Biolabs), 200 µM dNTPs, 0.5 µM forward primer, 0.5 µM reverse primer and

6.5 ng genomic DNA) using the thermal cycle program (30 s at 98˚C, 30 cycles [30 s,

98˚C; 30 s, 62˚C; 30 s, 72˚C], and 2 min at 72˚C). PCR products were examined by

agarose gel electrophoresis and sequenced using the same primers.

mRNA library preparation and Illumina next-generation

sequencing

An equivalent amount of total RNAs (0.52 - 3.34 µg/µl) isolated from 11 tissue

segments of inner organs (3 segments), intestines (2 segments), muscles (2 segments),

skin (3 segments), and head (1 segment) were mixed to obtain 100 ng/µl total RNA.

The mixed total RNA was analyzed by Tape Station (RNA Screen tape, Agilent

Technologies Ltd., USA) and determined to RIN (RNA integrity number) = 9.0. The

poly (A)+ fraction was isolated from the total RNA, followed by its fragmentation. The

fragmented mRNA was then reverse-transcribed to double-stranded cDNA using

random hexamers. The double-stranded cDNA fragments were processed for adaptor

ligation, size selection for 200-bp inserts, and amplification of strand-specific cDNA

libraries. Prepared libraries were subjected to paired-end 2 x 100 bp sequencing on the

HiSeq 2500 platform with v4 chemistry.

De novo transcriptome assembly and bioinformatic analysis

All analyses were performed mainly using the CLC Genomics Workbench 11.0

(QIAGEN Bioinformatics, Germany). The 2 x 100 bp paired-end reads were checked in

terms of the sequencing quality and trimmed (removal of adaptor and duplication) with

quality score limit of 0.05 and a maximum number of two ambiguous nucleotides. The

clean reads were then assembled and aligned with a word size of 20, bubble size of 50,

and minimum transcript length of 200, based on the algorithm of the De Brujin graph.

The open reading frame (ORF) in the assembled *P. adspersus* transcripts was predicted

under the following conditions: start codon as ATG, search as Both Strand, open-ended

sequence, minimum length (codons) as 25, and genetic code as standard. The transcripts

were homology searched using local BLASTx (National Center for Biotechnology

Information, NCBI) against the Uniprot database (https://www.uniprot.org/)

(all-proteins or *Xenopus tropicalis*) or SmProt database

(http://bioinfo.ibp.ac.cn/SmProt/index.htm) (small proteins curated from literature

mining, known databases, ribosome profiling data, or MS data). Homologous proteins

found in the Uniprot database (*X. tropicalis*) with an E-value lower than 1E-5 were

subjected to gene ontology (GO) analysis (https://www.uniprot.org) [15] to assign the

GO terms of biologic processes, molecular functions, and cellular components.

Data availability

The data sets supporting the results of this article are available from the following

databases.

Mitochondrial 16S rRNA of the African bullfrog (583 bp): GenBank: LC440402

Mitochondrial 12S rRNA of the African bullfrog (424 bp): GenBank: LC440403

Cytochrome *b* of the African bullfrog (transcripts #275): GenBank: LC440404

Raw reads of 2 x 100 bp RNA sequence of the African bullfrog: DDBJ sequencing read

archive (SRA): DRR164258, BioProject: PRJDB7457, BioSample: SAMD00143251,

Experiment: DRX154877

Transcript sequences of the African bullfrog: GenBank:

ICLD01000001-ICLD01101682

203 Results

204 Confirmation of animal species used in this study

To confirm that the experimental animal used in this study was in fact the African

bullfrog, a captive bred frog was raised for 5 years and the appearance of the adult frog

was observed. The frog was approximately 20 cm in length and had olive-green skin

with longitudinal skin folds (**Fig 1A**). These morphologic characteristics were

consistent with those of the African bullfrog reported by Channing [16] and Bishop [17],

and clearly differed from the characteristics of *P. edulis*, a species closely related to the

African bullfrog that is 12 cm in length and has olive green-yellow skin with spotted

folds. Furthermore, we determined the nucleotide sequence of mitochondrial 16S and

12S rRNA of the frog used for subsequent *de novo* assembly analysis. NCBI BLAST

analysis revealed that the mitochondrial 16S rRNA (583 bp, GenBank: LC440402) was

100% identical to the mitochondrial 16S rRNA of *Pyxicephalus cf. adspersus* FB-2006

isolate 0954 (551 bp, GenBank: DQ347304) (**Fig 1B**). The mitochondrial 12S rRNA

(424 bp, GenBank: LC440403) was 99% identical to the mitochondrial 12S rRNA of *P.*

cf. adspersus FB-2006 isolate 0954 (1,364 bp, GenBank: DQ347013) (**S1 Fig**). These

morphologic characteristics and the results of the homology analysis of rRNA

sequences confirmed that the frog used in this study was *P. adspersus,* the African

bullfrog.

The two types of frog, *P. adspersus* and *P. edulis*, are sometimes confused with

each other [16,18]. Top hits against the 16S and 12S rRNA sequences of the frog used in

this study are found in both species in no particular order. Furthermore, the phylogenetic

tree analysis of the top hits 16S and 12S rRNA sequences revealed that the two species

were not clustered (**Fig 1B and S1 Fig**). We speculate that the nomenclatures of these

rRNA sequences were mistakenly registered in the database.

Figure 1C shows the phylogenetic tree of mitochondrial 16S rRNA partial

sequences of the African bullfrog used in this study and of representative frog species

belonging to the major family of Anura. The two aestivating frogs, *C. alboguttata* and *S.*

cauchii, are phylogenetically distant from *P. adspersus*, as suggested by Pyron's report

[19].

RNA sequencing and *de novo* assembly

RNA sequencing and *de novo* assembly data are summarized in Table 1. RNA

sequencing produced 75,604,146 raw reads. *De novo* transcriptome assembly of the

trimmed 75,320,390 reads generated 101,682 transcripts. The transcript size ranged

from 236 to 35,997 bases with a mean length of 866 bases and an N50 length of 1,701

bases. The number of transcripts ranging from 1000 to 1999 bases was highest among

that of transcripts ranging from 500 to 5,000 bases (**Fig 2**).

In silico analysis to identify ORF-encoding proteins larger than 25 amino acids

revealed that 100,093 transcripts (98.4% of total transcripts) were protein-coding genes

(**Table 2**). The longest ORF-encoded protein was 11,679 amino acids long. The average

ORF-encoded protein was 140.7 amino acids long. In contrast, 1590 transcripts (1.6%

of total transcripts) had no ORF, suggesting that these transcripts were non-coding

RNAs. When the ORF detection conditions were changed to 30, 50 or 100 amino acids

long, the number of transcripts carrying the ORF was decreased to 97,122 (95.5%),

71,305 (70.1%) or 29,296 (28.8%), respectively (**Table 2**).

Within these *de novo* assembled transcripts, we identified that transcript #275 had

high similarity with cytochrome *b* on the basis of the BLASTx search. The DNA

sequence of transcript #275 (GenBank: LC440404) was 100% identical to *P. adspersus*

isolate Brak01 cytochrome *b* gene (GenBank: FJ613662) (**S2 Fig**). This result

confirmed that that the frog used in this study was *P. adspersus*.

Table 1. Summary of the RNA sequence reads and the *de novo* transcriptome

assembly for the African bullfrog

	Length of raw reads (bp)	100
Sequencing results	Total number of raw reads	75,604,146
	Total number of clean reads	75,320,390
	Total number of transcripts	101,682
	Total length of all transcripts (bases)	88,093,202
De novo transcriptome assembly results	GC contents of all transcripts (%)	40.8
	N50 length of all transcripts (bases)	1,701
	Mean length	866

260 **Table 2. Detection of ORF in the African bullfrog transcripts**

	Minimum ORF length			
	25 aa	**30 aa**	**50 aa**	**100 aa**
Total transcripts				101,683
Detected ORF	729,832	562,570	219,579	45,086
Transcripts coding ORF	100,093 (98.4%)	97,122 (95.5%)	71,305 (70.1%)	29,296 (28.8%)
Maximum ORF length	11,679	11,679	11,679	11,679
Minimum ORF length	25	30	50	100
Average ORF length (aa)	140.7	144.1	181.4	341.2
Transcripts of non-coding RNA	1,590 (1.6%)	4,561 (4.6%)	30,378 (29.9%)	72,387 (71.2%)

Transcriptome annotation

We performed a BLASTx homology search of the African bullfrog transcripts

against the Uniprot protein database (all-proteins or *Xenopus tropicalis* proteins). In the

all-proteins database, 27,286 transcripts (26.8%) had an E-value lower than 1E-5,

53,064 transcripts (52.2%) had an E-value higher than 1E-5, and 21,332 transcripts

(21.0%) showed no hits with the database (**Fig 3A**, All proteins). In the *X. tropicalis*

protein database, 28,572 transcripts (28.1%) had an E-value lower than 1E-5, 64,963

transcripts (63.9%) had an E-value higher than 1E-5, and 8,147 transcripts (8.0%)

showed no hits with the database (**Fig 3A**, *X. tropicalis*). Figure 3B shows the

distribution of transcripts with an E-value lower than 1E-5 according to the amino acid

identity with the protein database. 22,670 transcripts (83.1%) and 25,407 transcripts

(88.9%) had an amino acid identity higher than 50% in the all-proteins database or in

the *X. tropicalis* protein database, respectively.

To identify small proteins that are absent in the Uniprot database, we performed a

BLASTx homology search of the African bullfrog transcripts against the SmProt small

protein databases (small proteins curated from literature mining, known databases,

ribosome profiling data, or MS data). In any of the SmProt small protein databases,

more than 80% transcripts had an E-value higher than 1E-5 or showed no hit (**Fig 4A**).

We further examined whether the African bullfrog transcripts having little similarity to

the Uniprot all protein database (53,064 transcripts, >1E-5; 21332 transcripts, no hits)

and the Uniprot X. tropicalis protein database (64,963 transcripts, >1E-5; 8,147

transcripts, no hits) shows little similarity against the SmProt databases. As a result,

more than 99% non-annotated transcripts in the Uniprot all protein database as well as

those in the Uniprot X. tropicalis protein database had E-value higher than 1E-5 or no

hits in the SmProt databases (**Fig 4B**).

288 Gene ontology analysis

The 28,572 transcripts with an E-value lower than 1E-5 in Uniprot *X. tropicalis*

protein database were subjected to gene ontology (GO) analysis and classified

according to three major GO categories: cellular components, molecular function, and

biologic processes (**Fig 5**). In the cellular component category, 11,634 (33.4%) and

7,689 (22.0%) genes were associated with the cellular components and membrane

components, respectively. In the molecular function category, 11,142 (45.6%) and 6,995

(28.6%) genes were associated with binding and catalytic activity, respectively. In the

biologic process category, 9,765 (30.5%), 6,213 (19.4%), and 5,553 (17.3%) genes were

associated with cellular processes, biologic regulation, and metabolic processes,

respectively.

Comparison with the aestivation-related genes reported in other organisms

To determine whether the African bullfrog transcripts contain genes homologous to

the aestivation-related genes of the two aestivating frogs and the African lungfish, we

performed a homology search of aestivation-related genes against the African bullfrog

transcripts (**Table 3**) [8,9,10]. The #2083 transcript in the African bullfrog had high

identity with riboflavin-binding protein precursor (RfBP) in Couch's spadefoot toad,

which has increased expression during aestivation and might be involved in egg

maturation in the frog during the aestivating stage. The African bullfrog genes had high

identity with genes of the Green striped burrowing frog, which are highly upregulated

during aestivation, such as adaptor related protein complex 3 (beta 2 subunit) that

functions in vesicle budding from the Golgi apparatus; an immunoglobulin heavy chain

that recognizes foreign antigens; and a serine/threonine protein kinase, Chk1, that

activates DNA repair in response to DNA damage. In addition, the African bullfrog

genes had high identity with African lungfish genes that are expressed during

aestivation, such as argininosuccinate synthetase 1 and carbamoylphosphate synthetase

III, which are key enzymes in the ornithine-urea cycle and function to detoxify

ammonia to urea; rhesus family C glycoprotein that maintains muscle function; and

superoxide dismutase 1, an anti-oxidant. These findings suggest that African bullfrog

has a set of genes that are highly expressed in other vertebrates during aestivation.

Table 3. Presence of aestivation-related genes in the African Bullfrog

Species	Gene	Predicted Function	Top Hit Contig of African Bullfrog	E-value	% Identity
Scaphiopus couchii (Couch's spadefoot toad)	Riboflavin binding protein precursor (AF102545)	Maturation of eggs in preparation for the explosive breeding	#2083 (ICLD012083)	8.63E-114	67
Cyclorana alboguttata (Green-striped burrowing frog)	Adaptor related protein complex 3, beta 2 subunit (contig #8496)	Budding of vesicles from the golgi membrane	#4968 (ICLD014968)	0	100
	Solute carrier family 28 member 3-like (contig #7653)	Homeostasis of endogenous nucleosides	#2599 (ICLD012599)	1.99E-45	100
	Nuclease HARBI1-like (contig #14756)	Nuclease activity	#59155 (ICLD0159155)	1.31E-122	88
	Ribosome biogenesis protein NSA2-like (contig #40)	Biogenesis of the 60S ribosomal subunit	#4109 (ICLD014109)	5.97E-103	99
	Coiled coil domain containing protein 81 (contig #15484)	Positive regulator of T-cell maturation and inflammatory function	#18527 (ICLD0118527)	0	75
	Pentraxin precursor (contig #27476)	Mediating uptake of synaptic material	#3473 (ICLD013473)	7.25E-171	90
	Immunoglobulin heavy prechain (contig #36672)	Recognization of foreign antigens and initiation of immune responses	#3055 (ICLD013055)	1.01E-118	82
	Receptor type tyrosine protein phosphatase T-like (contig #7294)	Signal transduction and cellular adhesion	#10027 (ICLD0110027)	0	99
	Immunoglobulin J chain precursor (contig #48432)	Serving to link two monomer units of either IgM or IgA	#1190 (ICLD011190)	1.82E-15	76
	UHRF1 binding protein 1-like (contig #1661)	Negative regulator of cell growth	#8912 (ICLD018912)	0	95
	Inositol 3 phosphate synthase 1 (contig #6523)	Key enzyme in myo-inositol biosynthesis pathway that catalyzes the conversion of glucose 6-phosphate to 1-myo-inositol 1-phosphate	#13219 (ICLD0113219)	0	93

	Serine/threonine protein kinase Chk1 (contig #13899)	Activation of DNA repair in response to the presence of DNA damage or unreplicated DNA	#24942 (ICLD0124942)	0	85
Protopterus annectens (African Lungfish)	Rhesus family C glycoprotein (KX583647)	Glycolysis and muscle development	#10697 (ICLD0110697)	4.05E-162	79
	Argininosuccinate synthetase 1 (JZ575533)	Ornithine-urea cycle capacity to detoxify ammonia to urea	#297 (ICLD01297)	1.72E-56	81
	Carbamoylphosphate synthetase III (JZ575539)	Ornithine-urea cycle capacity to detoxify ammonia to urea	#265 (ICLD01265)	3.70E-108	91
	Betain homocyctein-S-transferase 1 (JZ575536)	Prevent the accumulation of hepatic homocyctein	#243 (ICLD01243)	5.19E-86	94
	Superoxide dismutase 1 (JZ575606)	Anti ROX	#1087 (ICLD011087)	2.73E-62	100

321   tBLASTx search was performed to identify the African bullfrog contig having high identity to the genes that are up-regulated during the aestivation

322   in the three organisms. The ID numbers in parenthesis indicate Genbank accession numbers or contig numbers (*C. alboguttata*).

Discussion

The African bullfrog is well known as an aestivating frog and is bred all over the world. Molecular biologic analysis of this frog, however, has not been performed and no genetic information has been available. In this study, we performed the first *de novo* transcriptome assembly and identified 101,682 transcripts. Of the 101,682 transcripts identified, 53,064 (52.2%) had an E-value higher than 1E-5 and 21,332 transcripts (21.0%) had no hits in a BLASTx search in the Uniprot all protein database. In addition, 64,963 (63.9%) had an E-value higher than 1E-5 and 8,147 transcripts (8.0%) had no hits in the Uniprot *X. tropicalis* protein database. These results suggest that more than 70% of total transcripts are novel genes in the African bullfrog. Of these novel genes, more than 99% had little similarity to the SmProt small protein database. The result suggests that the high proportion of novel genes of the African bullfrog are not explained by the genes coding small proteins, which are available in the database. The high proportion of the novel genes in total transcripts is also observed in other organisms whose *de novo* transcriptome assembly was recently performed, such as the Caecilian (*Caecilia tentaculate*), the Sodom apple (*Calotropis gigantea*), and the Black ant (*Formica fusca*) [20, 21, 22], and confirms that *de novo* transcriptome assembly is a powerful tool to identify novel genes. The novel genes of the African bullfrog might encode proteins without known domains, or non-coding RNAs. With regard to the very limited genomic and transcriptomic information in amphibians, the novel genes of the African bullfrog are the significant biologic resources to perform molecular biologic analysis of the physiologic functions of the African bullfrog, including its aestivation ability.

In contrast, 28,572 transcripts (29.1%) had an E-value lower than 1E-5 in the *X.*

tropicalis database, suggesting that these genes are conserved in *X. tropicalis*. These

genes were subjected to gene ontology analysis and their molecular functions were

estimated. They are assumed to have functions conserved between *P. adspersus* and *X.*

tropicalis. Furthermore, we revealed that the African bullfrog has a set of

aestivation-related genes, which has been reported in two aestivating frogs and the

African lungfish. These genes presumably function in the aestivation mechanism

conserved between different aestivating vertebrates. The aestivation-related genes are

not specific to the aestivating organisms, but are conserved in other non-aestivating

organisms, and must therefore be transcriptionally controlled by factor(s) specific to the

aestivating environment or the aestivating organisms. How the conserved genes are

regulated during aestivation remains an important unanswered question.

The genetic information of the African bullfrog obtained in this study enables

future comparative transcriptome analysis between the active stage and the aestivating

stage of the frog and contributes to uncover the molecular mechanisms of aestivation.

Supporting Information

S1 Fig. The phylogenetic tree analysis of 12S rRNA sequence between the frog used in this study and several *P. adspersus* and *P. edulis* species deposited in database. Sequences were aligned using MUSCLE algorithm implemented in MEGA X. Phylogenetic tree was constructed using neighbor-joining algorithm with bootstrap test of 1000 replicates.

S2 Fig. The phylogenetic tree analysis of cytochrome b partial sequence between the frog used in this study and several frogs deposited in database. Sequence of cytochrome *b* of the African bullfrog (GenBank: LC440404) was derived from *de novo* assembled transcript #275. Sequences were aligned using MUSCLE algorithm implemented in MEGA X. Phylogenetic tree was constructed using neighbor-joining algorithm with bootstrap test of 1000 replicates.

REFERENCES

1. Michael H, Hastings Francis JPE. Hibernation Proteins: Preparing for Life in the Freezer. Cell. 2006 125(1): 21-23. doi: 10.1016/j.cell.2006.03.019. PMID: 16615885

2. Kondo N, Sekijima T, Kondo J, Takamatsu N, Tohya K, Ohtsu T. Circannual Control of Hibernation by HP Complex in the Brain. Cell. 2006 125(1):161-72. doi: 10.1016/j.cell.2006.03.017. PMID: 16615897

3. Costanzo JP, Amaral MC, Rosendale AJ, Lee RE. Hibernation physiology, freezing adaptation and extreme freeze tolerance in a northern population of the wood frog. J Exp Biol. 2013; 216(18): 3461-73. doi: 10.1242/jeb.089342. PMID: 23966588

4. Christenson MK, Trease AJ, Potluri LP, Jezewski AJ, Davis VM, Knight LA, et al. De novo Assembly and Analysis of the Northern Leopard Frog *Rana pipiens* Transcriptome. J Genomics. 2014; 2:141-9. doi: 10.7150/jgen.9760. PMID: 25371763

5. Hadj-Moussa H, Storey KB. Micromanaging Freeze Tolerance: The Biogenesis and Regulation of microRNAs in Frozen Frogs. Cell Mol Life Sci. 2018; 75(19): 3635-3647. doi: 10.1007/s00018-018-2821-0, PMID: 29681008

6. Storey KB. Turning Down The Fires Of Life: Metabolic Regulation Of Hibernation And Estivation. Comp Biochem Physiol B Biochem Mol Biol. 2001; 126. doi: 10.1016/S0305-0491(00)80178-9

7. Reilly BD, Schlipalius DI, Cramp RL, Ebert PR, Franklin CE. Frogs and estivation: transcriptional insights into metabolism and cell survival in a natural model of extended muscle disuse. Physiol Genomics. 2013; 45(10): 377-88. doi: 10.1152/physiolgenomics.00163.2012. PMID: 23548685

8. Storey KB, Dent ME, Storey JM. Gene expression during estivation in spadefoot toads, *Scaphiopus couchii*: upregulation of riboflavin binding protein in liver. J Exp Zool. 1999; 284(3): 325-33. PMID: 10404124

9.   Hiong KC, Ip YK, Wong WP, Chew SF. Differential gene expression in the liver of the

African lungfish, *Protopterus annectens*, after 6 months of aestivation in air or 1 day of

arousal from 6 months of aestivation. PLoS One. 2015; 10(3): e0121224. doi:

10.1371/journal.pone.0121224. PMID: 25822522

10. Loveridge JP, Withers PC. Metabolism and Water Balance of Active and Cocooned African

Bullfrogs *Pyxicephalus adspersus*. Physiol Biochem Zoo. 1981; 54(2). doi:

210.1086/physzool.54.2.30155821

11. Secor SM. Physiological responses to feeding, fasting and estivation for anurans. J Exp

Biol. 2005; 208(13): 2595-608. doi: 10.1242/jeb.01659. PMID: 15961745

12. Strauss WM. Preparation of Genomic DNA from Mammalian Tissue. Curr Protoc Immunol.

2001; Chapter 10 (10.2). doi: 10.1002/0471142735.im1002s08. PMID: 18432684

13. De Lisle RC. Isolation of Pancreatic RNA. Pancreapedia: Exocrine Pancreas Knowledge

Base. doi: 10.3998/panc.2014.9

14. Sumida M, Kondo Y, Kanamori Y, Nishioka M. Inter-and intraspecific evolutionary

relationships of the rice frog *Rana limnocharis* and the allied species *R. cancrivora* inferred

from crossing experiments and mitochondorial DNA sequences of the 12S and 16S rRNA

genes. Mol Phylogenet Evol. 2002; 25(2): 293-305. doi: 10.1016/S1055-7903(02)00243-9.

PMID: 12414311

15. Birol I, Behsaz B, Hammond SA, Kucuk E, Veldhoen N, Helbing CC. *De novo*

Transcriptome Assemblies of *Rana (Lithobates) catesbeiana* and *Xenopus laevis* Tadpole

Livers for Comparative Genomics without Reference Genomes. PLoS One. 2015; 10(6):

e0130720. doi: 10.1371/journal.pone.0130720. PMID: 26121473

16. Channing A, DU Preez L, Passmore N. Status, vocalization and breeding biology of two

species of African bullfrogs (Ranidae: *Pyxicephalus*). J Zoo. 1994; 234(1): 141-148. doi:

10.1111/j.1469-7998

17. Bishop P. Announcing AmphibiaWeb (University of California Berkeley, CA, USA). 2004

Oct 19 [Edited by Janzen P, Tunstall T. 2005 Sep 21]. Available from:

https://amphibiaweb.org/species/4966

18. Engelbrecht D, Mashao M, Halajian A. Notes on the breeding behavior and ecology of

Edible Bullfrogs *Pyxicephalus edulis* Peters, 1854 in the Limpopo Province, South Africa.

Herpetology Notes. 2015; 8: 365-369.

19. Pyron RA, Wiens JJ. A large -scale phylogeny of Amphibia including over 2,800 species,

and a revised classification of extant frogs, salamanders, and caecilians. Mol Phylogenet

Evol. 2011; 61(2): 543-583. doi: 10.1016/j.ympev.2011.06.012. PMID: 21723399

20. Torres M, Creevey CJ, Kornobis E, Gower DJ, Wilkinson M, San D. Multi-tissue

transcriptomes of caecilian amphibians highlight incomplete knowledge of vertebrate gene

families. DNA Res. 2018; 0(0): 1-8. doi: 10.1093/dnares/dsy034. PMID: 30351380

21. Muriira N, Xu W, Muchugi A, Xu J, Liu A. *De novo* sequencing and assembly analysis of

transcriptome in the Sodom apple (*Calotropis gigantea*). BMC Genomics. 2015 (16):723.

doi: 10.1186/s12864-015-1908-3. PMID: 26395839

22. Morandin C, Pulliainen U, Bos N, Schultner E. *De novo* transcriptome assembly and its

annotation for the black ant *Formica fusca* at the larval stage. Scientific Data. 2018.

180282. doi: 10.1038/sdata.2018.282. PMID 30561435

Figure legend

Fig 1. The African bullfrog used in this study.

A) The adult male African bullfrog used in this study is shown.

B) Phylogenetic tree analysis of the 16S rRNA sequence between the frog used in this study and several *P. adspersus* and *P. edulis* species deposited in the database. Sequences were aligned using MUSCLE algorithm in MEGA X. Phylogenetic tree was constructed using neighbor-joining algorithm and the node support values were assessed by bootstrap test of 1000 replicates.

C) The phylogenetic tree using the 16S rRNA sequence between the representative frog species belonging to the major family of Anura. *P. adspersus* used in this study is shown in red and that of the other two aestivating frogs are shown in blue. Phylogenetic tree was constructed in the same manner in Fig 1B.

Fig 2. Distribution of the length of the African bullfrog transcripts.

The number of transcripts was count and the transcript length was determined.

Fig 3. Similarity of the African bullfrog transcripts against the Uniprot protein databases.

A) BLASTx search of the African bullfrog transcripts against the Uniprot database (all-proteins or *X. tropicalis* proteins), and the number of transcripts were counted according to E-values.

B) Distribution of the identity of the African bullfrog transcripts with an E-value lower than 1E-05 against the *X. tropicalis* protein database.

Fig 4. Similarity of the African bullfrog transcripts against the SmProt small protein

databases.

A) BLASTx search of the African bullfrog transcripts against the SmProt databases

including literature mining, ribosome profiling data, known databases, and MS data.

The number of transcripts were counted according to E-values.

B) The African bullfrog 74,396 transcripts having little similarity to Uniprot all protein

database (53,064 transcripts, >1E-5; 21,332 transcripts, no hit) were subjected to

BLASTx search against SmProt database (Literature mining, Known database,

Ribosome profiling, and MS data). Red dotted line represents the total number of

transcripts subjected to this analysis.

Fig 5. Gene ontology terms assigned to the African bullfrog genes.

The 28,570 genes with high identity (E-value lower than 1E-5) in a BLASTx search of

the *X. tropicalis* protein database were subjected to gene ontology analysis.

Figure 1

B

P. edulis (DQ022366), 99%
51
P. adspersus (DQ347304), 100%
61
This study
P. edulis (KF991277), 94%
P. adspersus (KY177061), 95%
89
P. adspersus (AF215505), 94%
P. edulis (EF107211), 93%
100
P. edulis (DQ283157), 93%
58
P. adspersus (AF206472), 92%

0.010

C

72
Pyxicephalus adspersus
59
Pelophylax nigromaculatus
Odorrana margaretae
57
58
Rana catesbeiana
98
Rana pipiens
36
Nanorana parkeri
Microhyla fissipes
70
37
Polypedates megacephalus
80
Rhacophorus dennysi
99
Bufotes viridis
Rhinella marina
87
Dryophytes cinereus
58
Cyclorana alboguttata
Oreobates cruralis
43
Leptobrachium boringii
Xenophrys sangzhiensis
100
Xenopus laevis
61
57
Xenopus tropicalis
Bonbina Bonbina
48
Scaphiopus cauchii

0.050

Figure 2

Figure 3

Figure 4

Figure 5