Familiar neighbours, but not relatives, enhance fitness in a territorial mammal

Authors: Erin R. Siracusa[1,2*], Stan Boutin[3], Ben Dantzer[4,5], Jeffrey E. Lane[6], David W.
Coltman[3], Andrew G. McAdam [1,7]

Affiliations:
[1]Department of Integrative Biology, University of Guelph, 50 Stone Rd E, Guelph, ON, N1G
2W1, CA

[2]Centre for Research in Animal Behaviour, School of Psychology, University of Exeter, Stocker
Rd, Exeter, EX4 4PY, UK

[3]Department of Biological Sciences, University of Alberta, 116 St & 85 Ave, Edmonton, AB,
T6G 2R3, CA

[4]Department of Psychology, University of Michigan, 500 S State St, Ann Arbor, MI, 48109,
USA

[5]Department of Ecology & Evolutionary Biology, University of Michigan, 500 S State St, Ann
Arbor, MI, 48109, USA

[6]Department of Biology, University of Saskatchewan, Science Pl, Saskatoon, SK, S7N 5C8, CA

[7]Department of Ecology and Evolutionary Biology, University of Colorado, 1900 Pleasant Street,
Boulder, CO, 80309, USA

*Lead Contact: Erin R. Siracusa, Center for Research in Animal Behaviour, School of
Psychology, University of Exeter, Exeter, EX4 4QG, UK | email: erinsiracusa@gmail.com |
Twitter: @erin_sira

Keywords: dear-enemy, familiarity, fitness, kin selection, mutualism, reciprocity, red squirrel,
reproductive success, senescence, social behaviour, survival, territoriality

 Graphical Abstract

Summary

One of the outstanding questions in evolutionary biology is the extent to which mutually

beneficial interactions and kin-selection can facilitate the evolution of cooperation by mitigating

conflict between interacting organisms. The indirect fitness benefits gained from associating with

kin are an important pathway to conflict resolution [1], but conflict can also be resolved if

individuals gain direct benefits from cooperating with one another (e.g. mutualism or reciprocity)

[2]. Owing to the kin-structured nature of many animal societies, it has been difficult for

previous research to assess the relative importance of these mechanisms [3-5]. However, one

area that might allow for the relative roles of kin-selection and mutualistic benefits to be

disentangled is in the resolution of conflict over territorial space [6]. While much research has

focused on group-living species, the question of how cooperation can first be favoured in solitary,

territorial species remains a key question. Using 22 years of data from a population of North

American red squirrels, we assessed how kinship and familiarity with neighbours affected fitness

in a territorial mammal. While living near kin did not enhance fitness, familiarity with

neighbours increased survival and annual reproductive success. These fitness benefits were

strong enough to compensate for the effects of aging later in life, with potential consequences for

the evolution of senescence. We suggest that such substantial fitness benefits provide the

opportunity for the evolution of cooperation between adversarial neighbours, offering insight

into the role that mutually beneficial behaviours might play in facilitating and stabilizing social

systems.

Results and Discussion

Stable social relationships among group-living species are known to play an important role in mediating intragroup conflict and enhancing fitness [3,7,8]. For territorial species, maintaining territorial boundaries can be a costly endeavor, and while kin can help to mitigate territorial aggression and enhance fitness [9], there is also potential for stable interactions between neighbours to play an important role in helping to resolve conflict over territorial space [6,10]. It is well understood that familiarity among long-term neighbours allows for the formation of stable social relationships that can facilitate reduced aggression and territory defence, a phenomenon known as the 'dear enemy' effect [11,12]. Although it is not clear whether dear-enemy relationships are stabilized by reciprocity or mutualism, by minimizing negotiation of territory boundaries and alleviating costs of territoriality for both individuals, social familiarity provides a pathway to conflict resolution through mutual direct benefits.

To assess the relative importance of kinship and familiarity for resolving conflict over territorial space and thereby enhancing reproductive success and survival, we used 22 years of data from a natural population of North American red squirrels (*Tamiasciurus hudsonicus*). Red squirrels are arboreal rodents in which both sexes defend exclusive territories year-round [13]. Territories are important for overwinter survival [14] and are defended through vocalizations called 'rattles' [15], which are uniquely identifiable [16]. Juveniles typically disperse 100 m on average and females sometimes bequeath all or part of their territory to offspring [17]. As a result, neighbours are sometimes closely related and have been shown to exhibit affiliative behaviour with kin [18,19]. Red squirrels also have potential to establish long-term familiarity with neighbours since individuals rarely relocate to a vacant territory following natal dispersal

[20]. Familiarity with neighbours is demonstrated to have important benefits for red squirrels

including reduced risk of territory intrusion [21] and reduced time spent on territory defence [22].

In this study we tested whether red squirrels that were more familiar with neighbouring

conspecifics had increased annual reproductive success (ARS) and a higher probability of

surviving to the following year, than squirrels that were less familiar with their social

neighbourhood (i.e. squirrels within a 130 m radius). Through genetic analysis and monitoring of

red squirrel pups in the natal nest we were able to establish a long-term pedigree, allowing us to

assess effects of social familiarity while simultaneously accounting for relatedness. We used

average neighbourhood familiarity and relatedness as our metrics of interest, as previous

research has demonstrated that these metrics have important effects on territory dynamics and

behaviour in red squirrels [21,22]. Given that stable social connections have been demonstrated

to reduce mortality risk and enhance longevity in other species [23], we also assessed whether

maintaining familiar social relationships into later life might help to buffer squirrels against age-

related declines in fitness. To do this, we looked for effects of familiarity on survival and

reproductive success in red squirrels specifically during the 'senescent' period (squirrels $\geq$ 4

years old) [24,25].

Effects of familiarity

Over the 22-year period (1994-2015) in which we analyzed survival and ARS, red squirrels, on

average, had 13 neighbours. Average neighbourhood familiarity ranged between 0 and 1186

days (mean = 191 $\pm$ 4 days; CV = 1.04) and average relatedness ranged between 0 and 0.5 (mean

= 0.05 $\pm$ 0.001; CV = 1.09). Familiarity was not strongly correlated with either relatedness (r = -

0.09) or grid-wide density (r = -0.17). We found no effect of average relatedness of the social

neighbourhood on male or female survival, number of pups sired annually, or number of pups

recruited annually in either the full models or senescent models (all $|\beta| < 0.14$, all $|z| < 1.48$, all P

> 0.13; Table 1). In contrast, familiarity with neighbouring individuals was associated with an

increase in male and female survival ($\beta = 0.18 \pm 0.06$, $z = 2.79$, $P = 0.005$; Table 1) and an

increase in the annual number of pups sired ($\beta = 0.18 \pm 0.09$, $z = 2.04$, $P = 0.04$; Table 1),

although there was no overall effect of social familiarity on the annual number of pups recruited

by females ($\beta = 0.06 \pm 0.05$, $z = 1.38$, $P = 0.17$; Table 1). Familiarity is, however, strongly

correlated with age (and therefore survival) in early life ($r = 0.60$), making it difficult to

disentangle whether familiarity drives survival or survival drives familiarity. However, for

senescent squirrels (≥ 4 years), age and familiarity are disassociated ($r = -0.04$; Figure S1), such

that familiarity is not driven by an individual's own survival but by the survival of its neighbours,

allowing us to better disentangle these effects. When analyzing effects of familiarity in the

senescent period (≥ 4 years old), we found that the benefits of social familiarity for all fitness

measures were even more substantial, and were consistently at least three times greater than the

effects of relatedness in the same models (Table 1). For squirrels aged 4 and older, living near

familiar neighbours was associated with an increase in the probability of annual survival for both

sexes ($\beta = 0.45 \pm 0.11$, $z = 3.40$, $P < 0.001$; Figure 1). For example, senescent squirrels with high

familiarity with their neighbours (500 days; third quartile) had a 69% chance of surviving to the

next year, while squirrels with low familiarity with their neighbours (130 days; first quartile) had

a 52% chance of surviving to the next year. In the senescent period, living near familiar

neighbours was associated with a substantial increase in the annual number of pups sired ($\beta =$

0.39 ± 0.15, $z = 2.67$, $P = 0.008$; Figure 1). Males with high familiarity with their neighbours

(500 days, third quartile) sired 1.7 pups annually while males with low familiarity with their

neighbours (130 days, first quartile) sired 0.9 pups annually. For females aged 4 and older,

familiarity with neighbours was also associated with an increase in the annual number of pups

recruited ($\beta = 0.23 \pm 0.09$, $z = 2.52$, $P = 0.01$; Figure 1). Senescent females with high familiarity

with their neighbours (500 days; third quartile) recruited 0.7 pups annually compared to females

with low average familiarity with their neighbours (130 days; first quartile) that recruited 0.5

pups annually.

An important consideration is whether variation in individual quality or habitat quality

might lead to a spurious correlation between familiarity and survival when the causation is

actually reversed. Familiarity, however, is not a trait of an individual but a trait of an interaction

between individuals. Therefore, an individual with 'good genes' that survives well cannot obtain

high familiarity without also having neighbours that survive well. For this to be possible there

would need to be spatial clustering of 'high quality' individuals or spatial variation in resource

availability or predation risk that leads to local areas of consistently high survival (and therefore

high familiarity). Furthermore, in order for local areas of high habitat or individual quality to

confound the familiarity-fitness relationship, spatial autocorrelation in these fitness measures

must persist across multiple years (because familiarity is a product of past survival). To assess

this possibility, we calculated Moran's I for each study grid and year to test whether similarity in

ARS or probability of survival was related to the spatial proximity of any two individuals.

Evidence of spatial autocorrelation in any of these measures was both rare and inconsistent

(Table S4). We found only one instance of spatial correlation in consecutive years (Table S4),

suggesting that a spurious correlation between ARS or survival and familiarity is unlikely. When

we removed the year/grid combinations with positive spatial autocorrelation from our data and

reran the analyses our conclusions remained the same (Table S5). In addition, while survival and

143 familiarity might be confounded in younger squirrels where familiarity increases with age, we

144 demonstrated that familiarity still has substantial effects on fitness in older squirrels where age

145 and familiarity are independent and variation in familiarity is driven by the survival of a

146 squirrel's neighbours rather than their own survival (Figure 1; Table 1). Finally, there is no

147 mechanism by which reproductive success might affect familiarity. Therefore, there is

148 substantial evidence to suggest that familiarity is a cause rather than a consequence of high

149 fitness.

150  Although we do not have direct evidence for the mechanism by which familiarity with

151 neighbours might lead to increased reproductive success, previous work in this study system has

152 shown that social familiarity results in reduced risk of pilfering [21] and reduced time spent on

153 territory defence [22]. Conceivably then, these direct fitness benefits may result from cooperative

154 'agreements' of non-intrusion among long-term neighbours, which might increase food stores or

155 reduce time and energy allocated to territory defence. Given the particularly substantial effects of

156 familiarity on male siring success, we sought to provide some mechanistic understanding of how

157 familiarity contributes to fitness by exploring two hypotheses in male red squirrels. We

158 hypothesized first that social familiarity might directly benefit males through increased mating

159 success with familiar females in their social neighbourhood. Alternatively, we hypothesized that

160 if familiarity enhances energetic resources through reduced pilferage [21,22] that this might

161 allow male squirrels in familiar social neighbourhoods to travel farther to obtain mating

162 opportunities and thereby increase their siring success [26,27]. Our analyses revealed that there

163 was no association between familiarity and a male's siring success within his social

164 neighbourhood ($\leq$ 130 m away; $\beta = 0.12 \pm 0.10$, $t = 1.25$, $P = 0.21$; Table 2). Instead, we found

165 evidence that males traveled farther to mate ($\beta = 19.93 \pm 8.83$, $t = 2.26$, $P = 0.03$) and sired more

pups ($\beta = 0.39 \pm 0.10$, $z = 3.79$, $P <0.001$) outside their social neighbourhood (>130 m away)

when more familiar with their neighbours (Figure 2; Table 2). This does not appear to be due to

inbreeding avoidance (see Table 2). Together, these results suggest that stable social

relationships with neighbours enhance male fitness by allowing them to travel farther and

increase their siring success [27]. This might result from familiarity with neighbours increasing

the energetic resources available to males through reduced territory defence [22] or reduced

cache pilfering by neighbours [21]. While the mechanisms by which social familiarity might

affect female recruitment remain unexplored, there are several pathways by which this might

occur, including familiarity increasing food resources, which might lead to earlier parturition

[28] or faster growth rates [29], known to be associated with enhanced juvenile survival.

Alternatively the reduced time spent on defence in familiar neighbourhoods might allow females

to spend more time in nest [22], and more attentive mothers have been found to have enhanced

reproductive success [30]. Exploring the mechanisms by which social familiarity contributes to

fitness is a viable area for future research.

Effects on senescence

Our findings also demonstrate a previously unappreciated benefit of maintaining social

relationships into later life. The magnitudes of the benefits of social familiarity in the senescent

period were sufficient to offset age-related declines in survival and reproductive success.

Specifically, for an average 4-year old squirrel, aging by one year decreases the probability of

survival from 68% to 59%. But if that individual were to maintain all of its neighbours, such that

average familiarity increased by one year (from the mean familiarity of 318 days to 683 days),

the maintenance of those stable social relationships would more than compensate for the change

in age, resulting in an increase in survival probability from 68% to 74%. However, despite the

large individual benefits of increased social familiarity late in life, few squirrels currently enjoy

this fitness advantage. For example, only 4% of 4-year olds surviving to age 5 maintain all of

their neighbours, and mean familiarity does not continue to increase with age through the

senescent period (Figure S1). As a result, our observed fitness increases in later life associated

with social stability are not currently widespread enough to affect the decline in the force of

natural selection with age and the arrival of the mutational 'wall of death' [31] (Figure S2).

Nevertheless, such individual fitness benefits of familiarity provide strong incentive for

squirrels to reduce turnover in their social neighbourhood. While squirrels compete with

territorial neighbours for resources [32], space [13], and reproductive success [33], here we have

shown that individuals also benefit from the enhanced survival of neighbours. The idea that

territorial neighbours might engage in cooperative behaviours to prevent neighbourhood turnover

has been theorized [34]. Here our documented benefits of social familiarity provide the

opportunity for the evolution of cooperative behaviour toward otherwise adversarial neighbours.

Specifically, while the loss of an unfamiliar neighbour would have no effect on survival

probability, the loss of a neighbour with 6 years of familiarity, although rare, would decrease the

owner's probability of survival by 7%. A squirrel should, therefore, be willing to engage in

behaviours that ensure their neighbour's survival as long as this does not reduce the actor's

survival by more than 7%. This scope for cooperation is comparable to kin selection interactions

between first cousins once removed (r = 0.06) [1]. It is possible then that reduced aggression

when interacting with familiar neighbours (i.e. the dear-enemy phenomenon) [12] is a

mechanism that not only serves to reduce an individual's own time and energy spent on defence,

but also enhances the survival of familiar neighbours, which provides indirect fitness returns

through increased social familiarity. In red squirrels, reduced territory defence toward familiar

individuals [22] may convey survival benefits to neighbours through reduced aggression or

lowered defense of food resources. Red squirrels have also been known to nest communally

during cold temperatures [18], which might enhance neighbour survival. Other mechanisms by

which red squirrels might reduce neighbourhood turnover such as predator alarm calling, or

forming defensive coalitions [e.g. 34] to prevent a territory takeover, have yet to be tested.

Ultimately, if cooperative behaviours enhancing the survival of familiar neighbours were to

become widespread enough that mean social familiarity increased with age across the population,

then this socially driven increase in fitness late in life could influence the force of selection and

consequently the evolution of senescence (Figure S2).

Effects of relatedness

Given that there is evidence of affiliative behaviour among kin in this study system [18,19], in

addition to localized dispersal and bequeathal of territory space to offspring [17], we expected

that relatedness among neighbours might play an important role in red squirrel survival and

reproductive success. Additionally, we were interested in the possibility that relatedness with

neighbours might help to mitigate the consequences of living with unfamiliar neighbours, as seen

in other systems [6], However, we found no evidence to suggest that kin provide fitness benefits

for red squirrels (Table 1), and there were no significant interactions between relatedness and

familiarity in any of our models. It is possible that the observed low relatedness among

neighbours (average neighbourhood relatedness $= 0.05 \pm 0.001$), which likely occurs as the result

of low overwinter juvenile survival [35], could have constrained the potential for kin selection to

act at this scale. However, our findings do not mean that neighbour relatedness has no

biologically meaningful effects on red squirrel fitness, but simply that the scale at which we

measured these effects did not exhibit any influence of kinship. If, for instance, the primary

benefits of kinship come from sharing resources such as food [21] or nests [18], relatedness of

nearest neighbours (rather than all neighbours within 130 m) may be a more relevant measure to

assess whether red squirrels obtain benefits from kin. Future work should therefore attempt to

use other measures of the social environment to offer additional insight as to the relative

importance of kin-selection in this system. Regardless, our results demonstrate that stable social

relationships are associated with increased fitness in red squirrels, independent of kinship,

suggesting that inclusive fitness benefits are not a necessary prerequisite to facilitate cooperation

between social partners.

Conclusion

Although interactions among territorial individuals are fundamentally defined by competition for

food, mates, or other resources, here we have shown that stable social relationships can provide

important fitness benefits that might feed back to limit these competitive interactions and

maintain strong affiliative relationships among unrelated territorial neighbours. Similar fitness

benefits of established relationships with neighbours have also been demonstrated in territorial

birds such as red-winged blackbirds [36], song sparrows [37], and great tits [10]. Our findings, in

conjunction with these previous studies, demonstrate that the benefits of stable social

relationships are not exclusive to group-living organisms but can extend to solitary species,

suggesting a shared evolutionary importance of stable relationships across a range of social

complexities. The strength of the effects of familiarity, particularly in the senescent period, was

surprising. We suggest that by increasing the force of selection later in life, stable social

relationships have the potential to lead to the evolution of slower rates of aging, although these

late-life social benefits are not currently widespread enough to affect senescence in red squirrels.

Importantly, our results indicate that intraspecific conflict can be mitigated through mutualistic

pathways, with resulting fitness benefits, independent of kinship effects [3,6]. Consequently,

stable social relationships with neighbours may be an important and overlooked contributor to

fitness in territorial species, and may offer insight into the relative importance of mutually

beneficial behaviours for facilitating conflict resolution.

Acknowledgments: We thank the Champagne and Aishihik First Nations for allowing us to

conduct our research within their traditional territory and thank Agnes MacDonald and her

family for long-term access to her trap line. We are grateful to all of the field technicians who

have contributed to the long-term KRSP database over the years, and J. Kapitain at KAP Design

for assistance with figure design. This research was supported by Discovery Grants and Northern

Research Supplements from the Natural Sciences and Engineering Research Council of Canada

(A.G.M.: RGPIN-2015-04707; S.B.: RGPIN-2014-05874; J.E.L.: RGPIN-2014-04093, RGPNS-

2014-459038; D.W.C.: RGPIN-2011-312207), as well as funding from the University of

Michigan (B.D.), and Grants-in-Aid of research from the American Society of Mammalogists

and the Arctic Institute of North America (E.R.S.).

Author Contributions: E.R.S. and A.G.M. conceived of and designed the study. E.R.S.

conducted statistical analyses and wrote the manuscript with support from A.G.M. All authors

contributed to field logistics, data collection and maintenance, and assisted with review and

editing of the manuscript.

Declaration of Interests: The authors declare no competing interests.

Figure 1. Effects of average neighbourhood familiarity on A) annual survival (N = 512), B) male annual reproductive success (ARS; N = 78), and C) female ARS (N = 268) during the senescent period (≥ 4 years old). Shaded grey bars indicate 95% confidence intervals. Values on x-axis are standardized measures of average familiarity. Points indicate raw data with a small amount of jitter introduced to show overlapping points.

Figure 2. Effects of average neighbourhood familiarity on A) the distance that males traveled to mate and B) number of pups sired outside the social neighbourhood (i.e. 130 m radius; N = 199). Shaded grey bars indicate 95% confidence intervals. Values on x-axis are standardized measures of average familiarity. Points indicate raw data. See also Table 2 and S2.

297 **Table 1. Fixed effects from annual survival, male annual reproductive success (ARS), and**

298 **female ARS generalized linear mixed-effects models.** Models are based on the full and

299 senescent (≥ 4 years old) datasets are shown with significant effects indicated in bold.

300 Regression coefficients are standardized. See also Table S1 for random effects.

Data set	N	Response	Fixed effect	Parameter ± SE	z	P
Full data						
	2346	Probability of survival	**Familiarity**	**0.18 ± 0.06**	**2.79**	**0.005**
			Age	**-0.27 ± 0.08**	**-3.47**	**0.001**
			Age^2	**-0.11 ± 0.04**	**-2.63**	**0.01**
			Relatedness	0.01 ± 0.05	0.16	0.88
			Density	0.10 ± 0.10	0.92	0.36
			Grid-SU	0.17 ± 0.10	1.67	0.09
	412	No. pups sired	**Familiarity**	**0.18 ± 0.09**	**2.04**	**0.04**
		(Male ARS)	**Age**	**0.68 ± 0.14**	**4.75**	**<0.001**
			Age2	**-0.41 ± 0.08**	**-5.32**	**<0.001**
			Relatedness	-0.12 ± 0.08	-1.47	0.14
			Density	0.04 ± 0.12	0.37	0.71
			Grid-SU	-0.07 ± 0.19	-0.38	0.70
	981	No. pups recruited	Familiarity	0.06 ± 0.05	1.38	0.17
		(Female ARS)	Age	0.08 ± 0.05	1.42	0.16
			Age^2	**-0.10 ± 0.04**	**-2.72**	**0.007**
			Relatedness	0.03 ± 0.04	0.63	0.53
			Density	**-0.37 ± 0.13**	**-2.78**	**0.005**
			Grid-SU	-0.06 ± 0.08	-0.78	0.43
Senescent data						
	512	Probability of survival	**Familiarity**	**0.45 ± 0.11**	**4.00**	**<0.001**
			Age	**-0.36 ± 0.10**	**-3.42**	**0.001**
			Relatedness	0.13 ± 0.11	1.20	0.23
			Density	0.15 ± 0.11	1.33	0.18
			Grid-SU	0.14 ± 0.21	0.68	0.49
	78	No. pups sired	**Familiarity**	**0.39 ± 0.15**	**2.67**	**0.008**
		(Male ARS)	**Age**	**-0.39 ± 0.19**	**-2.01**	**0.04**
			Relatedness	0.05 ± 0.14	0.37	0.72
			Density	0.11 ± 0.19	0.60	0.55
			Grid-SU	0.26 ± 0.32	0.82	0.41
	268	No. pups recruited	**Familiarity**	**0.23 ± 0.09**	**2.52**	**0.01**
		(Female ARS)	**Age**	**-0.29 ± 0.10**	**-2.79**	**0.005**
			Relatedness	0.06 ± 0.08	0.79	0.43

Density	**-0.38 ± 0.17**	**-2.20**	**0.03**
Grid-SU	-0.32 ± 0.17	-1.85	0.07

303 **Table 2. Fixed effects from models assessing effects of familiarity on (i) the farthest**

304 **distance that males traveled to mate, (ii) the number of pups sired inside the**

305 **neighbourhood, and (iii) the number of pups sired outside the neighbourhood (N = 199**

306 **observations over 129 males).** Regression coefficients are standardized and significant effects

307 are indicated in bold. See also Table S2 for random effects.

Response	Fixed Effect	Parameter ± SE	Test-statistic	P
Distance of farthest mate	**Familiarity**	**19.93 ± 8.83**	**2.26**	**0.03**
	Relatedness	6.91 ± 6.82	1.01	0.31
	Age	-4.67 ± 9.56	-0.49	0.63
	Density	**-24.86 ± 7.49**	**-3.32**	**0.001**
No. pups sired inside	Familiarity	0.12 ± 0.10	1.24	0.21
	Relatedness	0.01 ± 0.07	0.18	0.85
	Age	0.03 ± 0.11	0.24	0.81
	Density	**0.27 ± 0.12**	**2.32**	**0.02**
No. pups sired outside	**Familiarity**	**0.39 ± 0.10**	**3.78**	**<0.001**
	Relatedness	-0.003 ± 0.09	-0.04	0.97
	Age	-0.09 ± 0.12	-0.77	0.44
	Density	**-0.25 ± 0.09**	**-2.66**	**0.008**

STAR Methods

Resource Availability

Lead Contact

Further information and requests for resources and reagents should be directed to and will be

fulfilled by the Lead Contact, Erin Siracusa (erinsiracusa@gmail.com).

Materials Availability

This study did not generate new unique reagents.

Data and Code Availability

Data are available from the Figshare Repository (https://doi.org/10.6084/m9.figshare.8813564).

Experimental Model and Subject Details

We studied a wild population of North American red squirrels located in the southwest Yukon,

Canada (61° N, 138° W). This population has been followed since 1987 on two 40-ha study grids

separated by the Alaska Highway ('Kloo' and 'Sulphur'). We monitored squirrels annually from

March to August and used a combination of live-trapping and behavioural observations to assess

territory ownership, track reproduction and survival, and determine offspring recruitment [17,25].

We trapped squirrels using Tomahawk live-traps (Tomahawk Live Trap Co., Tomahawk,

Wisconsin, USA) baited with peanut butter. During their single-day oestrous, female red

squirrels mate with multiple males and produce multiply-sired litters [38], with an average of

three pups per litter [39]. We monitored the reproductive statuses and parturition dates of

females through palpation, mass changes, and evidence of lactation. After parturition, we used

radio telemetry and/or behavioural observations to follow females to their nests. We fitted pups

with unique alphanumeric metal ear tags (Monel #1; National Band and Tag, Newport, KY,

USA) at 25 days old, allowing us to follow individuals throughout their lifetimes.

Ethical guidelines

This study required trapping individuals using Tomahawk live traps to determine territory

ownership and assess reproductive status. Traps were checked every 60-90 min and squirrels

were never left in a trap for longer than 120 min. We also entered the natal nest when pups were

1-2 days old and 25 days old to collect DNA, measure pup growth, and tag individuals. We

returned the pups to the natal nest immediately after processing to minimize time spent away

from the dam. These procedures had no detectable negative effects on the survival or welfare of

the study animals. This research was approved by the University of Guelph Animal Care

Committee (AUP number 1807).

Method Details

Measuring familiarity

We completed a full census of the population twice annually in May and August and determined

territory ownership using the aforementioned methods. Both male and female red squirrels

defend exclusive territories year round, which are centered around a larder hoard of food

resources called a 'midden' [13]. We defined each squirrel's social neighbourhood to be all

conspecifics whose middens (i.e. the center of the territory) were within a 130 m radius of the

focal squirrel's midden. One hundred and thirty meters is the farthest distance that red squirrel

territorial vocalizations ('rattles') are known to be detectable to the human ear [40], and is

similar to the distance at which red squirrels have been demonstrated to be most responsive to

changes in local density (150 m) [41]. This suggests that 130 m is a reasonable measure of the

distance at which squirrels can receive and respond to information about their social environment.

We measured the pairwise familiarity between the territory owner and each neighbour as the

number of days that both individuals had occupied their current territories within 130 m of each

other. We then averaged all pairwise measures of familiarity to obtain a measure of each

individual's average familiarity with its neighbours. For all analyses we used average familiarity

values calculated in May, as this aligned most closed with the reproductive season. The timing of

our semi-annual censuses meant that we could only update each squirrel's familiarity with its

neighbours twice per year. Therefore, pairwise familiarities could increase from one year to the

next in increments of either 273 days (new neighbour in August survived until May), or 365 days

(neighbour survived from previous May until current May). Average familiarity, however, could

take on many possible values because pairwise familiarities were averaged across all neighbours.

Measuring relatedness

We temporarily removed juveniles from their natal nest immediately following parturition to

weigh, sex, and take ear tissue samples. Observing pups in the natal nest also allowed us to

assign maternity with certainty. Starting in 2003, we determined paternity by genotyping all

individuals at 16 microsatellite loci [42] and assigned paternity with 99% confidence using

CERVUS 3.0 [43]. Genotyping error based on known mother-offspring pairs was 2%. Additional

details on the genetic methods can be found in Lane et al. (2008). Using the established

multigenerational pedigree, we calculated the coefficient of relatedness between the territory

owner and each neighbour. We averaged all pairwise measures of relatedness to provide a

measure of the average relatedness between each squirrel and all of its neighbours.

Fitness measures

For individuals alive between 1994 and 2015, we measured annual survival as the probability of

surviving to the following year ($N = 2346$ records over 1009 individuals). Given that long-term

monitoring of individuals in this population first began in 1987, we excluded data prior to 1994

to ensure accurate measurement of familiarity between neighbours. Our ability to successfully

redetect adults in the population each year is estimated to be 1.0 [44].

We measured female annual reproductive success (ARS) as the number of pups recruited

each year (i.e. surviving overwinter). We included data from 450 breeding females between 1994

and 2015 ($N = 981$ records). All females with a known parturition date were considered to have

bred; all other non-breeding females were excluded from analysis. Most non-breeding females

were yearlings that were too young to breed. Since red squirrel males do not contribute to the

raising of pups and the male's social environment is, therefore, unlikely to affect pup recruitment,

we defined male ARS as the number of pups sired each year. We used data from 207 males

between 2003 and 2014 ($N = 412$ records). We excluded male reproductive success outside of

this timeframe because paternity data was not available prior to 2003 and after 2014. For all

analyses we only used data for adults (≥ 1 year old) whose age could be assigned with certainty

(i.e. individuals tagged in nest as juveniles).

It is possible that our ability to accurately assess recruitment is somewhat confounded by

dispersal beyond the borders of our study grids [45]. On average, juveniles typically disperse less

than 100 m [17,46], and estimates of juvenile survival do not differ between the center and edge

of our study grids, which might be expected if dispersing juveniles were mistakenly considered

to have died [25]. However, 37% of our population is comprised of immigrants from outside our

study grids [45], which suggests that some juvenile recruitment outside the study area is likely to

have been missed by our methods.

Quantification and Statistical Analysis

Annual survival and ARS

To assess the effects of familiarity on ARS and survival, we used generalized linear mixed

effects models (GLMMs) with a BOBYQA optimizer. We used a binomial distribution (logit-

link) to assess probability of annual survival and a Poisson distribution (log-link) for male and

female ARS. For all models we fitted average familiarity, average relatedness, a linear and

quadratic term for age, and grid-wide density as continuous predictors and included study grid as

a categorical fixed effect (Table 1). For the survival models we checked for an effect of sex, but

sex was not significant and so was not considered further. We included a random effect of

squirrel ID and year to account for repeated measures of individuals and temporal variation in

resource availability, respectively (Table S1). To account for inherent spatial structure in our

data, we grouped squirrels into 150 m squares (within each year) based on their known spatial

locations and included this 'spatial ID' as a random effect in all of our models. Our results

remained unchanged even when this spatial grouping term was defined to be a 75 m or 300 m

square (see Table S3). Fitting separate spatial autocorrelation terms for each year would have

been ideal, however, it was not possible to obtain convergence from our data using this method.

To account for the fact that familiarity and age are strongly correlated in early, but not later, life

(Figure S1), we also fit models for just the senescent period ($\geq$ 4 years old). Models for the

senescent period included the same covariates as the full models described above, but only

included a linear effect of age. Finally, we conducted an additional analysis to test for spatial

autocorrelation by calculating Moran's I (ape package version 5.1) [47] for each study grid and

year (Table S4). We removed the year/grid combinations with positive spatial autocorrelation

from our data and reran the analyses above (Table S5).

Mechanisms underlying male reproductive success

We conducted a post-hoc analysis to reveal potential mechanisms by which familiarity with

neighbours might lead to increased male reproductive success. We had two hypotheses: 1) that

familiarity with neighbours leads to increased energetic resources, allowing males to travel

farther to mate or 2) that familiarity with females provides males with more mating opportunities

within the males' social neighbourhood. To test hypothesis one, we assessed whether familiarity

with neighbours affected the farthest distance that a male traveled to mate using a linear mixed

effect model (LMM) with average familiarity, average relatedness, age, and grid density as

continuous fixed effects and squirrel ID and year as random effects (Table 2, S2). To test

hypothesis two, we measured whether average familiarity affected the number of pups that a

male sired inside or outside his neighbourhood. We fitted both these models using a Poisson

GLMM with the same fixed and random effects structure as above (Table 2, S2).

Force of selection

If the fitness benefits of stable social relationships were sufficiently widespread in the red

squirrel system then an increase in the mean number of reproductive opportunities later in life

associated with social familiarity would lead to a less rapid decline in the force of natural

selection with age. This would delay the arrival of the 'wall of death', a spike in mortality caused

by the accumulation of deleterious alleles, leading to the evolution of slower rates of aging [31].

We conducted a post-hoc analysis to explore this possibility in red squirrels by creating a

'simulated' population where social interactions were excluded from the calculation of age-

specific survival and fecundity (Figure S2). To do this, we fitted the same models as above,

except that we included both breeding and non-breeding females and the response variable for

females was now number of pups born rather than number recruited. For the first set of models

we included age, a quadratic term for age, density and grid as fixed effects, as well as squirrel ID

and year as random effects. We extracted the parameter estimates for age in each of these models

and used those parameters to estimate age-specific fecundity and survival for females and males

separately. This represented our 'observed' population, and provided us with the estimates of

age-specific fecundity and survival as observed and as potentially confounded by social

familiarity among squirrels. We then measured what the effects of age would have been had

there not be been variation in familiarity by including familiarity as a continuous predictor in the

same models described above, and thus statistically controlling for its effects. Our parameter

estimates for age now represented the effects of age in the absence of social effects. We again

used these parameters to estimate age-specific fecundity and survival for both sexes. We

calculated force of selection separately for males and females in our 'observed' population and

'simulated' population (where the benefits of social relationships were statistically excluded) by

using the following formula [48]:

$$S(x) = \sum_{y=x+1} e^{-ry} l(y)m(y)$$

Here $S(x)$ represents the force of selection against a deleterious allele in the population, where r

is the intrinsic rate of increase, $l(y)$ and $m(y)$ are the survivorship and fecundity functions and y

is used to sum up the net expected reproduction over all ages after age x.

Data analysis

We conducted statistical analyses using R version 3.4.1 [49] and fitted models using the lme4

package (version 1.1-13) [50]. We obtained P values for the LMMs using the package lmerTest

(version 3.0-1) [51]. For all models, we checked for significant non-linearities between the

predictor and response variables by fitting generalized additive models. For GLMMs we checked

for overdispersion by assessing whether the sum of squared Pearson residuals approximated a

Chi-squared distribution with *N-P* degrees of freedom [52]. Of the 1009 squirrels used in this

analysis 12 individuals were not in the pedigree, meaning there were 25 neighbourhoods for

which we could not calculate relatedness with neighbours. We assigned these neighbourhoods

the mean average neighbourhood relatedness (r = 0.05). (Results remained the same even when

these 25 neighbourhoods were removed from the analysis). To facilitate direct comparison of

effect sizes [53] we standardized all continuous predictors to a mean of zero and unit variance.

We estimated all effect sizes by using first and third quartile familiarity values while holding all

other variables constant at their mean and using 'Kloo' as the study grid. We present all means $\pm$

SE, unless otherwise stated, and consider all differences statistically significant at $\alpha < 0.05$.

References:

[1]   Hamilton WD. (1964). The genetical evolution of social behaviour. I. Journal of
        Theoretical Biology. 7, 1–16. doi:10.1016/0022-5193(64)90038-4.
[2]   Clutton-Brock T. (2002). Breeding Together: Kin Selection and Mutualism in
        Cooperative Vertebrates. Science. 296, 69–72. doi:10.1126/science.296.5565.69.
[3]   Cameron EZ, Setsaas TH, Linklater WL. (2009). Social bonds between unrelated
        females increase reproductive success in feral horses. PNAS. 106, 13850–3.
        doi:10.1073/pnas.0900639106.
[4]   Schino G, Aureli F. (2010). The relative roles of kinship and reciprocity in explaining
        primate altruism. Ecology Letters. 13, 45–50. doi:10.1111/j.1461-0248.2009.01396.x.
[5]   Carter GG, Wilkinson GS. (2013). Food sharing in vampire bats: reciprocal help predicts
        donations more than relatedness or harassment. Proceedings of the Royal Society B:
        Biological Sciences. 280, 20122573–8. doi:10.1098/rspb.2012.2573.
[6]   Bebbington K, Kingma SA, Fairfield EA, Dugdale HL, Komdeur J, Spurgin LG,
        Richardson DS. (2017). Kinship and familiarity mitigate costs of social conflict between
        Seychelles warbler neighbors. PNAS. 46, 201704350–15. doi:10.1073/pnas.1704350114.
[7]   Silk JB. (2003). Social Bonds of Female Baboons Enhance Infant Survival. Science. 302,
        1231–4. doi:10.1126/science.1088580.
[8]   Vander Wal E, Festa-Bianchet M, Reale D, Coltman D, Pelletier F. (2015). Sex-based
        differences in the adaptive value of social behavior contrasted against morphology and
        environment. Ecology. 96, 631–41.
[9]   Viblanc VA, Arnaud CM, Dobson FS, Murie JO. (2010). Kin selection in Columbian
        ground squirrels (*Urocitellus columbianus*): littermate kin provide individual fitness
        benefits. Proc Biol Sci. 277, 989–94. doi:10.1098/rspb.2009.1960.
[10]   Grabowska-Zhang AM, Wilkin TA, Sheldon BC. (2012). Effects of neighbor familiarity
        on reproductive success in the great tit (*Parus major*). Behavioral Ecology. 23, 322–33.
        doi:10.1093/beheco/arr189.
[11]   Fisher J. (1954). Evolution and bird sociality. In: Huxley J, Hardy AC, Ford EB, editors.
        Evolution as a Process, London, U.K. pp. 71–83.
[12]   Temeles E. (1994). The role of neighbours in territorial systems: when are they'dear
        enemies'? Animal Behaviour. 47, 339–50.
[13]   Smith CC. (1968). The Adaptive Nature of Social Organization in the Genus of Three
        Squirrels Tamiasciurus. Ecological Monographs. 38, 31–64. doi:10.2307/1948536.
[14]   LaMontagne JM, Williams CT, Donald JL, Humphries MM, McAdam AG, Boutin S.
        (2013). Linking intraspecific variation in territory size, cone supply, and survival of
        North American red squirrels. Journal of Mammalogy. 94, 1048–58. doi:10.1644/12-
        MAMM-A-245.1.
[15]   Siracusa E, Morandini M, Boutin S, Humphries MM, Dantzer B, Lane JE, McAdam AG.
        Red squirrel territorial vocalizations deter intrusions by conspecific rivals. (2017).
        Behaviour. 154, 1259–73. doi:10.1163/1568539X-00003467.
[16]   Wilson DR, Goble AR, Boutin S, Humphries MM, Coltman DW, Gorrell JC, Shonfield
        J, McAdam AG. (2015). Red squirrels use territorial vocalizations for kin discrimination.
        Animal Behaviour. 107, 79–85. doi:10.1016/j.anbehav.2015.06.011.
[17]   Berteaux D, Boutin S. (2000). Breeding dispersal in female North American red
        squirrels. Ecology. 81, 1311–26. doi:10.1890/0012-

9658(2000)081[1311:BDIFNA]2.0.CO;2.

[18]   Williams CT, Gorrell JC, Lane JE, McAdam AG, Humphries MM, Boutin S. (2013).
Communal nesting in an "asocial" mammal: social thermoregulation among spatially
dispersed kin. Behav Ecol Sociobiol. 67, 757–63. doi:10.1007/s00265-013-1499-4.

[19]   Gorrell JC, McAdam AG, Coltman DW, Humphries MM, Boutin S. (2010). Adopting
kin enhances inclusive fitness in asocial red squirrels. Nature Communications. 1, 1–4.
doi:10.1038/ncomms1022.

[20]   Larsen KW, Boutin S. (1995). Exploring territory quality in the North American red
squirrel through removal experiments. Canadian Journal of Zoology. 73, 1115–22.
doi:10.1139/z95-133.

[21]   Siracusa E, Boutin S, Humphries MM, Gorrell JC, Coltman DW, Dantzer B, Lane JE,
McAdam AG. (2017). Familiarity with neighbours affects intrusion risk in territorial red
squirrels. Animal Behaviour. 133, 11–20. doi:10.1016/j.anbehav.2017.08.024.

[22]   Siracusa ER, Wilson DR, Studd EK, Boutin S, Humphries MM, Dantzer B, Lane JE,
McAdam AG. (2019). North American red squirrels mitigate costs of territory defence
through social plasticity. Animal Behaviour. 151, 29–42.
doi:10.1016/j.anbehav.2019.02.014.

[23]   Snyder-Mackler N, Burger JR, Gaydosh L, Belsky DW, Noppert GA, Campos FA,
Bartolomucci A, Yang YC, Aiello AE, O'Rand A, et al. (2020). Social determinants of
health and survival in humans and other animals. Science. 368, eaax9553–14.
doi:10.1126/science.aax9553.

[24]   Descamps S, Boutin S, Berteaux D, Gaillard J-M. (2008). Age-specific variation in
survival, reproductive success and offspring quality in red squirrels: evidence of
senescence. Oikos. 117, 1406–16. doi:10.1111/j.0030-1299.2008.16545.x.

[25]   McAdam A, Boutin S, Sykes A, Humphries M. (2007). Life histories of female red
squirrels and their contributions to population growth and lifetime fitness. Ecoscience.
14, 362–9.

[26]   Lane JE, Boutin S, Speakman JR, Humphries MM. (2010). Energetic costs of male
reproduction in a scramble competition mating system. Journal of Animal Ecology. 79,
27–34. doi:10.1111/j.1365-2656.2009.01592.x.

[27]   Lane JE, Boutin S, Gunn MR, Coltman DW. (2009). Sexually selected behaviour: red
squirrel males search for reproductive success. Journal of Animal Ecology. 78, 296–304.
doi:10.1111/j.1365-2656.2008.01502.x.

[28]   Kerr T, Boutin S, LaMontagne J, McAdam A, Humphries M. (2007). Persistent maternal
effects on juvenile survival in North American red squirrels. Biology Letters. 3, 289–91.

[29]   Dantzer B, Newman AEM, Boonstra R, Palme R, Boutin S, Humphries MM, McAdam
AG. (2013). Density triggers maternal hormones that increase adaptive offspring growth
in a wild mammal. Science. 340, 1215–7. doi:10.1126/science.1235765.

[30]   Westrick SE, Taylor RW, Boutin S, Lane JE, McAdam AG, Dantzer B. (2020).
Attentive red squirrel mothers have faster growing pups and higher lifetime reproductive
success. Behavioural Ecology and Sociobiology. 74, 1–13. doi:10.1007/s00265-020-
02856-7.

[31]   Hamilton WD. (1966). The moulding of senescence by natural selection. Journal of
Theoretical Biology. 12, 12–45. doi:10.1016/0022-5193(66)90184-6.

[32]   Donald JL, Boutin S. (2011). Intraspecific cache pilferage by larder-hoarding red
squirrels (T*amiasciurus hudsonicus*). Journal of Mammalogy. 92, 1013–20.

doi:10.1644/10-MAMM-A-340.1.

[33]   Fisher DN, Boutin S, Dantzer B, Humphries MM, Lane JE, McAdam AG. (2017).
Multilevel and sex-specific selection on competitive traits in North American red
squirrels. Evolution. 71, 1841–54. doi:10.1111/evo.13270.

[34]   Getty T. (1987). Dear enemies and the prisoner's dilemma: why should territorial
neighbors form defensive coalitions? Integrative and Comparative Biology. 27, 327–36.
doi:10.1093/icb/27.2.327.

[35]   Hendrix JG, Fisher DN, Martinig AR, Boutin S, Dantzer B, Lane JE, McAdam AG.
(2020). Territory acquisition mediates the influence of predators and climate on juvenile
red squirrel survival. Journal of Animal Ecology. 89, 1408–18. doi:10.1111/1365-
2656.13209.

[36]   Beletsky LD, Orians GH. (1989). Familiar neighbors enhance breeding success in birds.
PNAS. 86, 7933–6. doi:10.1073/pnas.86.20.7933.

[37]   Beecher MD, Akcay C, Campbell SE. (2020). Birdsong learning is mutually beneficial
for tutee and tutor in song sparrows. Animal Behaviour. 166, 281–8.
doi:10.1016/j.anbehav.2020.05.015.

[38]   Lane J, Boutin S, Gunn M, Slate J, Coltman D. (2008). Female multiple mating and
paternity in free-ranging North American red squirrels. Animal Behaviour. 75, 1927–37.

[39]   Humphries M, Boutin S. (2000). The determinants of optimal litter size in free-ranging
red squirrels. Ecology. 81, 2867–77.

[40]   Smith CC. (1978). Structure and Function of the Vocalizations of Tree Squirrels
(*Tamiasciurus*). Journal of Mammalogy. 59, 793–808. doi:10.2307/1380144.

[41]   Dantzer B, Boutin S, Humphries MM, McAdam AG. (2012). Behavioral responses of
territorial red squirrels to natural and experimental variation in population density.
Behavioural Ecology and Sociobiology. 66, 865–78. doi:10.1007/s00265-012-1335-2.

[42]   Gunn MR, Dawson DA, Leviston A, Hartnup K, Davis CS, Strobeck C, Slate J, Coltman
DW. (2005). Isolation of 18 polymorphic microsatellite loci from the North American
red squirrel, *Tamiasciurus hudsonicus* (Sciuridae, Rodentia), and their cross-utility in
other species. Mol Ecol Notes. 5, 650–3. doi:10.1111/j.1471-8286.2005.01022.x.

[43]   Kalinowski ST, Taper ML, Marshall TC. (2007). Revising how the computer program
cervus accommodates genotyping error increases success in paternity assignment.
Molecular Ecology. 6, 1099–106. doi:10.1111/j.1365-294X.2007.03089.x.

[44]   Descamps S, Boutin S, McAdam AG, Berteaux D, Gaillard JM. (2009). Survival costs
of reproduction vary with age in North American red squirrels. Proceedings of the Royal
Society B: Biological Sciences. 276, 1129–35. doi:10.1098/rspb.2008.1401.

[45]   Martinig AR, McAdam AG, Dantzer B, Lane JE, Coltman DW, Boutin S. (2020). The
new kid on the block: immigrant males win big whereas females pay fitness cost after
dispersal. Ecology Letters. 23, 430–8. doi:10.1111/ele.13436.

[46]   Cooper EB, Taylor RW, Kelley AD, Martinig AR, Boutin S, Humphries MM, Dantzer B,
Lane JE, McAdam AG. (2017). Personality is correlated with natal dispersal in North
American red squirrels (*Tamiasciurus hudsonicus*). Behaviour. 154, 939–61.
doi:10.1163/1568539X-00003450.

[47]   Paradis E, Schliep K. (2019). ape 5.0: an environment for modern phylogenetics and
evolutionary analyses in R. Bioinformatics. 35, 526–8.
doi:10.1093/bioinformatics/bty633.

[48]   Rose MR, Rauser CL, Benford G, Matos M, Mueller LD. (2007). Hamilton's forces of

natural selection after forty years. Evolution. 61, 1265–76. doi:10.1111/j.1558-
5646.2007.00120.x.
[49] R Core Team. (2017). R: A language and environment for statistical computing .
https://www.R-project.org/.
[50] Bates D, Maechler M, Bolker BM, Walker S. (2015). Fitting linear mixed-effects models
using lme4. Journal of Statistical Software. 67, 1–48. doi:doi:10.18637/jss.v067.i01.
[51] Kuznetsova A, Brockhoff PB, Christensen RHB. (2017). lmerTestPackage: Tests in
Linear Mixed Effects Models. Journal of Statistical Software. 82, 1–26.
doi:10.18637/jss.v082.i13.
[52] Bolker BM, Brooks ME, Clark CJ, Geange SW, Poulsen JR, Stevens MHH, White JSS.
(2009) Generalized linear mixed models: a practical guide for ecology and evolution.
Trends in Ecology & Evolution. 24, 127–35. doi:10.1016/j.tree.2008.10.008.
[53] Schielzeth H. (2010). Simple means to improve the interpretability of regression
coefficients. Methods in Ecology and Evolution. 1, 103–13. doi:10.1111/j.2041-
210X.2010.00012.x.
634