1 **Human Foveal Cone Photoreceptor Topography and its Dependence on Eye Length**
2

3 **Authors:**
4 Wang, Yiyi[1]; Bensaid, Nicolas[3]; Tiruveedhula, Pavan[1,2]; Ma, Jianqiang[4]; Ravikumar, Sowmya[1,2];
5 Roorda, Austin[1,2]
6

7 **Affiliations:**
8   1. School of Optometry, University of California, Berkeley, Berkeley, CA, USA
9   2. Vision Science Graduate Group, University of California, Berkeley, Berkeley, CA, USA
10   3. Carl Zeiss Meditec AG, Berlin Germany
11   4. Department of Mechanical Engineering, Ningbo University, Ningbo, Zhejiang, China
12

Abstract:
We provide the first measures of foveal cone density as a function of axial length in living eyes
and discuss the physical and visual implications of our findings. We used a new generation
Adaptive Optics Scanning Laser Ophthalmoscope to image cones at and near the fovea in 28 eyes
of 16 subjects. Cone density and other metrics were computed in units of visual angle and linear
retinal units. The foveal cone mosaic in longer eyes is expanded at the fovea, but not in proportion
to eye length. Despite retinal stretching (decrease in cones/mm^2), myopes generally have a higher
angular sampling density (increase in cones/deg^2) in and around the fovea compared to
emmetropes, offering the potential for better visual acuity. Reports of deficits in best-corrected
foveal vision in myopes compared to emmetropes cannot be explained by increased spacing
between photoreceptors caused by retinal stretching during myopic progression.
27
28
29

Introduction

There has been a rapid increase in prevalence of myopia, of all magnitudes, in the period between 1971-1972 and 1999-2004 (Vitale, Sperduto, & Ferris, 2009). Across sub-populations grouped by race, ethnicity and gender, several studies report axial length of the eye to be the primary variable related to myopia (Gonzalez Blanco, Sanz Ferńandez, & Muńoz Sanz, 2008; He et al., 2015; Iyamu, Iyamu, & Obiakor, 2011). Increased axial length is associated with retinal stretching and thinning of posterior segment layers and the choroid (Fujiwara, Imamura, Margolis, Slakter, & Spaide, 2009; Harb et al., 2015) and is associated with sight-threatening, often irreversible pathologies of the retina (Morgan, Ohno-Matsui, & Saw, 2012; Verkicharla, Ohno-Matsui, & Saw, 2015). Even without any detectable pathology, the structural changes associated with eye growth ought to have functional consequences for vision.

What Do We Know About Functional Deficits in Myopia?

One might expect that eye growth would stretch the photoreceptor layer and would increase the spacing between cones, causing a longer eye to more coarsely sample an image relative to a shorter eye. However the situation is not that simple; the axial elongation associated with eye growth is accompanied by magnification of the retinal image (Strang, Winn, & Bradley, 1998). If the enlargement of the retinal image exactly matched the stretching of the cone mosaic, then eyes of different lengths would sample the visual field similarly. In fact, in large scale studies, myopes generally attain reasonably good visual acuity with optical correction (He et al., 2004; Jong et al., 2018).

However, more careful inspection reveals that myopes generally (6 out of 9 studies) have poorer angular resolution and have uniformly (3 out of 3 studies) poorer retinal resolution. **Table 1** summarizes published results from psychophysical foveal tasks.

Table 1: Summary of studies investigating foveal spatial vision and sensitivity tasks in myopia.

Author	Refractive error range of myopic cohort [D]	Functional tests	Results for myopes at foveal center	Suggested cause
Fiorentini & Maffei, 1976	-5.5 to -10 (n=10)	CSF	Reduced CSF	Neural insensitivity (myopic amblyopia)
Thorn, Corwin, & Comerford, 1986	-6 to -9.75 (n=13)	CSF	No difference in CSF	Global expansion
Collins & Carney, 1990	-2 to -11 (n=16)	VA, CSF	No difference in VA or CSF between low and high myopic groups with contact lens correction	NA
Strang et al., 1998	0 to -14 (n=34)	VA	Reduced VA (MAR) with increasing myopia after controlling for spectacle magnification	Retinal expansion specifically at the posterior pole; increased aberrations
Liou & Chiu, 2001	0 to >-12 (n=105 eyes)	CSF	Reduced CSF with increasing myopia	Retinal stretching and disruption, neural

				insensitivity (myopic amblyopia)
Chui, Yap, Chan, & Thibos, 2005	-0.5 to -14 (n=60)	Grating resolution	Decreased resolution acuity in cyc/mm	Retinal expansion specifically at the posterior pole; global expansion along with ganglion cell loss
Coletta & Watson, 2006	+2 to -15 (n=17)	Interferometric grating resolution	Decreased resolution acuity in cyc/mm but not in cyc/deg	Retinal expansion specifically at the posterior pole
Atchison, Schmid, & Pritchard, 2006	+0.75 to -12.4 (n=121)	Spatial summation; interferometric grating resolution	Increased critical summation area in linear area, but not in angular area; Decreased resolution acuity in cyc/mm but not in cyc/deg	Retinal expansion specifically at the posterior pole; global expansion along with ganglion cell loss
Stoimenov, 2007	-1 to -8 (n=60)	Contrast thresholds of 20/120 letters	Lower sensitivity to contrast for letters with a fixed angular size	Morphologic changes in the retina
Rossi, Weiser, Tarrant, & Roorda, 2007	−0.5 to −3.75 (n=10)	AO-corrected VA	Reduced acuity (MAR) compared to emmetropes	Retinal expansion, neural insensitivity; neural insensitivity (myopic amblyopia)
Jaworski, Gentle, Zele, Vingrys, & McBrien, 2006	-8.5 to -11.5 (n=10)	Foveal summation thresholds; CSF	Increased critical summation area (angular) Decreased luminance sensitivity Reduced contrast sensitivity at high frequencies (cyc/deg)	Reduction in photoreceptor sensitivity; postreceptoral changes; increased aberrations
Ehsaei, Chisholm, Pacey, & Mallen, 2013	-2.00 to -9.62 (n=60)	Size threshold of high and low contrast letter targets	No difference in threshold retinal image size between myopes and emmetropes.	NA

Most notably, Atchison et al. (2006) and Coletta & Watson (2006) show clear deficits in
retinal resolution (cyc/mm) with increasing myopia using interferometric methods which bypass
the optics of the eye and Rossi et al. (2007) show significant deficits in angular resolution (cyc/deg)
in low myopes, even after using adaptive optics to correct for optical blur. All studies that find
myopic visual deficits implicate retinal stretching as a possible cause, but what is actually
happening structurally at the foveal center during myopic progression is not known. Therefore, the
aim of the current study is to more carefully investigate how the length of the eye affects cone
density at and near the foveal center.

Models for How Photoreceptors Change with Eye Growth

Two types of cone densities will be discussed in this study. Linear density quantifies how
many cones are within a fixed area, in square mm, and serves as a way to evaluate physical retinal
stretching caused by eye growth. Angular density quantifies how many cones are within one degree
visual angle, (the visual angle is measured from the secondary nodal point of the eye). Angular

density serves as a way to evaluate the visual implications of eye growth as it governs the sampling
resolution of the eye.
Figure 1 illustrates three models, along the lines of Strang et al. (1998), of how
photoreceptor structure might be affected by myopic eye growth. In the first model, called the
global expansion model, the retina is proportionally stretched with increasing axial length - cones
are more spaced out in longer eyes - and linear density decreases with eye length. Assuming that
the secondary nodal point remains at a fixed position relative to the anterior segment, the number
of cones within a fixed angular area will remain constant. Therefore, angular cone density will be
constant with eye length. In the second model, called the **equatorial stretching model,** the
posterior retina simply moves axially further from the anterior segment of the eye so that the linear
density does not change with eye length. Since the retina is moving further from the secondary
nodal point, more cones will fall within a fixed angular area and the angular cone density will
increase with eye length. The final model, called the **over-development model**, describes a
structural photoreceptor change that mimics the changes that occur during development (Springer
& Hendrickson, 2004) whereby the photoreceptors continue to migrate towards the fovea as the
eye grows. In this scenario, longer eyes will show both increased linear cone density and an even
steeper increase in angular cone density. The model is motivated by observations of increased
linear cone density in the foveas of marmosets that underwent lens-induced eye growth (Troilo,
1998).
94

95
96 **Figure 1**: 3 models of myopic eye growth: (A) Global expansion shows an eyeball that is
97 proportionally stretched. (B) The equatorial stretching model indicates a growth model where the
98 fovea stays rigid and unaffected as the eye grows. (C) The over-development model shows that
99 myopic eye growth is similar with developmental eye growth where photoreceptors continue to
100 migrate towards the fovea as the eye grows.
101

Previous Studies of Cone Spacing with Axial Length

The most definitive studies of cone spacing as a function of axial length are done through direct imaging of the retina – wherein sharp images of the cones are enabled through the use of adaptive optics, a set of technologies that actively compensate the blur caused by aberrations of the eye (Liang, Williams, & Miller, 1997). Combined with confocal scanning laser ophthalmoscopy (Webb, Hughes, & Delori, 1987), adaptive optics offers the highest contrast *en face* images of the foveal photoreceptor mosaic ever recorded in vivo (Dubra et al., 2011; Roorda et al., 2002).

Despite continued advances in image quality, previous studies investigating cone packing and eye length have not made their measurements at the foveal center, the most important region for spatial vision but the most difficult to image owing to the small size of photoreceptors. There are a number of studies on cone packing and eye length (Chui, Song, & Burns, 2008; Elsner et al., 2017; Kitaguchi et al., 2007; Li, Tiruveedhula, & Roorda, 2010; Obata & Yanagi, 2014; Park, Chung, Greenstein, Tsang, & Chang, 2013) and here we summarize the published results that are most relevant to our study. Chui et al. (2008) investigated angular and linear cone density at 1 mm and 3 degrees eccentricity. They found a significant decrease (P<0.05) in linear cone density as a function of eye length at 1mm (which, by angular distance, is closer to the fovea in a longer eye than in a shorter eye) in all directions except in the nasal retina. They found that the angular cone density at 3 degrees (which, by linear distance, is closer to the fovea, in a shorter eye than in a longer eye) increased with eye length, but the trends were not significant. Li, et al. (2010) made similar measures, but closer to the fovea (from 0.10 mm to 0.30 mm eccentricity). They found that linear cone density decreased with eye length, but the trends were not significant at the smallest eccentricities (0.1 and 0.2 mm). When the data were plotted in angular units and angular distance from the fovea, they found that angular cone density trended toward an increase with eye length but none of the trends were significant. A more recent study measured peak cone densities in the fovea as well as axial length for 22 eyes of 22 subjects (Wilk et al., 2017) but they did not plot peak cone density as a function of axial length, as it was not the aim of their study. We plotted the data they provided in their paper and found that the linear cone density at the foveal center dropped significantly with increases in axial length, similar to what was found by Li et al. (2010) and Chui et al. (2008), but the angular cone density had no dependency on eye length. Summary plots from previous literature are shown in **Figures 2ab.**

Wilk et al. (2017)'s data were consistent with a global expansion model and Li et al. (2010) and Chui et al. (2008)'s data only leaned toward a model that falls between the global expansion and equatorial stretching models. If the trends found by Li et al. (2010) and Chui et al. (2008) near the fovea were to extend to the foveal center, then myopes would have higher foveal photoreceptor sampling resolution with a consequent potential for better performance on visual tasks compared to emmetropes. As such, the simplest explanation for visual deficits in myopes –increased separation between cones caused by retinal stretching – would have to be ruled out.

With the improvements in resolution of adaptive optics ophthalmoscopes, imaging the smallest cones at the foveal center is now possible in many eyes, enabling a definitive analysis of the cone density at the fovea as a function of eye length.

144
145 **Figure 2.** Summary of published data from Li et al. (2010), Chui et al. (2008) and Wilk et al.
146 (2017). In both plots, the linear fits with the solid lines indicate the data that have significant trends.
147 **(a)** Linear cone density has a decreasing trend with axial length near the fovea. **(b)** Angular cone
148 density (sampling resolution) of the eye generally increases with axial length although none of the
149 data show a significant linear relationship.
150
151

Results

The experiments were approved by the University of California, Berkeley Committee for the Protection of Human Subjects. All subjects provided informed consent prior to any experimental procedures. Subjects self-reported their eye health so that only healthy individuals with no ocular conditions were included in the study. All eyes were dilated and cyclopleged with 1% Tropicamide and 2.5% Phenylephrine before imaging. We report data from 28 eyes of 16 subjects with a wide range of refractive error and axial length. Age, sex and ethnicity are listed on **Table 2**.

Biometry Data

All the biometric measures used to convert angular dimensions to linear retinal dimensions are listed on **Table 2.** The strong correlation between refractive error and eye length ($P < 0.0001$) indicates that the myopia was predominantly as a result of axial length.

Imaging Data

Images of the foveal region, the preferred retinal locus for fixation (PRL) and the fixation stability were recorded with an adaptive optics scanning laser ophthalmoscope (see **Methods and Materials**). The image of one subject (10003L) is shown in **Figure 3a**. All the cones were resolved with our imaging system. The scatter plot indicates the scatter plot of fixation over the course of a 10-sec video. **Figure 3b** shows the same image with all cones labeled and a color-coded overlay indicating the density. 16,184 labeled cones are shown on the figure. The point of maximum density is indicated by the blue cross and the average location of the PRL is indicated by the yellow cross (mean of the scatter plot locations in **Figure 3a**). This eye has a peak linear density of 200,482 cones/mm², and a peak angular density of 15,584 cones/deg². Cone density plots in linear and angular units for all eyes are shown on **supplemental figures 1 and 2.** Original images and a list of the cone locations for each can be downloaded from the Resources section of the Roordalab website (roorda.vision.berkeley.edu).

Figure 4 shows the linear cone density as a function of linear eccentricity, where the average linear cone density was computed in 25-micron wide annuli centered around the point of peak density.

Table 2. Each subject's refractive error was self-reported at the time of the study. Axial Length, corneal curvature and anterior chamber depth were measure by IOL Master, and retinal magnification factor (microns/deg) was calculated from biometry data.

Subject ID	Eye	Gender	Age	Ethnicity	Spherical equivalent refraction (D)	Axial length (mm)	Corneal curvature (mm)	Anterior chamber depth (mm)	Retinal magnification factor (microns/deg)	Angular cone density (cones/deg²)	Linear Cone Density (cones/mm²)	PRL distance from fovea (minutes)	PRL distance from fovea (microns)	PRL angular cone density (cones/deg²)	PRL linear cone density (cones/mm²)
20165	L	F	28	Caucasian	0.500	22.26	7.37	3.86	261.79	13247	193288	3.80	16.60	12650	184600
	R	F	28	Caucasian	0.500	22.64	7.44	3.80	267.79	12468	173857	5.48	24.45	11870	165500
20177	L	F	18	Mixed	0.000	23.04	7.80	3.24	273.59	12055	161053	7.12	32.48	11730	156800
	R	F	18	Mixed	0.000	23.23	7.91	3.20	275.85	11780	154810	4.60	21.16	11550	151800
10003	L	M	50	Caucasian	1.000	23.30	7.80	3.12	278.81	15584	200482	7.11	33.02	14070	181000
	R	M	50	Caucasian	1.000	23.50	7.81	3.14	282.00	15172	190784	4.40	20.68	14670	184400
20176	L	F	18	Asian	0.000	23.45	7.98	3.65	276.50	12513	163676	15.82	72.90	8984	117500
	R	F	18	Asian	0.000	23.58	8.01	3.62	278.52	12193	157174	3.97	18.42	11960	154200
20172	L	F	25	Caucasian	-0.750	23.56	7.71	3.90	280.13	15264	194508	2.16	10.06	15170	193300
	R	F	25	Caucasian	-0.500	23.65	7.72	3.96	281.33	14668	185324	3.43	16.08	14760	186500
20147	R	M	26	Caucasian	-0.375	24.16	7.73	2.36	298.73	15401	172581	6.17	30.70	14670	164400
	L	M	26	Caucasian	0.000	24.17	7.81	4.03	288.94	14805	177337	11.70	56.36	13570	162500
20124	L	F	26	Asian	-3.000	24.67	7.70	4.05	298.82	13843	155024	5.15	25.63	13380	149900
	R	F	26	Asian	-4.250	25.29	7.68	4.07	309.88	13659	142247	1.76	9.08	13800	143700
20174	L	F	43	Caucasian	-1.750	24.80	7.79	3.57	302.57	13476	147200	7.67	38.65	11550	126200
	R	F	43	Caucasian	-2.750	25.37	7.83	3.62	311.85	12697	130557	5.90	30.66	11640	119700
20173	R	F	22	Caucasian	-2.750	24.96	7.81	3.68	304.64	16547	178298	7.24	36.73	15910	136000
20170	R	M	26	Asian	-2.250	25.00	7.69	3.90	305.54	14393	154172	8.77	44.65	12740	136500
	L	M	26	Asian	-3.750	25.66	7.65	4.15	316.25	14759	147573	1.50	7.90	14990	149900
20138	R	F	29	Caucasian	-5.000	25.26	7.95	3.14	311.22	13568	140078	6.37	33.05	12830	132500
	L	F	29	Caucasian	-5.000	25.28	7.91	3.15	311.92	14347	147459	5.23	27.20	14300	147000
20114	R	F	24	Asian	-5.500	25.83	8.72	3.47	310.94	14393	148864	8.44	43.72	14070	145500
	L	F	24	Asian	-6.000	26.16	8.98	3.58	313.31	15584	158761	2.71	14.13	15490	157800
20160	R	F	25	Asian	-5.375	25.83	7.81	3.60	320.25	15539	151507	8.97	47.86	14810	144400
20143	R	F	23	Asian	-6.875	25.91	7.42	2.10	334.12	17051	152739	3.07	17.07	16640	149000
20158	R	F	34	Asian	-6.500	26.60	7.84	3.51	333.78	13018	116845	10.58	58.88	10630	95400
20163	R	F	25	Asian	-7.125	26.84	7.89	3.65	336.60	17922	158183	4.16	23.31	17510	154500
	L	F	25	Asian	-7.125	27.06	7.89	3.65	340.44	18793	162149	5.03	28.52	17650	152300

191
Figure 3. **(a)** AOSLO image of the fovea one subject (10003L). Only the central 1.5 degrees are
shown here (810 X 810 pixels), which contains 16,184 cones. The white dots are a scatter plot
showing the PRL, or position of the fixated stimulus over the course of a 10-second video. The red
dot is the centroid of the scatter plot. **(b)** Same image with a color overlay indicating the density.
Linear and angular cone densities are indicated on the right colorbar. Peak cones densities in this
eye are 200,482 cones/mm^2 and 15,584 cones/deg^2. The yellow ellipse is the best fitting ellipse
containing ~ 68% of the points in the scatterplot and indicates the PRL. The black cross indicates
the position of peak cone density. Scale bar is 0.5 degrees, which in this eye corresponds to 139.4
microns.
201

202
203 **Figure 4.** Cone density as a function of eccentricity for all eyes. The axial length ranges of the
204 subjects are color coded, with warmer colors for shorter eyes and cooler colors for longer eyes. In
205 this plot, it is apparent that shorter eyes generally have higher peak cone densities.
206

In order to show the trends of density with axial length **Figure 5a&b** plot linear and angular
cone density as a function of axial length where the colors indicate different eccentricity - red to
purple indicate distance from the from fovea towards more parafoveal locations. **Figure 5a** reveals
that peak linear density decreases significantly with axial length and the trend persists and remains
significant from the fovea out to 100 microns eccentricity. Axial length accounts for 38% of the
variance in the changes in linear cone density. **Figure 5b** shows the opposite trends when plotted
in angular units. Peak angular density increases significantly with axial length and the trend
persists and remains significant out to 40 arcminutes eccentricity. Axial length accounts for 32%
of the variance in the changes in angular cone density. The plots clearly indicate that although
stretching does occur (**Figure 5a**) it is not a simple global expansion and longer eyes have higher
sampling density. The trends hold at and around fovea with statistical significance.

218
Figure 5. (a) Linear cone densities as a function of axial length. Longer eyes have lower linear
cone density than shorter eyes. The trend remains significant out to 100 microns eccentricity and
(b) Angular cone densities as a function of axial length. The peak angular cone density increases

significantly with increasing axial length and this trend remains significant out to 40 arcminutes
eccentricity. Relationships with P-values <0.05 are labelled with asterisks and trendlines are shown
as solid lines. Relationships with P-values ≥0.05 have dashed trendlines.
225
A more relevant measure of the impact of eye length on vision is how the angular cone
density changes at the PRL, which is often displaced from the location of peak cone density (Li et
al., 2010; Putnam et al., 2005; Wilk et al., 2017). If, for example, longer eyes had more displaced
PRLs then that could diminish, or even reverse, the trend of increased angular density with eye
length reported in **Figure 5b**. We found that the average displacement between PRL and maximum
cone density was 5.82 arcminutes and 28.94 microns. There was no significant linear relationship
found between PRL displacement in either angular or linear units vs. axial length. Therefore, the
PRL was not more displaced in myopes than in emmetropes from the point of peak cone density.
Plots of the cone density at the PRL with axial length show the same trend at the PRL as at the
point of maximum cone density (**Figure 6 a&b**).

236
237   **Figure 6 ab.** The relationship between cone density and axial length shows the same pattern at
238   the PRL as for the peak cone density. The slopes in both (a) and (b) are significant (P=0.00975 &
239   P=0.00432 respectively) and axial length accounts for 23% and 27% of the variance in linear and
240   angular cone density, respectively.
241

Finally, we explored whether fixational eye movements might have a dependency on axial
length. Fixation stability around the PRL had an average standard deviation of 3.94 arcminutes
and 19.84 microns. The average area of the best fitting ellipse containing ~ 68% of the points in
the scatterplot (defined as the bivariate contour ellipse area, or BCEA) was 50.7 square arcminutes
and 1303 square microns. The plot of BCEA in square microns v.s. axial length v.s. showed a trend
that approached significance (P=0.0596) (**Figure 7a**), but when we plotted BCEA in square
arcminutes v.s. axial length, the trend was no longer apparent (P=0.364) (**Figure 7b**). In other
words, if there is any increase in fixational eye movements in microns, it is just a symptom of
having a longer eye.

251
252 **Figure 7. (a)** The plot of BCEA in linear units (square microns) v.s. axial length shows a trend
253 that approaches significance (P=0.0596) **(b)** There is no significant relationship between BCEA in
254 angular units (square arcminutes) and axial length (P=0.364).

255
256 **Discussion**

257
In this paper we measure the cone density at and near the foveal center and investigate how
it changes as a function of axial length. This is the first comprehensive study of cones in living
eyes at the foveal center, the area solely responsible for a human's fine spatial vision. Our results

show that although some expansion does occur (linear cone density decreases with axial length)
the angular sampling resolution actually increases, on average, with axial length. Prior to this study,
the relationships between cone density and axial length were only made outside of the fovea, the
closest being 0.1 mm, or 0.3 degrees (Li et al., 2010). Although an eccentricity of 0.3 deg might
seem close, it is noted that the cone density drops precipitously just outside of the location of peak
density (Curcio, Sloan, Kalina, & Hendrickson, 1990) as does human vision (Poletti, Listorti, &
Rucci, 2013)(Rossi & Roorda, 2010b). There are other factors that govern peak cone density,
however; eye length accounts for anywhere between 27% and 38% of the variance in cone density.
Our finding that the slopes of cone density vs. axial length are in opposite directions when
plotted in linear (negative slope) and angular (positive slope) units, supports an eye growth model
that lies between the global expansion model and an equatorial stretching model. Previous studies
from our lab (Li et al., 2010) and also from Chui et al. (2008) leaned in the same direction. None
of the cone density studies provide insight into the reasons why the photoreceptor density would
behave this way with eye growth, but the results do align with other observations reported in the
literature. Specifically, Atchison et al. (2004) used magnetic resonance imaging and found that
eyeball dimensions in axial myopes are variable but are generally larger in all directions with a
weak tendency to be preferentially greater in the axial direction. Their reported eye growth patterns
lie between that illustrated for the global expansion and equatorial stretching models in **figure 1**.
Our results differ from Wilk et al. (2017) whose data support a global expansion model (i.e.
there is no detectable change in angular cone density with axial length; **figure 2b**). But it is
important to point out that their study did not set out to address the same question and the number
of subjects with long axial lengths was disproportionately low.
Our results also differ from Troilo (1998) who studied retinal cell topography in a
marmoset animal myopia model. Higher cone packing densities were observed in the
experimentally enlarged eyes compared to normal eyes in the fovea. Their result followed the
overdevelopment model, which is the reason why we included it as one of the possible outcomes
of our study. In fact, the overdevelopment model is an extension of Springer's model of
development (Springer & Hendrickson, 2004), which offers a biomechanical explanation for how
cone packing increases at the foveal center in a developing eye. While our data do not support the
overdevelopment model, it does not preclude the existence of biomechanical factors working in
opposition to simple global expansion.
The fact that angular cone density (visual sampling resolution) increases with eye length
(myopia), at the peak density and at the PRL, means that poorer performance by myopes on
resolution tasks cannot be explained by a decrease in photoreceptor sampling. The deficit musts
arise at a post-receptoral level.
Low-level causes for myopic visual deficits might arise from differences in the
connectivity between cones and ganglion cells. Atchison et al. (2006) suggested that abnormal eye
growth may be associated with a loss of ganglion cells. Alternately, if ganglion cells pool signals
from multiple cones, then they will impose the retinal sampling limit and reduce certain aspects of
visual performance (acuity, for example). Recent electron microscopy studies of a human fovea
have revealed extensive convergence and divergence connections between photoreceptors and
ganglion cells, albeit in an eye from an individual who was born prematurely (Dacey, 2018). These
discoveries challenge our current understanding of neural connectivity in the foveal center and
force us to consider the possibility of interindividual differences in foveal cone wiring. More
experiments are necessary to explore these ideas.

To explain why low myopes did not perform as well on an acuity task as emmetropes, even
after correction or bypassing of high order aberrations, Rossi et al. (2007) and Coletta & Watson
(2006) both raised the possibility that myopes might have become desensitized to high frequency
information (low level myopic amblyopia) as a result of having less exposure to a high contrast
visual environment. In this case, it might be possible to train myopes to take advantage of their
higher sampling resolution, but one myope in a follow up study by Rossi & Roorda (2010a) never
reached the acuity levels of emmetropes in the same study.
313
314　***Comparisons with Previous Studies***
315
Peak cone densities: Curcio et al. 1990 measured spatial density of cones and rods in eight
explanted whole-mounted human retinas. They found a large range of peak foveal cone densities
with an average of 199,000 cones/mm^2. When we averaged the peak cone density over a circular
aperture of 7.5 arcminutes which was similar to the 29 x 45 micron window that Curcio et al. (1990)
used to compute density, we measured peak linear cone densities ranging from 123,611 to 214,895
with an average of 168,047 cones/mm^2. Zhang et al. (2015) reported an average peak density of
168,162 cones/mm^2 in 40 eyes although they used a much smaller 5 x 5 micron sampling window
to measure the peak. Wilk et al. (2017) reported an average peak density of 145,900 cones/mm^2 in
22 eyes using a 37 x 37 micron sampling window and Li et al. (2010) reported an average peak
density of 150,412 cones/mm^2 in 4 eyes over a sampling window encompassing 150 cones
(approximately 37 micron diameter at the foveal center). All reports of cone densities from
adaptive optics studies in living eyes are lower than reports from histology. Two possible reasons
for this are (i) the excised tissue in Curcio et al. (1990) underwent more shrinkage than estimated
or (ii) the adaptive optics reports are subject to selection bias, where individuals with the highest
angular cone densities might have been excluded because the image were less well resolved
rendering the cones images too difficult to label with confidence. In our study, we attempted to
image 73 eyes from 46 subjects and only succeeded in resolving cones across a sufficiently large
region at and around the fovea in 28 of them. The reason the images from 45 eyes were not
analyzed was due to poor or inconsistent image quality arising from a number of factors: Images
from 4 eyes (3 subjects) were not analyzed because their refractive errors were too high (all above
–8D) and we ran into the limits of the deformable mirror's dynamic range. Images from 18 eyes
(13 subjects) that were taken early on in the study were not analyzed because the optics of AOSLO
were not tuned well enough to resolve foveal cones. Images from 4 eyes (2 subjects) were not
analyzed because of uncorrectable image degradation caused by keratoconus and corneal scarring.
Images from 2 eyes (1 subject) were not analyzed because of excessive aberrations caused by an
orthokeratology refractive correction. The cause of poor or inconsistent image quality among the
remaining 17 eyes were varied, including ocular surface dryness, excessive eye motion and small
pupils. The average refractive error among these remaining 17 eyes was about the same as the
successful eyes.
Anisotropic density distribution: Like Curcio et al. (1990) and Zhang et al. (2015) we
found steeper drops in cone density in the superior and inferior directions compared to the nasal
and temporal directions. Plots of density along the two cardinal directions are shown on
Supplemental Figure 3.
PRL displacements: The distance of the PRL from the foveal center for our study (mean
29 microns; range 8 – 73; n = 28) roughly agrees with those of Wilk et al. (2017) (mean 63 microns;
range 20 – 263; n = 22), Li et al. (2010) (mean 34 microns; range 3 – 92; n = 18) and Putnam et

al. (2005) (mean 17; range 11 – 23; n = 5). The differences in cone density between the peak and
the PRL were small and the trends (**Figures 5 and 6**) persisted at both locations.
Spatial vision estimates: The cone array imposes the first retinal sampling limit to human
spatial vision (MacLeod, Williams, & Makous, 1992; Williams, 1985) and the photoreceptor row-
to-row spacing (assuming an hexagonal packing structure) imposes the maximum frequencies that
can be relayed to later stages without aliasing. We can compute the sampling limit and estimate
the cone center-to-center spacing using the following formulas:
359

$$Sampling\ Limit = \frac{1}{2}\sqrt{\frac{2}{\sqrt{3}}\ AngularDensity}$$

$$Cone\ Spacing = Sampling\ Limit^{-1} \times 60 \times \frac{1}{\sqrt{3}}$$

For the densities reported here, the potential spatial frequency resolution limits range from 58.3 to
73.6 cyc/deg (average: 64.5 cyc/deg) at the peak density and 50.9 to 71.4 cyc/deg (average: 62.7
cyc/deg) at the PRL. These correspond to potential acuities ranging from 20/11.8 to 20/8.2 (based
on the primary spatial frequency of the three bars of a Snellen E). The cone frequency cut-offs are
higher than almost all the interferometric acuity limits reported by Coletta & Watson (2006), even
for the emmetropic subjects. The acuities are, however, in the range of those measured from
emmetropic subjects after adaptive optics correction by Rossi et al. (2007). The cone center-to-
center spacing ranges from 0.59 to 0.47 arcminutes at the peak density and from 0.60 to 0.49
arcminutes at the PRL. A direct comparison of foveal structure and function for each of our
subjects was not the scope of this study but will be the topic of future investigation.
374

Measuring structure and function of cone photoreceptors at the foveal center – the most
important region of the human retina – has been one of the more challenging endeavors in vision
science. Fortunately, the latest generation of adaptive optics ophthalmoscopes are making it easier
and are facilitating new discoveries within this retinal region. The pattern of how cone density
changes with eye growth lands somewhere between the global expansion and equatorial stretching
models. The cone mosaic in longer eyes is expanded at the fovea, but not in proportion to eye
length. Despite retinal stretching, myopes generally have a higher angular sampling density in and
around the fovea compared to emmetropes. Reports of reduced best-corrected central visual acuity
in myopes compared to emmetropes cannot be explained by decreased photoreceptor density
caused by retinal stretching during myopic progression.
385

386 **Materials and Methods**
387

Foveal Imaging
We used our latest generation adaptive optics scanning laser ophthalmoscope (AOSLO)
for foveal imaging. The system used a mirror-based, out-of-plane optical design (Dubra et al.,
2011), and employed a deformable mirror with a continuous membrane surface and shaped with
97 actuators (DM97, ALPAO, Montbonnot-Saint-Martin, France). The system scans multiple
wavelengths simultaneously. Each wavelength was drawn from the same broadband
supercontinuum source (SuperK EXTREME, NKT Photonics, Birkerod, Denmark) using a

custom-built fiber coupler. Wave aberrations were measured with a custom-built Shack Hartmann
wavefront sensor using the 940 nm channel. Images were recorded using the 680 nm channel. 512
x 512 pixel videos were recorded over a 0.9 x 0.9 degree square field for an average sampling
resolution of 9.48 pixels per arcminute. Eye alignment and head stabilization was achieved by
using either a bite bar or a chin rest with temple pads. At least one 10-second video was recorded
at the fovea and at 8 more locations where the subjects were instructed to fixate on the corners and
sides of the raster, to image an entire foveal region spanning about 1.8 X 1.8 degrees. In order to
ensure the best possible focus of the foveal cones, multiple videos were taken over a range of 0.05
D defocus steps to find the sharpest foveal cones. Focus steps were generated by adding a focus
shape onto the deformable mirror. Online stabilization and registration algorithms were used to
facilitate rapid feedback on the image quality.
406
Locating the Preferred Retinal Locus of Fixation (PRL)
Steady fixation was achieved at the fovea center by having the subjects fixate on a dark,
circular, blinking dot with a diameter of 3.16 arcminutes (30 pixels) in the center of the AOSLO
scanning raster. The fixation target was generated by modulating the same 680 nm scanning beam
used for imaging and, as such, the target's location was encoded directly into each frame of the
video (Poonja, Patel, Henry, & Roorda, 2005). A scatter plot of the positions of the blinking dot
relative to the retina was generated and was fit with a bivariate ellipse using free online Matlab
scripts downloaded from [http://www.visiondummy.com/wp-](http://www.visiondummy.com/wp-content/uploads/2014/04/error_ellipse.m)
[content/uploads/2014/04/error_ellipse.m](http://www.visiondummy.com/wp-content/uploads/2014/04/error_ellipse.m). The bivariate contour ellipse area (BCEA), which is the
area of the best-fitting ellipse encompassing 68% of the points in the scatterplot (Castet &
Crossland, 2012) was used to quantify the fixation stability (**figure 7**) and the exact location of the
PRL within the imaged cone mosaic (**Table 2, Figure 3, Supplemental figures 1 and 2**).
419
Image Processing and Analysis
High quality images were generated from the recorded videos offline using custom
software (Matlab, The MathWorks, Inc., Natick, MA) to measure and correct for distortions caused
by eye movements (Stevenson & Roorda, 2005). Poor-quality frames were manually excluded and
registered frames were averaged into a single high signal-to-noise image. The processed images
were stitched together (Photoshop; Adobe Systems, Inc., Mountain View, CA) to create an
approximately 1.8-degree montage of the foveal cone mosaic.
We used custom software to identify and label individual cones in the AO retinal images.
The program allows the user to select a region of interest and manually add and delete cone labels.
A combination of both manual and automated methods (Li & Roorda, 2007) were used to identify
cone locations as the current version of the program does not adequately recognize cones in the
foveal center where they are dim and smaller (Li et al., 2010). All the cone coordinates were
selected and reviewed by two of the authors. In some cases cones were too dim to be seen but there
was only a gap in the mosaic (Bruce et al., 2015). If a space that might have been occupied by a
cone was dim or dark, we would assume it was a cone and mark its location. We rationalize this
for two reasons: First, if there is a gap in the mosaic, then it is likely that a cell is occupying that
space, otherwise the adjacent cells would migrate to fill it in (Scoles et al., 2014). Second, in our
experience and of others (Pallikaris, Williams, & Hofer, 2003), cones that appear dark in one visit,
can often appear bright in the next. In other cases (uncommon) the contrast was low in some
regions or there were interference artifacts in the images (Meadway & Sincich, 2018; Putnam,
Hammer, Zhang, Merino, & Roorda, 2010), making the cone locations slightly ambiguous. In these

instances, we made manual cone selections based on the assumption that the cones were all similar
in size and close-packed into a nearly hexagonal array (Curcio et.al., 1990).
Continuous density maps were generated by computing cone density within a circle of 10
arcminutes in diameter around every pixel location across the image. We kept the area large
enough to generate smooth maps, but small enough to resolve local changes. Changes in density
with eccentricity were generated by computing the density in 5 arcminute annuli surrounding the
point of peak cone density. For linear density measures we used annuli with 25 micron widths.
448
Retinal Magnification Factor Calculation
The exact angular dimensions of the AOSLO images were computed by imaging a
calibrated model eye in the AOSLO system, but the conversion to linear dimensions on the retinal
image requires additional measurements, since the dimensions of each eye governs the actual size
of the image on its retina. The conversion from visual angle to retinal distance requires a
measurement of the axial length of the eye and an estimation of the location of the secondary nodal
point. We used a four-surface schematic eye model, originally proposed by Li et al., 2010 to
estimate the location of the secondary nodal point. The corneal first surface radius of curvature,
the anterior chamber depth and the axial length were for measured for each subject with an IOL
Master (Zeiss Meditec, Dublin, CA). The radius of the curvature of the back surface of the cornea
was computed as 88.31% of the front surface (Bennett, Rudnicka, & Edgar, 1994). The indices of
refraction of the media and the radii of curvature of the front and back lens surface were taken
from the Gullstrand schematic eye (Vojnikovic & Tamajo, 2013). Once determined, retinal image
size is related to visual angle by the equation:
463
464 $$I = \tan(1°)(x - AN')\theta$$
465
466 Where I is retinal image size, x is axial length, AN' is the distance from the corneal apex to the
467 eye's second nodal point, and θ is the visual angle. As can be seen in **Table 2**, myopic eyes, which
468 generally have longer focal lengths, have proportionally larger retinal images.
469
Statistical Analysis
Given the trends of increased angular density as a function of axial length that Li et al
(2011) observed at the location closest to the fovea (slope = 531 cones/deg^2 for each mm increase
in axial length; standard deviation of the regression errors = 1377 cones/deg^2), we estimated that
data from approximately 32 eyes, evenly distributed across a range of axial lengths would be
sufficient to show if there was a true effect at the fovea. The targeted number was computed using
methods outlined by Dupont & Plummer (1998) implemented using free online software (Power
and Sample Size Program Version 3.0, January 2009, downloaded from
http://biostat.mc.vanderbilt.edu/wiki/Main/PowerSampleSize) with type 1 error probability of
0.05 and a power of 0.95.
All data collected in this study were analyzed using simple linear regression models in
Excel. P-values for all linear regressions are reported and linear trendlines with P-values less than
0.05 are plotted as solid lines and P-values greater than 0.05 as dashed lines.
483
484
485 **Competing interests:**
486

A.R. has a patent (USPTO#7118216) assigned to the University of Houston and the University of Rochester which is currently licensed to Boston Micromachines Corp (Watertown, MA, USA). Both he and the company stand to gain financially from the publication of these results. No other authors have competing interests.

References

Atchison, D. A., Jones, C. E., Schmid, K. L., Pritchard, N., Pope, J. M., Strugnell, W. E., & Riley, R. A. (2004). Eye shape in emmetropia and myopia. *Investigative Ophthalmology and Visual Science*, *45*(10), 3380–3386. https://doi.org/10.1167/iovs.04-0292

Atchison, D. A., Schmid, K. L., & Pritchard, N. (2006). Neural and optical limits to visual performance in myopia. *Vision Research*, *46*(21), 3707–3722. https://doi.org/10.1016/j.visres.2006.05.005

Bennett, A. G., Rudnicka, A. R., & Edgar, D. F. (1994). Improvements on Littmann's method of determining the size of retinal features by fundus photography. *Graefe's Archive for Clinical and Experimental Ophthalmology*, *232*(6), 361–367. https://doi.org/10.1007/BF00175988

Bruce, K. S., Harmening, W. M., Langston, B. R., Tuten, W. S., Roorda, A., & Sincich, L. C. (2015). Normal perceptual sensitivity arising from weakly reflective cone photoreceptors. *Investigative Ophthalmology and Visual Science*, *56*(8), 4431–4438. https://doi.org/10.1167/iovs.15-16547

Castet, E., & Crossland, M. (2012). Quantifying Eye Stability During a Fixation Task: A Review of Definitions and Methods. *Seeing and Perceiving*, *25*(5), 449–469. https://doi.org/10.1163/187847611x620955

Chui, T. Y. P., Song, H., & Burns, S. A. (2008). Individual Variations in Human Cone Photoreceptor Packing Density: Variations with Refractive Error. *Investigative Ophthalmology and Visual Science*, *49*(10), 4679–4687. https://doi.org/10.1167/iovs.08-2135

Chui, T. Y., Yap, M. K., Chan, H. H., & Thibos, L. N. (2005). Retinal stretching limits peripheral visual acuity in myopia. *Vision Research*, *45*(5), 593–605. https://doi.org/10.1016/j.visres.2004.09.016

Coletta, N. J., & Watson, T. (2006). Effect of myopia on visual acuity measured with laser interference fringes. *Vision Research*, *46*(5), 636–651. https://doi.org/S0042-6989(05)00268-3 [pii]\r10.1016/j.visres.2005.05.025

Collins, J. W., & Carney, L. G. (1990). Visual performance in high myopia. *Current Eye Research*, *9*(3), 217–224. https://doi.org/10.3109/02713689009044516

Curcio, C. A., Sloan, K. R., Kalina, R. E., & Hendrickson, A. E. (1990). Human photoreceptor topography. *Journal of Comparative Neurology*, *292*(4), 497–523. https://doi.org/10.1002/cne.902920402

Dacey, D. M. (2018). Discovering visual pathway origins in the center of the human foveola with connectomics. *Investigative Ophthalmology and Visual Science*, *59*(9), ARVO Abstract #14.

Dubra, A., Sulai, Y., Norris, J. L., Cooper, R. F., Dubis, A. M., Williams, D. R., & Carroll, J. (2011). Noninvasive imaging of the human rod photoreceptor mosaic using a confocal

adaptive optics scanning ophthalmoscope. *Biomedical Optics Express*, *2*(7), 1864–1876. https://doi.org/10.1364/BOE.2.001864

Dupont, W. D., & Plummer, W. D. (1998). Power and sample size calculations for studies involving linear regression. *Controlled Clinical Trials*, *19*(6), 589–601. https://doi.org/10.1016/S0197-2456(98)00037-3

Ehsaei, A., Chisholm, C. M., Pacey, I. E., & Mallen, E. A. H. (2013). Visual performance fall-off with eccentricity in myopes versus emmetropes. *Journal of Optometry*, *6*(1), 36–44. https://doi.org/10.1016/j.optom.2012.07.001

Elsner, A. E., Chui, T. Y. P., Feng, L., Song, H. X., Papay, J. A., & Burns, S. A. (2017). Distribution differences of macular cones measured by AOSLO: Variation in slope from fovea to periphery more pronounced than differences in total cones. *Vision Research*, *132*, 62–68. https://doi.org/10.1016/j.visres.2016.06.015

Fiorentini, A., & Maffei, L. (1976). Spatial contrast sensitivity of myopic subjects. *Vision Research*, *16*(4), 437–438. https://doi.org/10.1016/0042-6989(76)90214-5

Fujiwara, T., Imamura, Y., Margolis, R., Slakter, J. S., & Spaide, R. F. (2009). Enhanced Depth Imaging Optical Coherence Tomography of the Choroid in Highly Myopic Eyes. *American Journal of Ophthalmology*, *148*(3), 445–450. https://doi.org/10.1016/j.ajo.2009.04.029

Gonzalez Blanco, F., Sanz Fernández, J. C., & Muńoz Sanz, M. A. (2008). Axial length, corneal radius, and age of myopia onset. *Optometry and Vision Science*, *85*(2), 89–96. https://doi.org/10.1097/OPX.0b013e3181622602

Harb, E., Hyman, L., Gwiazda, J., Marsh-Tootle, W., Zhang, Q., Hou, W., … Taylor, C. (2015). Choroidal thickness profiles in myopic eyes of young adults in the correction of myopia evaluation trial cohort. *American Journal of Ophthalmology*, *160*(1), 62–71.e2. https://doi.org/10.1016/j.ajo.2015.04.018

He, M., Zeng, J., Liu, Y., Xu, J., Pokharel, G. P., & Ellwein, L. B. (2004). Refractive error and visual impairment in urban children in southern china. *Investigative Ophthalmology and Visual Science*, *45*(3), 793–799. https://doi.org/10.1167/iovs.03-1051

He, X., Zou, H., Lu, L., Zhao, R., Zhao, H., Li, Q., & Zhu, J. (2015). Axial length/corneal radius ratio: Association with refractive state and role on myopia detection combined with visual acuity in Chinese schoolchildren. *PLoS ONE*, *10*(2), e0111766. https://doi.org/10.1371/journal.pone.0111766

Iyamu, E., Iyamu, J., & Obiakor, C. I. (2011). The Role of Axial Length-Corneal Radius of Curvature Ratio in Refractive State Categorization in a Nigerian Population. *ISRN Ophthalmology*, *2011*, 1–6. https://doi.org/10.5402/2011/138941

Jaworski, A., Gentle, A., Zele, A. J., Vingrys, A. J., & McBrien, N. A. (2006). Altered visual sensitivity in axial high myopia: A local postreceptoral phenomenon? *Investigative Ophthalmology and Visual Science*, *47*(8), 3695–3702. https://doi.org/10.1167/iovs.05-1569

Jong, M., Sankaridurg, P., Li, W., Resnikoff, S., Naidoo, K., & He, M. (2018). Reduced vision in highly myopic eyes without ocular pathology: the ZOC-BHVI high myopia study. *Clinical and Experimental Optometry*, *101*(1), 77–83. https://doi.org/10.1111/cxo.12563

Kitaguchi, Y., Bessho, K., Yamaguchi, T., Nakazawa, N., Mihashi, T., & Fujikado, T. (2007). In vivo measurements of cone photoreceptor spacing in myopic eyes from images obtained by an adaptive optics fundus camera. *Japanese Journal of Ophthalmology*, *51*(6), 456–461. https://doi.org/10.1007/s10384-007-0477-7

Li, K. Y., & Roorda, A. (2007). Automated identification of cone photoreceptors in adaptive optics retinal images. *Journal of the Optical Society of America A*, *24*(5), 1358–1363.

https://doi.org/10.1364/JOSAA.24.001358

Li, K. Y., Tiruveedhula, P., & Roorda, A. (2010). Intersubject variability of foveal cone photoreceptor density in relation to eye length. *Investigative Ophthalmology and Visual Science*, *51*(12), 6858–6867. https://doi.org/10.1167/iovs.10-5499

Liang, J., Williams, D. R., & Miller, D. T. (1997). Supernormal vision and high-resolution retinal imaging through adaptive optics. *Journal of the Optical Society of America A*, *14*(11), 2884–2892. https://doi.org/10.1364/JOSAA.14.002884

Liou, S.-W., & Chiu, C.-J. (2001). Myopia and contrast sensitivity function. *Current Eye Research*, *22*(2), 81–84. https://doi.org/10.1076/ceyr.22.2.81.5530

MacLeod, D. I. A., Williams, D. R., & Makous, W. (1992). A visual nonlinearity fed by single cones. *Vision Research*, *32*, 347–363. https://doi.org/10.1016/0042-6989(92)90144-8

Meadway, A., & Sincich, L. C. (2018). Light propagation and capture in cone photoreceptors. *Biomedical Optics Express*, *9*(11), 5543–5565. https://doi.org/10.1364/boe.9.005543

Morgan, I. G., Ohno-Matsui, K., & Saw, S. M. (2012). Myopia. *The Lancet*, *379*(9827), 1739–1748. https://doi.org/10.1016/S0140-6736(12)60272-4

Obata, R., & Yanagi, Y. (2014). QuantItative analysis of cone photoreceptor distribution and its relationship with axial length, age, and early age-related macular degeneration. *PLoS ONE*, *9*(3), e91873. https://doi.org/10.1371/journal.pone.0091873

Pallikaris, A., Williams, D. R., & Hofer, H. (2003). The reflectance of single cones in the living human eye. *Investigative Ophthalmology and Visual Science*, *44*(10), 4580–4592. https://doi.org/10.1167/iovs.03-0094

Park, S. P., Chung, J. K., Greenstein, V., Tsang, S. H., & Chang, S. (2013). A study of factors affecting the human cone photoreceptor density measured by adaptive optics scanning laser ophthalmoscope. *Experimental Eye Research*, *108*, 1–9. https://doi.org/10.1016/j.exer.2012.12.011

Poletti, M., Listorti, C., & Rucci, M. (2013). Microscopic eye movements compensate for nonhomogeneous vision within the fovea. *Current Biology*, *23*(17), 1691–1695. https://doi.org/10.1016/j.cub.2013.07.007

Poonja, S., Patel, S., Henry, L., & Roorda, A. (2005). Dynamic visual stimulus presentation in an adaptive optics scanning laser ophthalmoscope. *Journal of Refractive Surgery*, *21*(5), S575–S580. https://doi.org/10.1364/fio.2004.fthv3

Putnam, N. M., Hammer, D. X., Zhang, Y., Merino, D., & Roorda, A. (2010). Modeling the foveal cone mosaic imaged with adaptive optics scanning laser ophthalmoscopy. *Optics Express*, *18*(24), 24902–24916. https://doi.org/10.1364/OE.18.024902

Putnam, N. M., Hofer, H., Doble, N., Chen, L., Carroll, J., & Williams, D. R. (2005). The locus of fixation and the foveal cone mosaic. *Journal of Vision*, *5*(7), 632–639. https://doi.org/10.1167/5.7.3

Roorda, A., Romero-Borja, F., Donnelly III, W. J., Queener, H., Hebert, T. J., & Campbell, M. C. W. (2002). Adaptive optics scanning laser ophthalmoscopy. *Optics Express*, *10*(9), 405–412. https://doi.org/10.1364/OE.10.000405

Rossi, E. A., & Roorda, A. (2010a). Is visual resolution after adaptive optics correction susceptible to perceptual learning? *Journal of Vision*, *10*(12), 1–11. https://doi.org/10.1167/10.12.11

Rossi, E. A., & Roorda, A. (2010b). The relationship between visual resolution and cone spacing in the human fovea. *Nature Neuroscience*, *13*(2), 156–157. https://doi.org/10.1038/nn.2465

Rossi, E. A., Weiser, P., Tarrant, J., & Roorda, A. (2007). Visual performance in emmetropia

and low myopia after correction of high-order aberrations. *Journal of Vision*, *7*(8), 1–14.
https://doi.org/10.1167/7.8.14
Scoles, D., Sulai, Y. N., Langlo, C. S., Fishman, G. A., Curcio, C. A., Carroll, J., & Dubra, A.
(2014). In vivo imaging of human cone photoreceptor inner segments. *Investigative*
Ophthalmology and Visual Science, *55*(7), 4244–4251. https://doi.org/10.1167/iovs.14-
14542
Springer, A. D., & Hendrickson, A. E. (2004). Development of the primate area of high acuity. 1.
Use of finite element analysis models to identify mechanical variables affecting pit
formation. *Visual Neuroscience*, *21*(1), 53–62. https://doi.org/10.1017/S0952523804041057
Stevenson, S. B., & Roorda, A. (2005). Correcting for miniature eye movements in high
resolution scanning laser ophthalmoscopy. *Ophthalmic Technologies XV*, *5688*, 145–151.
https://doi.org/10.1117/12.591190
Stoimenova, B. D. (2007). The Effect of Myopia on Contrast Thresholds. *Investigative*
Opthalmology and Visual Science, *48*(5), 2371–2734. https://doi.org/10.1167/iovs.05-1377
Strang, N. C., Winn, B., & Bradley, A. (1998). The role of neural and optical factors in limiting
visual resolution in myopia. *Vision Research*, *38*(11), 1713–1721.
https://doi.org/10.1016/S0042-6989(97)00303-9
Thorn, F., Corwin, T. R., & Comerford, J. P. (1986). High myopia does not affect contrast
sensitivity. *Current Eye Research*, *5*(9), 635–639.
https://doi.org/https://doi.org/10.3109/02713688609015130
Troilo, D. (1998). Changes in retinal morphology following experimentally induced myopia.
Vision Science and Its Applications: Technical Digest (OSA, Washington, D.C.), *1*(1), 206–
209.
Verkicharla, P. K., Ohno-Matsui, K., & Saw, S. M. (2015). Current and predicted demographics
of high myopia and an update of its associated pathological changes. *Ophthalmic and*
Physiological Optics, *35*(5), 465–475. https://doi.org/10.1111/opo.12238
Vitale, S., Sperduto, R. D., & Ferris, F. L. (2009). Increased prevalence of myopia in the United
States between 1971-1972 and 1999-2004. *Archives of Ophthalmology*, *127*(12), 1632–
1639. https://doi.org/10.1001/archophthalmol.2009.303
Vojnikovic, B., & Tamajo, E. (2013). Gullstrand's optical schematic system of the eye--modified
by Vojnikovic & Tamajo. *Coll Antropol*, *37 Suppl 1*, 41–45. Retrieved from
http://www.ncbi.nlm.nih.gov/pubmed/23841130
Webb, R. H., Hughes, G. W., & Delori, F. C. (1987). Confocal scanning laser ophthalmoscope.
Applied Optics, *26*(8), 1492–1499. https://doi.org/10.1364/AO.26.001492
Wilk, M. A., Dubis, A. M., Cooper, R. F., Summerfelt, P., Dubra, A., & Carroll, J. (2017).
Assessing the spatial relationship between fixation and foveal specializations. *Vision*
Research, *132*, 53–61. https://doi.org/10.1016/j.visres.2016.05.001
Williams, D. R. (1985). Aliasing in human foveal vision. *Vision Research*, *25*, 195–205.
https://doi.org/10.1016/0042-6989(85)90113-0
Zhang, T., Godara, P., Blanco, E. R., Griffin, R. L., Wang, X., Curcio, C. A., & Zhang, Y.
(2015). Variability in Human Cone Topography Assessed by Adaptive Optics Scanning
Laser Ophthalmoscopy. *American Journal of Ophthalmology*, *160*(2), 290–300 e1.
https://doi.org/10.1016/j.ajo.2015.04.034
668

Supplemental Figure 1: Linear cone density (cones/mm^2) plots over the central 450 microns for all 28 eyes. The black cross indicates the point of maximum cone density. The black ellipse is the best fitting ellipse about the fixation scatterplot indicating the PRL. Dark blue regions indicate where no cone density estimates were made.

1.5 deg

675
676 **Supplemental Figure 2:** Angular cone density (cones/deg^2) plots over the central 1.5 degrees for
677 all 28 eyes. The black cross indicates the point of maximum cone density. The black ellipse is the
678 best fitting ellipse about the fixation scatterplot indicating the PRL. Dark blue regions indicate
679 where no cone density estimates were made.
680

681
682 **Supplemental Figure 3:** Plots of density as a function of eccentricity in the vertical and horizontal
683 directions. (A) linear cone density (B) angular cone density. The dashed lines represent +/- 1
684 standard deviation from the mean.