1 **Human Foveal Cone Photoreceptor Topography and its Dependence on Eye Length**
2

3 **Authors:**
4 Wang, Yiyi[1]; Bensaid, Nicolas[3]; Tiruveedhula, Pavan[1,2]; Ma, Jianqiang[4]; Ravikumar, Sowmya[1,2];
5 Roorda, Austin[1,2]
6

7 **Affiliations:**
8   1. School of Optometry, University of California, Berkeley, Berkeley, CA, USA
9   2. Vision Science Graduate Group, University of California, Berkeley, Berkeley, CA, USA
10   3. Carl Zeiss Meditec AG, Berlin Germany
11   4. Department of Mechanical Engineering, Ningbo University, Ningbo, Zhejiang, China
12

Abstract:
We provide the first measures of foveal cone density as a function of axial length in living eyes
and discuss the physical and visual implications of our findings. We used a new generation
Adaptive Optics Scanning Laser Ophthalmoscope to image cones at and near the fovea in 28
eyes of 16 subjects. Cone density and other metrics were computed in units of visual angle and
linear retinal units. The foveal cone mosaic in longer eyes is expanded at the fovea, but not in
proportion to eye length. Despite retinal stretching (decrease in cones/mm^2), myopes generally
have a higher angular sampling density (increase in cones/deg^2) in and around the fovea
compared to emmetropes, offering the potential for better visual acuity. Reports of deficits in
best-corrected foveal vision in myopes compared to emmetropes cannot be explained by
increased spacing between photoreceptors caused by retinal stretching during myopic
progression.
28
29
30

Introduction

There has been a rapid increase in prevalence of myopia, of all magnitudes, in the period between 1971-1972 and 1999-2004 (Vitale, 2009). Across sub-populations grouped by race, ethnicity and gender, several studies report axial length of the eye to be the primary variable related to myopia (Gonzalez Blanco, Sanz Ferńandez, & Muńoz Sanz, 2008; X. He et al., 2015; Iyamu, Iyamu, & Obiakor, 2011). Increased axial length is associated with retinal stretching and thinning of posterior segment layers and the choroid (Fujiwara, Imamura, Margolis, Slakter, & Spaide, 2009; Harb et al., 2015) and is associated with sight-threatening, often irreversible pathologies of the retina (Morgan, Ohno-Matsui, & Saw, 2012; Verkicharla, Ohno-Matsui, & Saw, 2015). Even without any detectable pathology, the structural changes associated with eye growth ought to have functional consequences for vision.

What Do We Know About Functional Deficits in Myopia?

One might expect that eye growth would stretch the photoreceptor layer and would increase the spacing between cones, causing a longer eye to more coarsely sample an image relative to a shorter eye. However the situation is not that simple; the axial elongation associated with eye growth is accompanied by magnification of the retinal image (Strang, Winn, & Bradley, 1998). If the enlargement of the retinal image exactly matched the stretching of the cone mosaic, then eyes of different lengths would sample the visual field similarly. In fact, in large scale studies, myopes generally attain reasonably good visual acuity with optical correction (He et al., 2004; Jong et al., 2018).

However, more careful inspection reveals that that myopes generally (6 out of 9 studies) have poorer angular resolution and have uniformly (3 out of 3 studies) poorer retinal resolution. **Table 1** summarizes published results from psychophysical foveal tasks.

Table 1: Summary of studies investigating foveal spatial vision and sensitivity tasks in myopia.

Author	Refractive error range of myopic cohort [D]	Functional tests	Results for myopes at foveal center	Suggested cause
Fiorentini & Maffei, 1976	-5.5 to -10 (n=10)	CSF	Reduced CSF	Neural insensitivity (myopic amblyopia)
Thorn, Corwin, & Comerford, 1986	-6 to -9.75 (n=13)	CSF	No difference in CSF	Global expansion
Collins & Carney, 1990	-2 to -11 (n=16)	VA, CSF	No difference in VA or CSF between low and high myopic groups with contact lens correction	NA
Strang et al., 1998	0 to -14 (n=34)	VA	Reduced VA (MAR) with increasing myopia after controlling for spectacle magnification	Retinal expansion specifically at the posterior pole; increased aberrations
Liou & Chiu, 2001	0 to >-12 (n=105 eyes)	CSF	Reduced CSF with increasing myopia	Retinal stretching and disruption, neural insensitivity (myopic

				amblyopia)
Chui, Yap, Chan, & Thibos, 2005	-0.5 to -14 (n=60)	Grating resolution	Decreased resolution acuity in cyc/mm	Retinal expansion specifically at the posterior pole; global expansion along with ganglion cell loss
Coletta & Watson, 2006	+2 to -15 (n=17)	Interferometric grating resolution	Decreased resolution acuity in cyc/mm but not in cyc/deg	Retinal expansion specifically at the posterior pole
Atchison, Schmid, & Pritchard, 2006	+0.75 to -12.4 (n=121)	Spatial summation; interferometric grating resolution	Increased critical summation area in linear area, but not in angular area; Decreased resolution acuity in cyc/mm but not in cyc/deg	Retinal expansion specifically at the posterior pole; global expansion along with ganglion cell loss
Stoimenov, 2007	-1 to -8 (n=60)	Contrast thresholds of 20/120 letters	Lower sensitivity to contrast for letters with a fixed angular size	Morphologic changes in the retina
Rossi, Weiser, Tarrant, & Roorda, 2007	−0.5 to −3.75 (n=10)	AO-corrected VA	Reduced acuity (MAR) compared to emmetropes	Retinal expansion, neural insensitivity; neural insensitivity (myopic amblyopia)
Jaworski, Gentle, Zele, Vingrys, & McBrien, 2006	-8.5 to -11.5 (n=10)	Foveal summation thresholds; CSF	Increased critical summation area (angular) Decreased luminance sensitivity Reduced contrast sensitivity at high frequencies (cyc/deg)	Reduction in photoreceptor sensitivity; postreceptoral changes; increased aberrations
Ehsaei, Chisholm, Pacey, & Mallen, 2013	-2.00 to -9.62 (n=60)	Size threshold of high and low contrast letter targets	No difference in threshold retinal image size between myopes and emmetropes.	NA

Most notably, Atchison et al. (2006) and Coletta & Watson (2006) show clear deficits in
retinal resolution (cyc/mm) with increasing myopia using interferometric methods which bypass
the optics of the eye and Rossi et al. (2007) show significant deficits in angular resolution
(cyc/deg) in low myopes, even after using adaptive optics to correct for optical blur. All studies
the find myopic visual deficits implicate retinal stretching as a possible cause, but what is
actually happening structurally at the foveal center during myopic progression is not known.
Therefore, the aim of the current study is to more carefully investigate how the length of the eye
affects cone density at and near the foveal center.

Models for How Photoreceptors Change with Eye Growth

Two types of cone densities will be discussed in this study. Linear density quantifies how
many cones are within a fixed area, in square mm, and serves as a way to evaluate physical
retinal stretching caused by eye growth. Angular density quantifies how many cones are within
one degree visual angle, (the visual angle is measured from the secondary nodal point of the

eye). Angular density serves as a way to evaluate the visual implications of eye growth as it governs the sampling resolution of the eye.

Figure 1 illustrates three models, along the lines of Strang et al. (1998), of how photoreceptor structure might be affected by myopic eye growth. In the first model, called the **global expansion model**, the retina is proportionally stretched with increasing axial length - cones are more spaced out in longer eyes - and linear density decreases with eye length. Assuming that the secondary nodal point remains at a fixed position relative to the anterior segment, the number of cones within a fixed angular area will remain constant. Therefore, angular cone density will be constant with eye length. In the second model, called the **equatorial stretching model,** the posterior retina simply moves axially further from the anterior segment of the eye so that the linear density does not change with eye length. Since the retina is moving further from the secondary nodal point, more cones will fall within a fixed angular area and the angular cone density will increase with eye length. The final model, called the **over-development model**, describes a structural photoreceptor change that mimics the changes that occur during development (Springer & Hendrickson, 2004) whereby the photoreceptors continue to migrate towards the fovea as the eye grows. In this scenario, longer eyes will show both increased linear cone density and an even steeper increase in angular cone density. The model is motivated by observations of increased linear cone density in the foveas of marmosets that underwent lens-induced eye growth (Troilo, 1998).

Figure 1: 3 models of myopic eye growth: (A) Global expansion shows an eyeball that is proportionally stretched. (B) The equatorial stretching model indicates a growth model where the fovea stays rigid and unaffected as the eye grows. (C) The over-development model shows that myopic eye growth is similar with developmental eye growth where photoreceptors continue to migrate towards the fovea as the eye grows.

Previous Studies of Cone Spacing with Axial Length

The most definitive studies of cone spacing as a function of axial length are done through direct imaging of the retina – wherein sharp images of the cones are enabled through the use of adaptive optics, a set of technologies that actively compensate the blur caused by aberrations of the eye (Liang, Williams, & Miller, 1997). Combined with confocal scanning laser ophthalmoscopy (Webb, Hughes, & Delori, 1987), adaptive optics offers the highest contrast *en face* images of the foveal photoreceptor mosaic ever recorded in vivo (Dubra et al., 2011; Roorda et al., 2002).

Despite continued advances in image quality, previous studies investigating cone packing and eye length have not made their measurements at the foveal center, the most important region for spatial vision, but the most difficult to image, owing to the small size of photoreceptors. There are a number of studies on cone packing and eye length (Chui, Song, & Burns, 2008; Elsner et al., 2017; Kitaguchi et al., 2007; Li, Tiruveedhula, & Roorda, 2010; Obata & Yanagi, 2014; Park, Chung, Greenstein, Tsang, & Chang, 2013) and here we summarize the published results that are most relevant to our study. Chui et al. (2008) investigated angular and linear cone density at 1 mm and 3 degrees eccentricity. They found a significant decrease (P<0.05) in linear cone density as a function of eye length at 1mm (which, by angular distance, is closer to the fovea in a longer eye than in a shorter eye) in all directions except in the nasal retina. They found that the angular cone density at 3 degrees (which, by linear distance, is closer to the fovea, in a shorter eye than in a longer eye) increased with eye length, but the trends were not significant. Li, et al. (2010) made similar measures, but closer to the fovea (from 0.10 mm to 0.30 mm eccentricity). They found that linear cone density decreased with eye length, but the trends were not significant at the smallest eccentricities (0.1 and 0.2 mm). When the data were plotted in angular units and angular distance from the fovea, they found that angular cone density trended toward an increase with eye length but none of the trends were significant. A more recent study measured peak cone densities in the fovea as well as axial length for 22 eyes of 22 subjects (Wilk et al., 2017) but they did not plot peak cone density as a function of axial length, as it was not the aim of their study. We plotted the data they provided in their paper and found that the linear cone density at the foveal center dropped significantly with increases in axial length, similar to what was found by Li et al. (2010) and Chui et al. (2008), but the angular cone density had no dependency on eye length. Summary plots from previous literature are shown in **Figures 2ab.**

Wilk et al. (2017)'s data were consistent with a global expansion model and Li et al. (2010) and Chui et al. (2008)'s data only leaned toward a model that falls between the global expansion and equatorial stretching models. If the trends found by Li et al. (2010) and Chui et al. (2008) near the fovea were to extend to the foveal center, then myopes would have higher foveal photoreceptor sampling resolution with a consequent potential for better performance on visual tasks compared to emmetropes. As such, the simplest explanation for visual deficits in myopes – increased separation between cones caused by retinal stretching – would have to be ruled out.

With the improvements in resolution of adaptive optics ophthalmoscopes, imaging the smallest cones at the foveal center is now possible in many eyes, making it possible to complete a definitive analysis of the cone density at the fovea as a function of eye length.

Figure 2. Summary of published data from Li et al. (2010), Chui et al. (2008) and Wilk et al. (2017). In both plots, the linear fits with the solid lines indicate the data that have significant trends. **(a)** Linear cone density has a decreasing trend with axial length near the fovea. **(b)** Angular cone density (sampling resolution) of the eye generally increases with axial length although none of the data show a significant linear relationship.

Results

The experiments were approved by the University of California, Berkeley Committee for the Protection of Human Subjects. All subjects provided informed consent prior to any experimental procedures. Subjects self-reported their eye health so that only healthy individuals with no ocular conditions were included in the study. All eyes were dilated and cyclopleged with 1% Tropicamide and 2.5% Phenylephrine before imaging. We report data from 28 eyes of 16 subjects with a wide range of refractive error and axial length. Age, sex and ethnicity are listed on **Table 2**.

Biometry Data

All the biometric measures used to convert angular dimensions to linear retinal dimensions are listed on **Table 2.** The strong correlation of refractive error and eye length (P < 0.0001) indicates that the subjects were predominantly as a result of axial length.

Imaging Data

Images of the foveal region, the preferred retinal locus for fixation (PRL) and the fixation stability were recorded with an adaptive optics scanning laser ophthalmoscope (see **Methods and Materials**). The image of one subject (10003L) is shown in **Figure 3a**. All the cones were resolved with our imaging system. The scatter plot indicates the scatter plot of fixation over the course of a 10-sec video. **Figure 3b** shows the same image with all cones labeled and a color-coded overlay indicating the density. 16,184 labeled cones are shown on the figure. The point of maximum density is indicated by the blue cross and the average location of the PRL is indicated by the yellow cross (mean of the scatter plot locations in **Figure 3a**). This eye has a peak linear density of 200,482 cones/mm², and a peak angular density of 15,584 cones/deg². Cone density plots in linear and angular units for all eyes are shown on **supplemental figures 1 and 2.** Original images and a list of the cone locations for each can be downloaded from the Resources section of the Roordalab website (roorda.vision.berkeley.edu).

Figure 4 shows the linear cone density as a function of linear eccentricity, where the average linear cone density was computed in 25-micron wide annuli centered around the point of peak density.

Table 2. Each subject's refractive error was self-reported at the time of the study. Axial Length, corneal curvature and anterior chamber depth were measure by IOL Master, and retinal magnification factor (microns/deg) was calculated from biometry data.

Subject ID	Eye	Gender	Age	Ethnicity	Spherical equivalent refraction (D)	Axial length (mm)	Corneal curvature (mm)	Anterior chamber depth (mm)	Retinal magnification factor (microns/deg)	Angular cone density (cones/deg2)	Linear Cone Density (cones/mm2)	PRL distance from fovea (minutes)	PRL distance from fovea (microns)	PRL angular cone density (cones/deg2)	PRL linear cone density (cones/mm2)
20163	L	F	25	Asian	-7.125	27.06	7.89	3.65	340.44	18793	162149	5.03	28.52	17650	152300
20158	R	F	25	Asian	-7.125	26.84	7.89	3.65	336.60	17922	158183	4.16	23.31	17510	154500
20143	R	F	34	Asian	-6.500	26.60	7.84	3.51	333.78	13018	116845	3.21	17.88	12740	114400
20160	R	F	23	Asian	-6.875	25.91	7.42	2.10	334.12	17051	152739	2.92	16.26	17370	155600
20114	R	F	24	Asian	-5.375	25.83	7.81	3.60	320.25	15539	151507	8.97	47.86	14810	144400
	L	F	24	Asian	-6.000	26.16	8.98	3.58	313.31	15584	158761	5.63	29.38	14940	152200
20138	R	F	29	Caucasian	-5.500	25.83	8.72	3.47	310.94	14393	148864	7.34	38.05	13840	143200
	L	F	29	Caucasian	-5.000	25.28	7.91	3.15	311.92	14347	147459	5.23	27.20	14300	147000
20170	R	M	26	Asian	-5.000	25.26	7.95	3.14	311.22	13568	140078	6.37	33.05	12830	132500
	L	M	26	Asian	-3.750	25.66	7.65	4.15	316.25	14759	147573	1.50	7.90	14990	149900
20173	R	F	22	Caucasian	-2.250	25.00	7.69	3.90	305.54	14393	154172	8.77	44.65	12740	136500
	R	F	22	Caucasian	-2.750	24.96	7.81	3.68	304.64	16547	178298	7.24	36.73	15910	136000
20174	L	F	43	Caucasian	-2.750	25.37	7.83	3.62	311.85	12697	130557	5.90	30.66	11640	119700
	R	F	43	Caucasian	-1.750	24.80	7.79	3.57	302.57	13476	147200	7.67	38.65	11550	126200
20124	L	F	26	Asian	-4.250	25.29	7.68	4.07	309.88	13659	142247	1.76	9.08	13800	143700
	L	F	26	Asian	-3.000	24.67	7.70	4.05	298.82	13843	155024	5.15	25.63	13380	149900
20147	R	M	26	Caucasian	0.000	24.17	7.81	4.03	288.94	14805	177337	11.70	56.36	13570	162500
	L	M	26	Caucasian	-0.375	24.16	7.73	3.96	298.73	15401	172581	6.17	30.70	14670	164400
20172	R	F	25	Caucasian	-0.500	23.65	7.72	2.36	281.33	14668	185324	3.43	16.08	14760	186500
	L	F	25	Caucasian	-0.750	23.56	7.71	3.90	280.13	15264	194508	2.16	10.06	15170	193300
20176	R	F	18	Caucasian	0.000	23.58	8.01	3.62	278.52	12193	157174	3.97	18.42	11960	154200
	L	F	18	Caucasian	1.000	23.45	7.98	3.65	276.50	12513	163676	15.82	72.90	8984	117500
10003	R	M	50	Mixed	1.000	23.50	7.81	3.14	282.00	15172	190784	4.40	20.68	14670	184400
	R	M	50	Mixed	0.000	23.30	7.80	3.12	278.81	15584	200482	7.11	33.02	14070	181000
20177	L	F	18	Caucasian	0.000	23.23	7.91	3.20	275.85	11780	154810	4.60	21.16	11550	151800
	R	F	18	Caucasian	0.500	23.04	7.80	3.24	273.59	12055	161053	7.12	32.48	11730	156800
20165	L	F	28	Caucasian	0.500	22.64	7.44	3.80	267.79	12468	173857	5.48	24.45	11870	165500
	R	F	28	Caucasian	0.500	22.26	7.37	3.86	261.79	13247	193288	3.80	16.60	12650	184600

192
Figure 3. **(a)** AOSLO image of the fovea one subject (10003L). Only the central 1.5 degrees are
shown here (810 X 810 pixels), which contains 16,184 cones. The white dots are a scatter plot
showing the PRL, or position of the fixated stimulus over the course of a 10-second video. The
red dot is the centroid of the scatter plot. **(b)** Same image with a color overlay indicating the
density. Linear and angular cone densities are indicated on the right colorbar. Peak cones
densities in this eye are 200,482 cones/mm^2 and 15,584 cones/deg^2. The yellow and blue crosses
indicate the PRL and the position of peak cone density respectively. Scale bar is 0.5 degrees,
which in this eye corresponds to 139.4 microns.

202
Figure 4. The cone density changes over different eccentricities in all the eyes. The axial length
ranges of the subjects are color coded, with warmer colors for shorter eyes and cooler colors for
longer eyes. In this plot, it is apparent that shorter eyes generally have higher peak cone
densities.

In order to show the trends of density with axial length **Figure 5a&b** plot linear and
angular cone density as a function of axial length where the colors indicate different eccentricity
- red to purple indicate distance from the from fovea towards more parafoveal locations. **Figure**
5a reveals that peak linear density decreases significantly with axial length and the trend persists
and remains significant from the fovea out to 100 microns eccentricity. Axial length accounts for
38% of the variance in the changes in linear cone density. **Figure 5b** shows the opposite trends
when plotted in angular units. Peak angular density increases significantly with axial length and
the trend persists and remains significant out to 40 arcminutes eccentricity. Axial length accounts
for 32% of the variance in the changes in angular cone density. The plots clearly indicate that
although stretching does occur (**Figure 5a**) it is not a simple global expansion and longer eyes
have higher sampling density. The trends hold at and around fovea with statistical significance.

219
Figure 5. (a) Linear cone densities as a function of axial length. Longer eyes have lower linear
cone density than shorter eyes. The trend remains significant out to 100 microns eccentricity and
P values smaller than 0.05 are labelled with asterisks. **(b)** Angular cone densities as a function of
axial length. The peak angular cone density increases significantly with increasing axial length
and this trend remains significant out to 40 arcminutes eccentricity. The asterisks show all the
significant trends.
226

A more relevant measure of the impact of eye length on vision is how the angular cone
density changes at the PRL, which is often displaced from the location of peak cone density (Li
et al., 2010; Putnam et al., 2005; Wilk et al., 2017). If, for example, longer eyes had more
displaced PRLs then that could diminish, or even reverse, the trend of increased angular density
with eye length reported in **Figure 5b**. We found that the average displacement between PRL
and maximum cone density was 5.82 arcminutes and 28.94 microns. There was no significant
linear relationship found between PRL displacement in either angular or linear units vs. axial
length. Therefore, the PRL was not more displaced in myopes than in emmetropes from the point
of peak cone density. Plots of the cone density at PRL with axial length show the same trend at
the PRL as at the point of maximum cone density (**Figure 6 a&b**).

237
238   **Figure 6 ab.** The relationship between cone density and axial length shows the same pattern at
239   PRL as for the peak cone density. The slopes in both (a) and (b) are significant (P<0.005) and
240   axial length accounts for 27% and 30% of the changes in linear and angular cone density,
241   respectively.

242
Finally, we explored whether fixational eye movements might have a dependency on
axial length. Fixation stability around the PRL had an average standard deviation of 4.0
arcminutes and 20.2 microns. We found a small but significant increase in the standard deviation
of fixational eye movement in microns with axial length (**Figure 7a**). But when we plotted it in
arcminutes, the trend was no longer significant (**Figure 7b**). In another words, the increase in
fixational eye movements in microns was just a symptom of having a longer eye.

249
250 **Figure 7. (a)** A small but significant increase in standard deviation of fixational eye movement
251 in microns was found. **(b)** when plotted in arcminutes, the trend is no longer significant.
252
253 **Discussion**
254
In this paper we measure the cone density at and near the foveal center and investigate
how it changes as a function of axial length. This is the first comprehensive study of cones in
living eyes at the foveal center, the area solely responsible a for human's fine spatial vision. Our
results show that although some expansion does occur (linear cone density decreases with axial
length) the angular sampling resolution actually increases, on average, with axial length. Prior to
this study, the relationships between cone density and axial length were only made outside of the
fovea, the closest being 0.1 mm, or 0.3 degrees (Li et al., 2010). Although an eccentricity of 0.3
deg might seem close, it is noted that the cone density drops precipitously just outside of the
location of peak density (Curcio, Sloan, Kalina, & Hendrickson, 1990) as does human vision
(Poletti, Listorti, & Rucci, 2013)(Rossi & Roorda, 2010b). There are other factors that govern
peak cone density, however; eye length accounts for anywhere between 27% and 38% of the
variance in cone density.
Our finding that the slopes of cone density vs. axial length are in opposite directions
when plotted in linear (negative slope) and angular (positive slope) units, supports an eye growth
model that lies between the global expansion model and an equatorial stretching model. Previous
studies from our lab (Li et al., 2010) and also from (Chui, Song, & Burns, 2008) leaned in the
same direction. None of the cone density studies provide insight into the reasons why the
photoreceptor density would behave this way with eye growth, but the results do align with other

observations reported in the literature. Specifically, (Atchison et al., 2004) found that eyeball
dimensions in axial myopes are variable but are generally larger in all directions with a weak
tendency to be preferentially greater in the axial direction. These reported eye growth patterns lie
between that illustrated for the global expansion and equatorial stretching models in **figure 1**.
Our results differ from Wilk et al. (2017) whose data support a global expansion model
(i.e. there is no detectable change in angular cone density with axial length; **figure 2b**). But it is
important to point out that their study did not set out to address the same question and the
number of subjects with long axial lengths was disproportionately low.
Our results also differ from Troilo (1998) who studied retinal cell topography in a
marmoset animal myopia model. Higher cone packing densities were observed in the
experimentally enlarged eyes compared to normal eyes in the fovea. Their result followed the
overdevelopment model, which is the reason why we included it as one of the possible outcomes
of our study. In fact, the overdevelopment model is an extension of Springer's model of
development (Springer & Hendrickson, 2004), which offers a biomechanical explanation for
how cone packing increases at the foveal center in a developing eye. While our data do not
support the overdevelopment model, it does not preclude the existence of biomechanical factors
working in opposition to simple global expansion.
The fact that angular cone density (visual sampling resolution) increases with eye length
(myopia), at the peak density and at the PRL, means that poorer performance by myopes on
resolution tasks cannot be explained by a decrease in photoreceptor sampling. The deficit musts
arise at a post-receptoral level.
Low-level causes for myopic visual deficits might arise from differences in the
connectivity between cones and ganglion cells. Atchison et al. (2006) suggested that abnormal
eye growth may be associated with a loss of ganglion cells. Alternately, if ganglion cells pool
signals from multiple cones, then they will impose the retinal sampling limit and reduce certain
aspects of visual performance (acuity, for example). Recent electron microscopy studies of a
human fovea have revealed extensive convergence and divergence connections between
photoreceptors and ganglion cells, albeit in an eye from an individual who was born prematurely
(Dacey, 2018). These discoveries challenge our current understanding of neural connectivity in
the foveal center and force us to consider the possibility of interindividual differences in foveal
cone wiring. More experiments are necessary to explore these ideas.
To explain why low myopes did not perform as well on an acuity task as emmetropes,
even after correction or bypassing of high order aberrations, Rossi et al. (2007) and Coletta &
Watson (2006) both raised the possibility that myopes might have become desensitized to high
frequency information (low level myopic amblyopia) as a result of having less exposure to a high
contrast visual environment. In this case, it might be possible to train myopes to take advantage
of their higher sampling resolution, but one myope in a follow up study by Rossi & Roorda
(2010a) never reached the acuity levels of emmetropes in the same study.

Comparisons with Previous Studies

Peak cone densities: Curcio et al. 1990 measured spatial density of cones and rods in
eight explanted whole-mounted human retinas. They found a large range of peak foveal cone
densities with an average of 199,000 cones/mm^2. When we averaged the peak cone density over
a circular aperture of 7.5 arcminutes which was similar to the 29 x 45 micron window that
Curcio et al. (1990) used to compute density, we measured peak linear cone densities ranging

from 123,611 to 214,895 with an average of 168,047 cones/mm^2. Zhang et al. (2015) reported an average peak density of 168,162 cones/mm^2 in 40 eyes although they used a much smaller 5 x 5 micron sampling window to measure the peak. Wilk et al. (2017) reported an average peak density of 145,900 cones/mm^2 in 22 eyes using a 37 x 37 micron sampling window and Li et al. (2010) reported an average peak density of 150,412 cones/mm^2 in 4 eyes over a sampling window encompassing 150 cones (approximately 37 micron diameter at the foveal center). All reports of cone densities from adaptive optics studies in living eyes are lower than reports from histology. Two possible reasons for this are (i) the excised tissue in Curcio et al. (1990) underwent more shrinkage than estimated or (ii) the adaptive optics reports are subject to selection bias, where individuals with the highest angular cone densities might have been excluded because the image were less well resolved rendering the cones images too difficult to label with confidence. In our study, we attempted to image 73 eyes from 46 subjects and only succeeded in resolving cones across a sufficiently large region at and around the fovea in 28 of them. The reason the images from 45 eyes were not analyzed was due to poor or inconsistent image quality arising from a number of factors: Images from 4 eyes (3 subjects) were not analyzed because their refractive errors were too high (all above –8D) and we ran into to the limits of the deformable mirror's dynamic range. Images from 18 eyes (13 subjects) data were not analyzed because the optics of AOSLO was not tuned well enough to resolve foveal cones (those images were acquired early in the study). Images from 4 eyes (2 subjects) were not analyzed because of uncorrectable image degradation caused by keratoconus and corneal scarring. Images from 2 eyes (1 subject) were not analyzed because of excessive aberrations caused by an orthokeratology refractive correction. The cause of poor or inconsistent image quality among the remaining 17 eyes were varied, including ocular surface dryness, excessive eye motion and small pupils. The average refractive error among these remaining 17 eyes was about the same as the successful eyes.

Anisotropic density distribution: Like Curcio et al. (1990) and Zhang et al. (2015) we found steeper drops in cone density in the superior and inferior directions compared to the nasal and temporal directions. Plots of density along the two cardinal directions are shown on **Supplemental Figure 3.**

PRL displacements: The distance of the PRL from the foveal center for our study (mean 29 microns; range 8 – 73; n = 28) roughly agrees with those of Wilk et al. (2017) (mean 63 microns; range 20 – 263; n = 22), Li et al. (2010) (mean 34 microns; range 3 – 92; n = 18) and Putnam et al. (2005) (mean 17; range 11 – 23; n = 5). The differences in cone density between the peak and the PRL were small and the trends (**Figures 5 and 6**) persisted at both locations.

Spatial vision estimates: The cone array imposes the first retinal sampling limit to human spatial vision (MacLeod, Williams, & Makous, 1992; Williams, 1985) and the photoreceptor row-to-row spacing (assuming an hexagonal packing structure) imposes the maximum frequencies that can be relayed to later stages without aliasing. We can compute the sampling limit using the following formula:

$$Sampling\ Limit = \frac{1}{2}\sqrt{\frac{2}{\sqrt{3}}AngularDensity}$$

For the densities reported here, the potential spatial frequency resolution limits range from 58.3 to 73.6 cyc/deg at the peak density and 58.2 to 71.4 cyc/deg at the PRL. These correspond to

potential acuities ranging from 20/10.3 to 20/8.2 (based on the primary spatial frequency of the
three bars of a Snellen E). The cone frequency cut-offs are higher than almost all the
interferometric acuity limits reported by Coletta & Watson (2006), even for the emmetropic
subjects. The acuities are in the range of those measured from emmetropic subjects after
adaptive optics correction by Rossi et al. (2007). A direct comparison of foveal structure and
function for each of our subjects was not the scope of this study but will be the topic of future
investigation.
370
Measuring structure and function of cone photoreceptors at the foveal center – the most
important region of the human retina – has been one of the more challenging endeavors in vision
science. Fortunately, the latest generation of adaptive optics ophthalmoscopes are making it
easier and are facilitating new discoveries within this retinal region. The pattern of how cone
density changes with eye growth lands somewhere between the global expansion and equatorial
stretching models. The cone mosaic in longer eyes is expanded at the fovea, but not in proportion
to eye length. Despite retinal stretching, myopes generally have a higher angular sampling
density in and around the fovea compared to emmetropes. Reports of reduced best-corrected
central visual acuity in myopes compared to emmetropes cannot be explained by decreased
photoreceptor density caused by retinal stretching during myopic progression.
381
382 **Material and Methods**
383
Foveal Imaging:
We used our latest generation adaptive optics scanning laser ophthalmoscope (AOSLO)
for foveal imaging. The system used a mirror-based, out-of-plane optical design (Dubra et al.,
2011), and employed a deformable mirror with a continuous membrane surface and shaped with
97 actuators (DM97, ALPAO, Montbonnot-Saint-Martin, France). The system scans multiple
wavelengths simultaneously. Each wavelength was drawn from the same broadband
supercontinuum source (SuperK EXTREME, NKT Photonics, Birkerod, Denmark) using a
custom-built fiber coupler. Wave aberrations were measured with a custom-built Shack
Hartmann wavefront sensor using the 940 nm channel. Images were recorded using the 680 nm
channel. 512 x 512 pixel videos were recorded over a 0.9 x 0.9 degree square field for an average
sampling resolution of 9.48 pixels per arcminute. Eye alignment and head stabilization was
achieved by using either a bite bar or a chin rest with temple pads. At least one 10-second video
was recorded at the fovea and at 8 more locations where the subjects were instructed to fixate on
the corners and sides of the raster, to image an entire foveal region spanning about 1.8 X 1.8
degrees. In order to ensure the best possible focus of the foveal cones, multiple videos were
taken over a range of 0.05 D defocus steps to find the sharpest foveal cones. Focus steps were
generated by adding a focus shape onto the deformable mirror. Online stabilization and
registration algorithms were used to facilitate rapid feedback on the image quality.
402
Locating the Preferred Retinal Locus of Fixation (PRL)
Steady fixation was achieved at the fovea center by having the subjects fixate on a dark,
circular, blinking dot with a diameter of 3.16 arcminutes (30 pixels) in the center of the raster.
The fixation target was generated by modulating the same 680 nm scanning beam used for
imaging and, as such, the target's location was encoded directly into each frame of the video
(Poonja, Patel, Henry, & Roorda, 2005). A scatter plot of the position of the blinking dot relative

to the retina was generated and was used to determine the fixation stability (**figure 7**) and the
exact location of the PRL within the imaged cone mosaic (**Table 2, Figure 3, Supplemental**
figures 1 and 2).
412
Image Processing and Analysis
High quality images were generated from the recorded videos offline using custom
software (Matlab, The MathWorks, Inc., Natick, MA) to measure and correct for distortions
caused by eye movements (Stevenson & Roorda, 2005). Poor-quality frames were manually
excluded and registered frames were averaged into a single high signal-to-noise image. The
processed images were stitched together (Photoshop; Adobe Systems, Inc., Mountain View, CA)
to create an approximately 1.8-degree montage of the foveal cone mosaic.
We used custom software to identify and label individual cones in the AO retinal images.
The program allows the user to select a region of interest and manually add and delete cone
labels. A combination of both manual and automated methods (Li & Roorda, 2007) were used to
identify cone locations as the current version of the program does not adequately recognize cones
in the foveal center where they are dim and smaller (Li et al., 2010). All the cone coordinates
were selected and reviewed by two of the authors. In some cases cones were too dim to be seen
but there was only a gap in the mosaic (Bruce et al., 2015). If a space that might have been
occupied by a cone was dim or dark, we would assume it was a cone and mark its location. We
rationalize this for two reasons: First, if there is a gap in the mosaic, then it is likely that a cell is
occupying that space, otherwise the adjacent cells would migrate to fill it in (Scoles et al., 2014).
Second, in our experience and of others (Pallikaris, Williams, & Hofer, 2003), cones that appear
dark in one visit, can often appear bright in the next. In other cases (uncommon) the contrast was
low in some regions or there were interference artifacts in the images (Meadway & Sincich,
2018; Putnam, Hammer, Zhang, Merino, & Roorda, 2010), making the cone locations slightly
ambiguous. In these instances, we made manual cone selections based on the assumption that the
cones were all similar in size and close-packed into a nearly hexagonal array (Curcio et.al.,
1990).
Continuous density maps were generated by computing cone density within a circle of 10
arcminutes in diameter around every pixel location across the image. We kept the area large
enough to generate smooth maps, but small enough to resolve local changes. Changes in density
with eccentricity were generated by computing the density in 5 arcminute annuli surrounding the
point of peak cone density. For linear density measures we used annuli with 25 micron widths.
442
Retinal Magnification Factor Calculation
The exact angular dimensions of the AOSLO images were computed by imaging a
calibrated model eye in the AOSLO system, but the conversion to linear dimensions on the
retinal image requires additional measurements, since the dimensions of each eye governs the
actual size of the image on its retina. The conversion from visual angle to retinal distance
requires a measurement of the axial length of the eye and an estimation of the location of the
secondary nodal point. We used a four-surface schematic eye model, originally proposed by Li et
al., 2010 to estimate the location of the secondary nodal point. The corneal first surface radius of
curvature, the anterior chamber depth and the axial length were for measured for each subject
with an IOL Master (Zeiss Meditec, Dublin, CA). The radius of the curvature of the back surface
of the cornea was computed as 88.31% of the front surface (Bennett, Rudnicka, & Edgar, 1994).
The indices of refraction of the media and the radii of curvature of the front and back lens

surface were taken from the Gullstrand schematic eye (Vojnikovic & Tamajo, 2013). Once determined, retinal image size is related to visual angle by the equation:

$$I = \tan(1°)(x - AN')\theta$$

Where I is retinal image size, x is axial length, AN' is the distance from the corneal apex to the eye's second nodal point, and θ is the visual angle. As can be seen in **Table 2**, myopic eyes, which generally have longer focal lengths, have proportionally larger retinal images.

Competing interests:

A.R. has a patent (USPTO#7118216) assigned to the University of Houston and the University of Rochester which is currently licensed to Boston Micromachines Corp (Watertown, MA, USA). Both he and the company stand to gain financially from the publication of these results. No other authors have competing interests.

References

Atchison, D. A., Jones, C. E., Schmid, K. L., Pritchard, N., Pope, J. M., Strugnell, W. E., & Riley, R. A. (2004). Eye shape in emmetropia and myopia. *Investigative Ophthalmology and Visual Science*, *45*(10), 3380–3386. https://doi.org/10.1167/iovs.04-0292

Atchison, D. A., Schmid, K. L., & Pritchard, N. (2006). Neural and optical limits to visual performance in myopia. *Vision Res*, *46*(21), 3707–3722. https://doi.org/10.1016/j.visres.2006.05.005

Bennett, A. G., Rudnicka, A. R., & Edgar, D. F. (1994). Improvements on Littmann's method of determining the size of retinal features by fundus photography. *Graefe's Archive for Clinical and Experimental Ophthalmology*, *232*(6), 361–367. https://doi.org/10.1007/BF00175988

Bruce, K. S., Harmening, W. M., Langston, B. R., Tuten, W. S., Roorda, A., & Sincich, L. C. (2015). Normal perceptual sensitivity arising from weakly reflective cone photoreceptors. *Investigative Ophthalmology and Visual Science*, *56*(8), 4431–4438. https://doi.org/10.1167/iovs.15-16547

Chui, T. Y. P., Song, H., & Burns, S. A. (2008). Individual Variations in Human Cone Photoreceptor Packing Density: Variations with Refractive Error. *Investigative Opthalmology & Visual Science*, *49*(10), 4679. https://doi.org/10.1167/iovs.08-2135

Chui, T. Y., Yap, M. K., Chan, H. H., & Thibos, L. N. (2005). Retinal stretching limits peripheral visual acuity in myopia. *Vision Res*, *45*(5), 593–605. https://doi.org/10.1016/j.visres.2004.09.016

Coletta, N. J., & Watson, T. (2006). Effect of myopia on visual acuity measured with laser interference fringes. *Vision Res*, *46*(5), 636–651. https://doi.org/S0042-6989(05)00268-3

[pii]\r10.1016/j.visres.2005.05.025

Collins, J. W., & Carney, L. G. (1990). Visual performance in high myopia. *Current Eye*
Research, *9*(3), 217–224. https://doi.org/10.3109/02713689009044516

Curcio, C. A., Sloan, K. R., Kalina, R. E., & Hendrickson, A. E. (1990). Human photoreceptor
topography. *Journal of Comparative Neurology*, *292*(4), 497–523.
https://doi.org/10.1002/cne.902920402

Dacey, D. M. (2018). Discovering visual pathway origins in the center of the human foveola
with connectomics. *Investigative Ophthalmology and Visual Science*, *59*(9), ARVO
Abstract #14.

Dubra, A., Sulai, Y., Norris, J. L., Cooper, R. F., Dubis, A. M., Williams, D. R., & Carroll, J.
(2011). Noninvasive imaging of the human rod photoreceptor mosaic using a confocal
adaptive optics scanning ophthalmoscope. *Biomedical Optics Express*, *2*(7), 1864.
https://doi.org/10.1364/BOE.2.001864

Ehsaei, A., Chisholm, C. M., Pacey, I. E., & Mallen, E. A. H. (2013). Visual performance fall-off
with eccentricity in myopes versus emmetropes. *Journal of Optometry*, *6*(1), 36–44.
https://doi.org/10.1016/j.optom.2012.07.001

Elsner, A. E., Chui, T. Y. P., Feng, L., Song, H. X., Papay, J. A., & Burns, S. A. (2017).
Distribution differences of macular cones measured by AOSLO: Variation in slope from
fovea to periphery more pronounced than differences in total cones. *Vision Research*, *132*,
62–68. https://doi.org/10.1016/j.visres.2016.06.015

Fiorentini, A., & Maffei, L. (1976). Spatial contrast sensitivity of myopic subjects. *Vision*
Research, *16*(4), 437–438. https://doi.org/10.1016/0042-6989(76)90214-5

Fujiwara, T., Imamura, Y., Margolis, R., Slakter, J. S., & Spaide, R. F. (2009). Enhanced Depth
Imaging Optical Coherence Tomography of the Choroid in Highly Myopic Eyes. *American*
Journal of Ophthalmology, *148*(3), 445–450. https://doi.org/10.1016/j.ajo.2009.04.029

Gonzalez Blanco, F., Sanz Ferńandez, J. C., & Muńoz Sanz, M. A. (2008). Axial length, corneal
radius, and age of myopia onset. *Optometry and Vision Science*, *85*(2), 89–96.
https://doi.org/10.1097/OPX.0b013e3181622602

Harb, E., Hyman, L., Gwiazda, J., Marsh-Tootle, W., Zhang, Q., Hou, W., … Taylor, C. (2015).
Choroidal thickness profiles in myopic eyes of young adults in the correction of myopia
evaluation trial cohort. *American Journal of Ophthalmology*, *160*(1), 62–71.e2.
https://doi.org/10.1016/j.ajo.2015.04.018

He, M., Zeng, J., Liu, Y., Xu, J., Pokharel, G. P., & Ellwein, L. B. (2004). Refractive error and
visual impairment in urban children in southern china. *Invest Ophthalmol Vis Sci*, *45*(3),
793–799. https://doi.org/10.1167/iovs.03-1051

He, X., Zou, H., Lu, L., Zhao, R., Zhao, H., Li, Q., & Zhu, J. (2015). Axial length/corneal radius
ratio: Association with refractive state and role on myopia detection combined with visual
acuity in Chinese schoolchildren. *PLoS ONE*, *10*(2), e0111766.
https://doi.org/10.1371/journal.pone.0111766

Iyamu, E., Iyamu, J., & Obiakor, C. I. (2011). The Role of Axial Length-Corneal Radius of
Curvature Ratio in Refractive State Categorization in a Nigerian Population. *ISRN*
Ophthalmology, *2011*, 1–6. https://doi.org/10.5402/2011/138941

Jaworski, A., Gentle, A., Zele, A. J., Vingrys, A. J., & McBrien, N. A. (2006). Altered visual
sensitivity in axial high myopia: A local postreceptoral phenomenon? *Investigative*
Ophthalmology and Visual Science, *47*(8), 3695–3702. https://doi.org/10.1167/iovs.05-1569

Jong, M., Sankaridurg, P., Li, W., Resnikoff, S., Naidoo, K., & He, M. (2018). Reduced vision in

highly myopic eyes without ocular pathology: the ZOC-BHVI high myopia study. *Clin Exp Optom*, *101*(1), 77–83. https://doi.org/10.1111/cxo.12563

Kitaguchi, Y., Bessho, K., Yamaguchi, T., Nakazawa, N., Mihashi, T., & Fujikado, T. (2007). In vivo measurements of cone photoreceptor spacing in myopic eyes from images obtained by an adaptive optics fundus camera. *Japanese Journal of Ophthalmology*, *51*(6), 456–461. https://doi.org/10.1007/s10384-007-0477-7

Li, K. Y., & Roorda, A. (2007). Automated identification of cone photoreceptors in adaptive optics retinal images. *Journal of the Optical Society of America A*, *24*(5), 1358. https://doi.org/10.1364/JOSAA.24.001358

Li, K. Y., Tiruveedhula, P., & Roorda, A. (2010). Intersubject variability of foveal cone photoreceptor density in relation to eye length. *Investigative Ophthalmology and Visual Science*, *51*(12), 6858–6867. https://doi.org/10.1167/iovs.10-5499

Liang, J., Williams, D. R., & Miller, D. T. (1997). Supernormal vision and high-resolution retinal imaging through adaptive optics. *Journal of the Optical Society of America A*, *14*(11), 2884. https://doi.org/10.1364/JOSAA.14.002884

Liou, S.-W., & Chiu, C.-J. (2001). Myopia and contrast sensitivity function. *Current Eye Research*, *22*(2), 81–84. https://doi.org/10.1076/ceyr.22.2.81.5530

MacLeod, D. I. A., Williams, D. R., & Makous, W. (1992). A visual nonlinearity fed by single cones. *Vision Research*, *32*, 347–363. https://doi.org/10.1016/0042-6989(92)90144-8

Meadway, A., & Sincich, L. C. (2018). Light propagation and capture in cone photoreceptors. *Biomedical Optics Express*, *9*(11), 5543. https://doi.org/10.1364/boe.9.005543

Morgan, I. G., Ohno-Matsui, K., & Saw, S. M. (2012). Myopia. *The Lancet*, *379*(9827), 1739–1748. https://doi.org/10.1016/S0140-6736(12)60272-4

Obata, R., & Yanagi, Y. (2014). QuantItative analysis of cone photoreceptor distribution and its relationship with axial length, age, and early age-related macular degeneration. *PLoS ONE*, *9*(3), e91873. https://doi.org/10.1371/journal.pone.0091873

Pallikaris, A., Williams, D. R., & Hofer, H. (2003). The reflectance of single cones in the living human eye. *Investigative Ophthalmology and Visual Science*, *44*(10), 4580–4592. https://doi.org/10.1167/iovs.03-0094

Park, S. P., Chung, J. K., Greenstein, V., Tsang, S. H., & Chang, S. (2013). A study of factors affecting the human cone photoreceptor density measured by adaptive optics scanning laser ophthalmoscope. *Experimental Eye Research*, *108*, 1–9. https://doi.org/10.1016/j.exer.2012.12.011

Poletti, M., Listorti, C., & Rucci, M. (2013). Microscopic eye movements compensate for nonhomogeneous vision within the fovea. *Current Biology*, *23*(17), 1691–1695. https://doi.org/10.1016/j.cub.2013.07.007

Poonja, S., Patel, S., Henry, L., & Roorda, A. (2005). Dynamic visual stimulus presentation in an adaptive optics scanning laser ophthalmoscope. *Journal of Refractive Surgery*, *21*(5), S575–S580. https://doi.org/10.1364/fio.2004.fthv3

Putnam, N. M., Hammer, D. X., Zhang, Y., Merino, D., & Roorda, A. (2010). Modeling the foveal cone mosaic imaged with adaptive optics scanning laser ophthalmoscopy. *Optics Express*, *18*(24), 24902–24916. https://doi.org/10.1364/OE.18.024902

Putnam, N. M., Hofer, H., Doble, N., Chen, L., Carroll, J., & Williams, D. R. (2005). The locus of fixation and the foveal cone mosaic. *Journal of Vision*, *5*(7), 632–639. https://doi.org/10.1167/5.7.3

Roorda, A., Romero-Borja, F., Donnelly III, W. J., Queener, H., Hebert, T. J., & Campbell, M.

C. W. (2002). Adaptive optics scanning laser ophthalmoscopy. *Optics Express*, *10*(9), 405.
https://doi.org/10.1364/OE.10.000405

Rossi, E. A., & Roorda, A. (2010a). Is visual resolution after adaptive optics correction
susceptible to perceptual learning? *Journal of Vision*, *10*(12), 11.
https://doi.org/10.1167/10.12.11

Rossi, E. A., & Roorda, A. (2010b). The relationship between visual resolution and cone spacing
in the human fovea. *Nature Neuroscience*, *13*(2), 156–157. https://doi.org/10.1038/nn.2465

Rossi, E. A., Weiser, P., Tarrant, J., & Roorda, A. (2007). Visual performance in emmetropia
and low myopia after correction of high-order aberrations. *Journal of Vision*, *7*(8), 14.
https://doi.org/10.1167/7.8.14

Scoles, D., Sulai, Y. N., Langlo, C. S., Fishman, G. A., Curcio, C. A., Carroll, J., & Dubra, A.
(2014). In vivo imaging of human cone photoreceptor inner segments. *Investigative*
Ophthalmology and Visual Science, *55*(7), 4244–4251. https://doi.org/10.1167/iovs.14-
14542

Springer, A. D., & Hendrickson, A. E. (2004). Development of the primate area of high acuity. 1.
Use of finite element analysis models to identify mechanical variables affecting pit
formation. *Visual Neuroscience*, *21*(1), 53–62. https://doi.org/10.1017/S0952523804041057

Stevenson, S. B., & Roorda, A. (2005). Correcting for miniature eye movements in high
resolution scanning laser ophthalmoscopy. *Ophthalmic Technologies XV*, *5688*, 145.
https://doi.org/10.1117/12.591190

Stoimenova, B. D. (2007). The Effect of Myopia on Contrast Thresholds. *Investigative*
Opthalmology & Visual Science, *48*(5), 2371. https://doi.org/10.1167/iovs.05-1377

Strang, N. C., Winn, B., & Bradley, A. (1998). The role of neural and optical factors in limiting
visual resolution in myopia. *Vision Research*, *38*(11), 1713–1721.
https://doi.org/10.1016/S0042-6989(97)00303-9

Thorn, F., Corwin, T. R., & Comerford, J. P. (1986). High myopia does not affect contrast
sensitivity. *Curr Eye Res*, *5*(9), 635–639. Retrieved from
http://www.ncbi.nlm.nih.gov/pubmed/3769528

Troilo, D. (1998). Changes in retinal morphology following experimentally induced myopia.
Vision Science and Its Applications: Technical Digest (OSA, Washington, D.C.), *1*(1), 206–
209.

Verkicharla, P. K., Ohno-Matsui, K., & Saw, S. M. (2015). Current and predicted demographics
of high myopia and an update of its associated pathological changes. *Ophthalmic and*
Physiological Optics. https://doi.org/10.1111/opo.12238

Vitale, S. (2009). Increased Prevalence of Myopia in the United States Between 1971-1972 and
1999-2004. *Archives of Ophthalmology*, *127*(12), 1632.
https://doi.org/10.1001/archophthalmol.2009.303

Vojnikovic, B., & Tamajo, E. (2013). Gullstrand's optical schematic system of the eye--modified
by Vojnikovic & Tamajo. *Coll Antropol*, *37 Suppl 1*, 41–45. Retrieved from
http://www.ncbi.nlm.nih.gov/pubmed/23841130

Webb, R. H., Hughes, G. W., & Delori, F. C. (1987). Confocal scanning laser ophthalmoscope.
Applied Optics, *26*(8), 1492. https://doi.org/10.1364/AO.26.001492

Wilk, M. A., Dubis, A. M., Cooper, R. F., Summerfelt, P., Dubra, A., & Carroll, J. (2017).
Assessing the spatial relationship between fixation and foveal specializations. *Vision*
Research, *132*, 53–61. https://doi.org/10.1016/j.visres.2016.05.001

Williams, D. R. (1985). Aliasing in human foveal vision. *Vision Research*, *25*, 195–205.

https://doi.org/10.1016/0042-6989(85)90113-0

Zhang, T., Godara, P., Blanco, E. R., Griffin, R. L., Wang, X., Curcio, C. A., & Zhang, Y.

(2015). Variability in Human Cone Topography Assessed by Adaptive Optics Scanning

Laser Ophthalmoscopy. *American Journal of Ophthalmology*.

https://doi.org/10.1016/j.ajo.2015.04.034

Supplemental Figure 1: Linear cone density (cones/mm^2) plots over the central 450 microns for all 28 subjects. The black cross indicates the point of maximum cone density. The black circle indicates the PRL location. Dark blue regions indicate where no cone density estimates were made.

1.5 deg

652
653 **Supplemental Figure 2:** Angular cone density (cones/deg^2) plots over the central 1.5 degrees for
654 all 28 subjects. The black cross indicates the point of maximum cone density. The black circle
655 indicates the PRL location. Dark blue regions indicate where no cone density estimates were
656 made.
657

658
659 **Supplemental Figure 3:** Plots of density as a function of eccentricity in the vertical and
660 horizontal directions. (A) linear cone density (B) angular cone density. The dashed lines
661 represent +/- 1 standard deviation from the mean.