Phylogenetic inference of changes in amino acid propensities with single-position resolution

Galya V. Klink[1], Olga V. Kalinina[2], Georgii A. Bazykin[1,3]

[1]Institute for Information Transmission Problems (Kharkevich Institute) of the Russian Academy of Sciences, Moscow, Russia

[2]Helmholtz Institute for Pharmaceutical Research Saarland (HIPS), Helmholtz Centre for Infection Research (HZI), Saarbrücken, Germany; Medical Faculty, Saarland University, Homburg, Germany

[3]Skolkovo Institute of Science and Technology, Skolkovo, Russia

*Corresponding author: Georgii A. Bazykin, e-mail: g.bazykin@skoltech.ru

Abstract

Fitness conferred by the same allele may differ between genotypes, and these differences shape variation and evolution. Changes in amino acid propensities at protein sites over the course of evolution have been inferred from sequence alignments statistically, but the existing methods are data-intensive and aggregate multiple sites. Here, we develop an approach to detect individual amino acids that confer different fitness in different groups of species from combined sequence and phylogenetic data. Our method is based on the mutual distribution of homoplasic (convergent or parallel) and divergent substitutions over a phylogeny, and uses the fact that a higher rate of homoplasic evolution within a clade signifies high fitness of the derived allele. We apply it to the *env* gene of two HIV-1 subtypes, A and B, and to the HA gene of two influenza A subtypes, H1 and H3, and show that the inferred fitness changes are consistent with the fitness differences observed in deep mutational scanning experiments. Changes in fitness do not always trigger episodes of positive selection, and therefore cannot be identified by existing methods for its detection.

Introduction

A particular amino acid at a specific position of a protein may confer different fitness in different species. Changes in amino acid propensities can occur in the course of evolution due to substitutions at other epistatically interacting sites. Indeed, accounting for co-occurrence of amino acids improves predictions of the rates and patterns of substitutions (Kryazhimskiy et al. 2011) and polymorphism (Loewen and Ganetzky 2018; Zaidi and Makova 2019), as well as the pathogenic potential of mutations (Kondrashov et al. 2002; Jordan et al. 2015) and their functional effects in experiments (Mann et al. 2014; Figliuzzi et al. 2016; Hopf et al. 2017; Riesselman et al. 2018). Changes in amino acid propensities can also happen due to shifts in protein fitness landscape caused by an altering

environment, e.g. shifts in immune pressure on proteins of pathogens (Blackburne et al. 2008; Elena et al. 2009; Stray and Pittman 2012, Popova et al. 2019).

Variability of amino acid propensities has been inferred from statistical analyses of pooled data from numerous amino acid positions of large protein alignments (Bazykin 2015; Storz 2016). In particular, it has been postulated to bias the phylogenetic distribution of homoplasies – repeated substitutions giving rise to the same allele: parallelisms, convergences (Povolotskaya and Kondrashov 2010; Goldstein et al. 2015; Zou and Zhang 2015; Klink et al. 2017; Klink and Bazykin 2017) and reversals (Naumenko et al. 2012; Soylemez and Kondrashov 2012). The idea behind this approach is that the fixation probability of a mutation, and therefore the rate of substitution, giving rise to an allele increases with its conferred fitness (Kimura 1983). Therefore, substitutions producing the same allele are expected to be more frequent in groups of species where this allele confers high fitness. On a phylogeny, this can be observed as clustering of homoplasies in these groups of species, and such clustering has been interpreted as evidence for changes in fitness landscapes. To increase power, existing methods (Povolotskaya and Kondrashov 2010; Naumenko et al. 2012; Soylemez and Kondrashov 2012; Goldstein et al. 2015; Zou and Zhang 2015; Klink et al. 2017; Klink and Bazykin 2017) aggregate data across multiple sites, and therefore do not allow to track changes in amino acid propensities at individual sites.

Here, we develop an approach for identification of amino acid variants at a protein site that confer different fitness in two taxa. This approach is based on the phylogenetic distribution of substitutions at this site. Our method performs well on phylogenies that include hundreds of samples (species or strains) or more. Using this method, we infer sites in the gp160 protein (encoded by the *env* gene) of HIV-1 that have undergone a change in the relative fitness of different amino acids between two highly divergent HIV-1 subtypes, and sites in the hemagglutinin (HA) protein of influenza A that have undergone such a change between two influenza subtypes. We then validate these findings using the fitness differences measured directly in deep mutational scanning (DMS) experiments (Haddox et al. 2016; Haddox et al. 2018; Lee et al. 2018).

Results

Inferring variable fitness amino acids

Using the reconstructed phylogenetic positions of substitutions, we analyze how homoplasic (parallel or convergent, i.e., giving rise to the same amino acid) and divergent (giving rise to different amino acids) substitutions at the same amino acid site are distributed relative to each other over the phylogeny. We propose that aggregation of homoplasic substitutions at a particular fragment of the phylogenetic tree ("phylogenetic neighborhood") implies that the descendant allele confers higher fitness at this group of species, compared to the fitness conferred by it in other species. By contrast, a deficit of substitutions into the allele implies that it confers lower fitness at this group of species (Klink and Bazykin 2017).

Specifically, we aim to infer individual amino acids at specific sites that undergo significant changes in frequency of homoplasies between the phylogenetic neighborhoods of two given species. For this, we start by considering the first of the two species as the "focal" species. For each amino acid at each site, we then test two hypotheses: that the phylogenetic positions of substitutions into this amino acid are biased towards ("proximal amino acids") or away from ("distal amino acids") the phylogenetic neighborhood of the focal species (Fig. 1). Next, we consider the second species as the focal species, and repeat the procedure. Finally, we assume that the fitness conferred by an amino acid has changed between the two focal species if it was classified as proximal to one species and as distal from the other, and designate such amino acids as variable fitness amino acids.

Figure 1. Schematic representation of the approach for inference of changes in fitness of individual amino acids. Both A and B are variable fitness amino acids when species 1 and 2 are chosen as focal species, because each of them is proximal to one species and distal from the other. By contrast, A is distal, and B is proximal, for both species in the comparison of species 2 and 3, so neither A nor B is identified as a variable fitness amino acid in this comparison.

Method validation

To understand the power of our method to identify amino acids with variable fitness, we used SELVa (Nabieva and Bazykin 2019) to simulate the evolution of individual amino acid sites under variable fitness landscape. In our model, the substitution rates between different amino acids were determined by their relative fitness, which could differ between tree branches. Specifically, we assumed that at a given site a and at each position of the tree, exactly one of the 20 amino acids, AA_k, is favored by selection and has the scaled fitness $F_a^{AA_k} = 2N_e f_a^{AA_k} = X > 1$, where N_e is the effective population size; while the remaining 19 amino acids have $F_a^{AA_{i \neq k}} = 1$. The $AA_k \longrightarrow AA_{i \neq k}$ substitutions then occur at rate $\frac{1-X}{1-e^{X-1}}$, while the $AA_{i \neq k} \longrightarrow AA_k$ substitutions occur at rate $\frac{X-1}{1-e^{1-X}}$ (Yang and Nielsen 2008). To model fitness variability, we assumed that selection changes exactly once at a predefined phylogenetic branch b; at this point, the identity of the newly favored amino acid with the fitness X, AA_l, is picked out of the 20 amino acids randomly, and all remaining amino acids get fitness 1. In the process, the

fitness of amino acids AA_k and AA_l has changed, while the fitness of the remaining 18 variants has remained constant.

We asked if our method could retrieve the identity of AA_k and AA_l from simulated protein sequences. For simulation, we used the reconstructed phylogenetic tree of the gp160 protein encoded by the *env* gene of HIV-1 (see Methods). The tree contains 3300 tips and has two clades, corresponding to HIV-1 subtypes B and C (Supplementary Figure 1). We assumed that the change in the relative fitness values had occurred in the last common ancestor (LCA) of subtype C. To evaluate the dependency of the method performance on sampling density, we additionally created two artificial phylogenetic trees on the basis of the original one: "sparse tree" with roughly half of the tips and "dense tree" with twice as many tips as the original tree (see Methods).

The results of the simulations are shown in Figure 2. We find that under moderate selection ($X = 4$ to 9), we correctly identify ~20%, ~40% and ~80% of amino acids for which fitness has changed in sparse, original and dense trees, respectively. Under stronger selection ($X = 10$), many sites are monomorphic, and the sensitivity of our method decreases, constituting ~30% for the dense tree. Under weaker selection ($X = 3$), the power of the method also drops, probably because changes in selection of such a low magnitude are masked by the noise associated with drift and tree topology. The percent of amino acids with constant fitness that are identified as positives is low for all trees, never exceeding 0.6%.

Figure 2. Performance of our method for "sparse" (~1650 tips), original (3300 tips) and "dense" (6600 tips) phylogenetic trees in simulations for different scaled fitness values of the preferred variant. Red line, mean True Positive Rate (TPR); blue line, mean False Positive Rate (FPR). 90% confidence bands were obtained by random subsampling of 1000 amino acids in 100 trials. A zoom-in for low FPR values is shown in Supplementary Figure 2.

Many *env* sites change amino acid preferences in the course of evolution

As a test case, we applied our method to the evolution of the gp160 protein encoded by the *env* gene in HIV-1. This protein evolves under strong positive selection (Yang et al. 2003; Bazykin et al. 2006), suggesting that site-specific amino acid propensities are likely to change between strains. As focal strains, we considered the two strains tested in DMS experiments (see below): BF520 strain of the subtype A and LAI strain of the subtype B.

Indeed, among the 3491 amino acids with sufficient variability for our analysis (potentially variable fitness amino acids, see Methods), we detected 59 variable fitness amino acids (at FDR=11%; Supplementary figure 3; see Methods), distributed over 46 of the 724 sites of gp160 (Fig. 3).

Substitutions giving rise to the variable fitness amino acid were positioned on the phylogeny closer to the species for which they were proximal: the phylogenetic distance to that species comprised, on average, 0.87 (range: 0.57 to 0.99, standard deviation = 0.097) of that expected randomly. They also were positioned more remotely from the species for which they were distal: the phylogenetic distance to that species comprised, on average, 1.19 (range: 1.01 to 1.91, standard deviation = 0.19) of that expected randomly.

Of these 59 amino acids, 36 amino acids at 34 sites were proximal to the BF520 strain and distal from the LAI strain; and 23 amino acids at 23 sites were proximal to the LAI strain and distal from the BF520 strain. At 11 sites, changes in fitness were reciprocal: some amino acids were proximal to the BF520 strain and distal from the LAI strain, while others were proximal to the LAI strain and distal from the BF520 strain.

The sites carrying variable fitness amino acids were found at all domains and structural regions of gp120 and gp41, and were not biased towards or away from any functional domain (Fisher's exact test, mean two-sided p-value for functional domains > 0.6).

Figure 3. Amino acid sites carrying variable fitness amino acids in gp160 protein of HIV-1. For each position of the amino acid within the protein (horizontal axis), the red dot shows the number of amino acids that are simultaneously proximal to the BF520 strain and distal from the LAI strain; and the blue dot, the number of amino acids that are simultaneously proximal to the LAI strain and distal from the LBF520 strain, multiplied by -1. Grey dots represent the number of amino acids at a site that could potentially have significantly variable fitness (potentially variable fitness amino acids, see Methods). The positions without any dots are sites filtered out due to alignment ambiguity. Vertical grey lines connect dots for the same site. Blue background, gp120; green background, gp41. Variable loops V1-V5 and known functional domains are marked by horizontal black lines: *, CD4-binding loop; **, fusion peptide; ***, immunosuppressive region; ****, membrane proximal external region (MPER). Numbering of protein positions is hxb2-based.

Phylogenetically inferred changes in fitness between strains match those observed in DMS experiments

We compared the changes in phylogenetically inferred amino acid preferences between two strains to those inferred experimentally by DMS (Haddox et al. 2016; Haddox et al. 2018). To estimate the changes in DMS preferences, for each amino acid, we measured the ratio $r = \pi_x / \pi_y$, where π_x is the DMS preference for this amino acid in the strain for which it is phylogenetically more proximal, and π_y is the preference for it in the strain

for which it is more distal. As a control, we analyzed the distribution of r between the two strains for invariable fitness amino acids.

The values of r were significantly higher for the variable fitness amino acids than in the control (one-tailed Wilcoxon's test, p = 0.005; Fig. 4), indicating that the phylogenetically and experimentally inferred variable fitness amino acids match. This trend is pronounced both for amino acids that conferred higher fitness in BF520 and in LAI strains (Fig. 4). It is mainly driven by amino acids with modest preferences ($\pi < 0.2$; Fig. 4B), which are not the highest-ranking amino acids in DMS experiments in either of the strains (Fig. 4C).

Figure 4. *A*, The ratio of DMS preferences π for an amino acid between the strain for which this amino acid is phylogenetically more proximal and the strain for which it is more distal. Amino acids with significant (left, p<0.01 for both strains) or insignificant (right, p>0.1 for both strains) changes in conferred fitness between strains are shown; bold line is the median, box borders represent inter quartile range (IQR) between first (Q1) and third (Q3) quartiles, whiskers are Q1 - 1.5*IQR and Q3 + 1.5*IQR intervals, outliers are not shown; boxplots are made using all data; each variable amino acid close to subtype A or subtype B and a subsample of 100 other amino acids are shown as red, blue and grey dots respectively. Dashed line indicates no change in preferences (r=1). *B*, *C*, Experimentally measured fitness π (B) or fitness rank (C, with rank 1 corresponding to the fittest amino acid) of each amino acid in two strains of HIV1. Grey dots, amino acids that persist at least in one strain in our phylogenetic tree; red, variable fitness amino acids that are proximal to subtype A; blue, variable fitness amino acids that are proximal to subtype B; dashed line, diagonal.

Most sites with variable fitness amino acids show no evidence of recurrent positive selection

Changes in fitness conferred by an allele can trigger episodes of positive selection. To study the association between these processes, we identified sites under recurrent positive selection, as evidenced by an above-neutral ratio of nonsynonymous to synonymous substitutions (dN/dS) estimated by codeml ($\omega > 1$) in subtype A, subtype B or both. Among the 46 sites carrying variable fitness amino acids, only 20 (43%) have experienced positive selection in subtypes A or B, despite the fact that the power of both the site-specific dN/dS test for positive selection and our test increases with the number of substitutions. An even lower overlap was observed with the sets of positively selected sites described in previous studies (Yang et al. 2003: 17%; Wood et al. 2009: 9%). Although there was an excess of positive selection among sites with variable fitness amino acids (Methods; Fisher's exact test, two-sided $p < 0.001$, Table 1), the fact that no such selection was detected in over a half of variable sites suggests that these two phenomena are frequently uncoupled. Furthermore, r was not higher for variable fitness amino acids at positively selected sites than at sites without any signal of positive selection (one-tailed Wilcoxon's test, $p = 0.09$).

Table 1. Sites with variable fitness amino acids and recurrently positively selected sites.

	Sites with variable fitness amino acids	Sites without variable fitness amino acids	p-value*
Our analysis			
Positively selected sites	20	71	
			<0.001
Other sites	26	422	
(Wood et al. 2009)			
Positively selected sites	4	21	
Other sites	42	472	0.26
(Yang et al. 2003)			
Positively selected sites	8	37	
Other sites	38	456	0.04

A closer examination of the data helps understand the interplay between the dN/dS ratio and fitness variability. For example, according to the DMS data, Y159 is strongly preferred within subtype A, while F159 is strongly preferred in subtype B. In our analysis, Y159 is a variable fitness amino acid proximal to the subtype A. At site 159, codeml has detected positive selection in subtype A ($\omega = 3.47$), but not in subtype B ($\omega = 0.10$). Within subtype A, F159Y substitutions constitute nearly all (81%) observed substitutions at site 159, so the signal of positive selection picked up by codeml has been created by them, implying recurrent adaptation (Fig. 5A).

By contrast, while S273 was designated a variable fitness amino acid in our analysis, no positive selection has been detected at site 273 by codeml. R273S substitutions constitute only 5.7% of all substitutions at this site in subtype B, while they account for 46.5% of substitutions in the remaining subtypes (including subtype A). Consistently, S273 confers higher fitness in DMS experiments in subtype A than in subtype B. Still, even within subtype A, S273 is just one of the several variants with near-equal fitness (Fig. 5B). Therefore, the increased frequency of R273S substitutions in subtype A is due to relaxation of negative selection against 273S, rather than positive selection in its favour.

A

Figure 5. Substitutions to variable fitness amino acids (red) and to other amino acids (black) at sites 159 (A) and 273 (B) of gp-160, and amino acid propensities in DMS experiments in strains BF520 and LAI. Phylogenetic distances are in numbers of amino acid substitutions per site. Numbering of protein positions is hxb2-based. Logos of experimentally derived amino acid propensities were built with Weblogo (https://weblogo.berkeley.edu/logo.cgi).

Differences in fitness constraints between 3 subtypes

Our approach can be extended to detect amino acids with fitness differing in a predetermined pattern between multiple subtypes. To illustrate this, we study the amino acids that change the conferred fitness between the three major subtypes of HIV-1, A, B and C, which are the most prevalent subtypes in Asia, Northern America/Europe and Africa, respectively. Besides the 59 variable fitness amino acids detected at 46 different sites in the A-B pairwise analysis (see above), we also found 268 additional amino acids that had changed fitness between C and at least one of the subtypes A and B. Overall, we detected 7 amino acids at 5 sites in the A-C comparison, and 285 amino acids at 152 sites in the B-C comparison. These amino acids are listed in Table S1; the web script that lists the variable fitness amino acids for an arbitrarily chosen pattern of preferences between subtypes is available at http://makarich.fbb.msu.ru/galkaklink/hiv_landscape/.

Changes in amino acid propensities may be associated with changes in immunogenicity

To illustrate the potential biological basis of the observed changes in amino acid propensities, we studied how broadly neutralizing antibodies (bnAbs) bind the epitopic sites of gp160. We hypothesized that changes in binding between the major HIV-1 subtypes may be associated with changes in propensities at corresponding sites.

To test this, we considered 9 bnAbs; among those, 6 target the amino acids of gp120 (including VRC01, b12, NIH45-46 which target the CD4-binding site, PG9 which has an epitope in the V1/V2 loop region, and PGT121 and PGT128 that contact the outer domain of gp120 and the V3 loop base); 1 (2G12) targets glycans in the outer domain of gp120; and 2 (2F5 and 4E10) contact with gp41 (Fig. 6A). For each bnAb, we asked if it differed

in inhibition capacity between subtypes. For this, we compared its experimentally measured half maximal inhibitory concentration (IC50) values between subtypes (Doria-Rose et al. 2012; Georgiev et al. 2013). For 5 of the 9 bnAbs, the differences were significant in at least one of the comparisons (Supplementary Table 1).

We then mapped the sites carrying variable fitness amino acids in the A-B (46 sites) or the B-C (152 sites) comparison onto the three-dimensional structure of the gp120/gp41 trimer and its complexes with bnAbs. Among these sites, 7 and 28 respectively were epitopic, i.e., were involved in binding with bnAbs according to (Chuang et al. 2019). (In the A-C comparison, only 5 sites carried variable fitness amino acids, and only one of them was epitopic; these were not analyzed further).

For many of the bnAbs that substantially differed in binding between two subtypes, we found that the associated epitope carried some sites with variable fitness amino acids between these subtypes (Supplementary table 1). For example, the two epitopes that are responsible for differential binding between subtypes A and B (those in the V1/V2 region and the CD4-binding site) both carry amino acids with variable fitness between these two subtypes, while the remaining of the gp120 epitopes (the one in the V3 loop) carries no such amino acids and experiences the same level of binding between the two subtypes (Fig. 6B).

Figure 6. (A) Overview of the structure of the gp120 trimer and its complexes with bnAbs (see Methods). gp120 is colored green and shown as cartoons, one subunit is also shown as a semi-translucent surface. bnAbs are in various color and shown as cartoons. Glycan molecules are shown as sticks. bnAbs binding only one gp120 subunit are shown for clarity. (B) Close-up of one gp120 subunit, with epitopes signed; sites with variable fitness amino acids between subtypes A and B shown as red spheres. Epotopic sites are defined according to (Chuang et al. 2019); CD4-binding site is colored pink, V1/V2 loop region cyan, and V3 loop region purple.

This suggests that shifts in fitness landscapes of epitopic sites may be associated with changes of antigenic properties of these epitopes. Remarkably, in 16 out of 18 sites that carry variable fitness amino acids and reside in epitopes differentially bound by antibodies, the identity of the amino acid that actually occupies the site remains the same between the two subtypes, despite changes in propensities (Supplementary Table 2). This

suggests that the differences in binding may be associated with changes in fitness conferred by the incumbent and/or other amino acid variants without an actual amino acid replacement.

Variable fitness amino acids in HA of Influenza A

We applied our method to the HA gene of H1 and H3 subtypes of human influenza A, and detected 9 variable fitness amino acids at 7 sites (FDR = 39%; Fig. 7).

Figure 7. Amino acid sites carrying variable fitness amino acids in HA of influenza A virus. For each protein position (horizontal axis), the red dot shows the number of amino acids that are simultaneously proximal to the H1 strain and distal from the H3 strain, and the blue dot, the number of amino acids that are simultaneously proximal to the H3 strain and distal from the H1 strain, multiplied by -1. Grey dots represent the number of amino acids at a site that could potentially have significantly variable fitness (see Methods). The positions without any dots are sites filtered out due to alignment ambiguity. Vertical grey lines connect dots for the same site. Protein domains are marked by horizontal lines: *, signal peptide; HA1, HA1 ectodomain; HA2, HA2 ectodomain; TM, transmembrane domain; **, cytoplasmic tail; numeration of protein positions is A/Perth/16/2009 based.

The results for HA were again consistent with the DMS experiment: amino acids with an excess of substitutions into them in the phylogenetic neighborhood of one of the strains conferred higher fitness at this strain than in the other strain (one-tailed Wilcoxon's test, $p = 0.02$; Fig. 8).

Figure 8. The ratio between experimentally measured amino acid preferences of the strain for which an amino acid is more proximal and the strain for which it is more distal in HA of influenza A. Amino acids with significant (left, p<0.05 for both strains) or insignificant (right, p>0.1 for both strains) changes in the phylogenetic analysis are shown; notations are the same as in Figure 4.

For some of the sites, changes in fitness could be interpreted from the functional perspective. For example, site 88 (A/Puerto Rice/8/1934 numbering) is antigenic in H1, but not in H3 (Caton et al. 1982). We suggest that variability in fitness of amino acid L that we found reflects differences in the amino acid propensities at this site between the subtype where it is epitopic (H1; Fig. 9A) and subtype where it is not epitopic (H3; Fig. 9B). Our suggestion is supported by the fact that 7 amino acids can be observed at this site in H1, but only 2 in H3.

Another example is site 202 (in A/Perth/16/2009 numbering) that was previously shown to be antigenic in H3, but not in the H1 subtype (Wiley et al. 1981). At this site, S is the ancestral amino acid for both H1 and H3 clades. Both our analysis and the DMS experiment suggest that P202 confers higher relative fitness in H1 than in H3. Four S202P substitutions have occurred in the H1 clade, and 90% of the H1 isolates contain P202. Conversely, no substitutions into P have occurred in the H3 clade, and 95% of H3 isolates carry G202, which has originated from S in a total of 3 substitutions. The observed differences in the fitness landscapes between subtypes might reflect the differences in the function of site 202 between the two viral subtypes (Fig. 9).

Figure 9. Variable fitness sites in the structure of the Influenza hemagglutinin. Variable fitness amino acids are shown as red spheres except when they are located in one of the antigenic sites. (A) Subtype H1 trimer (PDB id 1ru7). Antigenic sites were mapped based on (Caton et al. 1982): Ca1 is shown in yellow, Ca2 in orange, Cb in magenta, Sa in blue, Sb in green. (B) Subtype H3 trimer (PDB id 2viu). Antigenic sites were mapped based on (Wiley et al. 1981): A is shown in orange, B in green, C in blue, D in yellow.

Discussion

Patterns of substitutions in the course of evolution of a protein are shaped by the selection pressure experienced by it, and changes in this pressure may lead to non-uniformity of substitution rates between evolving lineages. While the overall non-stationarity of

evolutionary rate (heterotachy, Lopez et al. 2002; Yang and Nielsen 2002; Murrell et al. 2012) as well as of individual substitution types (heteropecilly, Tamuri et al. 2009; Roure and Philippe 2011; Klink et al. 2017; Klink and Bazykin 2017) have been used to infer changes in underlying selection, these approaches require pooling multiple sites to obtain a statistical signal. Therefore, they are unable to pick up changes in amino acid propensities within individual sites, especially if the directions of these changes were discordant between sites.

Large sequence datasets with inferable phylogenetic relationships can be harvested to increase the resolution of these methods. Here, we develop an approach to infer, for individual amino acids at individual sites, the changes in relative fitness that they confer between two related lineages.

Our method makes use of changes in relative frequencies of convergent (or parallel) and divergent substitutions between evolving lineages. While the high frequency of substitutions giving rise to the same amino acid suggests a high fitness of that amino acid, usage of divergent substitutions provides an internal control, making the method robust and insensitive to other types of non-stationarity. In particular, first, it is not sensitive to differences in the overall substitution rate (across sites and mutation types) between lineages. Second, as it considers each site individually, it is not sensitive to differences in substitution rates or patterns between sites. Third, as it controls for the identity of the ancestral amino acids, it is not sensitive to differences in rates of different types of substitutions (e.g., transitions vs. transversions) within a site that are conserved throughout the considered phylogeny. Fourth, for the same reason, it is not sensitive to changes in the rate of the substitutions of the same type that are caused by the changes in the prevalence of a particular ancestral amino acid (Klink and Bazykin 2017).

Still, our method has some limitations (Klink and Bazykin 2017). First, as it is based on the analysis of phylogenetic distribution of amino acid substitutions, its power is limited by the number of substitutions observed, which in turn depends on the tree shape and the selection regime. In particular, it is inadequate for detection of changes in amino acid

fitness between closely related species or at slowly evolving sites. Here, we explicitly study the power of our method for different selection strengths. Simulations show that our method works best within a window of selection coefficients corresponding to moderate selection. This is to be expected. If selection is too weak, both weakly advantageous and weakly deleterious substitutions occur throughout the phylogeny, making our approach impotent. Conversely, if selection is very strong, the number of variable sites becomes low: the favored amino acid is usually nearly fixed in one of the clades, so that there are no substitutions giving rise to it to analyze. In particular, our method is inadequate for detection of selection in favour of a highly conserved ancestral variant, since multiple substitutions into this variant are unlikely. Furthermore, the power of our method increases when sequences of more species are available (Fig. 2). This happens for two reasons: first, an increase in the number of substitutions gives more power; second, even with the same number of substitutions at a site, the power is higher for sites with more distinct ancestral amino acids, as at such sites, there are more combinations of substitutions in the null distribution, and thus a lower minimal p-value can be obtained.

Second, our test detects any change in the frequency of an individual substitution at a site independent of its cause. While at the amino acid level, such changes are probably always caused by changes in amino acid propensities, if a similar approach is applied to nucleotide substitutions, rate changes can also occur due to differences in mutation biases or in non-amino-acid level selection, e.g. selection for nucleotide composition. In theory, synonymous sites can be used to control for non-selective changes in substitution biases, although this is not straightforward for viruses where many of such sites can be constrained by selection on structure or overlapping reading frames. However, mutation spectra are similar between HIV-1 subtypes (Rawson et al. 2017), strongly suggesting that the changes we detect are caused by differences in selection rather than mutation.

As the probability of a substitution to an amino acid depends on its N_e-scaled relative fitness, theoretically, our method could be sensitive to differences in N_e between subtypes. We are unaware of any data showing systematic differences in N_e between HIV

or influenza A subtypes. More generally, changes in N_e should affect all sites concordantly, and it is difficult to imagine a scenario whereby they will lead to changes in density of homoplasies of opposite directions between sites, as observed in our analysis.

To illustrate and experimentally validate our approach, we make use of two DMS datasets for viral surface proteins, in each of which the amino acid preferences have been measured experimentally for two different strains: the gp160 protein in two HIV-1 strains, and the HA protein in two influenza A strains. Overall, we find that the results of the phylogenetic test are consistent with experimental results. If a particular amino acid is gained more frequently by a species closely related to a particular strain, it also tends to confer higher fitness in experimental assays in that same strain.

We find that the inferred variable fitness amino acids mostly have modest experimentally measured preference π even in the strain for which they are proximal. There can be multiple reasons for this. First, our phylogenetic test is not guaranteed to assign high preference to the fittest amino acid at a site. In particular, it is only able to assign differential fitness to derived, but not ancestral, variants. For example, 159F confers high experimentally measured fitness in the gp160 of the LAI strain, but its fitness is lower in the BF520 strain (Fig. 5A); still, it cannot be picked up by the phylogenetic test, as it is the ancestral variant, and the substitutions giving rise to it are rare in both strains (Fig. 5A). As the variant that is much more fit than all other variants is likely to be conserved throughout the clade, the phylogenetic method is less likely to pick it up, compared to a variant with fitness comparable to that of at least one other variant.

Second, many of the fitness shifts between A and B subtypes of HIV-1 did not affect the fittest amino acid. Indeed, among the 42 amino acids with significant phylogenetically inferred fitness shifts between these subtypes, only 8 (19%) had the highest DMS-inferred fitness rank in either strain (Fig. 4C). The remaining 34 amino acids were suboptimal in both strains. Therefore, our method allows to infer the changes in fitness between

suboptimal amino acids, which is relevant for understanding the overall fitness landscape governing the evolution of the protein.

An evolutionary model that incorporates the results of a DMS scan (ExpCM; Hilton et al. 2017) has been shown to outperform other substitution models, and, in particular, detect positive selection more robustly. This is achieved by obtaining the strength of functional constraint, indicative of negative selection, for individual sites from the DMS data (Haddox et al. 2018; Lee et al. 2018). In line with that model, the changes in substitution rates observed here can be caused by changes in the strength of negative selection between strains. Additionally, changes in amino acid preferences can trigger episodes of positive selection if the newly preferred amino acid is rare in the ancestral population. We find that less than a half of sites with variable fitness amino acids also had signatures of positive selection that could be detected by a standard dN/dS approach. Therefore, changes in site-specific fitness landscape are not necessarily coupled with strong positive selection favouring new optimal variants.

Our method can be applied to more than two focal species to find amino acids that have the same relative fitness in some lineages but change it in other lineages. We illustrate this using the three subtypes of HIV-1. As the number of strains substantially differs between these three clades (223, 2035 and 1265 for A, B and C correspondingly), the power of our method differs between them. Still, our results show that with enough data we are able to compare fitness of an amino acid between more than two clades. Importantly, unlike other methods for detection of changes in selection preferences, our test can be applied without prior identification of candidate clades differing in selection pressure (Tamuri et al. 2009), and makes no assumption regarding model of epistatic interactions or fitness shift (Tamuri et al. 2009; Louie et al. 2018).

Predicting the effects of genetic variation on fitness is often based on the extent of conservation of a position (Sim et al. 2012; Adzhubei et al. 2013; Kircher et al. 2014), but functionally important residues are not necessarily conserved (Wu et al. 2015). Our method can be used to distinguish between rapidly evolving sites with relaxed and

variable constraints. For example, in gp160, there are 10 sites each with 18 different amino acids that originated at some branch(es) of the phylogenetic tree. While this is suggestive of complete neutrality of these sites, we find that, among these 10 sites, 5 carry variable fitness amino acids.

Despite the recent surge in DMS experiments that allow high-throughput fitness measurements, such experiments can still only be performed for a relatively few model systems, and are still impractical for large numbers of strains. Moreover, the fitness measurements in experimental conditions are not necessarily reflective of the selective pressure experienced by a lineage over the course of its evolution (Haddox et al. 2016). By contrast, comparative genomics allows to assess the fitness differences that have shaped evolution for a large number of non-model species simultaneously.

Materials and Methods

Inference of phylogenetic biases

We define the phylogenetic distance d between the focal species and each substitution into amino acid A as the sum of branch lengths along the path between them on the phylogeny, assuming that a substitution occurs at the middle of the corresponding branch, and $\bar{d}$ as the mean of those distances. If the substitutions preferentially occur in the phylogenetic neighborhood of the focal species, $\bar{d}$ will be lower than expected. By contrast, if substitutions are underrepresented in the phylogenetic neighborhood of the focal species, $\bar{d}$ will be higher than expected. To obtain the null distribution of $\bar{d}$, we make use of the substitutions into non-A amino acids. $\bar{d}$ depends both on the phylogenetic distribution of the amino acids ancestral to A and on the number of substitutions into A (Klink and Bazykin 2017). To control for this, we subsampled the substitutions from the same ancestral amino acids as follows. We listed all "ancestral" amino acids that were substituted by A at least once. For each such ancestral amino acid, we then considered all substitutions into any (A or non-A) amino acid, and subsampled, out of these substitutions, the number of substitutions equal to that of substitutions into A. We then pooled these

substitutions across all ancestral amino acids together, and measured the mean phylogenetic distance $\tilde{d}$ for them. This subsampling was repeated 10,000 times to obtain the null distribution of $\bar{d}$.

The p-value for the bias of substitutions towards the focal species was defined as the percentile of $\bar{d}$ in this distribution, and for their bias away from the focal species, as 1 minus this value. We required p < α for proximal amino acids, and p > (1-α) for distal amino acids, where α is the significance threshold. An amino acid was considered "potentially significant" (proximal or distal) if the minimal attainable p-value (i.e., the fraction of the values of $\bar{d}$ in the most extreme bin) was lower than α. For each significance threshold α, we calculated the expected fraction of false positives (false discovery rate, FDR) as argmin ((α*PotVar)/VarFit, 1), where PotVar is the number of potentially significant amino acids at this α, and VarFit is the number of significant amino acids at this α. We used α=0.01 for gp160 throughout; the percent of expected false positives at this p-value is less than 11% (Supplementary figure 3). For HA, where the substitution rate is lower, we used α=0.05, yielding the FDR<39%.

Finally, for a pair of species 1 and 2, we defined an amino acid as "variable fitness amino acid" if it was proximal for one species and distal for the other one. All amino acids that were potentially significantly proximal for one species and potentially significantly distal for the other were considered as "potentially variable fitness amino acids".

Data

gp-160

We downloaded 4241 gp160 DNA and protein sequences belonging to nine subtypes of HIV-1 from the Los Alamos HIV sequence database (http://www.hiv.lanl.gov/) using the following settings: alignment type=filtered web, organism=HIV-1/SIVcpz, region=Env, subtype=M group without recombinants, year: 2016. The subtypes were represented unequally, with the majority of sequences from subtypes B (>2000 isolates) and C (>1000 isolates). We discarded sequences with lengths not divisible by 3 and internal stop codons, leaving 3789 sequences. We then added two experimentally studied sequences

LAI and BF520, and an outgroup type O (isolate_ANT70), for a total of 3792 sequences. Using Pal2Nal (Suyama et al. 2006) and Mafft (Katoh et al. 2002), we obtained amino acid alignments and codon-informed nucleotide sequence alignments. Using nucleotide alignments, we built a maximum likelihood phylogenetic tree using RAxML under the GTRGAMMA model. We then optimized branch lengths on the basis of the amino acid alignment under the PROTGAMMAGTR model (Stamatakis 2014) and reconstructed ancestral states with codeml program of the PAML package (Yang 2007).

The analysis was performed on amino acid alignments, using LAI or BF520 as focal species. To ensure that our results are robust to uncertainty of phylogenetic reconstruction, for all analyses involving one "proximal-distal" comparison, we constructed maximum likelihood phylogenetic trees on the basis of the same alignment in 10 independent runs, identified the variable fitness amino acids for each tree, and performed downstream analyses with just those amino acids that had significant shifts in fitness in all 10 replicates.

Functional regions were defined as annotated in UniProt (The UniProt Consortium 2017; entry P04578). We obtained solvent accessibility data for the LAI strain from (Haddox et al. 2016). To define buried and exposed residues, we used the threshold relative solvent accessibility = 0.25.

Hemagglutinin

We obtained 2238 and 50,109 DNA sequences of HA from seasonal H1N1 and H3N2 influenza A subtypes correspondingly from the GISAID database (Shu and McCauley 2017). We discarded sequences shorter than 1680 nucleotides, with lengths not divisible by 3 or with internal stop codons, leaving 2142 sequences of H1 and 49543 sequences of H3. We then used cd-hit program (Li and Godzik 2006) to discard identical sequences from the dataset, leaving 1557 sequences of H1 an 20191 sequences of H3. Finally, we randomly subsampled 1557 sequences of H3 to make the H1 and H3 datasets equal in size. Protein sequences were obtained from DNA sequences with BioPerl scripts. All subsequent procedures were conducted as for the HIV-1 dataset. We obtained

experimentally derived amino acid propensities for A/WSN/1933 (H1N1) and A/Perth/16/2009 (H3N2) strains from (Lee et al. 2018).

Simulated evolution

We simulated evolution of the protein across the phylogeny of two HIV-1 subtypes, B and C. The phylogeny was constructed as above, but only sequences from B and C subtypes were used. The "sparse" tree was constructed by removing a fraction of terminal branches from the real tree, and the "dense" tree, by adding, to each terminal node of the initial tree, two new branches ending with new terminal nodes, with lengths uniformly picked from the range (0;0.2]. The trees were processed using custom BioPerl scripts.

Simulations were performed with the Simulator of Evolution with Landscape Variation (SELVa) program (Nabieva and Bazykin 2019). SELVa allows to specify the branches of the phylogenetic tree where a fitness shift occurs, and the vectors of amino acid fitness before and after the shift. It simulates substitutions between amino acids according to the model described in (Yang and Nielsen 2008). The mutation rates are assumed to be uniform, so the differences in substitution rates are solely determined by differences in fitness.

We simulated evolution of 500 sites along the phylogenetic tree of HIV-1 with the amino acid fitness vector $\{X,1,1,1,1,1,1,1,1,1,1,1,1,1,1,1,1,1,1,1\}$, for each integer X between 3 and 10 . The identity of the amino acid with fitness X differed between subtypes B and C; the corresponding fitness shift was assumed to have occurred in the last common ancestor of subtype C.

For each X, we calculated two values: the true positive rate (TPR) – the proportion of amino acids, among those that have changed fitness in the course of evolution, that were correctly identified as positives; and the false positive rate (FPR) – the proportion of amino acids, among those that have not changed fitness, that were incorrectly identified as positives. These values were calculated for 1000 randomly sampled amino acids for 100 times to obtain the mean and the 90% confidence interval.

Comparison with DMS data

To test whether our results match the results of the DMS experiments, for each amino acid, we measured the ratio r between the DMS preference for this amino acid in the strain for which it is phylogenetically more proximal (π_x) and in the strain for which it is more distal (π_y). The proximity to one or the other strain was determined by comparison of the corresponding p-values, independent of their significance, so that all amino acids could be thus classified. We then compared the mean r for variable fitness and other amino acids with Wilcoxon's rank sum test.

Identifying sites under positive selection

Sites of gp160 were tested for positive selection using the codeml program of the PAML package with site model. The presence of positive selection was tested by comparison of models M1a and M2a with the likelihood ratio test. Probabilities that positive selection acts at each site were calculated by BEB (Bayes Empirical Bayes) method as implemented in codeml. We separately performed the analysis for subtype A subtree, subtype B subtree and A+B tree. The trees for these analysis were built as described above, but only sequences from subtypes A and/or B were used.

Studying association of variable fitness amino acids with differences of antigenic properties between HIV1 subtypes

We used IC50 values for 19 subtype A strains, 31 subtype B strains, and 28 subtype C strains for bnAbs VRC01, NIH45-46, b12, PG9, 2G12, 2F5, and 4E10 from (Georgiev et al. 2013); for 8 subtype A strains, 13 subtype B strains, and 12 subtype C strains for bnAbs PGT121 and PGT128 also from (Georgiev et al. 2013); and for 31 subtype A strains, 48 subtype B strains, and 59 subtype C strains for bnAbs VRC01 and PG9 from (Doria-Rose et al. 2012). For each bnAb, we performed a two-sided Mann-Whitney test to understand whether its neutralising capacity differs between subtypes A and B or subtypes B and C. Then we mapped sites with the variable fitness amino acids onto the overlay of multiple gp120/bnAb complexes: with VRC01 (PDB id 3ngb), PG9 (PDB id 3u4e), b12 (PDB id

2ny7), NIH45-46 (PDB id 4jkp), 2G12 (PDB id 6e5p), PGT121 (PDB id 5c7k), and PGT128 (PDB id 5cez). Epitopic sites were defined according to (Chuang et al. 2019). To show all considered antibodies in connection with gp120 on one picture (Fig. 6A), we combined structures 6e5p, 4jkp, 3ngb, 2nz7, 5cez and 5c7k.

Acknowledgments

This work was supported by the Russian Foundation for Basic Research grant no. 18-34-00358 and by the Molecular and Cellular Biology program of the Russian Academy of Sciences. We thank Juhye Lee, Hugh Haddox and Jesse Bloom for sharing their DMS data pre-publication and for valuable comments. We also thank Alexey Neverov for helpful discussions.

References

Adzhubei I, Jordan DM, Sunyaev SR. 2013. Predicting functional effect of human missense mutations using PolyPhen-2. Curr. Protoc. Hum. Genet. Chapter 7:Unit7.20.

Bazykin GA. 2015. Changing preferences: deformation of single position amino acid fitness landscapes and evolution of proteins. Biol. Lett. [Internet] 11:20150315. Available from: http://rsbl.royalsocietypublishing.org/lookup/doi/10.1098/rsbl.2015.0315

Bazykin GA, Dushoff J, Levin SA, Kondrashov AS. 2006. Bursts of nonsynonymous substitutions in HIV-1 evolution reveal instances of positive selection at conservative protein sites. Proc. Natl. Acad. Sci. U. S. A. 103:19396–19401.

Blackburne BP, Hay AJ, Goldstein RA. 2008. Changing selective pressure during antigenic changes in human influenza H3. PLoS Pathog. 4:e1000058.

Caton AJ, Brownlee GG, Yewdell JW, Gerhard W. 1982. The antigenic structure of the influenza virus A/PR/8/34 hemagglutinin (H1 subtype). Cell 31:417–427.

Chuang G-Y, Zhou J, Acharya P, Rawi R, Shen C-H, Sheng Z, Zhang B, Zhou T, Bailer RT, Dandey VP, et al. 2019. Structural Survey of Broadly Neutralizing Antibodies Targeting the HIV-1 Env Trimer Delineates Epitope Categories and Characteristics of Recognition. Struct. Lond. Engl. 1993 27:196-206.e6.

Doria-Rose NA, Louder MK, Yang Z, O'Dell S, Nason M, Schmidt SD, McKee K, Seaman MS, Bailer RT, Mascola JR. 2012. HIV-1 neutralization coverage is improved by combining monoclonal antibodies that target independent epitopes. J. Virol. 86:3393–3397.

Elena SF, Agudelo-Romero P, Lalić J. 2009. The evolution of viruses in multi-host fitness landscapes. Open Virol. J. 3:1–6.

Figliuzzi M, Jacquier H, Schug A, Tenaillon O, Weigt M. 2016. Coevolutionary Landscape Inference and the Context-Dependence of Mutations in Beta-Lactamase TEM-1. Mol. Biol. Evol. 33:268–280.

Georgiev IS, Doria-Rose NA, Zhou T, Kwon YD, Staupe RP, Moquin S, Chuang G-Y, Louder MK, Schmidt SD, Altae-Tran HR, et al. 2013. Delineating antibody recognition in polyclonal sera from patterns of HIV-1 isolate neutralization. Science 340:751–756.

Goldstein RA, Pollard ST, Shah SD, Pollock DD. 2015. Nonadaptive Amino Acid Convergence Rates Decrease over Time. Mol. Biol. Evol. [Internet] 32:1373–1381. Available from: https://academic.oup.com/mbe/article-lookup/doi/10.1093/molbev/msv041

Haddox HK, Dingens AS, Bloom JD. 2016. Experimental Estimation of the Effects of All Amino-Acid Mutations to HIV's Envelope Protein on Viral Replication in Cell Culture.Swanstrom R, editor. PLOS Pathog. [Internet] 12:e1006114. Available from: http://dx.plos.org/10.1371/journal.ppat.1006114

Haddox HK, Dingens AS, Hilton SK, Overbaugh J, Bloom JD. 2018. Mapping mutational effects along the evolutionary landscape of HIV envelope. eLife [Internet] 7. Available from: https://elifesciences.org/articles/34420

Hilton SK, Doud MB, Bloom JD. 2017. phydms: software for phylogenetic analyses informed by deep mutational scanning. PeerJ 5:e3657.

Hopf TA, Ingraham JB, Poelwijk FJ, Schärfe CPI, Springer M, Sander C, Marks DS. 2017. Mutation effects predicted from sequence co-variation. Nat. Biotechnol. 35:128–135.

Jordan DM, Frangakis SG, Golzio C, Cassa CA, Kurtzberg J, Task Force for Neonatal Genomics, Davis EE, Sunyaev SR, Katsanis N. 2015. Identification of cis-suppression of human disease mutations by comparative genomics. Nature 524:225–229.

Katoh K, Misawa K, Kuma K, Miyata T. 2002. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. Nucleic Acids Res. 30:3059–3066.

Kimura, M. 1983. The Neutral Theory of Molecular Evolution. Cambridge: Cambridge University Press

Kircher M, Witten DM, Jain P, O'Roak BJ, Cooper GM, Shendure J. 2014. A general framework for estimating the relative pathogenicity of human genetic variants. Nat. Genet. 46:310–315.

Klink GV, Bazykin GA. 2017. Parallel Evolution of Metazoan Mitochondrial Proteins. Genome Biol. Evol. [Internet] 9:1341–1350. Available from: https://academic.oup.com/gbe/article-lookup/doi/10.1093/gbe/evx025

Klink GV, Golovin AV, Bazykin GA. 2017. Substitutions into amino acids that are pathogenic in human mitochondrial proteins are more frequent in lineages closely related to human than in distant lineages. PeerJ [Internet] 5:e4143. Available from: https://peerj.com/articles/4143

Kondrashov AS, Sunyaev S, Kondrashov FA. 2002. Dobzhansky-Muller incompatibilities in protein evolution. Proc. Natl. Acad. Sci. U. S. A. 99:14878–14883.

Kryazhimskiy S, Dushoff J, Bazykin GA, Plotkin JB. 2011. Prevalence of epistasis in the evolution of influenza A surface proteins. PLoS Genet. 7:e1001301.

Lee JM, Huddleston J, Doud MB, Hooper KA, Wu NC, Bedford T, Bloom JD. 2018. Deep mutational scanning of hemagglutinin helps predict evolutionary fates of human H3N2 influenza variants. Proc. Natl. Acad. Sci. U. S. A. 115:E8276–E8285.

Li W, Godzik A. 2006. Cd-hit: a fast program for clustering and comparing large sets of protein or nucleotide sequences. Bioinformatics [Internet] 22:1658–1659. Available from: https://academic.oup.com/bioinformatics/article-lookup/doi/10.1093/bioinformatics/btl158

Loewen CA, Ganetzky B. 2018. Mito-Nuclear Interactions Affecting Lifespan and Neurodegeneration in a Drosophila Model of Leigh Syndrome. Genetics 208:1535–1552.

Lopez P, Casane D, Philippe H. 2002. Heterotachy, an important process of protein evolution. Mol. Biol. Evol. 19:1–7.

Louie RHY, Kaczorowski KJ, Barton JP, Chakraborty AK, McKay MR. 2018. Fitness landscape of the human immunodeficiency virus envelope protein that is targeted by antibodies. Proc. Natl. Acad. Sci. U. S. A. 115:E564–E573.

Mann JK, Barton JP, Ferguson AL, Omarjee S, Walker BD, Chakraborty A, Ndung'u T. 2014. The fitness landscape of HIV-1 gag: advanced modeling approaches and validation of model predictions by in vitro testing. PLoS Comput. Biol. 10:e1003776.

Murrell B, Wertheim JO, Moola S, Weighill T, Scheffler K, Kosakovsky Pond SL. 2012. Detecting individual sites subject to episodic diversifying selection. PLoS Genet. 8:e1002764.

Nabieva E, Bazykin GA. 2019. SELVa: Simulator of Evolution with Landscape Variation. bioRxiv [Internet]. Available from: http://biorxiv.org/lookup/doi/10.1101/647834

Naumenko SA, Kondrashov AS, Bazykin GA. 2012. Fitness conferred by replaced amino acids declines with time. Biol. Lett. [Internet] 8:825–828. Available from: http://rsbl.royalsocietypublishing.org/cgi/doi/10.1098/rsbl.2012.0356

Popova, A. V., Safina, K. R., Ptushenko, V. V., Stolyarova, A. V., Favorov, A. V., Neverov, A. D., & Bazykin, G. A. 2019. Allele-specific nonstationarity in evolution of influenza A virus surface proteins. Proceedings of the National Academy of Sciences, 116(42), 21104-21112.

Povolotskaya IS, Kondrashov FA. 2010. Sequence space and the ongoing expansion of the protein universe. Nature [Internet] 465:922–926. Available from: http://www.nature.com/articles/nature09105

Rawson JMO, Gohl DM, Landman SR, Roth ME, Meissner ME, Peterson TS, Hodges JS, Beckman KB, Mansky LM. 2017. Single-Strand Consensus Sequencing Reveals that HIV Type but not Subtype Significantly Impacts Viral Mutation Frequencies and Spectra. J. Mol. Biol. 429:2290–2307.

Riesselman AJ, Ingraham JB, Marks DS. 2018. Deep generative models of genetic variation capture the effects of mutations. Nat. Methods 15:816–822.

Roure B, Philippe H. 2011. Site-specific time heterogeneity of the substitution process and its impact on phylogenetic inference. BMC Evol. Biol. 11:17.

Shu Y, McCauley J. 2017. GISAID: Global initiative on sharing all influenza data – from vision to reality. Eurosurveillance [Internet] 22. Available from: http://www.eurosurveillance.org/ViewArticle.aspx?ArticleId=22750

Sim N-L, Kumar P, Hu J, Henikoff S, Schneider G, Ng PC. 2012. SIFT web server: predicting effects of amino acid substitutions on proteins. Nucleic Acids Res. 40:W452-457.

Soylemez O, Kondrashov FA. 2012. Estimating the rate of irreversibility in protein evolution. Genome Biol. Evol. 4:1213–1222.

Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. Bioinformatics [Internet] 30:1312–1313. Available from: https://academic.oup.com/bioinformatics/article-lookup/doi/10.1093/bioinformatics/btu033

Storz JF. 2016. Causes of molecular convergence and parallelism in protein evolution. Nat. Rev. Genet. [Internet] 17:239–250. Available from: http://www.nature.com/doifinder/10.1038/nrg.2016.11

Stray SJ, Pittman LB. 2012. Subtype- and antigenic site-specific differences in biophysical influences on evolution of influenza virus hemagglutinin. Virol. J. 9:91.

Suyama M, Torrents D, Bork P. 2006. PAL2NAL: robust conversion of protein sequence alignments into the corresponding codon alignments. Nucleic Acids Res. 34:W609-612.

Tamuri AU, Dos Reis M, Hay AJ, Goldstein RA. 2009. Identifying changes in selective constraints: host shifts in influenza. PLoS Comput. Biol. 5:e1000564.

The UniProt Consortium. 2017. UniProt: the universal protein knowledgebase. Nucleic Acids Res. 45:D158–D169.

Wiley DC, Wilson IA, Skehel JJ. 1981. Structural identification of the antibody-binding sites of Hong Kong influenza haemagglutinin and their involvement in antigenic variation. Nature 289:373–378.

Wood N, Bhattacharya T, Keele BF, Giorgi E, Liu M, Gaschen B, Daniels M, Ferrari G, Haynes BF, McMichael A, et al. 2009. HIV evolution in early infection: selection pressures, patterns of insertion and deletion, and the impact of APOBEC. PLoS Pathog. 5:e1000414.

Wu NC, Olson CA, Du Y, Le S, Tran K, Remenyi R, Gong D, Al-Mawsawi LQ, Qi H, Wu T-T, et al. 2015. Functional Constraint Profiling of a Viral Protein Reveals Discordance of Evolutionary Conservation and Functionality. PLoS Genet. 11:e1005310.

Yang W, Bielawski JP, Yang Z. 2003. Widespread Adaptive Evolution in the Human Immunodeficiency Virus Type 1 Genome. J. Mol. Evol. [Internet] 57:212–221. Available from: http://link.springer.com/10.1007/s00239-003-2467-9

Yang Z. 2007. PAML 4: phylogenetic analysis by maximum likelihood. Mol. Biol. Evol. 24:1586–1591.

Yang Z, Nielsen R. 2002. Codon-substitution models for detecting molecular adaptation at individual sites along specific lineages. Mol. Biol. Evol. 19:908–917.

Yang Z, Nielsen R. 2008. Mutation-selection models of codon substitution and their use to estimate selective strengths on codon usage. Mol. Biol. Evol. 25:568–579.

Zaidi AA, Makova KD. 2019. Investigating mitonuclear interactions in human admixed populations. Nat. Ecol. Evol. 3:213–222.

Zou Z, Zhang J. 2015. Are Convergent and Parallel Amino Acid Substitutions in Protein Evolution More Prevalent Than Neutral Expectations? Mol. Biol. Evol. [Internet] 32:2085–2096. Available from: https://academic.oup.com/mbe/article-lookup/doi/10.1093/molbev/msv091