Stoichiometric constraints modulate the effects of temperature and nutrients on biomass

distribution and community stability

Arnaud Sentis[1,2*], Bart Haegeman[1], and José M. Montoya[1]

[1]Centre for Biodiversity Theory and Modelling, Theoretical and Experimental Ecology Station,

CNRS and Paul Sabatier University, Moulis, France.

[2] IRSTEA, Aix Marseille Univ., UMR RECOVER, 3275 route Cézanne, 13182 Aix-en-Provence,

France.

*: Corresponding author. Arnaud Sentis. IRSTEA, Aix Marseille Univ., UMR RECOVER, 3275

route Cézanne, 13182 Aix-en-Provence, France.

Email: arnaud.sentis@irstea.fr;

Author contributions

A.S., B.H., and J.M.M. conceived the study. B.H. and A.S. developed and analysed the models.

A.S. wrote the first draft of the manuscript. All authors contributed substantially to revisions.

Running title: Stoichiometry alters thermal effects in food webs

Key-words: temperature, stoichiometry, temporal variability, paradox of enrichment, trophic

interactions, nutrient quota, consumer-resource dynamics, biomass structure.

Legal and ethical requirements

The authors declare that they have no conflict of interest.

Abstract

Temperature and nutrients are two of the most important drivers of global change. Both can

modify the elemental composition (i.e. stoichiometry) of primary producers and consumers. Yet

their combined effect on the stoichiometry, dynamics, and stability of ecological communities

remains largely unexplored. To fill this gap, we extended the Rosenzweig-MacArthur consumer-

resource model by including thermal dependencies, nutrient dynamics, and stoichiometric

constraints on both the primary producer and the consumer. We found that stoichiometric

constraints dampen the paradox of enrichment and increased persistence at high nutrient levels.

Nevertheless, they also reduced consumer persistence at extreme temperatures. Finally, we also

found that stoichiometric constraints can strongly influence biomass distribution across trophic

levels by modulating consumer assimilation efficiency and resource growth rates along the

environmental gradients. Our findings highlight the importance of accounting for stoichiometric

constraints as they can mediate the temperature and nutrient impact on the dynamics and

functioning of ecological communities.

Key words: temperature, stoichiometry, temporal variability, paradox of enrichment, trophic

interactions, nutrient quota, consumer-resource dynamics, biomass structure.

Introduction

Temperature and nutrients regulate many biological processes, including species geographical

distribution, primary production, species interactions, and energy and material fluxes (Falkowski

et al. 1998; Enquist *et al.* 1999; Elser *et al.* 2007; Thomas *et al.* 2017). They are at the core of

several ecological theories. While temperature is a fundamental component of metabolic scaling

theory (Brown *et al.* 2004), nutrients are at the core of resource competition theory (Tilman 1982)

and ecological stoichiometry (i.e. the element composition of organisms) theory (Sterner & Elser

2002). Cross et al. (2015) suggested that a better understanding of the interactions between

temperature and nutrients is crucial for developing realistic predictions about ecological

responses to multiple drivers of global change, including climate warming and elevated nutrient

supply. Nutrients can modulate the effects of warming on communities directly by altering

primary production, and/or indirectly by changing the elemental composition of primary

producers. Conversely, thermal effects on trophic interaction strengths (i.e. the *per capita* effect

of predators on prey population densities) and on consumer energetic efficiencies (i.e. ingestion

relative to metabolic demand) depend on both the quantity and quality of their resources. While

Cross et al. (2015) provided a road map on how to investigate the combined effects of

temperature and nutrient on ecological processes, we still lack an integrative theory to better

understand how the links between stoichiometry, nutrient enrichment, and temperature influence

the dynamics and stability of multispecies communities. Such a theory will allow us to

understand how and when stoichiometric variation modulates the consequences of single and

combined components of global change on trophic interactions, community dynamics, and

ecosystem functioning.

66
Predicting the effects of global warming and nutrient changes on ecosystems is challenging as

species are embedded within communities of multiple interacting species (Petchey *et al.* 1999;

Tylianakis *et al.* 2008; Montoya & Raffaelli 2010; Gilbert *et al.* 2014). Increased resource

availability (hereafter: enrichment) and warming can jointly affect food-web stability and

structure by modifying the strength of trophic interactions (O'Connor *et al.* 2009; Binzer *et al.*

2012; Kratina *et al.* 2012; Sentis *et al.* 2014; Binzer *et al.* 2016). Enrichment typically increases

energy flux from resources to higher trophic levels which often leads to the well-known paradox

of enrichment where the amplitude of population fluctuations increase with nutrients, leading to

extinctions at high nutrient concentrations (Rosenzweig 1971; Rip & McCann 2011; Gilbert *et al.*

2014). Nevertheless, most consumer species become less efficient at processing matter and

energy at warmer temperatures as their metabolic rates often increase faster with temperature

than their feeding rates (Vucic-Pestic *et al.* 2011; Fussmann *et al.* 2014; Iles 2014). This

reduction of energetic efficiency lessens energy flow between trophic levels and hence stabilizes

food-web dynamics by reducing population fluctuations (Rip & McCann 2011; Binzer *et al.*

2012; Gilbert *et al.* 2014). As a result, mild warming may alleviate the paradox of enrichment by

decreasing consumer energetic efficiency (Binzer *et al.* 2012; Sentis *et al.* 2017).

The theoretical expectations and results described above have already improved our ability to

understand and predict the effects of temperature and enrichment on food webs (Boit *et al.* 2012;

Tabi *et al.* 2019). However, most previous studies using metabolic scaling theory assumed that

nutrient enrichment lead to an increase in resource carrying capacity without influencing resource

elemental composition (Vasseur & McCann 2005; Binzer *et al.* 2012; Gilbert *et al.* 2014; Binzer

et al. 2016; Sentis *et al.* 2017). Yet nutrient enrichment effects are more complex. The elemental

composition of primary producers is likely to be altered, in response to the supplies of energy and

materials relative to their growth and nutrient intake rates (Rastetter *et al.* 1997;

Robert W. Sterner *et al.* 1997; Finkel *et al.* 2009) Sterner et al. 1997). This, in turn, can affect the

dynamics of the producer population and the herbivores feeding on it. For instance, previous

modelling studies showed that introducing stoichiometric heterogeneity in predator-prey

population dynamic models can dampen the negative effect of nutrient enrichment on system

persistence by reducing population biomass fluctuations (Andersen 1997; Loladze *et al.* 2000;

Andersen *et al.* 2004; Elser *et al.* 2012). More generally, the stoichiometric flexibility of primary

producers, in particular the flexibility in carbon to nutrient ratios (e.g. C:N or C:P), has important

implications for animal feeding behaviour (White 1993), consumer population stability (White

1993; Sterner & Hessen 1994; Hessen *et al.* 2002), community structure (Andersen 1997), and

ecosystem processes such as biogeochemical cycling (Andersen 1997; Hessen *et al.* 2004).

Previous theoretical and empirical studies reported that stoichiometric variations can have a

strong influence on the stability of consumer-resource interactions (Andersen 1997; Andersen *et*

al. 2004; Diehl *et al.* 2005; Elser *et al.* 2012). For instance, populations of crustacean *Daphnia*

feeding on low quality (i.e. low nutrient: carbon ratio) algae cannot persist even when resource

quantity is not a limiting factor (Elser *et al.* 2007). Consumer extinction is explained by the fact

that the consumer assimilation efficiency is, for most organisms, a function of resource quality

(Elser *et al.* 2000). When resource quality is low, the consumers assimilate only few nutrients

relative to the biomass they ingest, which limits their growth and reproduction (Elser *et al.* 2000;

Elser *et al.* 2012). Temporal variations in resource quality can stabilize the system by weakening

interaction strength and dampening population fluctuations (Andersen *et al.* 2004; Diehl *et al.*

2005)but see(Loladze *et al.* 2000; Elser *et al.* 2012). However, it remains unclear whether and

how temporal variations in the elemental composition of primary producers and consumers can

modulate the effects of temperature and nutrients on important community features such as

stability and biomass distribution across trophic levels. Previous studies indicated that the spatial

and temporal intraspecific variations in the elemental composition of primary producers are

expected to increase in response to global change drivers such as temperature, CO_2, and nutrient

availability (Bezemer & Jones 1998; Woods *et al.* 2003; Finkel *et al.* 2009). This increased

variation can be of importance for both primary producer and consumer populations as the

growth rate of primary producers is well known to depend on their elemental composition (Droop

1974) as is the assimilation efficiency of the consumers (Sterner & Elser 2002).

Altogether, previous studies indicated that both temperature and stoichiometric variations can

have important effects on species interactions and community dynamics (Andersen *et al.* 2004;

Diehl *et al.* 2005; Fussmann *et al.* 2014; Binzer *et al.* 2016; Sentis *et al.* 2017). However, the

effects of temperature and nutrient stoichiometry on food web dynamics and stability have only

been studied in isolation. Recent theory by Uszko et al. (2017) showed that considering nutrient

dynamics can help to better understand the influence of temperature on consumer-resource

population dynamics and resource carrying capacity. Nevertheless, they considered that the

elemental composition of both the resource and the consumer are constant and independent of

temperature and nutrient dynamics. This contrasts with the empirical observation that resource

elemental composition is flexible and can vary with both temperature and nutrient dynamics

(Droop 1974; Elser *et al.* 2000; Woods *et al.* 2003). Here we thus focused on the combined

effects of temperature and nutrients on the stoichiometry of primary producers and how this

affects community stability and biomass distribution across trophic levels in a consumer-resource

system. Understanding the determinants of stability and biomass distribution has been at the core

of ecology for a long time (Elton (1927), Lindeman (1942)). Recent theory aims at explaining

empirical observations of trophic pyramids (i.e. population biomass decreases with trophic

levels), inverted trophic pyramids (i.e. population biomass increases with trophic levels), trophic

cascades and the link between biomass distribution and stability (McCauley *et al.* 2018; Barbier

& Loreau 2019).

Here, we used the Rosenzweig-MacArthur model as a baseline non-stoichiometric model because

this model is one of the most studied models used to investigate the effects of temperature and

nutrient enrichment on community dynamics (Vasseur & McCann 2005; Binzer *et al.* 2012;

Fussmann *et al.* 2014; Sentis *et al.* 2017). Inspired by previous temperature-independent

stoichiometric consumer-resource models (Andersen 1997; Andersen *et al.* 2004; Diehl *et al.*

2005), we then extended the Rosenzweig-MacArthur model to account for nutrient dynamics as

well as for the simultaneous dependence of community dynamics on temperature and flexible

resource stoichiometry. Our objective here was not to develop a complex and very realistic

stoichiometric model that would include additional important abiotic and biotic features such as

light intensity (Diehl 2007) or compensatory feeding (Cruz-Rivera & Hay 2000). Instead, we

aimed at introducing two fundamental stoichiometric features (i.e. stoichiometric flexibility and

stoichiometric imbalance) and investigate how these stoichiometric considerations can change

predictions of the Rosenzweig-MacArthur model. We thus used our extended Rosenzweig-

MacArthur model to predict the effects of warming and nutrient enrichment on population

dynamics and biomass distribution across trophic levels and compared these predictions with the

predictions of the nonstoichiometric Rosenzweig-MacArthur model. We particularly addressed

two questions: (i) How do stoichiometric constraints modulate the effects of enrichment and

warming on community stability and persistence? and (ii) How do stoichiometric constraints

modulate the effects of enrichment and warming on biomass distribution across multiple trophic

levels?

Methods: Population dynamic models

The Rosenzweig-MacArthur (RM) model.

Rates of change of the consumer and resource biomass densities $\dot{C}$ and $\dot{R}$ depend on their

respective biomass densities C and R (g.m^{-3}):

$\dot{R} = r(1 - \frac{R}{K})R - \frac{aR}{1+ahR}C$ (1)

$\dot{C} = (e\frac{aR}{1+ahR} - m)C$ (2)

The population growth rate of the resource is given by the logistic equation where r is the

resource maximum growth rate and K is the resource carrying capacity. The population growth

rate of the consumer is equal to its feeding rate multiplied by its assimilation efficiency e (i.e. the

fraction of resource biomass converted into consumer biomass) minus a loss term associated to

metabolic losses m. The feeding rate of the consumer C depends on the density of its resource R

and follows a Holling type II functional response, with consumer-resource attack rate a and

handling time h. Our choice for a type II functional response is motivated by a meta-analysis

reporting that most consumers feed on their prey following a saturating type II functional

response rather than a linear type I or a sigmoidal type III functional response (Jeschke *et al.*

2004).

In the RM model, consumer and resource population growth rates are only limited by nutrient or

resource density. Nutrient enrichment is assumed to increase resource carrying capacity, which

often leads to the well-known paradox of enrichment where populations fluctuates up to

extinctions (Rosenzweig 1971). Nevertheless, this model neither considers nutrient dynamics nor

temporal variations of resource stoichiometry and their consequences on population dynamics.

To circumvent these limitations of the RM model, we extended it to better consider nutrient

dynamics, resource stoichiometry and the way they can affect resource and consumer population

dynamics.

The Stoichiometric Rosenzweig-MacArthur (SRM) model.

We derived a stoichiometric extension of the Rosenzweig-MacArthur consumer–resource model

with additional stoichiometric and temperature dependencies of several biological rates. We

considered two stoichiometric constraints: one on the resource population growth rate, and the

other on the consumer assimilation efficiency (see below for more details). These stoichiometric

constraints have been observed for several consumer-resource pairs suggesting that they are core

components of species growth and interactions (Sterner & Elser 2002).

Stoichiometric constraint on the resource population growth rate

Inspired by previous stoichiometric models (Andersen 1997; Loladze *et al.* 2000; Andersen *et al.*

2004; Diehl *et al.* 2005), we extended the RM model by considering explicit nutrient dynamics

and nutrient effects on resource population growth rate. The system is assumed to be closed for

nutrients. Thus, nutrient supply originates exclusively from biomass excretion and

remineralization. The total amount of nutrients in the system (N_{tot}) is then a measure of nutrient

enrichment. In contrast to the very high plasticity in C:N or C:P exhibited by autotrophs,

heterotrophs regulate elemental composition within narrower bounds, even when consuming food

with large variation in elemental composition (Andersen & Hessen 1991; Sterner & Hessen 1994;

Andersen 1997; Elser *et al.* 2000). In other words, the elemental homeostasis is much stronger for

consumers compared to primary producers. We thus assumed the nutrient quota (i.e. the nutrient

to carbon ratio) of the consumer Q_C to be conserved whereas the one of the resource Q_R is

flexible over time with the only constraint that $Q_R > 0$. As in the RM model, rates of change of

the consumer and resource biomass densities $\dot{C}$ and $\dot{R}$ depend on their respective carbon biomass

densities C and R (gC.m^{-3}), except that the resource population growth rate follows the Droop

equation (Droop 1974) given by $r(1-Q_{min}/Q_R)R$ and is now limited by Q_R relative to the

minimum nutrient quota Q_{min}:

$$N_{tot} = Q_R R + Q_C C \tag{3}$$

$$\dot{R} = r(1 - \frac{Qmin}{Q_R})R - \frac{aR}{1+ahR}C \tag{4}$$

$$\dot{C} = (e\frac{aR}{1+ahR} - m)C \tag{5}$$

From the nutrient conservation equation (eqn 3) we obtain that $Q_R = \frac{N_{tot} - Q_C C}{R}$. The intuitive

interpretation is that the resource nutrient quota Q_R changes instantaneously with the density of

the resource population R and with the density of the nutrient stored in the consumer biomass

$Q_C C$, to maintain nutrient balance (see Text S1 for details).

Stoichiometric constraint on the consumer population growth rate

In the RM model, the growth rate of the consumer population only depends on resource density.

We relaxed this assumption by making the population growth rate of the consumer dependent on

both resource quality (i.e. nutrient quota) and quantity (i.e. biomass density). In the SRM model,

consumer production is also limited by resource quality as the consumer assimilation efficiency e

is a saturating function of resource nutrient quota Q_R:

$$e(Q_R) = e_{max}\frac{Q_R}{Q_R + Q_C} \tag{6}$$

The intuitive interpretation of eqn. 6 is that resource quality is not a limiting factor for consumer

growth as long as the nutrient content of the resource is superior to the nutrient content of the

consumer (i.e. $Q_R > Q_C$). In other words, $e(Q_R)$ is at its maximum for $Q_R > Q_C$ and proportional to

Q_R for $Q_R < Q_C$. By replacing e by $e(Q_R)$ in eqn. 5, we obtain the SRM model.

Temperature dependence of model parameters

To investigate the effect of temperature and stoichiometric constraints on consumer-resource

dynamics, we next extended the RM and SRM models described above by adding thermal

dependencies of the parameters. Following Uszko et al. (2017), we assumed that the total amount

of nutrient N_{tot}, the maximum food conversion efficiency e_{max}, and fixed stoichiometric traits

(Q_C) are independent of temperature, as there is no evidence of systematic temperature

dependence for any of them (Peters 1983; Ahlgren 1987; Borer *et al.* 2013; Yvon-Durocher *et al.*

2015). Rate of maintenance respiration and natural background mortality m typically increases

exponentially with temperature (Fig. S1a and b). We thus used the Arrhenius equation to describe

the effect of temperature T (in Kelvin) on m:

$$m(T) = m_0 e^{\frac{-E_m}{kT}} \tag{7}$$

where m_0 is a parameter-specific constant calculated at temperature of 0°C (= 273.15 K). The

temperature dependence is characterized by the respective activation energy E_m (eV) and the

Boltzmann constant $k=8.62\times10^{-5}$ eVK^{-1}. As the temperature dependencies of resource intrinsic

growth rate r and functional response parameters (a, $1/h$) are often unimodal rather than

exponential (Englund *et al.* 2011; Rall *et al.* 2012; Sentis *et al.* 2012; Thomas *et al.* 2012), we

used Gaussian functions for r and a and an inverted Gaussian function for h:

$$g(T) = g_0 e^{\pm\frac{(T-Topt)^2}{2s^2}} \tag{8}$$

where T_{opt} is the temperature at which the rate g reaches its minimum or maximum, s is the

function width and g_0 is a parameter-specific constant calculated at T_{opt}. The minus-sign

corresponds to Gaussian functions and the plus-sign to inverted Gaussian functions.

Model parameterisation and simulations

To parameterise the models we assumed the resource and consumer species to be a unicellular

freshwater algae and a *Daphnia* grazer, respectively. The choice for this system was motivated by

the good characterization of both the stoichiometric parameters and thermal dependencies for this

system (Andersen 1997; Uszko *et al.* 2017). Uszko *et al.* (2017) recently estimated the thermal

dependencies for biological rates of the green algae *Monoraphidium minutum* and the grazer

Daphnia hyalina. We thus used their estimates of stoichiometric parameters and thermal

dependencies (See Table S1 and Fig. S1 for further details).

To investigate the individual and combined effects of enrichment, warming, and stoichiometric

constraints, we varied temperature (401 values ranging from 0 to 40°C by 0.1°C) and total

amount of nutrients (parameter N_{tot} in eqn. 10; 60 values ranging from 0.001 to 0.06 gP.m^{-3} by

0.001 gP.m^{-3}, overlapping with reported mean phosphorus concentration in European peri-alpine

lakes (Anneville *et al.* 2005)). For the RM model, we used the minimum nutrient quota to convert

nutrients into resource (i.e. $K = N_{tot}/Q_{min}$). This implies that carrying capacity is independent of

temperature which is expected for closed, nutrient-limited systems (Uszko *et al.* 2017) although

more experimental evidence are needed to verify this assumption. We then simulated the

consumer-resource dynamics for 1000 days to enable the system to reach an attractor (either an

equilibrium point or a limit cycle) before we assessed the final state. Therefore, for each model,

we simulated 24060 combinations of environmental conditions (401 temperatures by 60 nutrient

concentrations). Initial biomass density of each species was set to 0.98 times its equilibrium

density in the two-species system (calculated by solving for the two-species equilibrium, using

either eqns 1-2 for model RM or eqns 3-5 for model SRM). The value of 0.98 was chosen to be

(1) close enough to equilibria to avoid extinctions caused solely by transient dynamics and (2) not

exactly the equilibrium value to probe the stability of the equilibrium. Any population falling

below the extinction threshold of 10^{-9} g.m^{-3} during the simulation was deemed extinct and its

biomass set to zero to exclude ecologically unrealistic low biomass densities. For each model, we

calculated system persistence as the percentage of simulations with the two species remaining

extant at the end of the simulations. Population dynamics were simulated with R version 3.4.3 (R

Development Core Team 2017) using the "deSolve" package (Soetaert *et al.* 2012) with an

absolute error tolerance of 10^{-10} and a relative error tolerance of 10^{-6}.

Results

Stability: population fluctuations and persistence

Stoichiometric constraints dampened the paradox of enrichment, reducing fluctuations at high

nutrient levels and hence increasing persistence. However, stoichiometric constraints also

reduced the persistence of the consumer at low and high temperatures. As a result, the overall

effect of stoichiometric constraints on stability depends on its relative influence on population

fluctuations versus consumer persistence. In the two following paragraphs, we explain in more

detail these results and highlight key differences between the outcomes from RM and SRM

models.

The RM model predicts that increasing nutrient concentration is strongly destabilizing: the

system shifts from a stable equilibrium point to limit cycles (i.e. the system crosses a Hopf

bifurcation). This agrees with the paradox of enrichment. As population biomass fluctuations (i.e.

cycle amplitude) increase with nutrient concentration, minimal population densities are very low

at high nutrient concentrations leading to the extinction of both the consumer and resource once

the extinction threshold is crossed (Fig. 1). In the range of temperatures where the consumer

persists, warming does not have a strong influence on the nutrient concentration at which the

system shifts from the stable equilibrium point to limit cycles, although this qualitative shift is

absent at very high temperatures (i.e. 32°C) when the consumer is close to extinction. Warming

decreases fluctuation amplitude and thus dampens extinctions driven by the paradox of

enrichment, which results in warming enhancing the persistence of the consumer-resource system

at high nutrient concentrations. However, very warm and cold temperatures cause the extinction

of the consumer (see below for the mechanisms underlying extinctions), releasing resources from

top-down control. Overall, we found that, without considering the extinction threshold of 10^{-9}

g.m^{-3} (see Model parametrisation and simulations), both the consumer and the resource can

persist in 74% of the temperature-nutrient concentration scenarios (i.e. black + orange areas in

Fig 1C). Nevertheless, when considering the extinction threshold, they persist in only 21% of the

temperature-nutrient scenarios (i.e. black area in Fig. 1c) indicating that extinctions driven by

population fluctuations are highly prevalent in the RM model.

In contrast, the SRM model shows that increasing nutrient concentrations causes fewer

fluctuations than those observed for the RM model (Fig. 1). This is because: (1) more nutrients

are needed to shift the system from a stable equilibrium point to limit cycles—the system can

indeed persist without fluctuations up to 0.02 gP.m^{-3} whereas it was only up to 0.0005 gP.m^{-3} in

the RM model—and (2) when the system fluctuates, the amplitude of the fluctuations is smaller

in the SRM than in the RM model. As a result, stoichiometric constraints dampen the amplitude

of population fluctuations (i.e. the paradox of enrichment) and hence increase system persistence

at high nutrient levels. While the qualitative effect of temperature is similar to that observed in

the RM model, the thermal thresholds for consumer persistence are reduced at low and high

temperatures in the SRM predictions. Moreover, thermal thresholds remain almost constant along

the nutrient gradient in the RM model, whereas in the SRM model they depend on nutrient

concentration, with a smaller thermal range at low nutrient levels compared to high nutrient

levels (Fig. 1). The consumer is thus more likely to go extinct at low nutrient concentrations and

extreme temperatures in the SRM model than in the RM model. Overall, system persistence for

the SRM model was 44% without considering the extinction threshold and 37% when

considering it. In other words, without considering extinctions driven by very low biomass

densities, the SRM model predicts lower persistence of the consumer compared to RM model but

it is the opposite pattern when considering extinctions driven by very low biomass densities. We

thus conclude that the RM model predicts larger population fluctuations leading to high

probabilities of populations extinctions in comparison to the SRM model.

Biomass distribution

We next compared the predictions of both models for consumer-resource biomass ratios along the

temperature and nutrient gradients (Fig. 2). We found that the RM model systematically predicts

biomass ratio > 1 (i.e. consumer biomass is larger than resource biomass). In contrast, the SRM

model predicts biomass ratios both > or < than 1 depending on temperature and nutrient levels.

The RM model predicts that, as soon as the consumer can persist, its population biomass density

always exceeds the resource population biomass density (Fig. 2). With the SRM model, the

biomass ratios are below one at low nutrient levels (Fig. 2). However, at medium and high

nutrient levels, the ratios are above one as soon as the consumer can persist. We found

qualitatively similar results when considering unstable equilibrium points (Fig. S2). Finally, we

showed that, for equivalent parameter values, the RM model predicts biomass ratio that are

superior or equal to the ones predicted by the SRM model (text S2). This difference between the

two models is independent of the shape and position of the temperature function used to

parametrise the models.

Mechanisms underlying stability and biomass distribution patterns

Here, we detail the mechanisms underlying the stability and biomass distribution patterns to

better understand how and when stoichiometric constraints modulate the effects of temperature

and nutrients on consumer-resource dynamics. The first mechanism corresponds to the effect of

stoichiometric constraints on the consumer energetic efficiency that determines the consumer

persistence at extreme low and high temperatures. The second mechanism relates to the influence

of the stoichiometric constraints on population dynamical feedback that explains why the

stoichiometric model predicts more stability at high nutrient levels compared to the non-

stoichiometric model.

Consumer energetic efficiency

The persistence of the consumer at low and high temperatures is driven by the energetic

efficiency *EE* of the consumer (i.e. its feeding rate relative to metabolic losses) calculated as

follows:

$$EE = \frac{ef(R^*)}{m} \qquad (9)$$

Where $f(R^*)$ is the functional response of the consumer at resource density R^* (i.e. the resource

equilibrium density in absence of the consumer). We recall that the assimilation efficiency *e* is a

function of resource quality Q_R in the SRM model whereas it is constant in the RM model. The

intuitive interpretation of eqn. 9 is that *EE* should be above one for the consumer population to

grow and persist.

To better understand the influence of stoichiometric constraints on consumer persistence, we thus

investigated differences in the RM and SRM model predictions regarding the consumer energetic

efficiency *EE* along the temperature gradient at two nutrient concentrations (Fig. 3). For both

models, energetic efficiency at equilibrium has a hump-shaped relationship with temperature with

maximal efficiency values at medium temperatures. While this unimodal shape is conserved

across nutrient levels and models, the RM model systematically predicts higher consumer

energetic efficiency values than the SRM model because consumer assimilation efficiency is

lower in the SRM than in the RM model (Fig. S3). As a result, the temperatures at which

energetic efficiency falls below one and drives consumers extinct are more extreme in the RM

model compared to the SRM model (Fig. 3). In other words, energetic efficiency is above one for

a narrower thermal range in the SRM model.

Dynamical feedbacks due to the stoichiometric constraints

The second mechanism by which stoichiometric constraints influence consumer-resource stability

and biomass distribution are the dynamical feedbacks due to stoichiometric constraints on the

resource population growth rate and on the consumer energetic efficiency. In the SRM model, the

growth rate of the resource population depends on both the total nutrient load and the consumer

population density as $Q_R = (N_{tot} - Q_C C)/R$. In other words, when consumer population increases,

this decreases resource population growth by reducing both resource density (through predation)

and quality (through the total nutrient load) leading to a negative feedback on consumer

population growth rate. In contrast, for the RM model, the negative consumer feedback is only

driven by the reduction in resource density as resource quality is not considered. In addition to

this first dynamical feedback, there is a second dynamical feedback as the consumer population

growth rate also depends on Q_R and thus on its own biomass density. Thus, also this second

negative feedback loop limits the consumer population growth rate when its density increases.

Altogether, dynamical feedbacks reduce strongly the amplitude of population fluctuations, which

in turn increases resource and consumer persistence.

To reveal the dynamic effects of the stoichiometric constraints, we calculated the values of

assimilation efficiencies and carrying capacities predicted by the SRM model for each

temperature-nutrient scenario (Fig. S3) and used these effective parameter values to replace the

values of parameters e and K in the RM model for each temperature-nutrient scenario. In other

words, we calculated average values of e and K in the dynamic SRM model and used them as

constant input parameters in the RM model. The objective of using these effective parameter

values was to disentangle the static effect of stoichiometric constraints (i.e. changing the average

parameter values of consumer assimilation efficiency and of the resource carrying capacity) from

their population dynamical effect (i.e. the two dynamical feedback described above). We thus

simulated population dynamics along the temperature-nutrient gradient using the RM model with

these effective parameters; referred hereafter as effective RM model (Fig. 4). Comparing

predictions from the RM, effective RM, and SRM models allowed to disentangle the static

stoichiometric effects when going from the RM to the effective RM predictions (Fig. 4, panels a

to b) from the dynamical stoichiometric effects when going from the effective RM to the SRM

predictions (Fig. 4, panels b to c). In other words, the RM and effective RM only differ in their

parameter values because the effective RM takes into account the effect of stoichiometric

constraints on the average parameter values. On the other hand, the effective RM and SRM have

similar parameter values but different population dynamics, which helps understanding the

dynamical feedback induced by stoichiometric constraints.

We found that, at low nutrient concentrations, population fluctuations and consumer persistence

predicted by the effective RM model agreed with predictions of the SRM model. However, the

system shifted from a stable equilibrium point to a limit cycle at lower nutrient concentrations for

the effective RM model than for the SRM model. This suggests that more nutrients are needed to

destabilize the system with the SRM model. Moreover, the effective RM model predicts ampler

population fluctuations than the SRM model. As a result, the effective RM predicts high

extinction rates at high nutrient concentrations compared to the SRM model. Overall, we found

that the effective RM model cannot fully reproduce the dynamics predicted by the SRM, which

indicates that including stoichiometric constraints in the RM model involves more than only

changing parameter values.

Discussion

Temperature and nutrient enrichment are two of the most important drivers of global change

(Nelson 2005). However, most research on the effects of temperature and nutrients on community

dynamics assumes that the elemental composition of primary producers and consumers are

constant and independent of changes on energy and material fluxes (Binzer *et al.* 2012; Boit *et al.*

2012; Amarasekare & Coutinho 2014; Gilbert *et al.* 2014; Amarasekare 2015; Binzer *et al.* 2016;

Gilarranz *et al.* 2016). Yet, the elemental composition of primary producers is known to be

flexible, which can have important consequences for community dynamics and ecosystem

processes (Elser *et al.* 2000). We have shown how stoichiometric constraints that account for

flexible stoichiometry can affect predictions on how temperature and nutrients influence

community stability and biomass distribution across trophic levels. We thus argue that

considering stoichiometric constraints is an important step toward a better understanding of the

effects of global change on ecosystems.

Stoichiometric constraints and temperature can dampen the paradox of enrichment

We showed that both stoichiometric constraints and temperature dampen the negative effect of

nutrient enrichment on consumer-resource fluctuations and increase system persistence at high

nutrient levels. Temperature effects are driven by physiological mechanisms. In agreement with

previous empirical studies, our model parametrization reflects the observation that metabolic loss

rates increase faster with warming than consumer feeding rates (Vucic-Pestic *et al.* 2011; Sentis

et al. 2012; Fussmann *et al.* 2014; Iles 2014). Consumers are thereby less energetically efficient

at higher temperatures which stabilizes food-web dynamics by reducing energy flow between

trophic levels (Binzer *et al.* 2012; Kratina *et al.* 2012; Fussmann *et al.* 2014; Sentis *et al.* 2017).

In contrast, the effect of stoichiometric constraints is mainly linked to two mechanisms: a shift in

the position of the Hopf bifurcation and negative dynamical feedbacks of the consumer and

resource on their population growth rates. Both resources and consumers are composed of the

same essential elements (N, P, and C), which implies that, when consumer or resource population

biomass increases, it reduces the pool of free nutrients available for the growth of the resource

population. Therefore, more nutrients are needed to shift the system from a stable equilibrium to

population cycles. In other words, the paradox of enrichment is displaced to higher nutrient

concentrations (i.e., the position of the Hopf bifurcation is shifted to higher nutrient levels. In

contrast, the RM model does not take into account the storage of nutrients in both the resource

and consumer biomasses (i.e. the carrying capacity only depends on the total nutrient load). Less

enrichment is thus required to shift the system from a stable equilibrium point to limit cycles.

We found two dynamic effects that correspond to negative dynamical feedbacks of the consumer

and the resource on themselves. When consumer population increases, it decreases the population

growth rate of the resource by limiting nutrient availability, diminishing resource biomass, which,

in turn, decreases the consumer population growth rate. Conversely, when the resource biomass

increases, this decreases the nutrient content of the resource, which, in turn, limits the growth

rates of both the resource and consumer populations. These stoichiometric negative feedback

loops strongly decrease the amplitude of population fluctuations and thus dampen the paradox of

enrichment. Interestingly, our comparisons of the RM, effective RM and SRM model predictions

indicate that the dynamical effects contribute more to the reduction of fluctuations than the static

effects: population fluctuations are large in the effective RM model accounting for the static

effect only, whereas they are much smaller in SRM model accounting for both static and

dynamical effects (Fig. 4). This implies that the impact of stoichiometric constraints on

community dynamics goes beyond a simple modification of parameter values and encompass

more complex population feedbacks between the consumer and the resource.

Overall, these results demonstrate that both flexible stoichiometry and temperature can

synergistically dampen the paradox of enrichment by two different mechanisms: population

dynamic feedbacks and physiological constraints. Our consumer-resource model is simplified

compared to natural communities composed of numerous species. Moreover, in natural systems,

a large amount of nutrient can be stored in abiotic and slow biotic pools that have long turnover

times which, in turn, can influence the population dynamics. In particular, the amplitude of the

population fluctuations is expected to be smaller as abiotic pools can buffer the population

feedback. Nevertheless, considering the nutrient held in slow abiotic or biotic pools would not

change the equilibrium densities of primary producers and grazer if nutrients are released in the

environment proportionally to their density stored in the abiotic pool. Moreover, the predictions

of the stoichiometric model fit with empirical observations. In eutrophic lakes and experimental

mesocosms, populations can persist at relatively high nutrient concentrations even if fertilisation

enhance population fluctuations (O'Connor *et al.* 2009; Boit *et al.* 2012; Kratina *et al.* 2012), as

our stoichiometric model predicts. In contrast, the Rosenzweig-MacArthur model tends to

produce very large population fluctuations and extinctions at low nutrient concentrations which

can explain why these predictions are not well supported by empirical observations (McAllister *et*

al. 1972; Jensen & Ginzburg 2005).

Effects of stoichiometric constraints on system persistence across environmental gradients

While stoichiometric constraints dampen the paradox of enrichment and thus increase persistence

at high nutrient levels, they also reduce the persistence of the consumer at low and high

temperatures. Stoichiometric constraints affect the thermal thresholds for consumer extinctions.

Consumers can only persist over a narrower range of intermediate temperatures when they are

constrained by stoichiometry. This is due to the reduced biomass assimilation of the consumer at

low and high temperatures that, in turn, decreases its energetic efficiency and thus fastens

consumer extinction. In our stoichiometric model, the reduction of biomass assimilation

efficiency emerges from the effect of temperature on resource quality: extreme high and low

temperatures decrease resource quality and thus less resource biomass can be converted in

consumer biomass at these temperatures.　The emergence of a thermal dependency for

assimilation efficiency contrasts with previous theoretical studies that used the RM model and

assumed that the assimilation efficiency is temperature independent as resource quality is

assumed constant (Binzer *et al.* 2012; Gilbert *et al.* 2014; Sentis *et al.* 2017; Uszko *et al.* 2017).

In the SRM model, the thermal dependency of the consumer assimilation efficiency is fully

driven by the change in the resource stoichiometry induced by temperature. The SRM model thus

predicts an additional mechanism by which temperature can influence trophic interactions:

temperature changes resource stoichiometry, which, in turn, impacts the consumer assimilation

efficiency and its population growth rate. This prediction matches with empirical results showing

that primary producer stoichiometric composition can change with temperature (Woods *et al.*

2003) and that consumer assimilation efficiency is sensitive to resource stoichiometric

composition (Andersen 1997; Elser *et al.* 2000). To sum up, the overall effect of stoichiometric

constraints on system persistence thus depends on the temperature range considered and on their

relative influence on population fluctuations versus consumer persistence.

Effects of stoichiometric constraints on biomass distribution

We found that stoichiometric constraints can modulate the effects of temperature and nutrients on

biomass distribution across trophic levels. Without stoichiometric constraints (i.e. with the

Rosenzweig-MacArthur model), biomass ratios are above one for almost all temperatures or

nutrient levels as the biomass produced by the resource is efficiently transferred to the consumer

level consistently along the environmental gradients. This finding agrees with theoretical studies

reporting that Lotka-Volterra and RM models predict biomass ratios above one and fail to

reproduce biomass pyramids for a substantial region of parameter values (Jonsson 2017; Barbier

& Loreau 2019). However, in nature, consumer-resource biomass ratios are often below one

(McCauley & Kalff 1981; Del Giorgio & Gasol 1995; McCauley *et al.* 1999; Irigoien *et al.* 2004)

suggesting that additional mechanisms should be included to better understand and predict

biomass distribution patterns in natural food webs. Our stoichiometric model agrees with

experimental observations. It predicts that, at low nutrient concentrations (i.e. < 0.01 gP.m^{-3}), the

biomass ratio never exceeds one along the entire temperature gradient. This is observed in

oligotrophic aquatic systems where primary production is too low to sustain high consumer

populations (O'Connor *et al.* 2009). In addition, we also found that increasing nutrient levels

decreased the temperature ranges within which biomass ratio is below one. This corresponds to

results from manipulated nutrient concentrations and temperature in aquatic mesocosms, where

zooplankton to phytoplankton biomass ratio only exceeds one in the enriched mesocosms at

medium or warm temperatures (i.e. 27°C) (O'Connor *et al.* 2009). This suggests that the models

with stoichiometric constraints better reproduce the biomass patterns observed in experimental

and natural systems. Nevertheless, further experiments investigating the links between

stoichiometric flexibility and consumer-resource dynamics are needed to determine if these

stoichiometric mechanisms are underlying patterns of biomass distribution in nature.

Implications of our findings for global change

Temperature and nutrients do not act in isolation from each other. Climate warming, for example,

causes stronger water stratification which, in turn, can limit nutrient cycling (Sarmiento *et al.*

2004; Tranvik *et al.* 2009). Environmental policies such as the European water framework

directive (i.e. Directive 2000/60/EC of the European Parliament and of the Council establishing a

framework for the Community action in the field of water policy) effectively reduces input of

nutrients in aquatic ecosystems (Anneville *et al.* 2005) while the climate keeps warming. With

these two phenomena, water will often be warmer and contain fewer nutrients in aquatic systems.

Our models consistently predict that warmer temperatures should stabilise consumer-resource

dynamics but, if temperature further increases, the consumer goes extinct as energetic efficiency

decreases with warming. Moreover, we found that stoichiometric constraints can reduce this

thermal extinction threshold (i.e. the consumer persists in a narrower thermal range), especially at

low nutrient levels. Our stoichiometric model thus suggests that decreasing nutrient

concentrations alongside warmer temperatures should fasten the extinction of consumer

populations. This prediction matches empirical observations of consumer extinctions at warm

temperatures in oligotrophic aquatic systems (Petchey *et al.* 1999; O'Connor *et al.* 2009).

Altogether, these results indicate that considering stoichiometric constraints can be of importance

for the management of nutrient inputs and the conservation of natural populations and

communities under climate change.

Conclusion

Knowledge of how temperature and nutrient simultaneously influence the elemental composition

of primary producers and consumers is crucial to better understand and predict the effects of

global change on species interactions, community dynamics and fluxes of energy and material

within and among ecosystems. Here we showed that stoichiometric constraints dampen the

negative effect of enrichment on stability by reducing population fluctuations through population

dynamics feedbacks. However, stoichiometric constraints also decrease consumer energetic

efficiency, which increases consumer extinction risk at extreme temperatures and low nutrient

concentrations. Finally, stoichiometric constraints can reverse biomass distribution across trophic

level by modulating consumer efficiency and resource population growth rate along the

temperature and nutrient gradients. Our study provides a first step in the exploration of the

consequences of stoichiometric constraints and temperature on ecological communities. It

suggests that accounting for stoichiometric constraints can strongly influence our understanding

of how global change drivers impact important features of ecological communities such as

stability and biomass distribution patterns.

Acknowledgements: This work is funded by the FRAGCLIM Consolidator Grant (number

726176) to Jose M. Montoya from the European Research Council under the European Union's

Horizon 2020 Research and Innovation Program.

References

Ahlgren, G. (1987). Temperature functions in biology and their application to algal growth

constants. *Oikos*, 49, 177-190.

Amarasekare, P. (2015). Effects of temperature on consumer-resource interactions. *Journal of*

Animal Ecology, 84, 665-679.

Amarasekare, P. & Coutinho, R.M. (2014). Effects of temperature on intraspecific competition in

ectotherms. *The American Naturalist*, 184, E50-E65.

Andersen, T. (1997). *Pelagic nutrient cycles: herbivores as sources and sinks*. Springer-Verlag,

Berlin, Germany.

Andersen, T., Elser, J.J. & Hessen, D.O. (2004). Stoichiometry and population dynamics.

Ecology Letters, 7, 884-900.

Andersen, T. & Hessen, D.O. (1991). Carbon, nitrogen, and phosphorus content of freshwater

zooplankton. *Limnology and Oceanography*, 36, 807-814.

Anneville, O., Gammeter, S. & Straile, D. (2005). Phosphorus decrease and climate variability:

mediators of synchrony in phytoplankton changes among European peri-alpine lakes.

Freshwater Biology, 50, 1731-1746.

Barbier, M. & Loreau, M. (2019). Pyramids and cascades: a synthesis of food chain functioning

and stability. *Ecology Letters*, 22, 405-419.

Bezemer, T.M. & Jones, T.H. (1998). Plant-insect herbivore interactions in elevated atmospheric

C02: quantitative analyses and guild effects. *Oikos*, 82, 212-222.

Binzer, A., Guill, C., Brose, U. & Rall, B.C. (2012). The dynamics of food chains under climate

change and nutrient enrichment. *Philosophical Transactions of the Royal Society B:*

Biological Sciences, 367, 2935-2944.

Binzer, A., Guill, C., Rall, B.C. & Brose, U. (2016). Interactive effects of warming,

eutrophication and size structure: impacts on biodiversity and food☐web structure. *Global*

Change Biology, 22, 220-227.

Boit, A., Martinez, N.D., Williams, R.J. & Gaedke, U. (2012). Mechanistic theory and modelling

of complex food-web dynamics in Lake Constance. *Ecology Letters*, 15, 594-602.

Borer, E.T., Bracken, M.E., Seabloom, E.W., Smith, J.E., Cebrian, J., Cleland, E.E. *et al.* (2013).

Global biogeography of autotroph chemistry: is insolation a driving force? *Oikos*, 122,

1121-1130.

Brown, J.H., Gillooly, J.F., Allen, A.P., Savage, V.M. & West, G.B. (2004). Toward a metabolic

theory of ecology. *Ecology*, 85, 1771-1789.

Cruz-Rivera, E. & Hay, M.E. (2000). Can quantity replace quality? Food choice, compensatory

feeding, and fitness of marine mesograzers. *Ecology*, 81, 201-219.

Del Giorgio, P.A. & Gasol, J.M. (1995). Biomass Distribution in Freshwater Plankton

Communities. *The American Naturalist*, 146, 135-152.

Diehl, S. (2007). Paradoxes of enrichment: effects of increased light versus nutrient supply on

pelagic producer-grazer systems. *The American Naturalist*, 169, E173-E191.

Diehl, S., Berger, S. & Wöhrl, R. (2005). Flexible nutrient stoichiometry mediates environmental

influences on phytoplankton and its resources. *Ecology*, 86, 2931-2945.

Droop, M. (1974). The nutrient status of algal cells in continuous culture. *Journal of the Marine*

Biological Association of the United Kingdom, 54, 825-855.

Elser, J., Sterner, R., Gorokhova, E.a., Fagan, W., Markow, T., Cotner, J. *et al.* (2000). Biological

stoichiometry from genes to ecosystems. *Ecology Letters*, 3, 540-550.

Elser, J.J., Bracken, M.E., Cleland, E.E., Gruner, D.S., Harpole, W.S., Hillebrand, H. *et al.*

(2007). Global analysis of nitrogen and phosphorus limitation of primary producers in

freshwater, marine and terrestrial ecosystems. *Ecology Letters*, 10, 1135-1142.

Elser, J.J., Loladze, I., Peace, A.L. & Kuang, Y. (2012). Lotka re-loaded: modeling trophic

interactions under stoichiometric constraints. *Ecological Modelling*, 245, 3-11.

Elton, C. (1927). *Animal ecology*. Sidgwick & Jackson, LTD, London.

Englund, G., Ohlund, G., Hein, C.L. & Diehl, S. (2011). Temperature dependence of the

functional response. *Ecology Letters*, 14, 914-921.

Enquist, B.J., West, G.B., Charnov, E.L. & Brown, J.H. (1999). Allometric scaling of production

and life-history variation in vascular plants. *Nature*, 401, 907-911.

Falkowski, P.G., Barber, R.T. & Smetacek, V. (1998). Biogeochemical controls and feedbacks on

ocean primary production. *Science*, 281, 200-206.

Finkel, Z.V., Beardall, J., Flynn, K.J., Quigg, A., Rees, T.A.V. & Raven, J.A. (2009).

Phytoplankton in a changing world: cell size and elemental stoichiometry. *Journal of*

Plankton Research, 32, 119-137.

Fussmann, K.E., Schwarzmüller, F., Brose, U., Jousset, A. & Rall, B.C. (2014). Ecological

stability in response to warming. *Nature Climate Change*, 4, 206-210.

Gilarranz, L.J., Mora, C. & Bascompte, J. (2016). Anthropogenic effects are associated with a

lower persistence of marine food webs. *Nature communications*, 7, 10737.

Gilbert, B., Tunney, T.D., McCann, K.S., DeLong, J.P., Vasseur, D.A., Savage, V. *et al.* (2014).

A bioenergetic framework for the temperature dependence of trophic interactions.

Ecology Letters, 17, 902-9014.

Hessen, D.O., Ågren, G.I., Anderson, T.R., Elser, J.J. & De Ruiter, P.C. (2004). Carbon

sequestration in ecosystems: the role of stoichiometry. *Ecology*, 85, 1179-1192.

Hessen, D.O., Færøvig, P.J. & Andersen, T. (2002). Light, nutrients, and P:C ratios in algae:

grazer performance related to food quality and quantity. *Ecology*, 83, 1886-1898.

Iles, A.C. (2014). Towards predicting community level effects of climate: relative temperature

scaling of metabolic and ingestion rates. *Ecology*, 95, 2657–2668.

Irigoien, X., Huisman, J. & Harris, R.P. (2004). Global biodiversity patterns of marine

phytoplankton and zooplankton. *Nature*, 429, 863-867.

Jensen, C.X. & Ginzburg, L.R. (2005). Paradoxes or theoretical failures? The jury is still out.

Ecological Modelling, 188, 3-14.

Jeschke, J.M., Kopp, M. & Tollrian, R. (2004). Consumer☐food systems: why Type I functional

676   responses are exclusive to filter feeders. *Biological Reviews*, 79, 337-349.

677   Jonsson, T. (2017). Conditions for eltonian pyramids in Lotka-Volterra food chains. *Scientific*

678   *Reports*, 7, 10912.

679   Kratina, P., Greig, H.S., Thompson, P.L., Carvalho-Pereira, T.S.A. & Shurin, J.B. (2012).

680   Warming modifies trophic cascades and eutrophication in experimental freshwater

681   communities. *Ecology*, 93, 1421-1430.

682   Lindeman, R.L. (1942). The trophic☐dynamic aspect of ecology. *Ecology*, 23, 399-417.

683   Loladze, I., Kuang, Y. & Elser, J.J. (2000). Stoichiometry in producer-grazer systems: linking

684   energy flow with element cycling. *Bulletin of Mathematical Biology*, 62, 1137-1162.

685   McAllister, C., LeBrasseur, R., Parsons, T. & Rosenzweig, M. (1972). Stability of enriched

686   aquatic ecosystems. *Science*, 175, 562-565.

687   McCauley, D.J., Gellner, G., Martinez, N.D., Williams, R.J., Sandin, S.A., Micheli, F. *et al.*

688   (2018). On the prevalence and dynamics of inverted trophic pyramids and otherwise top-

689   heavy communities. *Ecology Letters*, 21, 439-454.

690   McCauley, E. & Kalff, J. (1981). Empirical relationships between phytoplankton and

691   zooplankton biomass in lakes. *Canadian Journal of Fisheries and Aquatic Sciences*, 38,

692   458-463.

693   McCauley, E., Nisbet, R.M., Murdoch, W.W., de Roos, A.M. & Gurney, W.S.C. (1999). Large-

694   amplitude cycles of *Daphnia* and its algal prey in enriched environments. *Nature*, 402,

695   653-656.

696   Montoya, J.M. & Raffaelli, D. (2010). Climate change, biotic interactions and ecosystem

697   services. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 365,

698   2013-2018.

Nelson, G.C. (2005). *Millennium ecosystem assessment: drivers of ecosystem change: summary*

chapter. World Resources Institute, Washington, DC.

O'Connor, M.I., Piehler, M.F., Leech, D.M., Anton, A. & Bruno, J.F. (2009). Warming and

resource availability shift food web structure and metabolism. *PLoS Biology*, 7,

e1000178.

Petchey, O.L., McPhearson, P.T., Casey, T.M. & Morin, P.J. (1999). Environmental warming

alters food-web structure and ecosystem function. *Nature*, 402, 69-72.

Peters, R.H. (1983). *The ecological implications of body size*. Cambridge University Press,

Cambridge.

Rall, B.C., Brose, U., Hartvig, M., Kalinkat, G., Schwarzmüller, F., Vucic-Pestic, O. *et al.*

(2012). Universal temperature and body-mass scaling of feeding rates. *Philosophical*

Transactions of the Royal Society B: Biological Sciences, 367, 2923-2934.

Rastetter, E.B., Ågren, G.I. & Shaver, G.R. (1997). Responses of N-limited ecosystems to

increased CO_2: a balanced-nutrition, coupled-element-cycles model. *Ecological*

Applications, 7, 444-460.

Rip, J.M.K. & McCann, K.S. (2011). Cross-ecosystem differences in stability and the principle of

energy flux. *Ecology Letters*, 14, 733-740.

Robert W. Sterner, James J. Elser, Everett J. Fee, Stephanie J. Guildford &

Thomas H. Chrzanowski (1997). The Light: Nutrient Ratio in Lakes: The Balance of

Energy and Materials Affects Ecosystem Structure and Process. *The American Naturalist*,

150, 663-684.

Rosenzweig, M.L. (1971). Paradox of enrichment: destabilization of exploitation ecosystems in

ecological time. *Science*, 171, 385-387.

Sarmiento, J.L., Slater, R., Barber, R., Bopp, L., Doney, S.C., Hirst, A. *et al.* (2004). Response of

ocean ecosystems to climate warming. *Global Biogeochemical Cycles*, 18, 1-23.

Sentis, A., Binzer, A. & Boukal, D.S. (2017). Temperature-size responses alter food chain

persistence across environmental gradients. *Ecology Letters*, 20, 852-862.

Sentis, A., Hemptinne, J.L. & Brodeur, J. (2012). Using functional response modeling to

investigate the effect of temperature on predator feeding rate and energetic efficiency.

Oecologia, 169, 1117-1125.

Sentis, A., Hemptinne, J.L. & Brodeur, J. (2014). Towards a mechanistic understanding of

temperature and enrichment effects on species interaction strength, omnivory and food-

web structure. *Ecology Letters*, 17, 785-793.

Sterner, R.W. & Elser, J.J. (2002). *Ecological stoichiometry: the biology of elements from*

molecules to the biosphere. Princeton University Press.

Sterner, R.W. & Hessen, D.O. (1994). Algal nutrient limitation and the nutrition of aquatic

herbivores. *Annual Review of Ecology and Systematics*, 25, 1-29.

Tabi, A., Petchey, O.L. & Pennekamp, F. (2019). Warming reduces the effects of enrichment on

stability and functioning across levels of organisation in an aquatic microbial ecosystem.

Ecology Letters, 22, 1061-1071.

Thomas, M.K., Aranguren☐Gassis, M., Kremer, C.T., Gould, M.R., Anderson, K., Klausmeier,

C.A. *et al.* (2017). Temperature–nutrient interactions exacerbate sensitivity to warming in

phytoplankton. *Global Change Biology*, 23, 3269-3280.

Thomas, M.K., Kremer, C.T., Klausmeier, C.A. & Litchman, E. (2012). A global pattern of

thermal adaptation in marine phytoplankton. *Science*, 338, 1085-1088.

Tilman, D. (1982). *Resource competition and community structure*. Princeton university press.

Tranvik, L.J., Downing, J.A., Cotner, J.B., Loiselle, S.A., Striegl, R.G., Ballatore, T.J. *et al.*

(2009). Lakes and reservoirs as regulators of carbon cycling and climate. *Limnology and*

Oceanography, 54, 2298-2314.

Tylianakis, J.M., Didham, R.K., Bascompte, J. & Wardle, D.A. (2008). Global change and

species interactions in terrestrial ecosystems. *Ecology Letters*, 11, 1351-1363.

Uszko, W., Diehl, S., Englund, G. & Amarasekare, P. (2017). Effects of warming on predator–

prey interactions – a resource-based approach and a theoretical synthesis. *Ecology Letters*,

20, 513-523.

Vasseur, D.A. & McCann, K.S. (2005). A mechanistic approach for modeling temperature-

dependent consumer–resource dynamics. *American Naturalist*, 166, 184-198.

Vucic-Pestic, O., Ehnes, R.B., Rall, B.C. & Brose, U. (2011). Warming up the system: higher

predator feeding rates but lower energetic efficiencies. *Global Change Biology*, 17, 1301-

1310.

White, T. (1993). The inadequate environment. Nitrogen and the abundance of animals. Springer

Verlag, Berlin.

Woods, H.A., Makino, W., Cotner, J.B., Hobbie, S.E., Harrison, J.F., Acharya, K. *et al.* (2003).

Temperature and the chemical composition of poikilothermic organisms. *Functional*

Ecology, 17, 237-245.

Yvon-Durocher, G., Dossena, M., Trimmer, M., Woodward, G. & Allen, A.P. (2015).

Temperature and the biogeography of algal stoichiometry. *Global Ecology and*

Biogeography, 24, 562-570.

Figure legends

Fig. 1. Population fluctuations (consumer biomass coefficient of variation; panels *a* and *b*) and

species persistence (number of species; panels *c* and *d*) across the temperature (*y* axis) and

nutrient (*x* axis) gradients as predicted by the Rosenzweig-MacArthur (RM; panels *a* and *c*) and

by the Stoichiometric Rosenzweig-MacArthur (SRM; panels *b* and *d*) models. In panels *a* and *b*,

the white colour corresponds to the temperature-nutrient scenario for which the consumer has

gone extinct whereas the orange to red to dark red represent population fluctuations of increasing

amplitude. In panels *c* and *d*, in black: both consumer and resource persist; in red: only the

resource persists; in orange: none persists. Resource biomass CV is not shown; it is qualitatively

similar to the consumer biomass CV as resource and consumer biomass fluctuation are strongly

coupled.

Fig. 2. Consumer-resource biomass ratio along the temperature gradient for the Rosenzweig-

MacArthur (RM, green lines) and the Stoichiometric Rosenzweig-MacArthur (SRM, black lines)

models at three nutrient concentrations (0.008, 0.02, and 0.032 gP.m^{-3}). In each panel, the dotted

line represents biomass ratio of one; i.e. the biomass densities of the resource and the consumer

are equal. Biomass values shown at equilibrium points. For unstable equilibrium points (i.e. limit

cycles), see Fig. S2.

Fig. 3. Consumer energetic efficiency along the temperature gradient for the Rosenzweig-

MacArthur (RM, in green) and the Stoichiometric Rosenzweig-MacArthur (SRM, in black)

models at two nutrient concentrations (0.008 and 0.02 gP/m^{3}). In each panel, the dotted line

represents energetic efficiency equal to one.

Fig. 4. Population fluctuations (consumer biomass coefficient of variation) across the temperature

(y axis) and nutrient (x axis) gradients as predicted by the Rosenzweig-MacArthur (RM; panel a),

the RM with effective parameters (panel b), and the Stoichiometric Rosenzweig-MacArthur

(SRM; panel c) models.

Figure 1

802   **Figure 2**

805   **Figure 3**

811 **Figure 4**

Supplementary information

Stoichiometric constraints modulate the effects of temperature and nutrients on biomass distribution and community stability

Arnaud Sentis, Bart Haegeman, and José M. Montoya

Text S1. Derivation of the Stoichiometric Rosenzweig-MacArthur (SRM) model

The model studied in the main text is very similar to previous stoichiometric consumer-resource models (Andersen 1997; Loladze *et al.* 2000; Andersen *et al.* 2004; Diehl *et al.* 2005). To make our paper self-contained, we here present the model assumptions and derive the model equations (eqs. 3-6 in main text). Our objective was not to develop a complex and very realistic stoichiometric model that would include additional important abiotic and biotic features such as light intensity (Diehl 2007) or compensatory feeding (Cruz-Rivera & Hay 2000). Instead, we aimed at introducing two fundamental stoichiometric features (i.e. stoichiometric flexibility and stoichiometric imbalance) and investigate how these stoichiometric considerations can change predictions of the Rosenzweig-MacArthur model. We assumed that resource and consumer production are limited by energy and a single mineral nutrient. Moreover, we assume the system is closed for nutrients. Thus, nutrient supply originates exclusively from excretion and remineralization of biomass. The total amount of nutrients in the system (N_{tot}) is then a measure of nutrient enrichment. As elemental homeostasis is much stronger for consumers compared to primary producers (Andersen 1997), we assumed the nutrient quota of the consumer Q_C to be constant whereas the nutrient quota of the resource Q_R is flexible. Four differential equations determine the dynamics of four state variables, that is, the concentrations of resource (R) and consumer (C) carbon biomasses and of dissolved mineral nutrients (N), and the nutrient quota of the resource (Q_R):

$$\dot{C} = \left(e\,\frac{aR}{1+ahR} - m\right)C \tag{S1}$$

$$\dot{R} = r\left(1 - \frac{Q_{min}}{Q_R}\right)R - \frac{aR}{1+ahR}C \tag{S2}$$

$\dot{Q}_R = h(N) - r(Q_R - Qmin)$ (S3)

$\dot{N} = (Q_R - Q_C e)\frac{aR}{1+ahR}C + Q_C mC - h(N)R$ (S4)

As in the RM model, rates of change of the consumer and resource biomass densities $\dot{C}$ and $\dot{R}$

depend on their respective carbon biomass densities C and R (gC.m^{-3}), except that the resource

population growth rate follows the Droop equation (Droop 1974) and is now limited by its

nutrient quota Q_R relative to the minimum nutrient quota Q_{min}. Rate of change of Q_R depends on

the nutrient uptake rate by the resource species $h(N)$ and the amount of nutrient invested in

growth (eqn S3). $h(N)$ is the specific resource nutrient uptake rate and can be represented by a

Michaelis-Menten model where the amount of nutrient uptake saturates at high nutrient

concentrations.

With the mass-balance equation, we get that the total amount of nutrient is the sum of the free

nutrient plus the nutrient fixed in the resource biomass plus the nutrient fixed in the consumer

biomass: $N_{tot} = N + Q_R R + Q_C C$. As Eqns S1-S4 conserve total biomass (the system is closed), the

time derivative of N_{tot} is zero. We can thus replace one of the four differential equations S1-S4

with the algebraic equation $N_{tot} = N + Q_R R + Q_C C$:

$\dot{C} = (e\frac{aR}{1+ahR} - m)C$

(S5)

$\dot{R} = r(1 - \frac{Qmin}{Q_R})R - \frac{aR}{1+ahR}C$

(S6)

$\dot{N} = (Q_R - Q_C e)\frac{aR}{1+ahR}C + Q_C mC - h(N)R$

(S7)

$N_{tot} = N + Q_R R + Q_C C$

(S8)

It is possible to derive a simpler model by reducing the number of dimensions in the above model

from three to two. This model reduction is based on the assumption that free nutrients are taken

up very quickly relative to the dynamics of the consumer and resource biomasses. This

corresponds to taking $h(N)$ large, $h(N) = \frac{1}{\varepsilon}\tilde{h}(N)$ for small ε. The fast dynamics (on the

timescale t ~ ε) are

$\dot{N} = -\frac{1}{\varepsilon}\tilde{h}(N)R + (slower\ contributions)$

(S9)

Which converge to N $\rightarrow$ 0, and $Q_R \rightarrow \frac{N_{tot}-Q_C C}{R}$ with N_{tot} the total nutrient in the system. In other

words, N in dead and excreted matter is immediately recycled and acquired by the resource

species. When substituting the quasi-steady-state in eqns. (S5, S6), we get the resulting dynamics

(on the timescale t ~1):

$\dot{C} = (e\frac{aR}{1+ahR} - m)C$ (S10)

$\dot{R} = r(1 - \frac{Qmin}{Q_R})R - \frac{aR}{1+ahR}C$ (S11)

$N_{tot} = Q_R R + Q_C C$ (S12)

From the nutrient conservation equation (eqn. S12) we obtain that $Q_R = \frac{N_{tot}-Q_C C}{R}$. The intuitive

interpretation is that the resource nutrient quota Q_R decreases with the density of the resource

population and with the density of nutrient stored in the consumer biomass. In contrast to eqns

S5-S8, the reduced model has only two differential equations and one algebraic equation. It can

be equivalently written as a set of three differential equations with $\dot{C}$ and $\dot{R}$ similar as equations

S10 and S11 and with $\dot{Q}_R = \frac{d\frac{N_{tot}-Q_C C}{R}}{dt} = (Q_R - Q_C e)\frac{aR}{1+ahR}C + \frac{Q_C mC}{R} - r(Q_R - Qmin).$

In the RM model, the growth rate of the consumer population is assumed to depend only on

resource density. We relaxed this assumption by making the population growth rate of the

consumer dependent on both the resource quality (i.e. nutrient quota) and quantity (i.e. density).

In the SRM model, consumer production is also limited by resource quality as the consumer

assimilation efficiency e is a saturating function of resource nutrient quota Q_R:

$$e(Q_R) = e_{max} \frac{Q_R}{Q_R + Q_C} \tag{S13}$$

The intuitive interpretation of eqn. S13 is that resource quality is not a limiting factor for

consumer growth as long as the nutrient content of the resource is superior to the nutrient content

of the consumer (i.e. $Q_R > Q_C$). In other words, when $Q_R \gg Q_C$, $e(Q_R) \rightarrow e_{max}$ and when $Q_R \ll$

Q_C, $e(Q_R) \rightarrow 0$. By replacing e by $e(Q_R)$ in eqn. S10, we obtain the SRM model.

Text S2. Differences in biomass ratios predicted by the two models

Here we show that the equilibrium consumer-to-resource biomass ratio in the model with
stoichiometric constraints (SRM model) is always smaller than the one in the model without
stoichiometric constraints (RM model), keeping the same parameter values. For simplicity we
assume for both models that the consumer and the resource persist at equilibrium, and we do not
consider the stability of the equilibrium point (in particular, the equilibrium might be unstable at
the center of a limit cycle). We indicate the equilibrium values of the non-stoichiometric model
by the superscript "ns" and the equilibrium values of the stoichiometric model by the superscript
"s". We use the same superscripts to distinguish the assimilation efficiencies of both models.

Model without stoichiometric constraints

The model is defined as

$\dot{R} = r(1 - \frac{R}{K})R - f(R)C$ (S14)

$\dot{C} = (e^{ns}f(R) - m)C$ (S15)

With $f(R) = \frac{aR}{1+ahR}$ and $K = N_{\text{tot}}/Q_{\text{min}}$.

By solving equation (S15) we get the resource biomass at equilibrium:

$R^{ns} = \frac{1}{a(\frac{e^{ns}}{m}-h)}$ (S16)

From equation (S14) we get the consumer biomass at equilibrium. It follows from $f(R^{ns})C^{ns} =$
$r(1 - \frac{R^{ns}}{K})R^{ns}$, or $C^{ns} = r(1 - \frac{R^{ns}}{K})\frac{R^{ns}}{f(R^{ns})}$

Hence, the consumer-to-resource biomass ratio is

$\frac{C^{ns}}{R^{ns}} = \frac{re^{ns}}{m}(1 - \frac{R^{ns}}{K}) = \frac{re^{ns}}{m}(1 - \frac{Q_{min}R^{ns}}{N_{tot}})$ (S17)

Model with stoichiometric constraints

The model is defined as

$\dot{R} = r(1 - \frac{Q_{min}}{Q_R})R - f(R)C$ (S18)

$\dot{C} = (e^s(Q_R)f(R) - m)C$ (S19)

With $e^s(Q_R) = e_{max}\frac{Q_R}{Q_C+Q_R}$, $f(R) = \frac{aR}{1+ahR}$ and $N_{tot} = Q_R R + Q_C C$.

From equation (S19) we have $e^s(Q_R^s)f(R^s) = m$, or

$$R^s = \frac{1}{a(\frac{e^s(Q_R^s)}{m}-h)}$$ (S20)

From equation (S18) we have from $(R^s)C^s = r(1 - \frac{Q_{min}}{Q_R^s})R^s$, or

$$C^s = \frac{re^s(Q_R^s)}{m}(1 - \frac{Q_{min}}{Q_R^s})R^s$$

Hence, the consumer-to-resource biomass ratio is

$$\frac{C^s}{R^s} = \frac{re^s(Q_R^s)}{m}(1 - \frac{Q_{min}}{Q_R^s}) = \frac{re^s(Q_R^s)}{m}(1 - \frac{R^s Q_{min}}{N_{tot}-Q_C C^s})$$ (S21)

We now compare the biomass ratios of equations (S17) and (S21). We have $e^{ns} = e_{max}$, and hence

$$e^{ns} \geq e^s(Q_R^s).$$ (S22)

Using this inequality, we get $a\left(\frac{e^{ns}}{m} - h\right) \geq a(\frac{e^s(Q_R^s)}{m} - h)$, and by equations (S16) and (S20), we

see that

$$R^{ns} \leq R^s$$ (S23)

Clearly, we always have $N_{tot} \geq N_{tot} - Q_C C^s$. Combining this with equation (S23), we get

$$\frac{R^{ns}}{N_{tot}} \leq \frac{R^s}{N_{tot}-Q_C C^s} \text{ and}$$

$$1 - \frac{Q_{min}R^{ns}}{N_{tot}} \geq 1 - \frac{R^s Q_{min}}{N_{tot}-Q_C C^s}$$ (S24)

Finally, from equations (S22) and (S24),

$$\frac{re^{ns}}{m}(1 - \frac{Q_{min}R^{ns}}{N_{tot}}) \geq \frac{re^s(Q_R^s)}{m}\left(1 - \frac{R^s Q_{min}}{N_{tot}-Q_C C^s}\right),$$ (S25)

showing that $\frac{C^{ns}}{R^{ns}} \geq \frac{C^s}{R^s}$.

Table S1. Definitions and units of model parameters, from Uszko *et al.* (2017). For temperature-

dependent parameters, we list the value of the scaling constant Q_0 (in units of the parameter) and

the values of either the activation energy E_Q (eV, when temperature dependence is monotonous,

eqn. 7) or of the temperature T_{opt} (Kelvin) at which the parameter value reaches a

maximum/minimum and the width s (Kelvin) of this bell-/U-shaped function (when temperature

dependence is non-monotonous, eqn. 8). Biomass and nutrients are expressed in units of carbon

(C) and phosphorus (P), respectively

Parameter	Temperature independent value	Unit	Definition	Reference
	Thermal parameters			
r	$r_0 = 2.2$; $T_{opt} = 298.15$; $s = 12.0$	1/d	Intrinsic rate of resource net production (gross production – biosynthesis costs)	Uszko et al. 2017
h	$h_0 = 0.17$; $T_{opt} = 294.1$; $s = 6.4$	d	Handling time	Uszko et al. 2017
a	$a_0 = 8.9$; $T_{opt} = 296.0$; $s = 9.4$	m^3/(gC d)	Attack rate	Uszko et al. 2017
m	$m_0 = 4.4 \times 10^8$; $E_m = 0.55$	1/d	Consumer mortality plus maintenance rate	
e_{max}	0.385	-	Maximum assimilation efficiency	Yodzis
Q_C	0.042	g P/g C	Consumer P:C ratio	Diehl 2005
Q_{min}	0.009	g P/g C	Minimum nutrient quota	Diehl 2005
Q_R	Variable	g P/g C	Resource P:C ratio	
N_{tot}	Variable	g P/m^3	Total nutrients in the system	
T	Variable	K	Temperature	

959 **Fig. S1.** Thermal functions used to parametrize the model (adapted from Uszko et al. 2017)

Fig. S2. Consumer-resource biomass ratio (log scale) along the temperature gradient for the

Rosenzweig-MacArthur (RM, green lines) and the Stoichiometric Rosenzweig-MacArthur (SRM,

black lines) models at three nutrient concentrations (0.008, 0.02, and 0.032 gP.m^{-3}). In each

panel, the dotted lines represent unstable solutions whereas full lines represent stable solutions.

The thin horizontal dotted line represents biomass ratio of one; i.e. the biomass densities of the

resource and the consumer are equal.

Fig. S3. Effective assimilation efficiency e_{ef} and carrying capacity K_{ef} from the

Stoichiometric Rosenzweig-MacArthur (SRM) model along the temperature gradient at

three nutrient levels (0.004, 0.015, and 0.032 gP.m^{-3}) with $Q_C = 0.042$. Full lines represent

temperature and nutrient scenarios for which both the resource and consumer persist

whereas dotted lines represent scenarios for which only the resource persists. Effective

assimilation efficiency was calculated as $e_{ef} = e_{max}Q_R/(Q_R+Q_C)$, with Q_R the equilibrium

solution of the SRM model and the effective carrying capacity as $K_{ef} = Q_R R/Q_{min} = (N_{tot}-$

$Q_C C)/Q_{min}$, with Q_R, R and C the equilibrium solutions of the SRM model.

References

Ahlgren, G. (1987). Temperature functions in biology and their application to algal growth constants. *Oikos*, 49, 177-190.

Amarasekare, P. (2015). Effects of temperature on consumer-resource interactions. *Journal of Animal Ecology*, 84, 665-679.

Amarasekare, P. & Coutinho, R.M. (2014). Effects of temperature on intraspecific competition in ectotherms. *The American Naturalist*, 184, E50-E65.

Andersen, T. (1997). *Pelagic nutrient cycles: herbivores as sources and sinks*. Springer-Verlag, Berlin, Germany.

Andersen, T., Elser, J.J. & Hessen, D.O. (2004). Stoichiometry and population dynamics. *Ecology Letters*, 7, 884-900.

Andersen, T. & Hessen, D.O. (1991). Carbon, nitrogen, and phosphorus content of freshwater zooplankton. *Limnology and Oceanography*, 36, 807-814.

Anneville, O., Gammeter, S. & Straile, D. (2005). Phosphorus decrease and climate variability: mediators of synchrony in phytoplankton changes among European peri-alpine lakes. *Freshwater Biology*, 50, 1731-1746.

Barbier, M. & Loreau, M. (2019). Pyramids and cascades: a synthesis of food chain functioning and stability. *Ecology Letters*, 22, 405-419.

Bezemer, T.M. & Jones, T.H. (1998). Plant-insect herbivore interactions in elevated atmospheric C02: quantitative analyses and guild effects. *Oikos*, 82, 212-222.

Binzer, A., Guill, C., Brose, U. & Rall, B.C. (2012). The dynamics of food chains under climate change and nutrient enrichment. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 367, 2935-2944.

Binzer, A., Guill, C., Rall, B.C. & Brose, U. (2016). Interactive effects of warming, eutrophication and size structure: impacts on biodiversity and food‐web structure. *Global Change Biology*, 22, 220-227.

Boit, A., Martinez, N.D., Williams, R.J. & Gaedke, U. (2012). Mechanistic theory and modelling of complex food-web dynamics in Lake Constance. *Ecology Letters*, 15, 594-602.

Borer, E.T., Bracken, M.E., Seabloom, E.W., Smith, J.E., Cebrian, J., Cleland, E.E. *et al.* (2013). Global biogeography of autotroph chemistry: is insolation a driving force? *Oikos*, 122, 1121-1130.

Brown, J.H., Gillooly, J.F., Allen, A.P., Savage, V.M. & West, G.B. (2004). Toward a metabolic theory of ecology. *Ecology*, 85, 1771-1789.

Cruz-Rivera, E. & Hay, M.E. (2000). Can quantity replace quality? Food choice, compensatory feeding, and fitness of marine mesograzers. *Ecology*, 81, 201-219.

Del Giorgio, P.A. & Gasol, J.M. (1995). Biomass Distribution in Freshwater Plankton Communities. *The American Naturalist*, 146, 135-152.

Diehl, S. (2007). Paradoxes of enrichment: effects of increased light versus nutrient supply on pelagic producer-grazer systems. *The American Naturalist*, 169, E173-E191.

Diehl, S., Berger, S. & Wöhrl, R. (2005). Flexible nutrient stoichiometry mediates environmental influences on phytoplankton and its resources. *Ecology*, 86, 2931-2945.

Droop, M. (1974). The nutrient status of algal cells in continuous culture. *Journal of the Marine Biological Association of the United Kingdom*, 54, 825-855.

Elser, J., Sterner, R., Gorokhova, E.a., Fagan, W., Markow, T., Cotner, J. *et al.* (2000). Biological stoichiometry from genes to ecosystems. *Ecology Letters*, 3, 540-550.

Elser, J.J., Bracken, M.E., Cleland, E.E., Gruner, D.S., Harpole, W.S., Hillebrand, H. *et al.* (2007). Global analysis of nitrogen and phosphorus limitation of primary producers in freshwater, marine and terrestrial ecosystems. *Ecology Letters*, 10, 1135-1142.

Elser, J.J., Loladze, I., Peace, A.L. & Kuang, Y. (2012). Lotka re-loaded: modeling trophic
interactions under stoichiometric constraints. *Ecological Modelling*, 245, 3-11.
Elton, C. (1927). *Animal ecology*. Sidgwick & Jackson, LTD, London.
Englund, G., Ohlund, G., Hein, C.L. & Diehl, S. (2011). Temperature dependence of the
functional response. *Ecology Letters*, 14, 914-921.
Enquist, B.J., West, G.B., Charnov, E.L. & Brown, J.H. (1999). Allometric scaling of production
and life-history variation in vascular plants. *Nature*, 401, 907-911.
Falkowski, P.G., Barber, R.T. & Smetacek, V. (1998). Biogeochemical controls and feedbacks on
ocean primary production. *Science*, 281, 200-206.
Finkel, Z.V., Beardall, J., Flynn, K.J., Quigg, A., Rees, T.A.V. & Raven, J.A. (2009).
Phytoplankton in a changing world: cell size and elemental stoichiometry. *Journal of*
Plankton Research, 32, 119-137.
Fussmann, K.E., Schwarzmüller, F., Brose, U., Jousset, A. & Rall, B.C. (2014). Ecological
stability in response to warming. *Nature Climate Change*, 4, 206-210.
Gilarranz, L.J., Mora, C. & Bascompte, J. (2016). Anthropogenic effects are associated with a
lower persistence of marine food webs. *Nature communications*, 7, 10737.
Gilbert, B., Tunney, T.D., McCann, K.S., DeLong, J.P., Vasseur, D.A., Savage, V. *et al.* (2014).
A bioenergetic framework for the temperature dependence of trophic interactions.
Ecology Letters, 17, 902-9014.
Hessen, D.O., Ågren, G.I., Anderson, T.R., Elser, J.J. & De Ruiter, P.C. (2004). Carbon
sequestration in ecosystems: the role of stoichiometry. *Ecology*, 85, 1179-1192.
Hessen, D.O., Færøvig, P.J. & Andersen, T. (2002). Light, nutrients, and P:C ratios in algae:
grazer performance related to food quality and quantity. *Ecology*, 83, 1886-1898.
Iles, A.C. (2014). Towards predicting community level effects of climate: relative temperature
scaling of metabolic and ingestion rates. *Ecology*, 95, 2657–2668.
Irigoien, X., Huisman, J. & Harris, R.P. (2004). Global biodiversity patterns of marine
phytoplankton and zooplankton. *Nature*, 429, 863-867.
Jensen, C.X. & Ginzburg, L.R. (2005). Paradoxes or theoretical failures? The jury is still out.
Ecological Modelling, 188, 3-14.
Jeschke, J.M., Kopp, M. & Tollrian, R. (2004). Consumer food systems: why Type I functional
responses are exclusive to filter feeders. *Biological Reviews*, 79, 337-349.
Jonsson, T. (2017). Conditions for eltonian pyramids in Lotka-Volterra food chains. *Scientific*
Reports, 7, 10912.
Kratina, P., Greig, H.S., Thompson, P.L., Carvalho-Pereira, T.S.A. & Shurin, J.B. (2012).
Warming modifies trophic cascades and eutrophication in experimental freshwater
communities. *Ecology*, 93, 1421-1430.
Lindeman, R.L. (1942). The trophic dynamic aspect of ecology. *Ecology*, 23, 399-417.
Loladze, I., Kuang, Y. & Elser, J.J. (2000). Stoichiometry in producer-grazer systems: linking
energy flow with element cycling. *Bulletin of Mathematical Biology*, 62, 1137-1162.
McAllister, C., LeBrasseur, R., Parsons, T. & Rosenzweig, M. (1972). Stability of enriched
aquatic ecosystems. *Science*, 175, 562-565.
McCauley, D.J., Gellner, G., Martinez, N.D., Williams, R.J., Sandin, S.A., Micheli, F. *et al.*
(2018). On the prevalence and dynamics of inverted trophic pyramids and otherwise top-
heavy communities. *Ecology Letters*, 21, 439-454.
McCauley, E. & Kalff, J. (1981). Empirical relationships between phytoplankton and
zooplankton biomass in lakes. *Canadian Journal of Fisheries and Aquatic Sciences*, 38,
458-463.

McCauley, E., Nisbet, R.M., Murdoch, W.W., de Roos, A.M. & Gurney, W.S.C. (1999). Large-amplitude cycles of *Daphnia* and its algal prey in enriched environments. *Nature*, 402, 653-656.

Montoya, J.M. & Raffaelli, D. (2010). Climate change, biotic interactions and ecosystem services. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 365, 2013-2018.

Nelson, G.C. (2005). *Millennium ecosystem assessment: drivers of ecosystem change: summary chapter*. World Resources Institute, Washington, DC.

O'Connor, M.I., Piehler, M.F., Leech, D.M., Anton, A. & Bruno, J.F. (2009). Warming and resource availability shift food web structure and metabolism. *PLoS Biology*, 7, e1000178.

Petchey, O.L., McPhearson, P.T., Casey, T.M. & Morin, P.J. (1999). Environmental warming alters food-web structure and ecosystem function. *Nature*, 402, 69-72.

Peters, R.H. (1983). *The ecological implications of body size*. Cambridge University Press, Cambridge.

R Development Core Team (2017). *R: a language and environment for statistical computing*, R Foundation for Statistical Computing, Vienna, Austria.

Rall, B.C., Brose, U., Hartvig, M., Kalinkat, G., Schwarzmüller, F., Vucic-Pestic, O. *et al.* (2012). Universal temperature and body-mass scaling of feeding rates. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 367, 2923-2934.

Rastetter, E.B., Ågren, G.I. & Shaver, G.R. (1997). Responses of N-limited ecosystems to increased CO_2: a balanced-nutrition, coupled-element-cycles model. *Ecological Applications*, 7, 444-460.

Rip, J.M.K. & McCann, K.S. (2011). Cross-ecosystem differences in stability and the principle of energy flux. *Ecology Letters*, 14, 733-740.

Robert W. Sterner, James J. Elser, Everett J. Fee, Stephanie J. Guildford & Thomas H. Chrzanowski (1997). The Light: Nutrient Ratio in Lakes: The Balance of Energy and Materials Affects Ecosystem Structure and Process. *The American Naturalist*, 150, 663-684.

Rosenzweig, M.L. (1971). Paradox of enrichment: destabilization of exploitation ecosystems in ecological time. *Science*, 171, 385-387.

Sarmiento, J.L., Slater, R., Barber, R., Bopp, L., Doney, S.C., Hirst, A. *et al.* (2004). Response of ocean ecosystems to climate warming. *Global Biogeochemical Cycles*, 18, 1-23.

Sentis, A., Binzer, A. & Boukal, D.S. (2017). Temperature-size responses alter food chain persistence across environmental gradients. *Ecology Letters*, 20, 852-862.

Sentis, A., Hemptinne, J.L. & Brodeur, J. (2012). Using functional response modeling to investigate the effect of temperature on predator feeding rate and energetic efficiency. *Oecologia*, 169, 1117-1125.

Sentis, A., Hemptinne, J.L. & Brodeur, J. (2014). Towards a mechanistic understanding of temperature and enrichment effects on species interaction strength, omnivory and food-web structure. *Ecology Letters*, 17, 785-793.

Soetaert, K., Cash, J. & Mazzia, F. (2012). *Solving differential equations in R*. Springer Science & Business Media.

Sterner, R.W. & Elser, J.J. (2002). *Ecological stoichiometry: the biology of elements from molecules to the biosphere*. Princeton University Press.

Sterner, R.W. & Hessen, D.O. (1994). Algal nutrient limitation and the nutrition of aquatic herbivores. *Annual Review of Ecology and Systematics*, 25, 1-29.

Tabi, A., Petchey, O.L. & Pennekamp, F. (2019). Warming reduces the effects of enrichment on
stability and functioning across levels of organisation in an aquatic microbial ecosystem.
Ecology Letters, 22, 1061-1071.
Thomas, M.K., Aranguren☐Gassis, M., Kremer, C.T., Gould, M.R., Anderson, K., Klausmeier,
C.A. *et al.* (2017). Temperature–nutrient interactions exacerbate sensitivity to warming in
phytoplankton. *Global Change Biology*, 23, 3269-3280.
Thomas, M.K., Kremer, C.T., Klausmeier, C.A. & Litchman, E. (2012). A global pattern of
thermal adaptation in marine phytoplankton. *Science*, 338, 1085-1088.
Tilman, D. (1982). *Resource competition and community structure*. Princeton university press.
Tranvik, L.J., Downing, J.A., Cotner, J.B., Loiselle, S.A., Striegl, R.G., Ballatore, T.J. *et al.*
(2009). Lakes and reservoirs as regulators of carbon cycling and climate. *Limnology and*
Oceanography, 54, 2298-2314.
Tylianakis, J.M., Didham, R.K., Bascompte, J. & Wardle, D.A. (2008). Global change and
species interactions in terrestrial ecosystems. *Ecology Letters*, 11, 1351-1363.
Uszko, W., Diehl, S., Englund, G. & Amarasekare, P. (2017). Effects of warming on predator–
prey interactions – a resource-based approach and a theoretical synthesis. *Ecology Letters*,
20, 513-523.
Vasseur, D.A. & McCann, K.S. (2005). A mechanistic approach for modeling temperature-
dependent consumer–resource dynamics. *American Naturalist*, 166, 184-198.
Vucic-Pestic, O., Ehnes, R.B., Rall, B.C. & Brose, U. (2011). Warming up the system: higher
predator feeding rates but lower energetic efficiencies. *Global Change Biology*, 17, 1301-
1310.
White, T. (1993). The inadequate environment. Nitrogen and the abundance of animals. Springer
Verlag, Berlin.
Woods, H.A., Makino, W., Cotner, J.B., Hobbie, S.E., Harrison, J.F., Acharya, K. *et al.* (2003).
Temperature and the chemical composition of poikilothermic organisms. *Functional*
Ecology, 17, 237-245.
Yvon-Durocher, G., Dossena, M., Trimmer, M., Woodward, G. & Allen, A.P. (2015).
Temperature and the biogeography of algal stoichiometry. *Global Ecology and*
Biogeography, 24, 562-570.
1158

1159