Balanced activity of multiple FtsW and RodA proteins is crucial for maintaining cell

shape and antibiotic resistance in *Listeria monocytogenes*

Jeanine Rismondo[1], Sven Halbedel[2] and Angelika Gründling[1#]

[1] Section of Microbiology and Medical Research Council Centre for Molecular Bacteriology

and Infection, Imperial College London, London SW7 2AZ.

[2] FG11 Division of Enteropathogenic bacteria and *Legionella*, Robert Koch-Institute,

Burgstrasse 37, 38855 Wernigerode, Germany.

[#]To whom correspondence should be addressed: Angelika Gründling –

a.grundling@imperial.ac.uk

22 Abstract word count: 247

23 Main text word count: 5957

Running Title: Functional analysis of multiple SEDS in *Listeria*

Abstract

Rod-shaped bacteria have two modes of peptidoglycan synthesis: lateral synthesis and

synthesis at the cell division site. These two processes are controlled by two macromolecular

protein complexes, the Rod complex and divisome, respectively. Recently, it has been shown

that the *Bacillus subtilis* RodA protein, which forms part of the Rod complex, has

peptidoglycan glycosyltransferase activity. The cell division specific RodA homolog FtsW

fulfils a similar role at the divisome. The human pathogen *Listeria monocytogenes* encodes

up to six FtsW/RodA homologs, however their functions have not yet been investigated.

Analysis of deletion and depletion strains led to the identification of the essential cell

division-specific FtsW protein, FtsW1. Interestingly, *L. monocytogenes* encodes a second

FtsW protein, FtsW2, which can compensate for the lack of FtsW1, when expressed from an

inducible promoter. *L. monocytogenes* also possesses three RodA homologs, RodA1, RodA2

and RodA3 and their combined absence is lethal. Cells of a *rodA1/rodA3* double mutant

strain are shorter and have increased antibiotic and lysozyme sensitivity, probably due to a

weakened cell wall. Results from promoter activity assays revealed that expression of *rodA3*

and *ftsW2* is induced in the presence of antibiotics targeting penicillin binding proteins.

Consistent with this, a *rodA3* mutant was also more susceptible to the ß-lactam antibiotic

cefuroxime. Taken together, our study highlights that *L. monocytogenes* encodes a multitude

of functional FtsW and RodA enzymes to produce its rigid cell wall and that their expression

needs to be tightly regulated to maintain growth, cell division and antibiotic resistance.

Importance

The human pathogen *Listeria monocytogenes* is usually treated with high doses of ß-lactam

antibiotics, often combined with gentamicin. However, these antibiotics only act

bacteriostatically on *L. monocytogenes* and the immune system is needed to clear the

infection. Therefore, individuals with a compromised immune system are at risk to develop a

severe form of *Listeria* infection, which can be fatal in up to 30% of cases. The development

of new strategies to treat *Listeria* infections is therefore necessary. Here we show that the

expression of some of the FtsW and RodA enzymes of *L. monocytogenes* is induced by the

presence of ß-lactam antibiotics and their combined absence makes bacteria more susceptible

to this class of antibiotics. The development of antimicrobials that inhibit the activity or

production of FtsW/RodA enzymes might therefore help to improve the treatment of *Listeria*

infections and thereby lead to a reduction in mortality.

Introduction

Bacterial cells are surrounded by a mesh of peptidoglycan (PG) that determines their shape and also protects the cells from lysis due to their high internal turgor pressure (Weidel and Pelzer, 1964, Vollmer et al., 2008). Peptidoglycan is comprised of glycan strands that are crosslinked by short peptides (Rogers et al., 1980). The glycan strands are composed of alternating N-acetylglucosamine and N-acetylmuramic acid residues that are connected by a ß-1,4 glycosidic bond (Ghuysen and Strominger, 1963). The synthesis of peptidoglycan begins in the cytoplasm with the production of the PG precursor lipid II by the proteins MurABCDEF, MraY and MurG (Blumberg and Strominger, 1974, van Heijenoort, 2001, Scheffers and Pinho, 2005, Pinho et al., 2013). Lipid II is then transported across the cytoplasmic membrane by the flippase MurJ and Amj (Meeske et al., 2015, Sham et al., 2014, Ruiz, 2008) and subsequently incorporated in the growing glycan strand by glycosyltransferases. The polymerization and the crosslinking of the glycan strands are facilitated by the activity of glycosyltransferases and transpeptidases, respectively. Class A penicillin binding proteins (PBPs) are bifunctional enzymes that possess glycosyltransferase and transpeptidase activity, whereas class B PBPs only contain a transpeptidase domain (Höltje, 1998, Sauvage et al., 2008, Goffin and Ghuysen, 1998). In addition, some species such as *Escherichia coli*, *Staphylococcus aureus* and *Streptococcus pneumoniae* encode monofunctional glycosyltransferases (MGTs) that can also incorporate lipid II into the growing glycan strand (Park and Matsuhashi, 1984, Park et al., 1985, Karinou et al., 2018, Wang et al., 2001, Hara and Suzuki, 1984).

B. subtilis encodes four class A PBPs and no MGT, however, deletion of all class A PBPs only manifests in small PG changes (McPherson and Popham, 2003). Recently, it has been shown that members of the SEDS (shape, elongation, division, sporulation) family of proteins, namely RodA and FtsW, also act as glycosyltransferases (Meeske et al., 2016, Cho et al., 2016, Emami et al., 2017, Taguchi et al., 2019). Both, RodA and FtsW, form

complexes with cognate class B PBPs to enable polymerization and crosslinking of glycan strands (Cho et al., 2016, Taguchi et al., 2019, Leclercq et al., 2017, Fraipont et al., 2011). Interestingly, SEDS proteins and the class B PBPs are more conserved among different bacterial species than class A PBPs (Meeske et al., 2016).

In rod-shaped bacteria, peptidoglycan is synthesized by two multiprotein complexes, the rod complex that is essential for the cell elongation and the divisome that is crucial for the formation of the division septum (Ricard and Hirota, 1973, Nanninga, 1991, Carballido-Lopez and Formstone, 2007). RodA is part of the rod complex and is essential in many bacteria including *B. subtilis* and *S. pneumoniae* (Liu et al., 2017, Henriques et al., 1998). Depletion of RodA results in the production of enlarged, spherical cells in *B. subtilis* (Henriques et al., 1998). In contrast, FtsW is essential for cell division and cells depleted for FtsW grow as long filaments (Kobayashi et al., 2003, Boyle et al., 1997, Gamba et al., 2016). *B. subtilis* harbors a sporulation specific member of the SEDS family, SpoVE in addition to RodA and FtsW. SpoVE is dispensable for growth, however, it is essential for the synthesis of the spore cortex peptidoglycan (Henriques et al., 1992, Ikeda et al., 1989). Other *Bacillus* species such as *B. cereus* and *B. anthracis* possess 4 to 5 FtsW/RodA proteins and strains of different serotypes of the human pathogen *Listeria monocytogenes* encode even up to 6 FtsW/RodA homologs in their genome. However, their functions have not yet been investigated.

Here, we determine the role of the different FtsW and RodA homologs for the growth and cell morphology of *L. monocytogenes*. Our results show that *L. monocytogenes* encodes two FtsW and three RodA enzymes. Absence of either FtsW1 or of all three RodA proteins is lethal under standard laboratory conditions. *L. monocytogenes* infections are usually treated with high doses of ß-lactam antibiotics such as ampicillin, which inhibit the transpeptidase activity of PBPs (Swaminathan and Gerner-Smidt, 2007). We demonstrate that the expression

of two SEDS proteins, FtsW2 and RodA3, is induced in the presence of ß-lactam antibiotics

likely to compensate for the inhibition of PBPs and that a *rodA3* mutant is more sensitive to

the ß-lactam antibiotic cefuroxime. Antimicrobials inhibiting the activity of proteins of the

SEDS family could therefore potentially improve the treatment of *Listeria* infections in the

future.

Materials and Methods

Bacterial strains and growth conditions

All strains and plasmids used in this study are listed in Table S1. Strain and plasmid

constructions are described in the supplemental materials and method section and all primers

used in this study are listed in Table S2. *E. coli* strains were grown in Luria-Bertani (LB)

medium and *L. monocytogenes* strains in brain heart infusion (BHI) medium at 37°C unless

otherwise stated. If necessary, antibiotics and supplements were added to the medium at the

following concentrations: for *E. coli* cultures, ampicillin (Amp) at 100 µg/ml and kanamycin

(Kan) at 30 µg/ml, and for *L. monocytogenes* cultures, chloramphenicol (Cam) at 10 µg/ml,

kanamycin (Kan) at 30 µg/ml and Isopropyl β-D-1-thiogalactopyranoside (IPTG) at 1 mM.

We used the *L. monocytogenes* strain 10403S and derivatives thereof. However, we refer to

L. monocytogenes EGD-e gene and locus tag numbers as this was the first fully sequenced *L.*

monocytogenes strain.

Growth curves

Overnight cultures of wildtype *L. monocytogenes* 10403S and the indicated deletion strains

were diluted to an OD_{600} of 0.01 or 0.05 in 15 ml BHI medium and the cultures were

incubated at 37°C with shaking. Growth was monitored by determining OD_{600} readings at

hourly intervals. For growth curves with the IPTG-inducible depletion strains 10403SΔ*ftsW1*

i*ftsW* (ANG4314), 10403SΔ*ftsW1* i*ftsW2* (ANG5119) and 10403SΔ*rodA1-3* i*rodA1*

(ANG5192), the strains were cultivated overnight in the presence of 1 mM IPTG. The next

day, cells were washed once with fresh medium, subsequently diluted 1:50 in 5 ml BHI

medium and grown for 8-10 h in the absence of the inducer. The cultures were diluted 1:100

into fresh BHI medium and grown until the next morning at 37°C. The depleted cells were

then diluted to an OD_{600} of 0.01 and grown in the presence or absence of 1 mM IPTG at

37°C. Averages and standard deviations from three independent experiments were plotted.

Determination of minimal inhibitory concentration (MIC)

The minimal inhibitory concentration for bacitracin, penicillin and moenomycin and

lysozyme was determined using a microbroth dilution assay in 96-well plates. Approximately

10^4 *L. monocytogenes* cells were used to inoculate 200 µl BHI containing two-fold dilutions

of the different antimicrobials. The starting antibiotic concentrations were: 1 mg/ml for

bacitracin A, 1 µg/ml for penicillin G, 0.8 or 1.6 µg/ml for moenomycin, 8 µg/ml cefuroxime

and 10 mg/ml for lysozyme. The OD_{600} was determined after incubating the 96-well plates

for 24 h at 37°C with shaking at 500 rpm in a plate incubator (Thermostar, BMG Labtech).

The MIC value refers to the antibiotic concentration at which bacterial growth was inhibited

by >90%.

Determination of antibiotic susceptibility using a spot plating assay

Overnight cultures of the indicated *L. monocytogenes* strains were adjusted to an OD_{600} of 1

and 5 µl of serial dilutions were spotted on BHI agar plates or BHI agar plates containing 1

µg/ml cefuroxime. Plates were photographed after incubation at 37°C for 24 h.

Fluorescence and phase contrast microscopy

For bacterial cell length measurements, 100 µl of mid-log cultures were mixed with 5 µl of

100 µg/ml nile red solution to stain the cell membrane. After an incubation of 20 min at

37°C, the cells were washed once with 1x PBS and resuspended in 50 µl 1x PBS. 1.5 µl of

the different samples were spotted on microscope slides covered with a thin agarose film (1.5

% agarose in distilled water), air-dried and covered with a cover slip. Phase contrast and

fluorescence images were taken at 100x magnification using the Zeiss Axio Imager.A1

microscope coupled to the AxioCam MRm and processed using the Zen 2012 software (blue

edition). For the detection of nile red fluorescence signals, the Zeiss filter set 00 was used.

For the determination of the cell lengths, 300 cells were measured for each experiment and

the median cell length was calculated. Averages and standard deviations of the median cell

length of three independent experiments were plotted.

Peptidoglycan isolation and analysis

Overnight cultures of *L. monocytogenes* 10403S, 10403SΔ*rodA1*Δ*rodA3* and

10403SΔ*rodA1*Δ*rodA3* pIMK3-*rodA1* were used to inoculate 1 L BHI broth (with 1 mM

IPTG for the complementation strain 10403SΔ*rodA1*Δ*rodA3* pIMK3-*rodA1*) to a starting

OD$_{600}$ of 0.06. The cultures were grown at 37°C until they reached an OD$_{600}$ of 1, at which

point the cultures were cooled on ice for 1 h. The bacteria were subsequently collected by

centrifugation and peptidoglycan was purified and digested with mutanolysin as described

previously (de Jonge et al., 1992, Corrigan et al., 2011). Digested muropeptides were

analyzed by high-performance liquid chromatography (HPLC) and recorded at an absorption

of 205 nm as described previously (de Jonge et al., 1992). For quantification, the areas of the

main muropeptide peaks were integrated using the Agilent Technology ChemStation

software. The sum of the peak areas was set to 100% and individual peak areas were

determined. Averages and standard deviations from three independent extractions were

calculated.

 β-galactosidase assay

 For the determination of the β-galactosidase activity, overnight cultures of strains 10403S

 pPL3e-P$_{lmo2689}$-*lacZ*, 10403SΔ*rodA1* pPL3e-P$_{lmo2689}$-*lacZ* and 10403SΔ*rodA1*Δ*rodA2* pPL3e-

 P$_{lmo2689}$-*lacZ* were diluted 1:100 in fresh BHI medium and grown for 6 h at 37°C. Sample

 collection and preparation were performed as described previously (Gründling et al., 2004).

 Briefly, OD$_{600}$ readings were determined (for the final β-galactosidase unit calculations) for

 the different cultures after 6 h of growth and cells from 1 ml culture were pelleted by

 centrifugation for 10 min at 13,200 x g, resuspended in 100 µl ABT buffer (60 mM K$_2$HPO$_4$,

 40 mM KH$_2$PO$_4$, 100 mM NaCl, 0.1% Triton X-100, pH 7.0, filter sterilized), snap frozen in

 liquid nitrogen and stored at -80°C until use. For the identification of substances inducing the

 expression of the *lmo2689-lmo2686* operon, an overnight culture of strain 10403S pPL3e-

 P$_{lmo2689}$-*lacZ* was diluted 1:100 in fresh BHI medium and the culture incubated with shaking

 at 37°C until an OD$_{600}$ of 0.5-0.6. The culture was divided into several flasks and incubated

 for two hours at 37°C in the presence or absence of the following substances: 0.5 µg/ml

 ampicillin, 0.05 µg/ml penicillin, 0.5 µg/ml vancomycin, 4 µg/ml cefuroxime, 0.05 µg/ml

 moenomycin, 0.5 mg/ml lysozyme, 1% ethanol, 300 µg/ml MgSO$_4$ or 300 µg/ml EDTA.

 After 2 h, bacterial cultures were harvested and frozen as described above.

 Samples were thawed and 1:10 dilutions were prepared in ABT buffer. 50 µl of the 1:10

 diluted samples were mixed with 10 µl of 0.4 mg/ml 4-methyl-umbelliferyl-ß-D-

 galactopyranoside (MUG) substrate prepared in DMSO and incubated for 60 min at room

 temperature (RT). A reaction with ABT buffer alone was used as negative control. Following

 this incubation step, 20 µl of each reaction was diluted into 180 µl of ABT buffer in a black

 96-well plate and fluorescence values were measured using an HIDEX Sense Microplate

 Reader at 355 nm excitation and 460 nm emission wavelengths. 0.125-20 µM of the

 fluorescent 4-methylumbelliferone (MU) standard were used to obtain a standard curve. β-

galactosidase units were calculated as (pmol of substrate hydrolyzed x dilution factor)/(ml

culture volume x OD_{600} x minute). The amount of hydrolyzed substrate was determined from

the standard curve as (emission reading – y intercept)/slope.

Results

***L. monocytogenes* 10403S encodes six FtsW/RodA homologs**

So far, FtsW and RodA proteins of the human pathogen *L. monocytogenes* have not been

studied. FtsW and RodA are members of the SEDS (shape, elongation, division, sporulation)

family of proteins and are multispanning membrane proteins with 8-10 transmembrane

helices and a large extracellular loop (Fig. 1A). Using BLAST, six proteins with homology to

the *B. subtilis* FtsW and RodA proteins could be identified in the genome of *L.*

monocytogenes 10403S (Tab. 1). The protein encoded by *lmo0421* has the weakest homology

to *B. subtilis* FtsW and RodA (Fig. S1, Tab. 1). *lmo0421* is part of the *sigC* operon, which is

comprised of *lmo0422* encoding the PadR-like repressor LstR and *lmo0423* coding for the

ECF-type sigma factor SigC (Fig. 1B). The *sigC* operon acts as a lineage II specific heat

shock system (Zhang et al., 2005) and is therefore not encoded in all *L. monocytogenes*

genomes. Due to the weak homology to FtsW and RodA and its absence in *L. monocytogenes*

strains of lineage I and III, Lmo0421 was excluded from further analysis.

The *L. monocytogenes* protein Lmo1071 is the closest homolog to *B. subtilis* FtsW with a

sequence identity of 48% (Table 1). Furthermore, *L. monocytogenes lmo1071* and *B. subtilis*

ftsW are found in the same chromosomal context. More specifically, *lmo1071* is located

between genes *lmo1070*, which encodes a protein with homology to the *B. subtilis* YlaN

protein, and *pycA* coding for the pyruvate carboxylase. This analysis suggests that gene

lmo1071 encodes the cell division protein FtsW. However, *L. monocytogenes* encodes a

second protein, Lmo2688, that shares a higher degree of homology to the *B. subtilis* FtsW as

compared to the *B. subtilis* RodA protein (Table 1). Due to these similarities and additional data presented in this study, we refer to Lmo1071 and Lmo2688 as FtsW1 and FtsW2, respectively.

The BLAST search with the *B. subtilis* RodA sequence as a query sequence yielded the *L. monocytogenes* protein Lmo2428 as the closest homolog with a sequence identity of 40% (Table 1). In addition to Lmo2428, two additional RodA homologs are present in *L. monocytogenes*, namely Lmo2427 and Lmo2687. As presented below, Lmo2427, Lmo2428 and Lmo2687 are likely bona-fide RodA homologs and were therefore renamed RodA1, RodA2 and RodA3, respectively. *rodA1* is located adjacent to *rodA2*, but despite their proximity, *rodA1* and *rodA2* are likely not transcribed as part of the same operon (Toledo-Arana et al., 2009). In contrast, *rodA3* and *ftsW2* are part of the four-gene operon *lmo2689-lmo2686*. Lmo2689 is similar to a Mg^{2+}-type ATPase, whereas *lmo2686* encodes a protein of unknown function.

lmo1071 encodes FtsW1 and is essential for the survival of *L. monocytogenes*

The cell division protein FtsW is essential for growth in the Gram-negative and Gram-positive model organisms *E. coli* and *B. subtilis* (Boyle et al., 1997, Ikeda et al., 1989, Khattar et al., 1994, Kobayashi et al., 2003). Depletion of FtsW in these organisms leads to a block in cell division and formation of elongated cells (Boyle et al., 1997, Gamba et al., 2016). All our attempts to delete the *ftsW1* gene in *L. monocytogenes* 10403S remained unsuccessful, suggesting that FtsW1 is also essential for growth in *Listeria*. Next, strain 10403SΔ*ftsW1* i*ftsW1* was constructed, in which the expression of *ftsW1* is controlled by an IPTG-inducible promoter. While no difference in the growth was observed between the wildtype and FtsW1-depletion strain (likely due to leakiness of the inducible promoter) (Fig. 2A), cells depleted for FtsW1 were significantly elongated (Fig. 2B-C). Bacteria depleted for

FtsW1 had a median cell length of 3.41±0.16 μm, while wildtype and 10403SΔ*ftsW1* i*ftsW1*

bacteria grown in the presence of 1 mM IPTG had a median cell length of 1.85±0.16 μm and

1.93±0.07 μm, respectively (Fig. 2C). These data indicate that *lmo1071* encodes the cell

division specific SEDS protein FtsW.

L. monocytogenes encodes a second FtsW protein

To our knowledge, all bacteria analyzed to date possess only one FtsW protein that is

essential for cell survival. We identified a second potential FtsW protein, Lmo2688, in *L.*

monocytogenes. In contrast to *ftsW1,* a *L. monocytogenes ftsW2* deletion strain could be

constructed and no significant growth or cell morphology phenotypes could be observed for

the Δ*ftsW2* deletion strain (data not shown). In a previous study, it was reported that the

operon comprised of genes *lmo2689-lmo2686* is only minimally expressed when *L.*

monocytogenes 10403S is grown in BHI broth (Lobel and Herskovits, 2016). We reasoned

that if *ftsW2* does indeed code for a second FtsW protein, it should be possible to delete *ftsW1*

in a strain in which *ftsW2* is artificially expressed from an IPTG-inducible promoter. Indeed,

strain 10403SΔ*ftsW1* i*ftsW2* could be generated in the presence of IPTG. In contrast, we were

unable to generate strain 10403SΔ*ftsW1* when any of the other FtsW/RodA homologs

Lmo2427 (RodA1), Lmo2428 (RodA2) or Lmo2687 (RodA3) were expressed from the same

IPTG-inducible promoter system. While prolonged depletion of FtsW2 in strain

10403SΔ*ftsW1* i*ftsW2* had again no impact on the growth, the cells were significantly

elongated in the absence of the inducer compared to wildtype or bacteria grown in the

presence of inducer (Fig. 3). These data strongly suggest that *ftsW2* encodes a second FtsW

enzyme while the remaining three proteins Lmo2427, Lmo2428 and Lmo2687 likely function

as RodA proteins.

***L. monocytogenes* encodes three RodA homologs**

We were able to assign roles for two of the FtsW/RodA homologs as FtsW-like proteins.

However, *L. monocytogenes* encodes three additional homologs, which show a higher

similarity to the *B. subtilis* RodA protein as compared to the *B. subtilis* FtsW protein (Tab. 1).

As described above, expression of none of the enzymes Lmo2427, Lmo2428 or Lmo2687

was able to rescue the growth of an *ftsW1* deletion strain, indicating that these enzymes likely

function as RodA proteins in *L. monocytogenes* and hence they were renamed RodA1,

RodA2 and RodA3, respectively. All attempts to construct a *rodA1-3* triple mutant failed

further corroborating that these proteins function as RodA proteins and at least one of them

needs to be present for cell viability. To determine whether the different RodA homologs

have distinct functions or whether they are merely duplications, single and double mutant

strains were generated. No significant differences with regards to growth and cell length

could be observed between the wildtype strain 10403S and single *rodA1*, *rodA2* or *rodA3*

deletion strains (Fig. S2). Similar observations were made with the *rodA* double mutant

strains 10403SΔ*rodA1*Δ*rodA2* and 10403SΔ*rodA2*Δ*rodA3* (Fig. 4). However, cells lacking

RodA1 and RodA3 were shorter (1.3±0.03 µm) compared to wildtype cells (1.9±0.06 µm,

Fig. 4B-C), indicating that either RodA1 or RodA3 needs to be present for *L. monocytogenes*

to maintain its rod shape. RodA3 is part of the *lmo2689-lmo2686* operon that is only

minimally expressed when *L. monocytogenes* is grown in BHI broth (Lobel and Herskovits,

2016). The fact that we observe differences in cell morphology between a *rodA1* single and

the *rodA1/rodA3* double mutant, suggests that *rodA3* expression might increase upon deletion

of *rodA1*. To test this hypothesis, we fused the promoter upstream of *lmo2689* and driving

rodA3 expression to *lacZ* and inserted this fusion into the chromosome of wildtype 10403S,

the *rodA1* and the *rodA1/rodA2* deletion strains. The promoter activity was indeed 1.5- to 2-

fold increased in the *rodA1* and *rodA1/rodA2* mutant strains as compared to the wildtype, as

assessed by the increase in the ß-galactosidase activity (Fig. 4G). This result indicates that

expression of the *lmo2689-lmo2686* operon, which encodes FtsW2 and RodA3, is induced in

the absence of RodA1, suggesting a coordination of the expression of the different RodA

homologs.

To confirm that the decrease in cell length of the double mutant strain 10403S*ΔrodA1ΔrodA3*

depends on the absence of RodA1 and RodA3, complementation strains with IPTG-inducible

expression of *rodA1* or *rodA3* were constructed. Expression of RodA1 restored the cell length

to 1.84±0.1 μm, which is comparable to the cell length of wildtype cells (1.76±0.15 μm, Fig.

4E). On the other hand, expression of RodA3 from the ectopic locus in strain

10403S*ΔrodA1ΔrodA3* led to the formation of longer cells with an average cell length of

2.47±0.01 μm (Fig. 4D-E). These results indicate that induction of RodA3 from the IPTG-

inducible promoter likely results in an overproduction of the protein as compared to the

expression from the native promoter, highlighting that in particular fine-tuning of RodA3

production is essential for cell-length determination in *L. monocytogenes*.

The observation that the *rodA1/rodA3* double mutant forms shorter cells suggests that RodA2

is not sufficient to maintain the cell length of *L. monocytogenes*. There are several possible

explanations for this: 1. RodA2 might have a reduced activity as compared to RodA1 or

RodA3. 2. RodA2 might have a function that is different from RodA1 and RodA3. 3. The

protein levels of RodA2 might be sufficient to maintain cell viability but too low to maintain

the rod shape. To determine which of these possibilities might be the case, a strain was

constructed which lacks *rodA1* and *rodA3*, but carries pIMK3-*rodA2* to allow for IPTG-

inducible expression of *rodA2* in addition to the expression of *rodA2* from its native locus

(10403S*ΔrodA1ΔrodA3* i*rodA2*). In the absence of the inducer, the cells had a median cell

length of 1.2±0.03 μm (data not shown). However, the cell length of strain

10403S*ΔrodA1ΔrodA3* i*rodA2* increased to 1.71±0.8 μm when the strain was grown in the

presence of IPTG (Fig. 4F). Therefore, additional expression of RodA2 can partially

complement the cell length phenotype of the *rodA1/rodA3* deletion strain, suggesting that

RodA2 has a similar function as RodA1 and RodA3, but that it has either a lower activity or

is not expressed in sufficient amounts.

As stated above, several attempts to construct a strain inactivated for all three RodA

homologs remained unsuccessful, suggesting that at least one of the proteins RodA1, RodA2

or RodA3 needs to be present for the viability of *L. monocytogenes*. The results presented so

far indicate that RodA1 is the most important RodA homolog considering that RodA2 alone

is not sufficient to maintain the rod shape and that RodA3 is only minimally expressed under

standard laboratory conditions (Lobel and Herskovits, 2016). To understand the impact of

RodA enzymes on cell growth and cell division in *L. monocytogenes*, a strain was

constructed which lacks all three *rodA* genes from its genome, but harbors pIMK3-*rodA1* to

enable IPTG-inducible expression of RodA1. Prolonged depletion of RodA1 in strain

10403SΔ*rodA1-3* i*rodA1* led to a growth defect that was not seen when the strain was grown

in the presence of the inducer (Fig. 5A). However, the depletion was not efficient enough to

see a complete growth inhibition, which would be expected for a strain lacking all three

RodA homologs. Cells of the *L. monocytogenes* strain 10403SΔ*rodA1-3* i*rodA1* that were

grown without IPTG were also significantly shorter with a cell length of 1.18±0.08 μm as

compared to cells of the double mutant 10403SΔ*rodA1*Δ*rodA3* or the wildtype strain 10403S

(Fig. 5B, D). Interestingly, different cell morphologies could be observed for strain

10403SΔ*rodA1-3* i*rodA1* after prolonged RodA1 depletion (Fig. 5C). The placement of the

division septum was affected in some cells and daughter cells of different size or cells with

two septa were observed (Fig. 5C). These morphological defects could be complemented and

the cell length increased to 1.95±0.04 μm upon addition of IPTG and expression of RodA1

(Fig. 5D). These data highlight that RodA1 alone is sufficient to maintain the cell shape of *L.*

monocytogenes.

Decreased moenomycin and lysozyme resistance in the absence of RodA homologs

Next, we wondered whether the absence of FtsW or RodA proteins affects the resistance of *L.*

monocytogenes towards the antibiotics penicillin, bacitracin and moenomycin, which target

different steps in the peptidoglycan biosynthesis process. Penicillin binds to the

transpeptidase domain of PBPs and inhibits their function, leading to a reduced crosslinking

of the peptidoglycan (Nakagawa et al., 1979, Korsak et al., 2010). Bacitracin inhibits the

dephosphorylation of the bactoprenol carrier leading to a block in lipid II synthesis (Stone

and Strominger, 1971). The phosphoglycolipid antibiotic moenomycin inhibits the

glycosyltransferase activity of bifunctional PBPs and thereby prevents the polymerization of

the glycan chain (van Heijenoort et al., 1987).

No significant differences could be observed in terms of resistance against penicillin,

bacitracin or moenomycin for the FtsW1 depletion strain 10403SΔ*ftsW1* i*ftsW1*. This is

presumably due to basal level expression of *ftsW1* even in the absence of the inducer (data

not shown). Simultaneous deletion of *rodA1* and *rodA3* resulted in a slight decrease in the

MIC for penicillin, however, this difference was not significant (Fig. 6A). However, strain

10403SΔ*rodA1*Δ*rodA3* was 2-4-fold more sensitive to the antibiotic bacitracin (Fig. 6B). This

phenotype could be complemented by expressing either RodA1, RodA2 or RodA3 from an

IPTG-inducible promoter (Fig. 6B).

As described above, moenomycin inhibits the transglycosylase activity of PBPs leading to a

decreased activity of these enzymes. In the absence of RodA1 and RodA3, cells are more

susceptible to a reduced activity of PBPs manifesting in a 4-fold reduced resistance to

moenomycin (Fig 6C). Induction of RodA1 expression in the 10403SΔ*rodA1*Δ*rodA3* deletion

strain resulted in a significantly higher resistance to moenomycin as compared to the

wildtype strain 10403S and expression of RodA2 or RodA3, led to partial or complete

complementation of the moenomycin sensitivity (Fig. 6C).

Moreover, resistance to lysozyme, an enzyme that cleaves the linkage between N-acetyl

muramic acid and N-acetylglucosamine residues of the peptidoglycan, was drastically

decreased in strain 10403S∆*rodA1*∆*rodA3* and could be fully restored by expression of

RodA1, RodA2 or RodA3 (Fig. 6D).

Lysozyme resistance in *L. monocytogenes* is mainly accomplished by two modifications of

the peptidoglycan; deacetylation of N-acetylglucosamine residues by PgdA or O-acetylation

of N-acetylmuramic acid residues by OatA (Boneca et al., 2007, Aubry et al., 2011). To

determine whether the activity of PgdA is changed in the absence of RodA1 and RodA3,

peptidoglycan was purified from the 10403S∆*rodA1*∆*rodA3* mutant strain, digested with

mutanolysin and the resulting muropeptides analyzed by HPLC. Peptidoglycan samples

isolated from the wildtype strain 10403S and the complementation strain

10403S∆*rodA1*∆*rodA3* i*rodA1*, that had been grown in the presence of IPTG, were analyzed

as controls (Fig. 6E). The main muropeptide peaks were assigned as described previously

(Rismondo et al., 2015, Burke et al., 2014). Peaks 1 and 2 correspond to the acetylated and

deacetylated monomeric muropeptides, respectively, whereas peak 3 and peaks 4-6 are

acetylated and deacetylated muropeptide dimers, respectively. Deletion of *rodA1* and *rodA3*

led to a reduction of both monomeric muropeptides and therefore to an increase in

crosslinked peptidoglycan fragments by approximately 2% as compared to the wildtype strain

10403S, in which 65% of the peptidoglycan was cross-linked (Fig. 6F). However, no

significant difference with regards to the deacetylated muropeptides could be observed

between the wildtype 10403S, the 10403S∆*rodA1*∆*rodA3* and the 10403S∆*rodA1*∆*rodA3*

i*rodA1* complementation strain (Fig. 6F). These results suggest that the lysozyme sensitivity

phenotype of strain 10403S*ΔrodA1ΔrodA3* is not caused by changes in the peptidoglycan

deacetylation, but rather due to general defects in the peptidoglycan structure.

Cell wall-acting antibiotics induce the promoter of *lmo2689*

The operon *lmo2689-lmo2686*, that contains the genes encoding FtsW2 and RodA3, is only

minimally expressed under standard laboratory conditions (Lobel and Herskovits, 2016).

However, a genome-wide transcriptional analysis performed in *L. monocytogenes* strain

LO28 has shown that *lmo2687, lmo2688* and *lmo2689* are part of the CesR regulon (Nielsen

et al., 2012). The <u>ce</u>phalosporin <u>s</u>ensitivity response regulator CesR is part of the CesRK two-

component system that regulates the transcription of several cell envelope-related genes in

response to changes in cell wall integrity, such as caused by the presence of cell wall-acting

antibiotics or alcohols such as ethanol (Gottschalk et al., 2008, Kallipolitis et al., 2003,

Nielsen et al., 2012). Therefore, we next used the *lmo2689* promoter-*lacZ* fusion described

above, to assess if expression of the *lmo2689-lmo2686* operon is induced in the presence of

antibiotics that target different processes of the PG biosynthesis or ethanol. Indeed, increased

β-galactosidase activity could be measured for cells that had been grown in the presence of

sub-inhibitory concentrations of the β-lactam antibiotics ampicillin, penicillin and cefuroxime

and the phosphoglycolipid moenomycin (Fig. 7A). In contrast, no increase in β-galactosidase

activity could be detected upon addition of vancomycin, lysozyme or ethanol as compared to

untreated control cells (Fig. 7A). We also tested whether the presence of MgSO$_4$ or EDTA

has an impact on the *lmo2689* promoter activity since *lmo2689* encodes a putative Mg^{2+}-type

ATPase. However, the β-galactosidase activity of cells grown in the presence of MgSO$_4$ or

EDTA was comparable to the β-galactosidase activity seen for untreated cells (Fig. 7A).

These results indicate that the expression of *ftsW2* and *rodA3*, that are part of the *lmo2689-*

lmo2686 operon, are induced in the presence of various cell wall-acting antibiotics,

suggesting that FtsW2 and RodA3 might be important for the intrinsic resistance of *L.*

monocytogenes against these antibiotics. However, no significant differences in MICs for

penicillin and moenomycin could be observed between wildtype 10403S, the *ftsW2* or *rodA3*

single mutant strains or the *ftsW2/rodA3* double mutant (Fig. S3). However, there was a slight

reduction in the resistance of the *rodA3* single mutant against cefuroxime as compared to the

wildtype (Fig. S3). To further assess whether there is a difference in the cefuroxime

resistance between the *L. monocytogenes* wildtype strain 10403S and the *rodA1*, *rodA2* and

rodA3 single mutant strains, dilutions of overnight cultures were spotted on BHI agar plates

with or without 1 µg/ml cefuroxime. Deletion of *rodA1* or *rodA2* results in a slightly reduced

ability of these strains to grow on BHI plates supplemented with 1 µg/ml cefuroxime as

compared to the wildtype 10403S strain (Fig. 7B). However, deletion of *rodA3* leads to a

stronger reduction of growth on BHI plates containing 1 µg/ml cefuroxime as compared to

the *rodA1* and *rodA2* single mutants (Fig. 7B). Our results therefore suggest that *L.*

monocytogenes induces the expression of *rodA3* and *ftsW2* in the presence of β-lactam

antibiotics and moenomycin to compensate for the inhibition of the glycosyltransferase and

transpeptidase activity of PBPs and that in particular the RodA homolog RodA3 has an

important function for the intrinsic antibiotic resistance of *L. monocytogenes*.

Discussion

Bacterial cell elongation and cell division need to be tightly regulated to maintain the cell

shape. This is accomplished by two multiprotein complexes, the rod complex and the

divisome, which are coordinated by the actin homolog MreB and the tubulin homolog FtsZ,

respectively (Ricard and Hirota, 1973, Nanninga, 1991, Carballido-Lopez and Formstone,

2007, Typas et al., 2012, Jones et al., 2001, Bi and Lutkenhaus, 1991). The SEDS protein

FtsW is part of the divisome and essential for growth as shown for many bacteria including

E. coli, B. subtilis and *S. aureus* (Boyle et al., 1997, Ikeda et al., 1989, Khattar et al., 1994,

Kobayashi et al., 2003, Reichmann et al., 2019). Our experiments suggested that FtsW1 is

also essential in *L. monocytogenes*, however, a second FtsW protein, FtsW2, can compensate

for the loss of FtsW1 if it is expressed from an inducible promoter. FtsW2 is encoded in the

lmo2689-lmo2686 operon that appears to be only minimally expressed when *L.*

monocytogenes 10403S is grown under standard laboratory conditions (Lobel and Herskovits,

2016). The expression of the *lmo2689-lmo2686* operon is regulated by the two-component

system CesRK that is activated by cell envelope stress (Kallipolitis et al., 2003, Gottschalk et

al., 2008, Nielsen et al., 2012). Using an *L. monocytogenes* strain carrying a P$_{lmo2689}$-*lacZ*

promoter fusion, we could detect increased β-galactosidase activity after incubation with sub-

inhibitory concentrations of different β-lactam antibiotics including penicillin, cefuroxime,

and moenomycin. However, the expression of the *lmo2689-lmo2686* operon was not induced

by other cell wall-targeting antibiotics such as vancomycin or the hydrolase lysozyme. This

suggests that inhibition of the glycosyltransferase or transpeptidase activity of PBPs leads to

activation of the *lmo2689-lmo2686* operon, and hence, to the expression of *ftsW2* as well as

rodA3.

The rod-shape determining protein RodA is part of the elongation machinery. The data

presented in this study suggest that *L. monocytogenes* encodes not one but three RodA

proteins and depletion of the three RodA enzymes leads to a decreased cell length (Fig. 5).

Simultaneous deletion of *rodA1* and *rodA3* already results in the formation of shorter cells,

whereas cells of strains deleted for *rodA1/rodA2* or *rodA2/rodA3* have a cell length that is

comparable to the wildtype strain 10403S. Taking into consideration that *rodA3* is only

minimally expressed under standard laboratory growth conditions in *L. monocytogenes*

10403S (Lobel and Herskovits, 2016), the results presented in this study suggest that *rodA3*

expression gets induced upon inactivation of RodA1, since we observed morphological

differences between the *rodA1* single and the *rodA1/3* double mutant strains. Indeed, β-

galactosidase assays confirmed that deletion of *rodA1* or *rodA1/2* increases the activity of the

promoter from which *rodA3* is expressed. The data presented in this study also indicate that

RodA1 is the "main" RodA enzyme in *L. monocytogenes* as no significant phenotypic

changes with regards to growth and cell division could be observed as long as RodA1 was

present. On the other hand, RodA2 was only able to compensate for the loss of RodA1 and

RodA3, when overproduced from an inducible promoter. This suggests that either RodA2 has

a reduced activity compared to RodA1 or RodA3, which can be overcome by overproducing

the enzyme, or the expression levels of RodA2 are too low to maintain the cell shape.

Interestingly, cells of strain 10403S*ΔrodA1ΔrodA3* in which *rodA3* is overexpressed from an

ectopic locus have an increased cell length as compared to the wildtype. An explanation for

this could be that elevated levels of RodA3 lead to the depletion of proteins needed at the cell

division site, resulting in an extended synthesis of PG on the lateral wall. Another possibility

could be that RodA3 directly inhibits FtsW1 or displaces FtsW1 at the cell division site,

leading to a block in cell division and therefore resulting in the formation of elongated cells.

Further experiments are needed to distinguish between these possibilities.

Recent studies have shown that SEDS proteins acts as glycosyltransferases (Meeske et al.,

2016, Emami et al., 2017). The glycosyltransferase activity of PBPs and MGT can be

inhibited by moenomycin, whereas RodA/FtsW enzymes are not affected by moenomycin

and are therefore important for moenomycin resistance (Tamura et al., 1980, Emami et al.,

2017). In good agreement with this, deletion of the genes encoding two of the three RodA

enzymes, RodA1 and RodA3, resulted in an increased moenomycin sensitivity of *L.*

monocytogenes (Fig. 7B).

In *B. subtilis*, RodA is in a complex with the class B PBP, PBP 2A (also named PBPH), and

these two proteins act together to polymerize and crosslink the glycan strands (Wei et al.,

2003, Henriques et al., 1998). Similarly, FtsW and PBP 2B form a subcomplex as part of the

divisome (Daniel et al., 1996, Gamba et al., 2009). Recently it was shown that RodA-PBP3

and FtsW-PBP1 act as cognate pairs in the coccoid bacterium *S. aureus* (Reichmann et al.,

2019). Depletion of all three RodA enzymes in *L. monocytogenes*, RodA1, RodA2 and

RodA3, leads to a drastic reduction in cell length (Fig. 5). A similar phenotype was observed

for a *L. monocytogenes* strain depleted for the essential class B PBP, PBP B1 (Rismondo et

al., 2015). In contrast, the absence of either FtsW1 (Fig.2) or the class B PBP, PBP B2, in *L.*

monocytogenes results in the formation of elongated cells (Rismondo et al., 2015). These

observations suggest that RodA and FtsW might work in a complex with the cognate PBPs

PBP B1 and PBP B2 during cell elongation and cell division, respectively. Further studies are

necessary to confirm this hypothesis.

It is also interesting to note that *L. monocytogenes* does not only encode multiple FtsW/RodA

enzymes, but also encodes two copies of the lipid II flippase MurJ, MurJ1 and MurJ2.

RNAseq data suggest that *murJ1* and *murJ2* are both expressed in *L. monocytogenes* 10403S

when grown in BHI broth (Lobel and Herskovits, 2016). *murJ1* and *murJ2* might therefore

solely be gene duplications, however, MurJ1 and MurJ2 might have specific roles in different

growth conditions, which needs to be assessed in future studies.

Taken together, *L. monocytogenes* has a repertoire of two lipid II flippases, several PBPs and

multiple members of the SEDS family of proteins to produce its rigid cell wall. The

expression and the activity of these enzymes need to be tightly regulated in *L. monocytogenes*

to maintain its cell shape. Our results suggest that *L. monocytogenes* adapts the expression of

two FtsW/RodA enzymes, FtsW2 and RodA3, to environmental stresses such as the presence

of β-lactam antibiotics, thereby preventing defects in the peptidoglycan synthesis and

subsequent cell lysis.

Acknowledgements

This work was funded by the Wellcome Trust grants 100289/Z/12/Z and 210671/Z/18/Z to

AG and the German research foundation (DFG) grant RI 2920/1-1 to JR.

References
547
AUBRY, C., GOULARD, C., NAHORI, M. A., CAYET, N., DECALF, J., SACHSE, M.,
BONECA, I. G., COSSART, P. & DUSSURGET, O. 2011. OatA, a peptidoglycan O-
acetyltransferase involved in *Listeria monocytogenes* immune escape, is critical for
virulence. *J Infect Dis,* 204**,** 731-40.
BI, E. F. & LUTKENHAUS, J. 1991. FtsZ ring structure associated with division in
Escherichia coli. Nature, 354**,** 161-4.
BLUMBERG, P. M. & STROMINGER, J. L. 1974. Interaction of penicillin with the
bacterial cell: penicillin-binding proteins and penicillin-sensitive enzymes. *Bacteriol*
Rev, 38**,** 291-335.
BONECA, I. G., DUSSURGET, O., CABANES, D., NAHORI, M. A., SOUSA, S., LECUIT,
M., PSYLINAKIS, E., BOURIOTIS, V., HUGOT, J. P., GIOVANNINI, M.,
COYLE, A., BERTIN, J., NAMANE, A., ROUSSELLE, J. C., CAYET, N.,
PREVOST, M. C., BALLOY, V., CHIGNARD, M., PHILPOTT, D. J., COSSART,
P. & GIRARDIN, S. E. 2007. A critical role for peptidoglycan N-deacetylation in
Listeria evasion from the host innate immune system. *Proc Natl Acad Sci U S A,* 104**,**
997-1002.
BOYLE, D. S., KHATTAR, M. M., ADDINALL, S. G., LUTKENHAUS, J. & DONACHIE,
W. D. 1997. *ftsW* is an essential cell-division gene in *Escherichia coli. Mol Microbiol,*
24**,** 1263-73.
BURKE, T. P., LOUKITCHEVA, A., ZEMANSKY, J., WHEELER, R., BONECA, I. G. &
PORTNOY, D. A. 2014. *Listeria monocytogenes* is resistant to lysozyme through the
regulation, not the acquisition, of cell wall-modifying enzymes. *J Bacteriol,* 196**,**
3756-67.
CARBALLIDO-LOPEZ, R. & FORMSTONE, A. 2007. Shape determination in *Bacillus*
subtilis. Curr Opin Microbiol, 10**,** 611-6.
CHO, H., WIVAGG, C. N., KAPOOR, M., BARRY, Z., ROHS, P. D. A., SUH, H.,
MARTO, J. A., GARNER, E. C. & BERNHARDT, T. G. 2016. Bacterial cell wall
biogenesis is mediated by SEDS and PBP polymerase families functioning semi-
autonomously. *Nature Microbiology,* 1**,** 16172.
CORRIGAN, R. M., ABBOTT, J. C., BURHENNE, H., KAEVER, V. & GRÜNDLING, A.
2011. c-di-AMP is a new second messenger in *Staphylococcus aureus* with a role in
controlling cell size and envelope stress. *PLoS Pathog,* 7**,** e1002217.
DANIEL, R. A., WILLIAMS, A. M. & ERRINGTON, J. 1996. A complex four-gene operon
containing essential cell division gene *pbpB* in *Bacillus subtilis. J Bacteriol,* 178**,**
2343-50.
DE JONGE, B. L., CHANG, Y. S., GAGE, D. & TOMASZ, A. 1992. Peptidoglycan
composition of a highly methicillin-resistant *Staphylococcus aureus* strain. The role
of penicillin binding protein 2A. *J Biol Chem,* 267**,** 11248-54.
EMAMI, K., GUYET, A., KAWAI, Y., DEVI, J., WU, L. J., ALLENBY, N., DANIEL, R.
A. & ERRINGTON, J. 2017. RodA as the missing glycosyltransferase in *Bacillus*
subtilis and antibiotic discovery for the peptidoglycan polymerase pathway. *Nat*
Microbiol, 2**,** 16253.
FRAIPONT, C., ALEXEEVA, S., WOLF, B., VAN DER PLOEG, R., SCHLOESSER, M.,
DEN BLAAUWEN, T. & NGUYEN-DISTECHE, M. 2011. The integral membrane
FtsW protein and peptidoglycan synthase PBP3 form a subcomplex in *Escherichia*
coli. Microbiology, 157**,** 251-9.
GAMBA, P., HAMOEN, L. W. & DANIEL, R. A. 2016. Cooperative Recruitment of FtsW
to the Division Site of *Bacillus subtilis. Front Microbiol,* 7**,** 1808.

GAMBA, P., VEENING, J. W., SAUNDERS, N. J., HAMOEN, L. W. & DANIEL, R. A. 2009. Two-step assembly dynamics of the *Bacillus subtilis* divisome. *J Bacteriol,* 191**,** 4186-94.

GHUYSEN, J. M. & STROMINGER, J. L. 1963. Structure of the Cell Wall of *Staphylococcus aureus*, Strain Copenhagen. Ii. Separation and Structure of Disaccharides. *Biochemistry, 2***,** 1119-25.

GOFFIN, C. & GHUYSEN, J. M. 1998. Multimodular penicillin-binding proteins: an enigmatic family of orthologs and paralogs. *Microbiol Mol Biol Rev,* 62**,** 1079-93.

GOTTSCHALK, S., BYGEBJERG-HOVE, I., BONDE, M., NIELSEN, P. K., NGUYEN, T. H., GRAVESEN, A. & KALLIPOLITIS, B. H. 2008. The two-component system CesRK controls the transcriptional induction of cell envelope-related genes in *Listeria monocytogenes* in response to cell wall-acting antibiotics. *J Bacteriol,* 190**,** 4772-6.

GRÜNDLING, A., BURRACK, L. S., BOUWER, H. G. & HIGGINS, D. E. 2004. *Listeria monocytogenes* regulates flagellar motility gene expression through MogR, a transcriptional repressor required for virulence. *Proc Natl Acad Sci U S A,* 101**,** 12318-23.

HARA, H. & SUZUKI, H. 1984. A novel glycan polymerase that synthesizes uncross-linked peptidoglycan in *Escherichia coli*. *FEBS Lett,* 168**,** 155-60.

HENRIQUES, A. O., DE LENCASTRE, H. & PIGGOT, P. J. 1992. A *Bacillus subtilis* morphogene cluster that includes *spoVE* is homologous to the *mra* region of *Escherichia coli*. *Biochimie,* 74**,** 735-48.

HENRIQUES, A. O., GLASER, P., PIGGOT, P. J. & MORAN, C. P., JR. 1998. Control of cell shape and elongation by the *rodA* gene in *Bacillus subtilis*. *Mol Microbiol,* 28**,** 235-47.

HÖLTJE, J. V. 1998. Growth of the stress-bearing and shape-maintaining murein sacculus of *Escherichia coli*. *Microbiol Mol Biol Rev,* 62**,** 181-203.

IKEDA, M., SATO, T., WACHI, M., JUNG, H. K., ISHINO, F., KOBAYASHI, Y. & MATSUHASHI, M. 1989. Structural similarity among *Escherichia coli* FtsW and RodA proteins and *Bacillus subtilis* SpoVE protein, which function in cell division, cell elongation, and spore formation, respectively. *J Bacteriol,* 171**,** 6375-8.

JONES, L. J., CARBALLIDO-LOPEZ, R. & ERRINGTON, J. 2001. Control of cell shape in bacteria: helical, actin-like filaments in *Bacillus subtilis*. *Cell,* 104**,** 913-22.

KALLIPOLITIS, B. H., INGMER, H., GAHAN, C. G., HILL, C. & SOGAARD-ANDERSEN, L. 2003. CesRK, a two-component signal transduction system in *Listeria monocytogenes*, responds to the presence of cell wall-acting antibiotics and affects beta-lactam resistance. *Antimicrob Agents Chemother,* 47**,** 3421-9.

KARINOU, E., SCHUSTER, C. F., PAZOS, M., VOLLMER, W. & GRUNDLING, A. 2018. Inactivation of the monofunctional peptidoglycan glycosyltransferase SgtB allows *Staphylococcus aureus* to survive in the absence of lipoteichoic acid. *J Bacteriol*.

KHATTAR, M. M., BEGG, K. J. & DONACHIE, W. D. 1994. Identification of FtsW and characterization of a new *ftsW* division mutant of *Escherichia coli*. *J Bacteriol,* 176**,** 7140-7.

KOBAYASHI, K., EHRLICH, S. D., ALBERTINI, A., AMATI, G., ANDERSEN, K. K., ARNAUD, M., ASAI, K., ASHIKAGA, S., AYMERICH, S., BESSIERES, P., BOLAND, F., BRIGNELL, S. C., BRON, S., BUNAI, K., CHAPUIS, J., CHRISTIANSEN, L. C., DANCHIN, A., DEBARBOUILLE, M., DERVYN, E., DEUERLING, E., DEVINE, K., DEVINE, S. K., DREESEN, O., ERRINGTON, J., FILLINGER, S., FOSTER, S. J., FUJITA, Y., GALIZZI, A., GARDAN, R., ESCHEVINS, C., FUKUSHIMA, T., HAGA, K., HARWOOD, C. R., HECKER, M.,

HOSOYA, D., HULLO, M. F., KAKESHITA, H., KARAMATA, D., KASAHARA, Y., KAWAMURA, F., KOGA, K., KOSKI, P., KUWANA, R., IMAMURA, D., ISHIMARU, M., ISHIKAWA, S., ISHIO, I., LE COQ, D., MASSON, A., MAUEL, C., MEIMA, R., MELLADO, R. P., MOIR, A., MORIYA, S., NAGAKAWA, E., NANAMIYA, H., NAKAI, S., NYGAARD, P., OGURA, M., OHANAN, T., O'REILLY, M., O'ROURKE, M., PRAGAI, Z., POOLEY, H. M., RAPOPORT, G., RAWLINS, J. P., RIVAS, L. A., RIVOLTA, C., SADAIE, A., SADAIE, Y., SARVAS, M., SATO, T., SAXILD, H. H., SCANLAN, E., SCHUMANN, W., SEEGERS, J. F., SEKIGUCHI, J., SEKOWSKA, A., SEROR, S. J., SIMON, M., STRAGIER, P., STUDER, R., TAKAMATSU, H., TANAKA, T., TAKEUCHI, M., THOMAIDES, H. B., VAGNER, V., VAN DIJL, J. M., WATABE, K., WIPAT, A., YAMAMOTO, H., YAMAMOTO, M., YAMAMOTO, Y., YAMANE, K., YATA, K., YOSHIDA, K., YOSHIKAWA, H., ZUBER, U. & OGASAWARA, N. 2003. Essential *Bacillus subtilis* genes. *Proc Natl Acad Sci U S A,* 100**,** 4678-83.

KORSAK, D., MARKIEWICZ, Z., GUTKIND, G. O. & AYALA, J. A. 2010. Identification of the full set of *Listeria monocytogenes* penicillin-binding proteins and characterization of PBPD2 (Lmo2812). *BMC Microbiol,* 10**,** 239.

LECLERCQ, S., DEROUAUX, A., OLATUNJI, S., FRAIPONT, C., EGAN, A. J., VOLLMER, W., BREUKINK, E. & TERRAK, M. 2017. Interplay between Penicillin-binding proteins and SEDS proteins promotes bacterial cell wall synthesis. *Sci Rep,* 7**,** 43306.

LIU, X., GALLAY, C., KJOS, M., DOMENECH, A., SLAGER, J., VAN KESSEL, S. P., KNOOPS, K., SORG, R. A., ZHANG, J. R. & VEENING, J. W. 2017. High-throughput CRISPRi phenotyping identifies new essential genes in *Streptococcus pneumoniae. Mol Syst Biol,* 13**,** 931.

LOBEL, L. & HERSKOVITS, A. A. 2016. Systems Level Analyses Reveal Multiple Regulatory Activities of CodY Controlling Metabolism, Motility and Virulence in *Listeria monocytogenes. PLoS Genet,* 12**,** e1005870.

MCPHERSON, D. C. & POPHAM, D. L. 2003. Peptidoglycan synthesis in the absence of class A penicillin-binding proteins in *Bacillus subtilis. J Bacteriol,* 185**,** 1423-31.

MEESKE, A. J., RILEY, E. P., ROBINS, W. P., UEHARA, T., MEKALANOS, J. J., KAHNE, D., WALKER, S., KRUSE, A. C., BERNHARDT, T. G. & RUDNER, D. Z. 2016. SEDS proteins are a widespread family of bacterial cell wall polymerases. *Nature,* 537**,** 634-638.

MEESKE, A. J., SHAM, L. T., KIMSEY, H., KOO, B. M., GROSS, C. A., BERNHARDT, T. G. & RUDNER, D. Z. 2015. MurJ and a novel lipid II flippase are required for cell wall biogenesis in *Bacillus subtilis. Proc Natl Acad Sci U S A,* 112**,** 6437-42.

NAKAGAWA, J., TAMAKI, S. & MATSUHASHI, M. 1979. Purified penicillin binding proteins 1Bs from *Escherichia coli* membrane showing activities of both peptidoglycan polymerase and peptidoglycan crosslinking enzyme. *Agric. Biol. Chem.,* 43**,** 1379-1380.

NANNINGA, N. 1991. Cell division and peptidoglycan assembly in *Escherichia coli. Mol Microbiol,* 5**,** 791-5.

NIELSEN, P. K., ANDERSEN, A. Z., MOLS, M., VAN DER VEEN, S., ABEE, T. & KALLIPOLITIS, B. H. 2012. Genome-wide transcriptional profiling of the cell envelope stress response and the role of LisRK and CesRK in *Listeria monocytogenes. Microbiology,* 158**,** 963-74.

PARK, W. & MATSUHASHI, M. 1984. *Staphylococcus aureus* and *Micrococcus luteus* peptidoglycan transglycosylases that are not penicillin-binding proteins. *J Bacteriol,* 157**,** 538-44.

696 PARK, W., SETO, H., HAKENBECK, R. & MATSUHASHI, M. 1985. Major peptidoglycan
697   transglycosylase activity in *Streptococcus pneumoniae* that is not a penicillin-binding
698   protein. *FEMS Microbiology Letters,* 27**,** 45-48.
699 PINHO, M. G., KJOS, M. & VEENING, J. W. 2013. How to get (a)round: mechanisms
700   controlling growth and division of coccoid bacteria. *Nat Rev Microbiol,* 11**,** 601-14.
701 REICHMANN, N. T., TAVARES, A. C., SARAIVA, B. M., JOUSSELIN, A., REED, P.,
702   PEREIRA, A. R., MONTEIRO, J. M., VANNIEUWENHZE, M. S., FERNANDES,
703   F. & PINHO, M. G. 2019. SEDS-bPBP pairs direct lateral and septal peptidoglycan
704   synthesis in *Staphylococcus aureus. bioRxiv,* 552034.
705 RICARD, M. & HIROTA, Y. 1973. Process of cellular division in *Escherichia coli*:
706   physiological study on thermosensitive mutants defective in cell division. *J Bacteriol,*
707   116**,** 314-22.
708 RISMONDO, J., MOLLER, L., ALDRIDGE, C., GRAY, J., VOLLMER, W. &
709   HALBEDEL, S. 2015. Discrete and overlapping functions of peptidoglycan synthases
710   in growth, cell division and virulence of *Listeria monocytogenes. Mol Microbiol,* 95**,**
711   332-51.
712 ROGERS, H. J., PERKINS, H. R. & WARD, J. B. 1980. Microbial Cell Walls and
713   Membranes. *Chapman and Hall New York, N.Y.,* 10**,** 33-33.
714 RUIZ, N. 2008. Bioinformatics identification of MurJ (MviN) as the peptidoglycan lipid II
715   flippase in *Escherichia coli. Proc Natl Acad Sci U S A,* 105**,** 15553-7.
716 SAUVAGE, E., KERFF, F., TERRAK, M., AYALA, J. A. & CHARLIER, P. 2008. The
717   penicillin-binding proteins: structure and role in peptidoglycan biosynthesis. *FEMS*
718   *Microbiol Rev,* 32**,** 234-58.
719 SCHEFFERS, D. J. & PINHO, M. G. 2005. Bacterial cell wall synthesis: new insights from
720   localization studies. *Microbiol Mol Biol Rev,* 69**,** 585-607.
721 SHAM, L. T., BUTLER, E. K., LEBAR, M. D., KAHNE, D., BERNHARDT, T. G. & RUIZ,
722   N. 2014. Bacterial cell wall. MurJ is the flippase of lipid-linked precursors for
723   peptidoglycan biogenesis. *Science,* 345**,** 220-2.
724 SIEVERS, F., WILM, A., DINEEN, D., GIBSON, T. J., KARPLUS, K., LI, W., LOPEZ, R.,
725   MCWILLIAM, H., REMMERT, M., SODING, J., THOMPSON, J. D. & HIGGINS,
726   D. G. 2011. Fast, scalable generation of high-quality protein multiple sequence
727   alignments using Clustal Omega. *Mol Syst Biol,* 7**,** 539.
728 SONNHAMMER, E. L., VON HEIJNE, G. & KROGH, A. 1998. A hidden Markov model
729   for predicting transmembrane helices in protein sequences. *Proc Int Conf Intell Syst*
730   *Mol Biol,* 6**,** 175-82.
731 STONE, K. J. & STROMINGER, J. L. 1971. Mechanism of action of bacitracin:
732   complexation with metal ion and C 55 -isoprenyl pyrophosphate. *Proc Natl Acad Sci*
733   *U S A,* 68**,** 3223-7.
734 SWAMINATHAN, B. & GERNER-SMIDT, P. 2007. The epidemiology of human listeriosis.
735   *Microbes Infect,* 9**,** 1236-43.
736 TAGUCHI, A., WELSH, M. A., MARMONT, L. S., LEE, W., SJODT, M., KRUSE, A. C.,
737   KAHNE, D., BERNHARDT, T. G. & WALKER, S. 2019. FtsW is a peptidoglycan
738   polymerase that is functional only in complex with its cognate penicillin-binding
739   protein. *Nat Microbiol.*
740 TAMURA, T., SUZUKI, H., NISHIMURA, Y., MIZOGUCHI, J. & HIROTA, Y. 1980. On
741   the process of cellular division in *Escherichia coli*: isolation and characterization of
742   penicillin-binding proteins 1a, 1b, and 3. *Proc Natl Acad Sci U S A,* 77**,** 4499-503.
743 TOLEDO-ARANA, A., DUSSURGET, O., NIKITAS, G., SESTO, N., GUET-REVILLET,
744   H., BALESTRINO, D., LOH, E., GRIPENLAND, J., TIENSUU, T.,
745   VAITKEVICIUS, K., BARTHELEMY, M., VERGASSOLA, M., NAHORI, M. A.,

SOUBIGOU, G., REGNAULT, B., COPPEE, J. Y., LECUIT, M., JOHANSSON, J.
& COSSART, P. 2009. The *Listeria* transcriptional landscape from saprophytism to
virulence. *Nature,* 459**,** 950-6.

TYPAS, A., BANZHAF, M., GROSS, C. A. & VOLLMER, W. 2012. From the regulation of
peptidoglycan synthesis to bacterial growth and morphology. *Nat Rev Microbiol,* 10**,**
123-36.

VAN HEIJENOORT, J. 2001. Formation of the glycan chains in the synthesis of bacterial
peptidoglycan. *Glycobiology,* 11**,** 25R-36R.

VAN HEIJENOORT, Y., LEDUC, M., SINGER, H. & VAN HEIJENOORT, J. 1987. Effects
of moenomycin on *Escherichia coli*. *J Gen Microbiol,* 133**,** 667-74.

VOLLMER, W., BLANOT, D. & DE PEDRO, M. A. 2008. Peptidoglycan structure and
architecture. *FEMS Microbiol Rev,* 32**,** 149-67.

WANG, Q. M., PEERY, R. B., JOHNSON, R. B., ALBORN, W. E., YEH, W. K. &
SKATRUD, P. L. 2001. Identification and characterization of a monofunctional
glycosyltransferase from *Staphylococcus aureus*. *J Bacteriol,* 183**,** 4779-85.

WEI, Y., HAVASY, T., MCPHERSON, D. C. & POPHAM, D. L. 2003. Rod shape
determination by the *Bacillus subtilis* class B penicillin-binding proteins encoded by
pbpA and *pbpH*. *J Bacteriol,* 185**,** 4717-26.

WEIDEL, W. & PELZER, H. 1964. Bagshaped Macromolecules--a New Outlook on
Bacterial Cell Walls. *Adv Enzymol Relat Areas Mol Biol,* 26**,** 193-232.

ZHANG, C., NIETFELDT, J., ZHANG, M. & BENSON, A. K. 2005. Functional
consequences of genome evolution in *Listeria monocytogenes*: the *lmo0423* and
lmo0422 genes encode sigmaC and LstR, a lineage II-specific heat shock system. *J*
Bacteriol, 187**,** 7243-53.

Table 1: Sequence homology between *B. subtilis* and *L. monocytogenes* FtsW/RodA proteins as determined by BLAST.

L. monocytogenes proteins	*B. subtilis* FtsW (403 aa)[*]	*B. subtilis* RodA (393 aa)[*]
Lmo0421	46/128 aa (36%)	77/350 aa (22%)
(416 aa)	7e-14	4e-09
FtsW1 (Lmo1071)	195/403 aa (48%)	100/333 aa (30%)
(402 aa)	1e-122	7e-35
RodA1 (Lmo2427)	124/403 aa (31%)	132/360 aa (37%)
(391 aa)	9e-29	1e-57
RodA2 (Lmo2428)	112/374 aa (30%)	159/402 aa (40%)
(389 aa)	1e-33	1e-66
RodA3 (Lmo2687)	95/324 aa (29%)	110/356 aa (31%)
(369 aa)	3e-25	3e-44
FtsW2 (Lmo2688)	147/342 aa (43%)	95/300 aa (32%)
(376 aa)	5e-67	6e-22

[*] *L. monocytogenes* proteins and their amino acid sizes (aa) are shown in bold in the left column and *B. subtilis* FtsW and RodA proteins used for the homology search are denoted in the top row. The number of amino acids, which were found to be identical between the respective *L. monocytogenes* and *B. subtilis* FtsW/RodA protein are denoted (left number) along with the amino acids that was utilized by the BLAST algorithm for this comparison (right number). The % identity is given in brackets and the corresponding e-values are indicated below.

Figure 1: Topology of FtsW and RodA enzymes and their genomic arrangement in *L. monocytogenes*. (A) Topology models for FtsW and RodA enzymes of *L. monocytogenes* as predicted using the TMHMM v. 2.0 server (Sonnhammer et al., 1998). (B) Genomic organization of *ftsW/rodA* genes in *L. monocytogenes*. Arrowheads indicate the gene orientation. Small black arrows indicate promoters based on the operon structures described in Toledo-Arana *et al*. (Toledo-Arana et al., 2009). Three different promoters have been identified for the *sigC* operon (Zhang et al., 2005).

Figure 2: Depletion of FtsW1 results in cell elongation. (A) Growth of the *L. monocytogenes* FtsW1 depletion strain. Strain 10403SΔ*ftsW1* i*ftsW1* was grown overnight in BHI medium in the presence of 1 mM IPTG. The following day, the cells were washed and used to inoculate fresh BHI medium without IPTG and incubated at 37°C for 8-10h, diluted again and grown until the next morning. The depletion culture was used to inoculate BHI medium without IPTG or with 1 mM IPTG and the growth was monitored by determining OD_{600} readings at hourly intervals. Wild type *L. monocytogenes* strain 10403S (wt) was used as control. Averages and standard deviations from three independent experiments were plotted. (B) Microscopy analysis of 10403S (wt) and the *ftsW1* depletion strain grown in the presence of absence of IPTG. For depletion of FtsW1, strain 10403SΔ*ftsW1* i*ftsW1* was grown as described in panel A. Cultures of 10403S and 10403SΔ*ftsW1* i*ftsW1* were diluted 1:100 in fresh BHI medium (with 1 mM IPTG where indicated), grown for 3 h at 37°C and the cells subsequently stained with the membrane dye nile red and analyzed by fluorescence and phase contrast microscopy. The scale bar is 2 μm. (C) Cell lengths measurement of 10403S (wt) and the *ftsW1* depletion strain. The cell length of 300 cells per strain was measured and the median cell length calculated and plotted. Three independent experiments were performed, and the average and standard deviation of the median cell length plotted. For statistical analysis, a one-way ANOVA coupled with a Dunnett's multiple comparisons test was performed (**** p≤0.0001).

Fig. 3: Overexpression of FtsW2 can compensate for the loss of FtsW1. (A) Growth comparison of strains 10403S (wt) and 10403SΔ*ftsW1* i*ftsW2*. Growth curves were performed as described in Figure 2. (B) Microscopy images of strains 10403S (wt) and 10403SΔ*ftsW1* i*ftsW2*. For depletion of FtsW2, strain 10403SΔ*ftsW1* i*ftsW2* was grown as described in Figure 2. The next morning, cells were diluted 1:100 and grown in BHI broth (with or without IPTG as indicated) until mid-logarithmic growth phase, stained with the membrane dye nile red and subjected to fluorescence and phase contrast microscopy analysis. Scale bar is 2 μm. (C) Cell lengths measurements of 10403S (wt) and 10403SΔ*ftsW1* i*ftsW2*. The cell length of 300 cells per strain was measured and the median cell length calculated and plotted. Three independent experiments were performed, and the average and standard deviation of the median cell length plotted. For statistical analysis, a one-way ANOVA coupled with a Dunnett's multiple comparisons test was performed (**** p≤0.0001).

Fig. 4: RodA1 and RodA3 are essential to maintain the rod shape of *L. monocytogenes*.

(A) Growth of *L. monocytogenes* 10403S (wt), 10403S*ΔrodA1ΔrodA2*, 10403S*ΔrodA1ΔrodA3* and 10403S*ΔrodA2ΔrodA3* strains in BHI broth at 37°C. (B) Microscopy images of wt and mutant *L. monocytogenes* strains. Cells from the strains described in panel A were stained with the membrane dye nile red and analyzed by fluorescence and phase contrast microscopy. Scale bar is 2 μm. (C) Cell lengths measurements of wt and mutant *L. monocytogenes* strains. The cell length of 300 cells per

strain was measured and the median cell length calculated and plotted. Three independent experiments were performed, and the average and standard deviation of the median cell length plotted. For statistical analysis, one-way ANOVA t-tests for multiple comparisons were used (** p≤0.01). (D) Microscopy images of wildtype and mutant *L. monocytogenes* strains. Cells from 10403S (wt), 10403S∆*rodA1*∆*rodA3* and 10403S∆*rodA1*∆*rodA3* expressing *rodA1*, *rodA2* or *rodA3* from an IPTG-inducible promoter were stained with the membrane dye nile red and analyzed by fluorescence and phase contrast microscopy. Scale bar is 2 μm. (E-F) Cell length analysis of 10403S∆*rodA1*∆*rodA3* complementation strains. Analysis was performed as described in panel C. (G) β-galactosidase assay using *L. monocytogenes* strains 10403S (wt), 10403S∆*rodA1* and 10403S∆*rodA1-2* carrying a P$_{lmo2689}$-*lacZ* fusion construct. Averages and standard deviations of three independent experiments were plotted. For statistical analysis, a one-way ANOVA coupled with a Dunnett's multiple comparisons test was used (*p≤0.1).

Figure 5: Absence of all three RodA proteins leads to a drastic cell shortening. (A) Growth of *L. monocytogenes* in the absence of all three RodA proteins. Growth curves were performed as described in Figure 2. (B and C) Microscopy analysis of strain 10403S*ΔrodA1-3* i*rodA1*. For depletion of RodA1, Strain 10403S*ΔrodA1-3* i*rodA1* was grown as described in Figure 2. (B) Wildtype 10403S and depleted 10403S*ΔrodA1-3* i*rodA1* cultures were diluted 1:100, grown for 3 h at 37°C (where indicated in the presence of 1 mM IPTG), stained with nile red and phase contrast and fluorescence microscopy images were taken. Scale bar is 2 μm. (C) Microscopy images showing examples of different cell morphologies observed for strain 10403S*ΔrodA1-3* i*rodA1* depleted for RodA1. Scale bar is 2 μm. (D) Cell lengths measurements of 10403S (wt) and strain 10403S*ΔrodA1-3* i*rodA1*. The cell length of 300 cells per strain was measured. Three independent experiments were performed, and the average and standard deviation of the median cell length plotted. For statistical analysis, a one-way ANOVA coupled with a Dunnett's multiple comparisons test was performed (* p≤0.1, **** p≤0.0001).

Figure 6: Impact of *rodA1* and *rodA3* deletion on the antibiotic and lysozyme resistance.

The minimal inhibitory concentration for the antibiotics penicillin G (A), bacitracin A (B), moenomycin (C) and lysozyme (D) was determined for the wildtype *L. monocytogenes* strain 10403S, the Δ*rodA1*Δ*rodA3* deletion strain and the complementation strains 10403SΔ*rodA1*Δ*rodA3* i*rodA1,* 10403SΔ*rodA1*Δ*rodA3* i*rodA2* and 10403SΔ*rodA1*Δ*rodA3* i*rodA3* using a microbroth dilution assay. The complementation strains were grown in the

presence of 1 mM IPTG. The result of three biological replicates are shown in A-D. For statistical analysis, a one-way ANOVA coupled with a Dunnett's multiple comparisons test was used (** p≤0.01, *** p≤0.001, **** p≤0.0001). (E) HPLC analysis of the muropeptide composition of 10403S (wt), 10403SΔ*rodA1*Δ*rodA3* and 10403SΔ*rodA1*Δ*rodA3* i*rodA1* (grown in the presence of IPTG to induce expression of RodA1). The major muropeptide peaks are numbered 1-6 as previously described (Rismondo et al., 2015, Burke et al., 2014). (F) Relative abundance of muropeptide peaks 1-6 in peptidoglycan isolated from strains 10403S (wt), 10403SΔ*rodA1*Δ*rodA3* and 10403SΔ*rodA1*Δ*rodA3* i*rodA1*. Average values and standard deviations were calculated from three independent peptidoglycan extractions.

Figure 7: Cell wall-acting antibiotics increase the *lmo2689* promoter activity. (A) Bacteria from mid-logarithmic cultures of strain 10403S pPL3e-P*lmo2689*-*lacZ* were exposed for 2 h at 37°C to different stressors. The activity of the *lmo2689* promoter was subsequently determined by performing β-galactosidase activity assays as described in the method section. Bacteria that had been grown in the absence of a stressor were included as negative (-) control. The averages of the β-galactosidase activity units and standard deviations from three independent experiments were plotted. For statistical analysis, a one-way ANOVA coupled with a Dunnett's multiple comparisons test was used (*** p≤0.001, **** p≤0.0001). (B) Absence of RodA3 results in decreased cefuroxime resistance. Dilutions of overnight cultures

of strains 10403S, 10403SΔ*rodA1*, 10403SΔ*rodA2* and 10403SΔ*rodA3* were spotted on BHI agar and BHI agar containing 1 µg/ml cefuroxime and incubated for 24 h at 37°C. A representative result from three independent experiments is shown.

Figure S1: Sequence alignment of FtsW and RodA proteins from *B. subtilis* and *L. monocytogenes.* The alignment was created using Clustal Omega (Sievers et al., 2011). Based on a study by Meeske *et al.,* amino acid residues of *B. subtilis* RodA that are essential for RodA function are colored in blue, where conservative replacements are possible in yellow and amino acids that can be changed to certain other amino acids are colored in orange (Meeske et al., 2016).

Figure S2: Single *rodA1, rodA2* or *rodA3 L. monocytogenes* strains do not show a growth or cell length defect. (A) Growth of *L. monocytogenes* 10403S (wt), 10403SΔ*rodA1*, 10403SΔ*rodA2* and 10403SΔ*rodA3* in BHI broth at 37°C. (B) Microscopy images of wildtype and mutant *L. monocytogenes* strains. Bacteria from cultures of the strains described in panel A were stained with the membrane dye nile red and analyzed by fluorescence and phase contrast microscopy. Scale bar is 2 μm. (C) Cell length of *L. monocytogenes* wt and *rodA* single mutants. The cell length of 300 cells per strain was measured and the median cell length calculated and plotted. Three independent experiments were performed, and the average and standard deviation of the median cell length plotted.

Figure S3: Deletion of *rodA3* but not *ftsW2* has a minor impact on the resistance towards a selected number of cell wall active antibiotics. The minimal inhibitory concentration for the antibiotics penicillin G (A), moenomycin (B) and cefuroxime (C) was determined for the wildtype *L. monocytogenes* strain 10403S, the 10403SΔ*rodA3* and 10403SΔ*ftsW2* single deletion strains and the 10403SΔ*ftsW2*Δ*rodA3* double mutant strain using a microbroth dilution assay. The result of three biological replicates are shown in A-C.