Transgenerational plasticity of inducible defenses: combined effects of grand-parental, parental and current environments

Juliette Tariel[1,*], Sandrine Plenet[1], Émilien Luquet[1]

Abstract

While an increasing number of studies highlights that parental environment shapes offspring phenotype (transgenerational plasticity TGP), TGP beyond the parental generation has received less attention. Studies suggest that TGP impacts population dynamics and evolution of phenotype, but these impacts will depend on how long an environmental effect can persist across generations and whether multigenerational effects are cumulative. Here we tested the impact of both grand-parental and parental environments on offspring reaction norm in a prey-predator system. We exposed three generations of *Physa acuta* snails without and with predator-cues according to a full factorial design and measured offspring inducible defenses. We found that both grand-parental and parental exposure to predator cues impacted offspring anti-predator defenses, but their effects were not cumulative and depended on the defenses considered. We also highlighted that grand-parental environment could alter reaction norm of offspring shell thickness, demonstrating an interaction between the grand-parental TGP and the within-generational WGP plasticity. We called for more studies covering the combine effects of multigenerational environments.

Keywords

transgenerational plasticity — grand-parental effect — inducible defenses — predator-prey interactions — *Physa acuta*

*Corresponding author: juliettetariel@gmail.com

Introduction

Organisms may respond to fluctuating environments by adapting through genetic

evolution over generations or through phenotypic plasticity. This last is traditionally defined as

the capacity of a given genotype to render alternative phenotypes under different

environmental conditions (within-generational plasticity, WGP) (Pigliucci, 2005; West-

Eberhard, 2003). Plasticity may also occur across generations (transgenerational plasticity,

TGP), when the phenotype of offspring is influenced by carry-over effects of past environments

experienced by the previous generation(s) (Agrawal et al., 1999; Galloway and Etterson, 2007;

Salinas et al., 2013). Ancestors can alter the phenotype of their offspring without involving

genetic changes through a range of non-genetic processes as parental effects, e.g. transmission

of nutrients, hormones, proteins (Crean and Bonduriansky, 2014; Mousseau, 1998), or by any

form of epigenetic inheritance, e.g. DNA methylation marks, histone protein modifications,

non-coding small RNAs (Holeski et al., 2012; Schlichting and Wund, 2014). TGP has been

shown for diverse organisms and traits in response to various environments (Bonduriansky and

Day, 2009; Donelson et al., 2018; Salinas et al., 2013). When past environmental cues provide

reliable proxies about future offspring conditions, TGP may enable organisms to cope with fast-

changing environments because it refines offspring phenotypes in anticipation of the

environmental conditions they are likely to experience (Bonduriansky and Day, 2009; Donelson

et al., 2018; Herman and Sultan, 2011).

Most studies on TGP focused on the effect of parental environment on offspring phenotype

(e.g. Donelson et al., 2018; Herman and Sultan, 2011; Mousseau, 1998; Wolf and Wade, 2009)

and on offspring reaction norm (interaction TGP x WGP; Salinas et al., 2013). But

transgenerational effects are not confined to the parental generation. A few examples have

shown that TGP can persist for multiple successive generations (Plaistow et al., 2006; Remy,

2010; Sarker and Peleg-Raibstein, 2019), even though the majority is restrained to three

generations (*i.e.* grand-parental effect) (e.g. Hafer et al., 2011; Herman and Sultan, 2011; Kou

et al., 2011; Lock, 2012; Shama et al., 2014; Walsh et al., 2014). However, it is unclear how

multigenerational effects can interact (Prizak et al., 2014) and experimental works controlling

for combination of multigenerational effects always showed complex patterns of phenotypic

offspring responses. For example, Hafer et al. (2011) and Walsh et al. (2014) in collembolan

(*Folsomia candida*) and cladoceran (*Daphnia ambigua*) respectively, demonstrated that some

life-history traits were affected by an interactive effect between grandparental and parental

environment, leading to phenotypic landscapes that do not fit simple adaptive scenarios.

Moreover, such complex patterns of phenotypic responses can depend on the offspring context.

Plaistow et al. (2006), for instance, showed in the soil mite *Sancassania berlesei* that the

persistence of past environments (across four generations) differed between high- and low-food

offspring contexts. Consequently, understanding how multigenerational combined environments

interact to shape the offspring phenotype is still limited and challenging.

Predator-induced plasticity is a well-known model in WGP study (e.g. Harvell, 1990;

Hoverman et al., 2005; Relyea, 2001) and allows an individual to fine-tune its phenotype facing

predation risk (Benard, 2004; Lima, 1998; Tollrian and Harvell, 1999). Predator-induced

defenses are also widely used to investigate TGP over two generations (parental and offspring

generations) (e.g. Agrawal et al., 1999; Bell and Stein, 2017; Colicchio, 2017). However, to our

knowledge, only two studies have been interested in TGP of predator-induced traits beyond

two consecutives generations (Sentis et al., 2018; Walsh et al., 2015).

We focused on a hermaphroditic gastropod *Physa acuta*. Physidae are well-known to develop

adaptive phenotypes in response to predation risk (Auld and Houser, 2015; Auld and Relyea,

2008, 2011; Beaty et al., 2016; DeWitt et al., 1999; Gustafson et al., 2014). Predator (crayfish)

cues induce WGP of *Physa sp.* life-history traits (larger age and size at first reproduction; Auld

and Relyea, 2008), shell thickness (thicker shell; Auld and Relyea, 2011), shell size (narrower

shape; DeWitt, 1998), and behavior (crawling-out the water; Alexander and Covich, 1991;

DeWitt et al., 1999). We exposed three successive generations of snails from hatching to sexual

maturity according to a full factorial design to environments without and with predator-cues.

The results concerning the first two generations have demonstrated a predator-induced TGP in

P. acuta (Luquet and Tariel, 2016) that has been confirmed in a concomitant study (Beaty et

al., 2016). Here, we focused on the F3 generation to investigate how the effects of grand-

parental, parental and offspring environments combine to influence behavior, shell thickness

and shell morphology.

Materials and Methods

Animal collection and experimental design

Adult *P. acuta* snails were collected (number of individuals are reported on Figure 1 and experimental design represented on Supplementary Figure 1) on March 2015 in a lentic backwater of the Rhône river (45° 48'6"N, 4° 55'33"E) in Lyon, France. *Physa acuta* is a freshwater and simultaneous hermaphroditic snail, invasive from North America (Lydeard et al., 2016). The wild-caught adult snails constituted our F_0 generation. We pooled them overnight in a 10L-aquarium to ensure that offspring result from outcrossing (*P. acuta* is a preferential outcrosser; Jarne et al., 2010). Then, we individually isolated all F_0 snails in 70 mL plastic vials filled with reconstituted water (2.4 g $NaHCO_3$, 3 g $CaSO_4$, 1.5 g $MgSO_4$, 0.1 g KCl to 25 L deionized water) in a 25°C experimental room with 12h light-dark photoperiod. After 24 hours, we removed the F_0 adults from the vials and we randomly choose 15 vials with one egg capsule each. These 15 egg capsules constituted our 15 maternal families (hereafter called only "families") of the F_1 generation and developed until hatching (~ 7 days). Two days after hatching, we randomly sampled 12 siblings per family and split them into two environments: 6 snails remained in a no-predator environment (control environment) while 6 others were moved in a predator-cue environment. These F_1 snails were reared in 70 mL plastic vials with their siblings until 28 days-old where they were isolated in the same type of plastic vials until the end of the experiment (35 days after hatching). We then generated the second F_2 generation in merging F_1 snails into copulation groups of 15 individuals (one snail from each of the 15 families). At this step, the snail identity was not recorded and thus the pedigree could not be further assessed. We made 6 copulation groups per treatment and each reproduced in a 5L-aquarium for 24h in no-predator water to ensure embryos are not exposed to predator environment. Then, individuals were isolated, and we randomly selected 18 F_1 snails that had laid eggs from each treatment. We then followed the same protocol as previously to rear F_2 snails in control and predator-cue environments according to a full factorial design until 49 days after hatching. The F_3 generation was then generated and reared using the procedure described above. As growth rate was slowing down every generation under our laboratory conditions and we wanted snails in a reasonably sufficient size, we reared the F_3 generation at a

113 later age (74 days after hatching). This F_3 generation was represented by eight combinations of

114 grand-parental (E_1), parental (E_2) and offspring (E_3) environments: CCC, CPC, PCC, PPC,

115 CCP, CPP, PCP and PPP with each time "C" for control environment and "P" for predator-

116 cue environment (Figure 1). The number of individuals and families per combination of

117 environments are reported on Figure 1.

Figure 1. Number of individuals (N) and families (Family) at each generation and in the two

119
treatments with "C" for control environment and "P" for predator-cue environment. For the 3rd

generation, two number of individuals are reported, one for behavioral measurements (first position) and

one for other measurements (second position).

Water and food (*ad libitum*, chopped and boiled lettuce) were renewed for all experimental

snails twice a week. Predator-conditioned water with predator cues was obtained by

individually rearing crayfishes (*Procambarus clarkii*) in 4L reconstituted water and feeding

with one *P. acuta* adult one day before water change (Auld and Relyea, 2011). This crayfish-

conditioned water was used for the predator-cue treatment while only reconstituted water was

used for the control treatment. This crayfish species coexists with *P. acuta* in its native

location in North America.

Measuring phenotypes

We assessed anti-predator behavior three times in 70-day-old F_3 snails through three

consecutive days starting one day after the water change. We measured anti-predator behavior

by recording the position above/on or below the water surface in the rearing boxes with

predator cues present or absent according to the treatment. Crawling-out of the water (position

above water surface) is considered as allowing to escape from benthic predators like crayfishes

(DeWitt et al., 1999).

At 74 days old, we gently dried snails with paper towel and measured the snail total wet

weight (body and shell) with an electronic scale at the nearest 0.001mg. A photograph of each

snail aperture upwards was taken with an Olympus SC50 camera installed on an Olympus

SZX9 binocular and its Olympus DF PLAPO 1X-2 objective at a x8 magnification. Shell and

aperture length and width (shell morphology) were measured on these photographs with the

software ImageJ (Schneider et al., 2012). Shell thickness was measured with an electronic

caliper at the nearest 0.01mm at the edge of the aperture. Shorter and narrower shell and

aperture dimensions (after adjusting for mass) and thicker shell are adaptive anti-predator

responses (Auld and Relyea, 2011).

Statistical analysis

The multigenerational effect of predator cues on anti-predator behavior (*i.e.* snail

position above/on or below the water surface) was analyzed using generalized linear mixed

models (GLMM) assuming a binomial distribution (logit link function). Shell and aperture

length and width were analyzed with a principal component analysis (PCA) with the package

FactoMineR (Lê et al., 2008) to assess shell morphology. They were highly correlated and the

first component (PC1) explained most of variance in shell morphology (95.8%; Supplementary

figure 2): a high value of PC1 was associated with a longer and wider shell and aperture,

whereas a low value was associated with a smaller and narrower shell and aperture. PC1 was

then used as a proxy for shell morphology and response variable in linear mixed model. The

multigenerational effect of predator cues on weight, shell thickness and shell morphology was

analyzed using linear mixed models (LMM) with restricted maximum likelihood estimation and

Kenward and Roger's approximation for degrees of freedom. These response variables were

scaled prior to statistical analysis. In all linear mixed models, grand-parental (E_1), parental

(E_2), offspring (E_3) environments, and all interactions were considered as fixed effects. Because

shell thickness and shell morphology were significantly correlated with weight, weight (cube

root transformed and then scaled) was added as a continuous fixed effect (analyses of

covariance) in these models. Family was considered as a random intercept effect in all models.

Individual identity was added as a random intercept model in the GLMM model (anti-predator

behavior analysis) to account for repeated measures on the same individual. We used type 2 F-

tests for significance of fixed effects in the LMM and likelihood ratio tests in the GLMM

(lmerTest package; Kuznetsova et al., 2017). We tested significance of random effects with

likelihood ratio tests comparing models with or without the tested random effect in the full

fixed effect structure. All statistical analyses were performed with R 3.4.1 (R Core Team, 2017)

and the glmer() and lmer() functions from the lme4 package (Bates et al., 2015, 4).

Results

Anti-predator behavior

The offspring exposure to predator cues (E3) significantly increased by 105% the

proportion of snails crawling-out the water (Table 1a; Fig. 2a). The parental environment (E2)

did not affect the proportion of crawling-out behavior (Table 1a; Fig. 2a). However, grand-

parental exposure to predator cues (E1) significantly increased by 28 % the proportion of

crawling-out behavior (Table 1; Fig. 2a). Family and individual random effects were significant

(Table 1a).

Snail weight

180   The offspring exposure to predator cues significantly decreased by 18% the snail weight

181 (Table 1b; Fig. 2b). Neither the parental environment nor the grand-parental environment

182 influenced snail weight (Table 1b; Fig. 2b). Family random effect was significant (Table 1b).

Shell thickness

After accounting for snail weight, grand-parental and parental environments interacted

both with the offspring environment (Table 1c; Fig. 2c), demonstrating that both could affect

the offspring response to predator cues. In the offspring control environment, the grand-

parental exposure to predator cues increased by 4% the offspring shell thickness whereas the

parental exposure to predator cues decreased by 9% the offspring shell thickness. And in the

offspring predator-cue environment, the grand-parental exposure to predator cues decreased by

11% the offspring shell thickness whereas the parental exposure to predator cues increased by

14% the offspring shell thickness. Regarding the direct effect of offspring environment, offspring

from predator-cue environment had a thicker shell than those from current control environment

(Table 1c; Fig. 2c). Family random effect was not significant (Table 1c).

Shell morphology

195   After accounting for snail weight, the offspring exposure to predator cues impacted the

196   shell morphology and induced a longer and narrower shell and aperture (Table 1d; Fig. 2d).

197   Neither the parental environment (interaction parental and offspring environments almost

198   significant $P < 0.10$) nor the grand-parental environment influenced the shell morphology

199   (Table 1d; Fig. 2d). Family random effect was not significant (Table 1d).

Figure 3. The effect of multi-generational exposure to predator-cues on offspring **a** crawling-out behavior (proportion of snails out the water in %), **b** total weight, **c** and **d** least-mean squares of shell thickness and shell morphology in the model accounting for weight described in Table 1. CCC, PCC, CPC, PCC, CCP, PCP, CPP and PPP represent the eight combinations of grand-parental (E1), parental (E2) and offspring (E3) environments with "C" for control environment and "P" for predator-cue environment for every generation. The vertical dashed line separates the two offspring treatment groups. White shapes are for grand-parental control environment and black shapes for grand-parental predator-cue environment. Circles are for parental control environment and squares for parental predator-cue environment. Data are means ± SE.

a. Crawling-out	*Fixed effects*	*Estimate (SE)*	*Df*	*Chisq*	*P*
	Grand-parental env. (E_1)	0.848 (0.6812)	1	8.29	**0.0040**
	Parental env. (E_2)	-0.090 (0.3864)	1	0.4	0.5274
	Offspring env. (E_3)	1.738 (0.3544)	1	114.46	**<0.0001**
	E1 x E2	-0.389 (0.7696)	1	0.01	0.9315
	E1 x E3	-0.168 (0.7882)	1	1.89	0.1687
	E2 x E3	-0.146 (0.4242)	1	0.01	0.9072
	E1 x E2 x E3	0.855 (0.8915)	1	0.91	0.3395
	Random effects	*Variance*	*Df*	*Chisq*	*P*
	Family	0.489	1	14.98	**0.0001**
	Individual	-0.047	1	8.26	**0.0040**
b. Weight	*Fixed effects*	*Estimate (SE)*	*Numdf, Dendf*	*F*	*P*
	Grand-parental env. (E_1)	-0.047 (0.1656)	1, 42.44	0.18	0.6766
	Parental env. (E_2)	-0.116 (0.1656)	1, 48.70	0.57	0.4545
	Offspring env. (E_3)	-0.366 (0.1329)	1, 354.26	14.76	**0.0001**
	E1 x E2	-0.041 (0.3312)	1, 55.77	0.01	0.9105
	E1 x E3	-0.027 (0.2659)	1, 348.77	0.001	0.9737
	E2 x E3	-0.104 (0.2659)	1, 362.32	0.30	0.5836
	E1 x E2 x E3	0.069 (0.5318)	1, 358.29	0.02	0.8969
	Random effects	*Variance*	*Df*	*Chisq*	*P*
	Family	0.085	1	6.6	**0.0102**
c. Shell thickness	*Fixed effects*	*Estimate (SE)*	*Numdf, Dendf*	*F*	*P*
	cuberoot(Weight)	0.656 (0.0404)	1, 364.6	257.77	**<0.0001**
	Grand-parental env. (E_1)	-0.092 (0.1152)	1, 40.41	0.19	0.6650
	Parental env. (E_2)	0.057 (0.1153)	1, 49.59	0.04	0.8446
	Offspring env. (E_3)	0.258 (0.1054)	1, 356.90	28.15	**<0.0001**
	E1 x E2	0.180 (0.2305)	1, 59.10	0.62	0.4329
	E1 x E3	-0.350 (0.2088)	1, 350.26	4.01	**0.0461**
	E2 x E3	0.497 (0.2088)	1, 363.53	7.68	**0.0059**
	E1 x E2 x E3	0.033 (0.4176)	1, 359.10	0.01	0.9366
	Random effects	*Variance*	*Df*	*Chisq*	*P*
	Family	0.020	1	0.78	0.3780
d. Shell morphology	*Fixed effects*	*Estimate (SE)*	*Numdf, Dendf*	*F*	*P*
	cuberoot(Weight)	1.901 (0.0167)	1, 366.61	12842.53	**<0.0001**
	Grand-parental env. (E_1)	-0.034 (0.0496)	1, 41.10	0.19	0.6613
	Parental env. (E_2)	-0.014 (0.0497)	1, 49.12	0.33	0.5677
	Offspring env. (E_3)	-0.287 (0.0433)	1, 355.56	73.25	**<0.0001**
	E1 x E2	-0.044 (0.0992)	1, 57.48	0.12	0.7288
	E1 x E3	0.053 (0.0858)	1, 349.39	0.02	0.8943
	E2 x E3	0.080 (0.0859)	1, 362.82	3.68	0.0558
	E1 x E2 x E3	-0.215 (0.1717)	1, 358.51	1.57	0.2114
	Random effects	*Variance*	*Df*	*Chisq*	*P*
	Family	0.005	1	1.88	0.1704

210 **Table 1.** Results of generalized mixed model (**a**) and linear mixed model (**b**, **c** and **d**) analysis on
211 offspring predator-induced traits. Bold values indicate significant P-values (P < 0.05).

Discussion

We first confirm that the exposure to predator cues induces well-known defenses

against crayfish predation in *P. acuta*. Our key finding is that predator cues alter also offspring

defenses two generations later but depending on the offspring environment (WGP x TGP) and

the defensive traits considered. Our experimental work highlights that TGP effects can be

complex beyond the parental generation and that the offspring phenotype results from a

combination of multigenerational effects.

Corroborating several studies on anti-predator defenses of *Physa sp.* (Auld and Relyea, 2011;

Dalesman et al., 2009; DeWitt et al., 1999, 2000; Turner et al., 1999), the offspring exposure to

predator cues induced higher crawling-out behavior, shell-crushing resistance (thicker shell) and

entry-resistant shell (narrower shell and aperture). Moreover, offspring exposed to predator

cues were lighter, suggesting a trade-off, *i.e.* a lower energetic investment in growth due to a

potential cost to produce these defenses (as showed in other gastropod species: (Brönmark et

al., 2012). This result stresses the fitness advantage of WGP which allow the production of

costly defenses only in case of predation (Harvell, 1990).

Grand-parental and parental effects on anti-predator defenses

TGP is expected to evolve when the ancestral environment is a good proxy of offspring

environment (Bonduriansky and Day, 2009; Dey et al., 2016; English et al., 2015; Harvell,

1990; Leimar and McNamara, 2015; Uller, 2008), allowing a pre-adaptation of offspring to

predation risk (Agrawal et al., 1999). In our predator-prey system, crayfish has a long lifespan

(ca. 3 years) compared to the generation time of *P. acuta* (ca. 50 days) and a relatively

sedentary lifestyle (Vioque-Fernández et al., 2009). This suggests that generational cues of

crayfish presence can be a good proxy of predation risk across several snail generations and

thus that TGP could have long-lasting effects on the anti-predator defenses. Consistently, in *P.*

acuta, parental exposure to predator cues induces a more crush-resistant shell and an higher

escape behavior in offspring (Beaty et al., 2016; Luquet and Tariel, 2016). In this study, as

expected, TGP went further than the parental generation: the grand-parental environment also

influenced the anti-predator behavior and the shell thickness of offspring.

How long can persist transgenerational effects on anti-predator responses is a remaining open

question. To our knowledge, the study of Sentis et al. (2018) on the pea aphid (*Acyrthosiphon*

pisum) is the only one to investigate predator-induced TGP over a large number of generations

(> 25). They found that the defensive phenotype – a high frequency of winged aphids in the

population – persists for one generation after removing predators whatever the induction time

is, *i.e.* the previous number of successive generations experiencing the novel environment

(predator presence). However, three generations are needed after removing predators for the

frequency of winged phenotypes to come back to the control level, and this number of

generations increases with the induction time. Together, these results suggest that

multigenerational environmental effects on inducible defenses are broader than just a parental

effect and could persist for many generations.

Combination of multigenerational effects on anti-predator defenses

We showed that the offspring phenotype results from a combination of

multigenerational effects (grand-parents, parents and offspring), similar to theoretical and other

experimental studies (Hafer et al., 2011; Kou et al., 2011; Lock, 2012; Prizak et al., 2014;

Shama and Wegner, 2014; Walsh et al., 2014). However, in our study, grand-parental and

parental effects acted independently (no significant interaction between grand-parental and

parental environmental effects): either only one affected the offspring environment (behavior),

or in interaction with the offspring environment (WGP x TGP) and in opposite directions

(shell thickness). This results in complex offspring phenotypic patterns that do not fit with

simple anti-predator scenarios. The adaptive relevance of such combinations of

multigenerational effects is far to be evident even studying, as in our study, fitness-related

traits. It would be thus interesting to compare the survival of snails from different past

environmental histories exposed to lethal predation challenges. The offspring crawling-out

behavior increased with offspring and grand-parental exposure to predator cues while the

parental environment did not alter this behavior. Shell thickness was influenced by both grand-

parental and parental environments, but in opposing directions and depending on the offspring

environment (grand-parental and parental WGP x TGP interactions). In offspring control

environment, grand-parental exposure to predator cues increased the offspring shell thickness

whereas parental exposure reduced the shell thickness. The effects were opposite in the

offspring predator-cue environment.

Firstly, these results confirm that offspring reaction norms can be altered by parental

environment (shell thickness; (Donelson et al., 2018; Luquet and Tariel, 2016; Salinas et al.,

2013)) but expand for the first time the WGP x TGP interaction to grand-parental

environmental cues (shell thickness). Secondly, opposing directions of grand-parental and

parental effects found on shell thickness are not rare in empirical studies (Magiafoglou and

Hoffmann, 2003; Shama and Wegner, 2014) and illustrate the complexity to determine the

adaptive significance of multigenerational effects. Such opposing effects may reflect different

mechanisms underlying the transfer of environmental information (Shea et al., 2011). This

complex opposing relationship between grand-parental and parental environmental effects could

be also theoretically beneficial by reducing the phenotypic variance which allow the population

to stay closer to the target phenotype (Prizak et al., 2014). Moreover, focusing on few

generations in short-term experiments artificially focuses the interpretations of such effects

while they could only be transient over larger timescales in a population dynamic framework.

For example, Sentis et al. (2018), after removing predators, observed that the frequency of

winged aphids remained high for one generation before dropping abruptly below the control

levels (grand-parental effect), and then converging with the winged aphid frequencies of the

control lines (great-grand-parental effect). Consequently, in focusing on only three consecutive

generations as in our study, these results could be interpreted as a negative grand-parental

effect (decrease of winged aphid frequency) opposing to a positive parental effect (increase of

winged individual frequency) on the offspring phenotype. These findings highlight the need to

develop empirical studies on larger timescales and controlling for the combination of

multigenerational effects.

Trait-dependence of transgenerational plasticity

Our results show that the pattern of TGP depends on the traits (anti-predator

behavior, shell thickness and shell morphology). Behavioral traits, which are often labile and

exhibiting reversible WGP within developmental or adult stages, are predicted to be influenced

by current environment rather than by past environmental experience (Dingemanse and Wolf,

2013; Piersma and Drent, 2003). Behavioral WGP in response to current environmental cues

should rapidly by-pass the behavioral TGP (Beaman et al., 2016). By contrast, the traits that

are more constrained during the development and exhibiting irreversible variations, as

morphological traits, are predicted to be relatively more influenced by past environments

(Kuijper and Hoyle, 2015). TGP on morphological traits could irreversibly engage the offspring

on developmental trajectories and could not be compensated by WGP. In *P. acuta*, crawling-

out behavior is indeed very flexible and reversible at a time scale of hours while a thicker shell

and a rounder or narrower shell shape are irreversible changes in the developmental trajectory

(DeWitt et al., 1999; Relyea, 2003). Surprisingly in our study, the escape behavior of offspring

is influenced by the grand-parental environment while shell morphology was not influenced by

parental or grand-parental environments. This highlights that transgenerational effects on

morphological traits may have a short persistence over generations while behavioral TGP may

be much more prevalent than currently realized. Parental TGP on behavioral traits has been

sometimes observed (Bestion et al., 2014; Donelan and Trussell, 2015; Giesing et al., 2011; e.g.

Storm and Lima, 2010) and few times with long-lasting effects over generations (Dias and

Ressler, 2014; Remy, 2010).

Conclusion

In our study, we demonstrated that the effects of multigenerational (grand-parental,

parental and offspring) exposure to predator cues on a variety of offspring defensive traits

(escape behavior, shell thickness and shape) do not fit simple anti-predator scenarios. The

multigenerational effects combined, sometimes in opposing directions and depending on the

traits, to shape the offspring anti-predator defenses. We also call for more theoretical and

empirical studies integrating the combined effects of multigenerational environments on larger

number of generations, investigating the underlying mechanisms (epigenetic inheritance) and

evaluating their evolutionary importance.

Author contributions

324   S.P. and E.L. conceived of and designed the study. All authors contributed to the data

325   collection. They all carried out the data analysis and interpretation. And they all drafted,

326   revised and approved the article for publication.

Acknowledgements

328   Thanks to P. Gibert for advice on the manuscript. We thank L. Bensousoussan for help
329   on data analysis.

Conflict of interest disclosure

331   The authors of this preprint declare that they have no financial conflict of interest with
332   the content of this article.

Data accessibility

334   The dataset and the R script for the linear mixed model analysis will be available soon
335   on Zenodo upon recommendation.

References

Agrawal, A. A., Laforsch, C., and Tollrian, R. (1999). Transgenerational induction of defences
in animals and plants. *Nature* 401, 60–63. doi:10.1038/43425.

Alexander, J. E., and Covich, A. P. (1991). Predator avoidance by the freshwater snail *Physella*
virgata in response to the crayfish *Procambarus simulans*. *Oecologia* 87, 435–442.

Auld, J. R., and Houser, R. (2015). Age-dependent effects of predation risk on reproductive
success in a freshwater snail. *Evolution* 69, 2793–2798. doi:10.1111/evo.12769.

Auld, J. R., and Relyea, R. A. (2008). Are there interactive effects of mate availability and
predation risk on life history and defence in a simultaneous hermaphrodite? *J.*
Evolution. Biol. 21, 1371–1378. doi:10.1111/j.1420-9101.2008.01562.x.

Auld, J. R., and Relyea, R. A. (2011). Adaptive plasticity in predator-induced defenses in a
common freshwater snail: altered selection and mode of predation due to prey
phenotype. *Evol. Ecol.* 25, 189–202. doi:10.1007/s10682-010-9394-1.

Bates, D., Mächler, M., Bolker, B., and Walker, S. (2015). Fitting linear mixed-effects models
using lme4. *J. Stat. Softw.* 67, 1–48. doi:10.18637/jss.v067.i01.

Beaman, J. E., White, C. R., and Seebacher, F. (2016). Evolution of plasticity: mechanistic
link between development and reversible acclimation. *Trends Ecol. Evol.* 31, 237–249.
doi:10.1016/j.tree.2016.01.004.

Beaty, L. E., Wormington, J. D., Kensinger, B. J., Bayley, K. N., Goeppner, S. R., Gustafson,
355   K. D., et al. (2016). Shaped by the past, acting in the present: transgenerational
356   plasticity of anti-predatory traits. *Oikos* 125, 1570–1576. doi:10.1111/oik.03114.

357   Bell, A. M., and Stein, L. R. (2017). Transgenerational and developmental plasticity at the
358   molecular level: lessons from *Daphnia*. *Mol. Ecol.* 26, 4859–4861.
359   doi:10.1111/mec.14327.

360   Benard, M. F. (2004). Predator-induced phenotypic plasticity in organisms with complex life
361   histories. *Annu. Rev. Ecol. Evol. S.* 35, 651–673.
362   doi:10.1146/annurev.ecolsys.35.021004.112426.

363   Bestion, E., Teyssier, A., Aubret, F., Clobert, J., and Cote, J. (2014). Maternal exposure to
364   predator scents: offspring phenotypic adjustment and dispersal. *Proc. R. Soc. B-Biol.*
365   *Sci.* 281, 20140701. doi:10.1098/rspb.2014.0701.

366   Bonduriansky, R., and Day, T. (2009). Nongenetic inheritance and its evolutionary
367   implications. *Annu. Rev. Ecol. Evol. S.* 40, 103–125.
368   doi:10.1146/annurev.ecolsys.39.110707.173441.

369   Brönmark, C., Lakowitz, T., Nilsson, P. A., Ahlgren, J., Lennartsdotter, C., and Hollander, J.
370   (2012). Costs of inducible defence along a resource gradient. *PLoS One* 7, e30467.
371   doi:10.1371/journal.pone.0030467.

372   Colicchio, J. (2017). Transgenerational effects alter plant defence and resistance in nature. *J.*
373   *Evolution. Biol.* 30, 664–680. doi:10.1111/jeb.13042.

374   Crean, A. J., and Bonduriansky, R. (2014). What is a paternal effect? *Trends Ecol. Evol.* 29,
375   554–559. doi:10.1016/j.tree.2014.07.009.

376   Dalesman, S., Rundle, S. D., and Cotton, P. A. (2009). Crawl-out behaviour in response to
377   predation cues in an aquatic gastropod: insights from artificial selection. *Evol. Ecol.* 23,
378   907–918. doi:10.1007/s10682-008-9280-2.

379   DeWitt, T. J. (1998). Costs and limits of phenotypic plasticity: tests with predator-induced
380   morphology and life history in a freshwater snail. *J. Evolution. Biol.* 11, 465–480.
381   doi:10.1046/j.1420-9101.1998.11040465.x.

382   DeWitt, T. J., Robinson, B. W., and Wilson, D. S. (2000). Functional diversity among
383   predators of a freshwater snail imposes an adaptive trade-off for shell morphology. *Evol.*
384   *Ecol. Res.* 2, 129–148.

385   DeWitt, T. J., Sih, A., and Hucko, J. A. (1999). Trait compensation and cospecialization in a
386   freshwater snail: size, shape and antipredator behaviour. *Anim. Behav.* 58, 397–407.
387   doi:10.1006/anbe.1999.1158.

Dey, S., Proulx, S. R., and Teotónio, H. (2016). Adaptation to temporally fluctuating
environments by the evolution of maternal effects. *PLOS Biol.* 14, e1002388.
doi:10.1371/journal.pbio.1002388.

Dingemanse, N. J., and Wolf, M. (2013). Between-individual differences in behavioural
plasticity within populations: causes and consequences. *Anim. Behav.* 85, 1031–1039.
doi:10.1016/j.anbehav.2012.12.032.

Donelan, S. C., and Trussell, G. C. (2015). Parental effects enhance risk tolerance and
performance in offspring. *Ecology* 96, 2049–2055. doi:10.1890/14-1773.1.

Donelson, J. M., Salinas, S., Munday, P. L., and Shama, L. N. S. (2018). Transgenerational
plasticity and climate change experiments: where do we go from here? *Glob. Change*
Biol. 24, 13–34. doi:10.1111/gcb.13903.

English, S., Pen, I., Shea, N., and Uller, T. (2015). The information value of non-genetic
inheritance in plants and animals. *Plos One* 10, e0116996.
doi:10.1371/journal.pone.0116996.

Galloway, L. F., and Etterson, J. R. (2007). Transgenerational plasticity is adaptive in the
wild. *Science* 318, 1134–1136. doi:10.1126/science.1148766.

Giesing, E. R., Suski, C. D., Warner, R. E., and Bell, A. M. (2011). Female sticklebacks
transfer information via eggs: effects of maternal experience with predators on offspring.
P. Roy. Soc. B-Biol. Sci. 278, 1753–1759. doi:10.1098/rspb.2010.1819.

Gustafson, K., Kensinger, B., Bolek, M., and Luttbeg, B. (2014). Distinct snail (*Physa*)
morphotypes from different habitats converge in shell shape and size under common
garden conditions. *Evol. Ecol. Res.* 16, 77–89.

Hafer, N., Ebil, S., Uller, T., and Pike, N. (2011). Transgenerational effects of food availability
on age at maturity and reproductive output in an asexual collembolan species. *Biol.*
Lett. 7, 755–758. doi:10.1098/rsbl.2011.0139.

Harvell, C. D. (1990). The ecology and evolution of inducible defenses. *Q. Rev. Biol.* 65, 323–
340.

Herman, J. J., and Sultan, S. E. (2011). Adaptive transgenerational plasticity in plants: case
studies, mechanisms, and implications for natural populations. *Front. Plant Sci.* 2, 102.
doi:10.3389/fpls.2011.00102.

Holeski, L. M., Jander, G., and Agrawal, A. A. (2012). Transgenerational defense induction
and epigenetic inheritance in plants. *Trends Ecol. Evol.* 27, 618–626.
doi:10.1016/j.tree.2012.07.011.

Hoverman, J. T., Auld, J. R., and Relyea, R. A. (2005). Putting prey back together again:
integrating predator-induced behavior, morphology, and life history. *Oecologia* 144,
481–491. doi:10.1007/s00442-005-0082-8.

Jarne, P., Pointier, J. P., David, P., and Koene, J. M. (2010). Basommatophoran gastropods.
The evolution of primary sexual characters in animals, 173–196.

Kou, H. P., Li, Y., Song, X. X., Ou, X. F., Xing, S. C., Ma, J., et al. (2011). Heritable
alteration in DNA methylation induced by nitrogen-deficiency stress accompanies
enhanced tolerance by progenies to the stress in rice (*Oryza sativa L.*). *J. Plant Physiol.*
168, 1685–1693. doi:10.1016/j.jplph.2011.03.017.

Kuijper, B., and Hoyle, R. B. (2015). When to rely on maternal effects and when on
phenotypic plasticity? *Evolution* 69, 950–968. doi:10.1111/evo.12635.

Kuznetsova, A., Brockhoff, P. B., and Christensen, R. H. B. (2017). lmerTest Package: Tests in
Linear Mixed Effects Models. *J. Stat. Softw.* 82, 1–26. doi:10.18637/jss.v082.i13.

Lê, S., Josse, J., and Husson, F. (2008). FactoMineR: A Package for Multivariate Analysis.
Journal of Statistical Software 25, 1–18. doi:10.18637/jss.v025.i01.

Leimar, O., and McNamara, J. M. (2015). The evolution of transgenerational integration of
information in heterogeneous environments. *Am. Nat.* 185, E55–E69.
doi:10.1086/679575.

Lima, S. L. (1998). Nonlethal effects in the ecology of predator-prey interactions. what are the
ecological effects of anti-predator decision-making? *BioScience* 48, 25–34.
doi:10.2307/1313225.

Lock, J. E. (2012). Transgenerational effects of parent and grandparent gender on offspring
development in a biparental beetle species. *Biol. Lett.* 8, 408–411.
doi:10.1098/rsbl.2011.0920.

Luquet, E., and Tariel, J. (2016). Offspring reaction norms shaped by parental environment:
interaction between within- and trans-generational plasticity of inducible defenses.
BMC Evol. Biol. 16, 209. doi:10.1186/s12862-016-0795-9.

Lydeard, C., Campbell, D., and Golz, M. (2016). Physa acuta Draparnaud, 1805 Should be
Treated as a Native of North America, Not Europe. *Malacologia* 59, 347–350.
doi:10.4002/040.059.0213.

Mousseau, T. (1998). The adaptive significance of maternal effects. *Trends Ecol. Evol.* 13, 403–
407. doi:10.1016/S0169-5347(98)01472-4.

Piersma, T., and Drent, J. (2003). Phenotypic flexibility and the evolution of organismal
design. *Trends Ecol. Evol.* 18, 228–233. doi:10.1016/S0169-5347(03)00036-3.

Pigliucci, M. (2005). Evolution of phenotypic plasticity: where are we going now? *Trends Ecol.*
Evol. 20, 481–486. doi:10.1016/j.tree.2005.06.001.

Plaistow, S. J., Lapsley, C. T., and Benton, T. G. (2006). Context-dependent intergenerational
effects: the interaction between past and present environments and its effect on
population dynamics. *Am. Nat.* 167, 206–215. doi:10.1086/499380.

Prizak, R., Ezard, T. H. G., and Hoyle, R. B. (2014). Fitness consequences of maternal and
grandmaternal effects. *Ecol. Evol.* 4, 3139–3145. doi:10.1002/ece3.1150.

R Core Team (2017). *R: A Language and Environment for Statistical Computing.* Vienna,
Austria: R Foundation for Statistical Computing Available at: https://www.R-
project.org/.

Relyea, R. A. (2001). Morphological and behavioral plasticity of larval anurans in response to
different predators. *Ecology* 82, 523–540. doi:10.1890/0012-
9658(2001)082[0523:MABPOL]2.0.CO;2.

Relyea, R. A. (2003). Predators come and predators go: the reversibility of predator-induced
traits. *Ecology* 84, 1840–1848. doi:10.1890/0012-9658(2003)084[1840:PCAPGT]2.0.CO;2.

Remy, J.-J. (2010). Stable inheritance of an acquired behavior in *Caenorhabditis elegans. Curr.*
Biol. 20, R877–R878. doi:10.1016/j.cub.2010.08.013.

Salinas, S., Brown, S. C., Mangel, M., and Munch, S. B. (2013). Non-genetic inheritance and
changing environments. *Non-Genetic Inheritance* 1, 38–50. doi:10.2478/ngi-2013-0005.

Sarker, G., and Peleg-Raibstein, D. (2019). Maternal overnutrition induces long-term cognitive
deficits across several generations. *Nutrients* 11, 7. doi:10.3390/nu11010007.

Schlichting, C. D., and Wund, M. A. (2014). Phenotypic plasticity and epigenetic marking: an
assessment of evidence for genetic accommodation. *Evolution* 68, 656–672.
doi:10.1111/evo.12348.

Schneider, C. A., Rasband, W. S., and Eliceiri, K. W. (2012). NIH Image to ImageJ: 25 years
of Image Analysis. *Nat Methods* 9, 671–675.

Sentis, A., Bertram, R., Dardenne, N., Ramon-Portugal, F., Espinasse, G., Louit, I., et al.
(2018). Evolution without standing genetic variation: change in transgenerational
plastic response under persistent predation pressure. *Heredity* 121, 266.
doi:10.1038/s41437-018-0108-8.

Shama, L. N. S., Strobel, A., Mark, F. C., and Wegner, K. M. (2014). Transgenerational
plasticity in marine sticklebacks: maternal effects mediate impacts of a warming ocean.
Funct. Ecol. 28, 1482–1493. doi:10.1111/1365-2435.12280.

Shama, L. N. S., and Wegner, K. M. (2014). Grandparental effects in marine sticklebacks:
transgenerational plasticity across multiple generations. *J. Evolution. Biol.* 27, 2297–
2307. doi:10.1111/jeb.12490.

Shea, N., Pen, I., and Uller, T. (2011). Three epigenetic information channels and their
different roles in evolution: epigenetic mechanisms and evolution. *J. Evolution. Biol.* 24,
1178–1187. doi:10.1111/j.1420-9101.2011.02235.x.

Storm, J. J., and Lima, S. L. (2010). Mothers forewarn offspring about predators: a
transgenerational maternal effect on behavior. *Am. Nat.* 175, 382–390.
doi:10.1086/650443.

Tollrian, R., and Harvell, C. D. (1999). *The Ecology and Evolution of Inducible Defenses.*
Princeton University Press.

Turner, A. M., Fetterolf, S. A., and Bernot, R. J. (1999). Predator identity and consumer
behavior: differential effects of fish and crayfish on the habitat use of a freshwater snail.
Oecologia 118, 242–247. doi:10.1007/s004420050724.

Uller, T. (2008). Developmental plasticity and the evolution of parental effects. *Trends Ecol.*
Evol. 23, 432–438. doi:10.1016/j.tree.2008.04.005.

Vioque-Fernández, A., Alves de Almeida, E., and López-Barea, J. (2009). Assessment of
Doñana National Park contamination in *Procambarus clarkii*: integration of
conventional biomarkers and proteomic approaches. *Sci. Total Env.* 407, 1784–1797.
doi:10.1016/j.scitotenv.2008.11.051.

Walsh, M. R., Cooley, F., Biles, K., and Munch, S. B. (2015). Predator-induced phenotypic
plasticity within- and across-generations: a challenge for theory? *P. Roy. Soc. B-Biol.*
Sci. 282, 20142205. doi:10.1098/rspb.2014.2205.

Walsh, M. R., Whittington, D., and Funkhouser, C. (2014). Thermal transgenerational
plasticity in natural populations of *Daphnia*. *Integr. Comp. Biol.* 54, 822–829.
doi:10.1093/icb/icu078.

West-Eberhard, M. J. (2003). *Developmental plasticity and evolution.* Oxford University Press.

Wolf, J. B., and Wade, M. J. (2009). What are maternal effects (and what are they not)? *Phil.*
Trans. R. Soc. Lond. B 364, 1107–1115. doi:10.1098/rstb.2008.0238.

517