Heterogeneous profiles of coupled sleep oscillations in human hippocampus

Roy Cox[1*], Theodor Rüber[1,2,3], Bernhard P Staresina[4], Juergen Fell[1]

1: Department of Epileptology, University of Bonn, Bonn, Germany
2: Epilepsy Center Frankfurt Rhine-Main, Department of Neurology, Goethe University Frankfurt, Frankfurt am Main, Germany
3: Center for Personalized Translational Epilepsy Research (CePTER), Goethe University Frankfurt, Frankfurt am Main, Germany
4: School of Psychology, University of Birmingham, Birmingham, United Kingdom

* corresponding author: roycox.roycox@gmail.com

Abstract

Cross-frequency coupling of sleep oscillations is thought to mediate memory consolidation. While the hippocampus is deemed central to this process, detailed knowledge of which oscillatory rhythms interact in the human hippocampus is lacking. Combining intracranial hippocampal and non-invasive electroencephalography from twelve neurosurgical patients, we characterized spectral power and coupling during non-rapid eye movement (NREM) and rapid eye movement (REM) sleep. Hippocampal coupling was extensive, with the majority of channels expressing spectral interactions. NREM consistently showed delta–ripple coupling, but ripples were also modulated by slow oscillations (SOs) and sleep spindles. SO–delta and SO–theta coupling, as well as interactions between delta/theta and spindle/beta frequencies also occurred. During REM, limited interactions between delta/theta and beta frequencies emerged. Moreover, oscillatory organization differed substantially between i) hippocampus and scalp, ii) sites along the anterior-posterior hippocampal axis, and iii) individuals. Overall, these results extend and refine our understanding of hippocampal sleep oscillations.

Keywords: sleep, hippocampus, electrophysiology, cross-frequency coupling, individual differences

Introduction

The hippocampus and wider medial temporal lobe system play a crucial role in episodic memory, the ability to recall past events and their individual components (1, 2). Besides its role in memory formation and retrieval, the hippocampus (HPC) is also a key player in systems consolidation, whereby initially labile memories are gradually transferred from HPC to neocortical sites for permanent storage (3–6). Intriguingly, consolidation depends heavily on sleep (7–9), with greatest importance usually ascribed to non-rapid eye movement (NREM) sleep, and a relatively poorly understood role for rapid eye movement (REM) sleep (10, 11).

NREM-dependent consolidation is thought to rely on the unique electrophysiological signatures of this brain state. These include 0.5–1 Hz neocortical slow oscillations (SOs), 12.5–16 Hz thalamocortical fast sleep spindles, and 80–100 Hz hippocampal ripples (frequency ranges in humans), all of which have been associated with human memory reactivation and consolidation processes (12–17). Although SOs and spindles are most prominent in neocortex, they are also present in human HPC (18–21), consistent with their postulated role in HPC-cortical memory transfer. Interestingly, SOs, spindles, and ripples are temporally organized, with faster oscillations preferentially expressed at a particular phase of the slower rhythm. In general terms, such cross-frequency phase-amplitude coupling (PAC) is thought to enable brain communication across multiple spatiotemporal scales (22, 23). During sleep, PAC occurs both within and across brain areas (24–28), including in human (para)hippocampus (21, 29, 30). Remarkably, animal findings have shown that stronger SO–spindle coupling enhances plasticity and memory retention (31, 32), suggesting that these nested oscillations are required for effective information transfer and storage (33).

In addition to these "classical" forms of NREM PAC among SOs, spindles, and ripples, SOs affect the expression of 1–4 Hz delta activity (34), which is physiologically distinct from SOs (35–37). Whether SO–delta coupling also exists in human HPC, and whether HPC SO and delta activity coordinate faster activity in a similar fashion, is presently unknown. Moreover, HPC ripples often occur in conjunction with ~4 Hz sharp waves (38), which could be expressed as delta–ripple PAC (21, 39). Finally, neocortical theta (~6 Hz) activity is also organized by the SO phase (40), and this coupling has been related to memory consolidation (41). Whether analogous SO–theta coupling exists in human HPC is unknown.

Contrary to the relatively organized structure of NREM oscillations, REM sleep exhibits an overall cortical "desynchronization" (i.e., fast, low-amplitude activity), although isolated REM SOs have been described in rodent cortex (42). Similarly, human HPC expresses both fast beta/gamma activity (43) and relatively slow activity during REM sleep. Yet, whether this slower component entails SO/delta (44) or theta (45) activity is still contentious. Moreover, with the exception of delta–gamma coupling in parahippocampal areas (46), the extent of interacting HPC oscillations in human REM sleep has not been explored.

Importantly, the HPC consists of anatomically and functionally distinct domains, with anterior and posterior regions showing differential relations with human memory performance (47–50). Given that sleep oscillations are often spatially restricted (20, 24, 25, 51), it is an open question whether these rhythms and their interactions vary along the longitudinal axis of the human HPC, possibly in relation to functional specialization. Another relevant factor concerns between-subject variability, with various aspects of scalp-recorded sleep oscillations showing large and reproducible individual differences (25, 52–55). Whether such variability extends to human HPC is unknown.

In sum, while electrophysiological HPC activity during sleep is deemed pivotal for memory processing, a comprehensive characterization of human HPC sleep oscillations and their interactions has not been performed. The wealth of fundamental animal knowledge notwithstanding (56–58), non-negligible species differences in the expression and frequency of sleep oscillations make it unclear whether theoretical accounts on the role of the sleeping HPC are consistent with human data. Taking advantage of the unique opportunity offered by neurosurgical monitoring in 12 epilepsy patients, we examined oscillatory activity along the longitudinal HPC axis during light NREM (N2), deep NREM (N3), and REM sleep. For comparison purposes, we also included scalp activity. We assessed spectral power and PAC across an extensive (0.5–200 Hz) frequency range, employing data-driven permutation-based approaches to characterize and contrast oscillatory activity in various ways. Our results indicate extensive PAC in human HPC, with the majority of channels expressing clear spectral interactions, but also a remarkable heterogeneity between HPC and scalp, HPC sites, and individuals.

Results

We examined overnight scalp and invasive hippocampal electroencephalography (EEG) in a sample of 12 epilepsy patients. A total of 61 HPC contacts were included (5.1 ± 2.3 per patient, range: 1–8). Only HPC contacts from the non-pathological hemisphere were used, as evidenced by clinical monitoring. HPC contacts were classified as anterior, middle, or posterior HPC based on individual anatomy (see Supp. Table 1 for all electrode locations). Polysomnography-based sleep architecture is shown in Table 1.

	mean	SD
N1(%)	29.8	21.8
N2 (%)	40.5	13.7
N3 (%)	13.2	9.0
REM (%)	16.5	8.0
N1 (min)	171.4	175.0
N2 (min)	203.9	74.0
N3 (min)	64.1	45.7
REM (min)	81.1	41.0
total sleep (min)	520.5	113.5
WASO (min)	82.9	77.2
sleep efficiency (%)	87.1	11.7

Table 1. Sleep architecture. WASO: wake after sleep onset.

<u>Spectral profiles</u>

Considering a total of 86.7 ± 41.3 (N2), 26.3 ± 25.5 (N3), and 55.3 ± 35.8 min (REM) of artifact-free data, power spectra were determined for each channel and sleep stage, adjusting for 1/f power scaling with a slope-fitting procedure to enhance the visibility of narrowband peaks. Averaged across subjects, the spectrum at the central scalp derivation (Cz) showed greatest SO/delta (0.5–4 Hz) power in N3, prominent fast spindle (12.5–16 Hz) and theta (4–8 Hz) peaks in both NREM stages, and theta and beta (~25 Hz) peaks in REM (Fig. 1A), consistent with typical sleep (53, 54). The HPC spectrum also displayed a strong fast spindle peak and additional delta/theta peaks during NREM (Fig. 1B). Most prominently, HPC showed a strong gamma (40–100 Hz) peak in all sleep stages, matching earlier observations in parahippocampal cortex (43) and consistent with the frequency range of human ripples (12, 21). Results of sleep stage comparisons are indicated at the top of each plot. Direct comparisons between scalp and HPC confirmed the specificity of HPC gamma while showing greater <20 Hz power at the scalp (Supp. Fig. 1).

Figure 1. Scalp and hippocampal sleep spectra. Top: Group-level spectra averaged across all 12 patients at the scalp (A) and in HPC (B). Error shading: standard error of the mean across patients. Horizontal color bars above plots indicate significant (P<0.05) pairwise stage differences (cluster-based permutation test). Filled color reflects stage with greater power. Gray vertical lines at 1.5, 4, 9, 12.5 and 16 Hz indicate approximate boundaries between SO, delta, theta, slow spindle, fast spindle, and faster activity. Right panels in (A) and (B) show same spectra without slope removal. The flattening of the raw spectrum at low (<4 Hz) frequencies results in comparatively low adjusted power after slope removal. Middle/bottom: Spectra of two example patients highlight differences between individuals, both at the

scalp (C and F) and in HPC (D and G; error shading: standard error of the mean across data segments; stage differences as above). A further breakdown by HPC channel (E and H) reveals spectral differences along the longitudinal HPC axis (anterior: lighter color; posterior: darker color; 7 HPC channels for both patients).

Because sleep spectra show substantial between-subject variability (53, 54), we examined individual patients' spectral profiles. While scalp NREM spectra commonly displayed strong peaks in the fast spindle range, the presence, prominence, and frequency of other spectral components varied (Fig. 1CF, Supp. Fig. 2). Theta and delta peaks, when present, could be largest in any sleep stage. Although slow spindles (9–12.5 Hz) are less prominent at central scalp positions (59), several patients showed distinct peaks in the slow spindle range. This peak was clearly separate from nearby theta and fast spindle peaks in some instances (Fig. 1C), suggesting they reflect distinct oscillatory phenomena.

Considering HPC, channel-averaged spectra exhibited quite consistent gamma peaks across individuals, but otherwise showed large variability, particularly in the delta and theta range (Fig. 1DG, Supp. Fig. 2). A fast spindle peak could be discerned in NREM in most individuals, but was absent in some. Separate slow/fast spindle peaks were not observed. Also noteworthy is that while fast spindle peaks were absent during REM at the scalp (in line with sleep scoring criteria), such peaks were sometimes observed in HPC (Fig. 1DG), possibly related to asynchronous sleep stage transitions between brain structures (60, 61). In contrast, REM beta power frequently showed a peak at the scalp, but generally not in HPC.

Next, we considered spectra of single HPC channels for individual patients (Fig. 1EH). HPC channel spectra from a single patient were much more similar compared to spectra from different patients, with spectral peaks at similar frequencies. Nonetheless, peak amplitudes varied considerably along the longitudinal HPC axis. Power typically changed gradually and monotonically from contact to contact. However, the direction of power increase (anterior-posterior or posterior-anterior) varied between individuals, and could occur in opposite directions for different frequency components in the same individual (e.g., N2 spindle and gamma power in Fig. 1E), ruling out non-specific channel differences in signal amplitude. As a consequence, group-level comparisons between anterior, middle, and posterior HPC did not indicate systematic regional differences in spectral power (Supp. Fig. 3).

In sum, HPC spectral profiles varied with sleep stage, but this pattern differed substantially from that seen at the scalp. Moreover, we observed surprisingly diverse spectral patterns between individuals, and for HPC, between channels. Notably, this variability in spectral composition is likely to affect their potential to engage in cross-frequency PAC, to which we turn next.

Single-channel PAC

We quantified two aspects of cross-frequency PAC. First, the strength of PAC (subsequently referred to as "coupling strength", "coupling", or simply "PAC") signifies the degree to which activity of a faster frequency is non-uniformly distributed across the phase of a slower frequency. Second, coupling phase indicates the phase of the

slower oscillation at which faster activity is preferentially expressed. Both metrics were assessed for every pair of frequencies in the 0.5–200 Hz range, separately for every one-minute artifact-free data segment, channel, sleep stage, and patient. Furthermore, coupling strengths for each segment were z-scored with respect to surrogate distributions. Given the large spectral variability reported in the previous section, we first sought to understand coupling dynamics for several example individual channels.

Figure 2. Phase-amplitude coupling for example channels. (A and D) Coupling strength at the scalp (A) and the most posterior HPC channel (D) for two example patients. White outlines indicate clusters of significantly higher than zero coupling across one-minute data segments (P<0.05, cluster-based permutation test). (B and E) Mean preferred coupling phases for significant frequency pairs. Insets:

normalized circular histograms of coupling phase across data segments for example frequency pairs, indicating consistent phase preferences. Phases defined with respect to sine wave, with peak at 90˚ and trough at 270˚. (C and F) Difference in coupling strength between sleep stages. White/red outlines indicate significant positive/negative clusters, respectively (P<0.05, cluster-based permutation test). (G) Difference in coupling strength between scalp and HPC. White/red outlines indicate significant positive/negative clusters, respectively (P<0.05, cluster-based permutation test).

Assessing the presence of PAC for the same example subjects as in Fig. 1, both scalp (Fig. 2A) and HPC (Fig. 2D) channels showed prominent hotspots of frequency pairs where coupling strengths were significantly higher than zero (P<0.05, cluster-based permutation test across data segments). Note that a significant cluster may comprise more than one coupling phenomenon due to bridging between subclusters. Interestingly, while some cluster maxima coincided with peaks in the power spectrum, shown in the lower and right margin of each panel, other coupling effects were not accompanied by corresponding spectral peaks. For frequency pairs that were part of a significant cluster, we determined the mean coupling phase (Fig. 2BE) and, for several example frequency pairs, we evaluated the distribution of coupling phases across data segments (Fig. 2BE, insets). We now consider several observations in detail.

Scalp PAC in both example patients was most clearly seen during NREM between SO and theta and between SO and fast spindle activity (Fig. 2A), with phases of maximal coupling towards the SO trough and peak, respectively (Fig. 2B), consistent with previous findings (25, 27, 40). Note that theta clusters extended into the slow spindle range without clear boundaries (see *Discussion*). Coupling in these ranges was significantly stronger in NREM than REM, and also differed between N3 and N2 (Fig. 2C; P<0.05, cluster-based permutation tests), indicating the stage-specificity of these effects. Scalp PAC during REM was generally less pronounced, although there was evidence for PAC between delta/theta and beta/gamma in both patients. Other effects differed between patients. For example, patient p12 exhibited N2 theta–beta coupling coinciding with a theta spectral peak (Fig. 2A, right). In contrast, patient p4 demonstrated strong PAC during NREM between gamma and >150 Hz activity (Fig. 2A, left), likely reflecting muscle activity (see *Discussion*).

HPC PAC was also clearest during NREM. Both example patients expressed strong evidence for the modulation of high gamma activity, with distinct clusters centered at ~80 Hz and ~140 Hz, consistent with ripple activity (Fig. 2DE). Ripple activity was coupled to the oscillatory phase of various slower frequency bands, including the SO, delta, theta, and fast spindle ranges, depending on NREM stage and patient. Note that ripple locking to the SO trough, rather than the peak, suggests an opposite polarity compared to the scalp, with the SO trough reflecting the physiological UP state for these channels (20). The observation of delta–ripple coupling is consistent both with sharp wave-ripple sequences (39), and with ripple coupling to a delta rhythm of different origin. Interestingly, whereas ripple activity was locked to the trough of both the SO and delta ranges for patient p12 (Fig. 2D, right), it was tied to opposite phases for SO (trough) and delta (peak) activity for patient p4 (Fig. 2D, left). Distinct phase preferences for SO and delta activity fit the idea that these slow components reflect separate phenomena (35). Spindle–ripple coupling was

tied to the spindle trough, as reported previously (21, 30). Direct stage comparisons indicated that ripple activity was most strongly coupled to slower activity during N3, followed by N2, and finally REM (Fig. 2F). In addition to the modulation of ripple activity, spindle activity was preferentially organized into the SO/delta trough in both patients (Fig. 2DE), again consistent with opposite polarities in HPC and scalp, although coupling was weak for p12. N2 SO–delta/theta coupling appeared in patient p12 (Fig. 2DE, right), but with a phase preference opposite to that seen at the scalp. The pattern of HPC PAC during REM was less clear, although some coordination of faster activity by the SO/delta phase was present, which could even surpass that seen during NREM (e.g., N2/N3 SO–theta in Fig. 2D, right). Finally, direct contrasts between HPC and scalp underscore the regional specificity of many of the aforementioned effects, including stronger N2/N3 ripple and N2 spindle modulation in HPC, but enhanced N2/N3 theta modulation at the scalp (Fig. 2G).

Combined, these results illustrate clear temporal coupling between various pairs of oscillatory rhythms. Although the expected coordination among SO, spindle, and ripple rhythms was present, coupling extended well beyond these frequency ranges, with different dynamics for HPC and scalp, and different patterns for NREM and REM sleep. Moreover, coupling dynamics differed between patients, potentially related to corresponding individual differences in spectral profiles.

Single-patient PAC along the hippocampal axis

Next, we considered PAC for all channels along the HPC axis within individuals. Fig. 3AB shows the complete profile for two example patients (including patient p4 described in the previous section). Although coupling patterns showed a certain degree of similarity across channels for both patients, clear differences were also present, such that frequency pairs showing large and above-chance coupling on one channel could show much reduced (and non-significant) coupling on other channels. Interestingly, while the degree of SO–spindle and spindle–ripple coupling corresponded to the height of the spindle peak in the spectrum for patient p4, this phenomenon was not observed for patient p11. (Also note the absence of clearly distinguishable spindle–ripple clusters in the latter patient.) In contrast, channels with largest ripple power were generally not the ones that were modulated most strongly by slower rhythms.

Interestingly, patients expressed distinct longitudinal profiles of HPC coupling, such that frequency pairs could be most strongly coupled at different HPC sites for different patients. These findings echo our earlier observations for spectral power (Fig. 1). To further examine longitudinal coupling profiles, we plotted coupling strengths along the HPC axis for five example frequency pairs (Fig. 3C). These plots not only illustrate distinct regional coupling patterns for the two patients, but also distinct patterns for different frequency pairs and sleep stages. Note the different profiles for SO–ripple and delta–ripple coupling, again suggesting that SOs and delta activity constitute distinct phenomena. Also noteworthy is that while coupling was generally weakest during REM, REM coupling was nonetheless apparent for most frequency pairs, and even surpassed the level seen during NREM on some channels (e.g., anterior

SO–ripple coupling for patient p4). The latter finding fits with the presence of SOs during REM sleep (42).

Figure 3. Phase-amplitude coupling along hippocampal axis for example patients. (A and B) Coupling strengths for all HPC contacts in two example patients. White outlines indicate clusters of significantly higher than zero coupling (P<0.05, cluster-based permutation test). (C) Coupling strengths for five frequency pairs, indicated in (A) with arrows. Presence of marker indicates frequency pair is part of significant cluster in (A) or (B).

In sum, while oscillatory interactions can be seen along the entire axis of human HPC, whether and how strongly specific frequency pairs interact varies from contact to contact, consistent with the notion of spatially localized sleep oscillations

(20, 42, 51). Moreover, HPC coupling profiles vary considerably between individuals, extending similar observations from non-invasive approaches (25, 53, 54).

Group-level PAC

We next examined scalp and HPC PAC across patients. As shown in Fig. 4A (left), scalp coupling was most evident during N3 between the SO and the theta/spindle bands, with a similar, albeit weaker, cluster present during N2. In contrast, REM showed evidence of weak coupling between delta/theta phase and beta/gamma power, consistent with the effects of individual patients (Fig. 2A). Both NREM stages also exhibited a cluster of gamma–high gamma coupling (maximum at 40 x 200 Hz), similar to that seen for one of the example patients (Fig. 2A). This finding is likely due to muscle rather than neural activity (see *Discussion*). Direct stage comparisons indicated significantly stronger PAC in N3 than REM for the SO-theta effect (Fig. 4B, left). Overall, these findings corroborate previous scalp findings (24, 25, 27), while also indicating the consistency of scalp-measured PAC across individuals.

Figure 4. Group-level phase-amplitude coupling. (A) Mean coupling strength for scalp (left) and HPC (right). White outlines indicate clusters of significantly greater than zero coupling (P<0.05, cluster-based permutation test). (B) Mean difference in coupling strength between sleep stages for scalp (left) and HPC (right). (C) Difference in coupling strength between scalp and HPC. White/red outlines in (B) and (C) indicate significant positive/negative clusters, respectively (P<0.05, cluster-based permutation test).

For HPC, coupling profiles were first averaged across available HPC contacts for each patient. Subsequent group analyses revealed prominent hotspots of delta–ripple coupling in both N2 and N3 (Fig. 4A, right). In contrast, clearly delineated foci for other coupling phenomena that might be expected (e.g., SO–spindle, SO–theta, spindle–ripple) were not apparent at the group level. REM PAC was much lower, showing weak hotspots in the SO–ripple and delta–ripple ranges. Nonetheless, statistical evaluation signaled massive clusters for every sleep stage, essentially comprising all frequency pairs. (An alternative statistical approach using False Discovery Rate-based (62) rather than cluster-based correction yielded essentially similar results (not shown)). Note that this observation should not be taken to imply that every frequency is consistently modulated by every other frequency. Rather,

these findings suggest that the aforementioned variability in coupling dynamics between patients and individual HPC channels results in above-zero group averages for every frequency pair, which then form a single big cluster. Direct stage comparisons indicated stronger coupling in both NREM stages relative to REM, for frequency pairs including, but not limited to, the delta–ripple range (Fig. 4B, right). Directly contrasting scalp and HPC coupling profiles revealed significantly stronger PAC in HPC for each stage, mainly between SO/delta and beta/gamma frequencies (Fig. 4C).

While the group analyses of Fig. 4A provide evidence for the general presence of PAC, they offer limited information about which exact frequency pairs are coupled, particularly in HPC. To obtain a qualitative view how often particular frequency pairs were coupled in our sample, we adopted an unsupervised clustering approach. Specifically, for every subject, sleep stage, and channel, we extracted all frequency pairs showing a local maximum in coupling strength. Of these, the frequency pairs corresponding to coupling strengths above a threshold were submitted to a density-based clustering algorithm (63). We employed both adaptive, data-derived thresholds (i.e., different thresholds for every subject, sleep stage, and region) and fixed thresholds (PAC > 2). This yielded, for every subject, sleep stage, and brain region, a relatively small number of clusters, whose centers were taken as frequency pairs expressing coupling. Clustering results for several example patients using adaptive thresholds are shown in Supp. Fig. 4.

Fig. 5 aggregates the clustering algorithm results across individuals as heat maps, with color indicating how many subjects exhibited evidence of coupling at each frequency pair, using either adaptive (Fig. 5A) or fixed (Fig. 5B) thresholds. Additional plots below and to the right of each panel indicate how many unique individuals showed coupling for every modulating and modulated frequency, respectively.

Using adaptive thresholds, scalp NREM showed strongest group effects for SO–spindle, SO–theta, and gamma–high gamma frequency pairs (Fig. 5A, left), similar to the effects using conventional group statistics in Fig. 4A (left). Scalp REM showed most consistent evidence for coupling between delta/theta modulating frequencies and spindle/beta/ripple modulated frequencies. In contrast, the clustering approach yielded a more clearly defined NREM profile for HPC (Fig. 5A, right), compared to the diffuse appearance of the conventional group approach (Fig. 4A, right), whereas HPC REM coupling continued to show a highly variable pattern (detailed description of coupling effects in next paragraph). However, it is important to note that overall coupling strengths were much lower in REM compared to NREM, and reduced at the scalp compared to HPC. Consequently, the fixed-threshold approach abolished the majority of scalp effects, and for REM, only a cluster of delta/theta–beta coupling remained in HPC (Fig. 5B).

Considering the adaptive threshold approach in detail, NREM scalp coupling was most consistently seen for N3 SO–spindle (9/12 individuals) frequency pairs, followed by N2 SO–spindle (5), N3 SO–theta (5), N2 SO–theta (4), N2 SO–theta/slow spindle (4), and N3 SO/delta (4) frequency pairs. In HPC, NREM coupling was highly

consistent for N2 (12) and N3 (11) delta–ripple frequencies, followed by coupling between modulating frequencies on the delta/theta border and their modulated frequencies covering the high spindle/low beta range (N3: 5; N2: 4). SO–ripple coupling was seen in both N2 and N3 (4), whereas SO–delta coupling only occurred in N2 (4). Spindle–ripple coupling emerged in only three patients in both NREM stages, whereas above-threshold SO–spindle PAC was not observed at all. A complete overview of all NREM effects present in at least 3 subjects (25%) can be found in Supp. Table 2. The even larger variability of REM coupling profiles, due to the inclusion of weaker (and likely noisier) effects, precludes meaningful construction of a similar table for REM.

Figure 5. Group-level counts of phase-amplitude coupling. Number of individuals showing PAC for each frequency pair using adaptive (A) and fixed (B) thresholds at the scalp (left) and HPC (right) for different sleep stages (rows). Line plots indicate number of unique individuals showing PAC at each modulating (bottom) or modulated frequency (right) bin.

In sum, while sleep oscillations are relatively consistently coupled at the scalp, group-level PAC within HPC was much less systematic, with many effects occurring in only a minority of individuals, further confirming the observations from individual patients (Fig. 2 and 3).

Group-level PAC in hippocampal regions

Finally, we considered group-level PAC for anterior, middle, and posterior HPC regions. Interestingly, direct comparisons between HPC regions revealed two systematic differences (Fig. 6). First, N2 coupling between SOs and activity in the theta/spindle range was significantly stronger in anterior than middle HPC. Second, SO/delta–ripple PAC showed a difference during REM, with greater coupling in

anterior HPC relative to both middle and posterior HPC. (This effect was also present at the subject-level for patient p4 in Fig. 3C.) Note that both effects occurred in the absence of systematic regional differences in spectral power (Supp. Fig. 3). Region-specific analyses considering the presence of coupling (Supp. Fig. 5) and sleep stage differences (Supp. Fig. 6) yielded results similar to those seen across all HPC channels, as presented in Fig. 4A (right) and Fig. 4B (right), respectively.

Figure. 6. Regional differences in hippocampal phase-amplitude coupling. Mean difference in coupling strength between anterior and middle, middle and posterior, and anterior and posterior HPC for each sleep stage. White outlines indicate significant positive clusters (P<0.05, cluster-based permutation test; no negative clusters present). Note different sample sizes for each regional comparison due to variable electrode coverage.

Discussion

The present study offers a comprehensive analysis of cross-frequency phase-amplitude coupling at the scalp and in the hippocampus of sleeping humans. Considering frequency pairs over an extensive 0.5–200 Hz range, we found strong evidence for the presence of spectral interactions, with the overall strength of coupling greater in HPC than at the scalp, and greater during NREM than REM sleep. Scalp PAC during NREM consisted of SO–theta and SO–spindle coupling, while theta–gamma coupling was seen during REM. In contrast, HPC PAC was seen for many frequency pairs, including both expected (e.g., NREM SO–ripple, SO–spindle, delta–ripple, spindle–ripple), and novel interactions (e.g., NREM SO–delta, SO–theta, REM delta/theta–beta). At the same time, coupling profiles differed substantially between individuals, and between HPC sites within individuals.

<u>Sleep oscillations at the scalp</u>

Scalp-level group analyses indicated well-known NREM coupling phenomena between SO phase and amplitudes of both spindle and theta activity (Fig. 4A) (24, 25, 27, 28, 40). These effects were stronger in N3 compared to N2, consistent with the lower density of SOs in N2 (mostly constituting K-complexes) (25). Although individuals expressed variability in both power spectra (Fig. 1CF, Supp. Fig. 2) (53, 54) and coupling profiles (Fig. 2A), SO–theta and SO–spindle effects were relatively

consistent across individuals, thus leading to circumscribed effects at the group level. Importantly, this corroboration of known effects establishes the general suitability of our analytical approach.

Interestingly, while coupling was generally weaker during REM, we observed significant coupling in this sleep stage between the phase of the delta/theta rhythm and amplitudes in the beta/gamma range, consistent with the presence of theta and beta peaks in the power spectrum. We are not aware of previous studies reporting such REM coupling in human scalp EEG, although this finding fits with similar findings in rodents (64, 65) and human parahippocampal cortex (46) (also see *Sleep oscillations in the hippocampus*).

During NREM, clusters of gamma–high gamma (>150 Hz) coupling were seen at both the group and individual levels. However, high-frequency (>~80 Hz) components of non-invasive EEG cannot be assumed to reflect neural activity without additional validation. In contrast, scalp-recorded electromyographic activity extends well beyond this range (66). For this reason, and because the observed coupling effect was absent during the atonic state of REM sleep (Fig. 4A), along with a REM-related reduction in high-gamma power (Fig. 1A), the most likely explanation for this effect is muscle activity.

Recent rodent work indicates that spindle oscillations coordinate the expression of ripple-like (~100 and ~200 Hz) activity in neocortex (67). However, our scalp recordings did not show modulation of frequencies >85 Hz by NREM spindles or SOs, in line with previous scalp analyses (21). Similarly, scalp spectra were not indicative of obvious ripple activity (Fig. 1ACF), suggesting that the possibility that ripples can be non-invasively recorded from the scalp is limited (68, 69).

A 9–12.5 Hz slow spindle peak was observed in only a minority of cases, likely due to suboptimal positioning of the central scalp electrode relative to the predominantly frontal generators of slow spindles (54, 59). As a consequence of this, as well as the limited spectral precision of our methods (see *Methodological considerations*), our PAC findings do not directly speak to the recently raised possibility that theta waves are mistaken for slow spindles (40). However, we note that distinct slow spindle and theta components were sometimes apparent in the power spectrum (Fig. 1C).

Sleep oscillations in the hippocampus

Considering HPC, we found strong evidence for various oscillatory interactions, particularly during NREM. Yet, coupling patterns were strikingly diverse across individuals and HPC contacts. Furthermore, the overall profile of oscillatory pairs engaging in HPC PAC deviated substantially from the one seen at the scalp, indicating the regional specificity of oscillatory organization.

Across individuals and channels, NREM delta-ripple PAC was the clearest and most consistent coupling phenomenon (Fig. 3, 4A, 5), corroborating earlier reports (12, 21). A likely possibility is that the delta aspect of this coupling effect reflects the

sharp wave component of sharp wave–ripple complexes, since the spectral representation of sharp waves peaks in the delta range (but extending up to 10 Hz) (39). Also noteworthy in this respect is the observation of delta–beta (20–25 Hz) coupling in several instances. This beta component could be akin to the "slow gamma" band modulation recently shown to constitute the rhythmicity of overlapping ripples (39). That said, the current findings do not rule out the possibility of alternative, non-sharp wave, delta oscillations contributing to delta-ripple PAC.

In this light, it is important to point out that HPC delta–ripple and SO–ripple coupling likely constitute distinct phenomena. First, above-threshold SO–ripple coupling appeared in only a third of our sample, compared to the full sample for delta–ripple PAC. Note that this observation corresponds to a previous report (partially overlapping with the data presented in the current paper) of ripples being most strongly modulated by delta rather than SO activity (21). Second, longitudinal profiles for these frequency pairs differed substantially within individuals (Fig. 3C). Third, separate SO–ripple and delta–ripple hotspots could be discerned within an individual, sometimes with opposite phase preferences (e.g., Fig. 2D, left, N2). Fourth, we observed SO and delta activity to be coupled themselves in several instances, similar to earlier findings in neocortex (34). Overall, these findings fit earlier arguments that <4 Hz activity does not constitute a single phenomenon, but may be separated into physiologically and functionally distinct entities (35–37).

Also noteworthy is the observation of N2 coupling in HPC between SOs and the high theta/slow spindle range (Fig. 3B, 5A). Given that we did not observe separate slow and fast spindle peaks in HPC, we tentatively interpret the modulated frequency to reflect theta activity. Recent non-invasive findings indicate a role for SO–theta PAC in memory consolidation (41). The presence of a similar dynamic in HPC lends further support to this idea, although it is striking that this phenomenon was not observed in N3, during which it was most prominently seen at the scalp.

While we observed unambiguous clusters of strong SO–spindle and spindle–ripple coupling for individual HPC channels (Fig. 2 and 3), these findings did not emerge consistently at the group level, using either standard (Fig. 4) or unsupervised clustering approaches (Fig. 5). This finding is somewhat surprising in light of recent demonstrations of SO–spindle–ripple coupling in human (para)hippocampus (21, 29, 30). However, an important methodological difference is that these previous studies mostly limited PAC calculations to detected waveforms, whereas we considered continuous data without imposing specific detection criteria (see *Methodological considerations*). Moreover, the study reporting hierarchical nesting of all three waveforms (partially overlapping with the current data) (21) included only the contact with highest spindle power, thereby making analyses contingent on the presence of clear spindles. Indeed, while we observed spindle spectral peaks on at least one HPC channel in most individuals, many channels were not suggestive of spindle activity (see spectral profiles inside panels of Fig. 3AB). This may also explain inconsistent reports regarding the presence of spectral spindle peaks in (para)hippocampus (70, 71). More generally, although absolute signal amplitudes are higher in HPC, both SOs and spindles are much less visually apparent in HPC compared to the scalp. In sum, while

our results confirm the existence of SO–spindle and spindle–ripple coupling in human HPC, our findings also indicate that this coupling may not be as ubiquitous as previously thought.

Similar to scalp findings, hippocampal coupling was much reduced during REM compared to NREM (Fig. 4AB). While conventional group analyses did not indicate clearly defined REM coupling, unsupervised clustering suggested the presence of delta/theta–beta (4 vs. 36 Hz) coupling in three patients (fixed thresholds, Fig. 5B). This result is reminiscent of previous findings in human parahippocampal cortex (46), although in that report the modulating frequency was slower (1–3 Hz) and modulation was strongest in the 60–100 Hz range. As such, our findings do little to answer whether the human functional analog of rodent hippocampal theta lies in the SO/delta (44, 46) or theta (45) range, since our 4 Hz modulating frequency falls right in between. Rather, our observations of variable spectral and coupling profiles during REM suggest that both SO/delta and theta frequencies can be observed (also see *Methodological considerations*). More generally, the ambiguous evidence for REM theta-like activity in human HPC underscores the need to consider species differences in oscillatory organization. Also noteworthy is that SO–ripple coupling was found to be stronger in HPC vs. scalp in all sleep stages including REM (Fig. 4C), suggesting that SOs may also occur in this sleep stage (also see *Regional and individual variability in hippocampus*). These findings are consistent with asynchronous sleep stage transitions between scalp and HPC (60, 61), as well as the existence of SOs during REM (42).

<u>Regional and individual variability in hippocampus</u>

A key observation of the present study is the large heterogeneity in oscillatory and coupling profiles between HPC contacts and individuals. What could underlie this variability? First, several technical factors could play a role. For example, electrode orientation relative to generating fields could render contacts differentially sensitive to certain waveforms. Related, HPC subfields (e.g., CA3, CA1, DG) are known to participate differently in sleep-related network activity (39, 72). Unfortunately, anatomical resolution in the current study cannot answer whether subfield identity could account for our findings. It also deserves mention that the observed regional variability makes it unlikely that HPC findings are driven by the mastoids reference, since such an influence should impact all HPC contacts equally.

While the preceding factors essentially argue that heterogeneity is a rather uninteresting consequence of data acquisition, another possibility is that the observed differences constitute true variability. Specifically, regional HPC variability is consistent with accumulating evidence of the local nature of sleep oscillations (20, 24, 25, 51). Particularly telling in this respect is that different spectral components could be strongest at opposite ends of HPC for the same individual (e.g., N2 spindle and ripple activity in Fig. 1E). Similarly, distinct coupling phenomena could be most pronounced at different contacts (e.g., N3 SO–ripple in anterior, and delta–ripple in posterior HPC; Fig. 3C).

At the group-level, HPC regional variability emerged as significantly enhanced SO–theta and SO–spindle PAC in anterior vs. middle HPC during N2, as well as

enhanced SO–ripple coupling in anterior vs. middle/posterior HPC during REM (Fig. 6). However, these effects should be interpreted with caution since these frequency pairs did not emerge as clearly defined clusters when assessing the presence of coupling (Fig. 4 and 5, Supp. Fig. 5). Nonetheless, these regional differences provide some evidence that the pertaining frequency bands were in fact coupled. Furthermore, regional variability in SO–ripple coupling during REM suggests that aspects of NREM electrophysiology can also be observed during REM in HPC (also see *Sleep oscillations in hippocampus*).

In contrast, no systematic regional coupling differences were observed for SO–ripple, delta–ripple, or spindle–ripple pairs during NREM, nor for SO–spindle coupling during N3. Similarly, we did not observe any systematic regional differences in spectral power for any frequency band during any sleep stage (Supp. Fig. 3). However, we note that the anatomy-based division into HPC thirds does not necessarily result in functionally homologous subregions across patients. Thus, defining regions functionally might have yielded different results. Although we did not have specific predictions regarding oscillatory structure in anterior, middle, and posterior HPC, we examined this possibility because of well-known intrinsic and extrinsic wiring differences between anterior and posterior HPC (or the analogous ventral/dorsal divide in rodent HPC) (73, 74), and evidence of differential regional relations with human memory (47–50). In sum, with the potential exception of N2 SO–spindle coupling, we conclude that the sleep oscillations commonly thought relevant for memory consolidation (i.e., SOs, spindles, ripples), are not differentially present or coupled along the hippocampal axis in a systematic fashion.

Nonetheless, we emphasize that this group-level conclusion does not reflect individual patients, who showed marked differences in spectral power and PAC along the HPC axis. We suggest that individual differences in HPC PAC constitute true variability representative of the population, similar to replicable individual differences of scalp sleep oscillations (25, 52–55). Whether such variability is further related to general (e.g., cognitive functioning, age) or disease-related (e.g., epilepsy history, medication) factors cannot be determined from the present sample.

Methodological considerations

Although findings from medicated epilepsy patients necessarily warrant caution when generalizing to healthy populations, sleep architecture (Table 1) and scalp power spectra (Fig. 1A, Supp. Fig. 2) were within normal ranges, only data from the non-pathological HPC was used, and a stringent artifact rejection procedure removed >50% of data, making it unlikely that our results are due to epileptogenic activity.

While our sample size (N=12) was sufficiently large to allow group-level inferences, its modest size, though not atypical for invasive studies, has implications for generalizability. In particular, while the general notion of individual and regional variability may be extrapolated to the population as a whole, the specific pattern of variability should not. For example, while NREM SO–theta, SO–spindle, delta–ripple, and spindle–ripple coupling were all observed, the present sample did not contain a

single individual/channel clearly expressing all of these effects. Nonetheless, it is not unreasonable to expect the co-existence of each of these effects in the larger population. Similarly, the population likely harbors individuals with idiosyncratic spectral peaks and coupling effects for frequencies not seen in the present sample. In this light, we also note that previous invasive studies may have reached quite different conclusions (e.g., presence vs. absence of spectral spindle peaks; SO/delta vs. theta activity during REM) due to taking small samples from a highly variable population, thus increasing the likelihood of capitalizing on chance.

In the present approach, PAC was calculated over continuous data. This contrasts with discrete approaches where PAC calculations are contingent on the presence of specific waveforms, detected according to specific criteria of frequency, amplitude, and/or duration. In line with our objective of assessing spectral interactions in a data-driven fashion without setting *a priori* restrictions on the forms of coupling that might be present, we favored the continuous approach. We also note that the employed wavelet approach is inherently limited in spectral (and temporal) resolution, hampering isolation of closely spaced spectral components (e.g., theta, slow spindle, and fast spindle components). Moreover, the use of a fixed set of wavelet frequencies does not allow subject-specific targeting of oscillatory phenomena that vary in frequency between individuals (54, 75). However, given that our approach identified various previously described coupling phenomena, we believe these methodological choices do not pose a major concern.

Metrics of PAC are typically sensitive to both genuine cross-frequency coupling and asymmetrical features of a single oscillator (76). However, specific and distinct generating circuits can reasonably be ascribed to several spectral components we found to be coupled (e.g., SO, spindle, ripple, sharp wave). Also worth mentioning is that the present PAC metric is not sensitive to multimodal phase preferences, for which other approaches may be preferred (77). Ultimately, detailed waveform analyses are required to fully understand the origin of each of the observed effects (76), particularly those that have little precedence in the literature.

In several instances we observed that the presence of coupling coincided with corresponding spectral peaks, and that coupling strength scaled with the height of the modulating peak (e.g., N3 spindle–ripple PAC in Fig. 3A). In other instances, this relation was not apparent (e.g., N3 spindle–ripple PAC in Fig. 3B), indicating no straightforward mapping between power and PAC. Moreover, it has been suggested that the presence of a narrowband spectral peak for the modulating frequency is required to meaningfully interpret PAC (22). However, the most robust coupling phenomenon in HPC, delta–ripple PAC, occurred without clear peaks in the delta range (Fig. 1B). Finally, there were instances where two frequencies showed clear spectral peaks, but were not significantly coupled, indicating that coupling does not invariably emerge when multiple oscillators are present.

Functional implications

The present findings are relevant to theories on the role of coupled sleep oscillations in memory consolidation. HPC PAC was seen for a remarkably diverse

range of frequency pairs during NREM, with coupling phenomena extending well beyond previously described interactions between SOs, spindles, and ripples (21, 29, 30). Indeed, these classical coupling patterns were less consistently present than we expected, whereas delta–ripple PAC was reliably present in the full sample. That said, SOs, spindles, and ripples are robustly tied to plasticity and consolidation processes (12–17, 31, 32), whereas the functional relevance for other coupling pairs remains to be demonstrated. It is also possible that the coordination of HPC ripple and spindle oscillations by spindles and SOs, respectively, becomes more evident when considering the phase of neocortical rather than HPC spindle/SO activity. Such cross-regional spectral interactions have been described in animals (78) and humans (21), but were outside the scope of the present study. In contrast to NREM, the functional relevance of REM sleep remains enigmatic (10, 11). Whatever its function, our findings indicate that the role of coupled sleep oscillations in REM-related processing is likely limited, since only (weak) coupling between delta/theta and beta/gamma components emerged during this sleep stage.

While variability is typically treated as a nuisance or noise factor in neuroscience, proper understanding of a phenomenon requires scrutiny of both the group mean and its natural variation (79, 80). (This is clearly illustrated by the HPC maps of Fig. 4A, where group means do not describe any individual particularly well.) More specifically, the degree of explanatory power that can be ascribed to a coupling phenomenon occurring in either 25 or 75% of individuals is decidedly different, even though group statistics might be similar. Analogously, observing a coupling phenomenon in either 1 or 9 out of 10 recording sites places different constraints on the functional role this coupling may play. While these additional layers of complexity present novel challenges, these obstacles can only be overcome by being aware of their existence in the first place.

Conclusion

While hippocampal sleep oscillations are thought to play a central role in systems memory consolidation, little is known about how coordinated rhythmic brain activity is organized in human HPC. Our findings reveal that such oscillatory interactions occur in remarkably diverse ways. While the causes of this heterogeneity remain to be determined, we believe that characterizing both the commonalities and variability of the sleep oscillatory landscape will contribute to a deeper understanding of sleep's role in memory consolidation, and the function of sleep more broadly.

Methods

Participants

We analyzed archival electrophysiological sleep data in a sample of 12 (6 male) patients suffering from pharmaco-resistant epilepsy (age: 36.5 ± 14.5 yrs, range: 22–62). This sample overlaps with ones presented previously (21, 81). Patients had been epileptic for 23.4 ± 12.0 yrs (range: 10–49) and were receiving anticonvulsive medication at the moment of recording. All patients gave informed consent, the study was conducted according to the Declaration of Helsinki, and was approved by the ethics committee of the Medical Faculty of the University of Bonn.

<u>Data acquisition</u>

Electrophysiological monitoring was performed with a combination of depth and subdural strip/grid electrodes placed according to clinical criteria. HPC depth electrodes (AD-Tech, Racine, WI, USA) containing 8–10 cylindrical platinum-iridium contacts (length: 1.6 mm; diameter: 1.3 mm diameter; center-to-center inter-contact distance: 4.5 mm) were stereotactically implanted. Implantations were done either bilaterally (n=9) or unilaterally (n=3), and either along the longitudinal HPC axis via the occipital lobe (n=11) or along a medial-lateral axis via temporal cortex (n=1).

Pre- and post-implantation 3D T1-weighted magnetic resonance image (MRI) scans were used to determine electrode locations. Pre-operative T1 (resolution = 0.8x0.8x0.8 mm^3, TR = 1,660 ms, TE = 2.54 ms, flip angle = 9°) was acquired using a 3.0 Tesla Magnetom Trio (Siemens Healthineers, Erlangen, Germany) with a 32-channel-coil. Post-operative T1 (resolution = 1x1x1 mm^3, TR = 11.09 ms, TE = 5.02 ms, flip angle = 8°) was conducted using an Achieva 3.0 Tesla Tx system (Philips Healthcare, Best, The Netherlands). Preprocessing and analyses of T1 volumes was done using FMRIB's Software Library 5.0 (FSL) (82). Brain extractions (83) were performed and followed by a bias-field correction (84). Post-operative volumes were linearly registered to the pre-operative volumes. Pre-operative volumes were normalized to the Montreal Neurological Institute template (MNI 152, 1 mm resolution) by means of a non-linear registration (85). The resulting normalization matrices were then applied to the registered post-operative volumes to normalize them to the MNI template. This way, the normalization process was not affected by electrode image artifacts. Anatomical labels of the electrodes were determined by an experienced physician (TR) based on the subject-specific co-registered T1 volumes. MNI coordinates were extracted from the normalized volumes.

We included a total of 61 HPC contacts (5.1 ± 2.3 per patient, range: 1–8). Only electrodes from the non-pathological side were considered (6 left), and electrode contacts were included only when they could be localized to HPC gray matter, the HPC gray/white matter border, or the border between HPC gray matter and the lateral ventricle. A further subdivision was made into anterior (total: 23, range: 0–4), middle (22, 0–3), and posterior (16, 0–3) thirds of the HPC. Note that not every patient had contacts inside each hippocampal subdivision (see Supp. Table 1 for an overview of anatomical labels and MNI coordinates for all electrodes). For sleep recordings, additional signals were recorded from the scalp (Cz, C3, C4, Oz, A1, A2; plus T5 and T6 in 10 patients), the outer canthi of the eyes for electrooculography (EOG), and chin for electromyography (EMG). All signals were sampled at 1 kHz (Stellate GmbH, Munich, Germany) with hardware low- and high-pass filters at 0.01 and 300 Hz respectively, using an average-mastoid reference.

<u>Sleep scoring</u>

Offline sleep scoring was done in 20 s epochs based on scalp EEG, EOG, and EMG signals in accordance with Rechtschaffen and Kales criteria (86). Stages S3 and S4 were combined into a single N3 stage following AASM criteria (87).

Preprocessing and artifact rejection

All data processing and analysis was performed in Matlab (the Mathworks, Natick, MA), using custom routines and EEGLAB functionality (88). For each patient, we included all epochs scored as sleep or wake and selected channels of interest (including channels not reported in the present paper). Data were filtered with a repeated notch filter (50 Hz and harmonics up to 300 Hz), and a 0.3 Hz high-pass. In a first round of artifact rejection, we visually identified and marked time segments during which any channel displayed obvious artifacts (e.g., excessive amplitudes, amplifier saturation). Next, remaining time segments were used to determine channel-specific thresholds for detection of epileptogenic artifacts. Specifically, each time point was converted to a z-score based on the mean and SD of i) the signal gradient (amplitude difference between adjacent time points), and ii) the signal amplitude after applying a 250 Hz high-pass filter. Artifacts were marked from 0.5 s before to 0.5 s after contiguous time points where the z-score of either of these measures exceeded a critical value. All records were visually reviewed to confirm that threshold settings resulted in appropriate artifact detection. We found that a z threshold of 6 offered an optimal balance between detection of epileptogenic spikes and unwarranted rejection of clean data. Next, we included only continuous data segments that were free of either type of artifact across all included channels for a minimum of 3 s. That is, if a single channel displayed an artifact, that data segment would not be included for any of the channels, further reducing the likelihood that data is contaminated by pathological activity. Data within 0.5 s from non-adjacent epochs was also excluded. All told, this approach removed 54.6 ± 16.6% of the data, leaving a total of 86.7 ± 41.3 (N2), 26.3 ± 25.5 (N3), and 55.3 ± 35.8 min (REM) of clean data. Remaining segments ("trials") were split if they spanned multiple sleep/wake stages and if the resulting new trials also exceeded a duration of 3 s. Each trial was assigned a stage label based on its midpoint. Average number and duration of included trials was 763 ± 383 and 8.3 ± 7.4 s (N2), 241 ± 210 and 6.6 ± 3.7 s (N3), and 272 ± 172 and 16.1 ± 12.7 s (REM).

Spectral analysis

For each trial and channel, we estimated power spectral density using Welch's method with 3 s windows and 80% overlap (0.244 Hz resolution). Mean stage spectra were determined with a weighted average approach using trial durations as weights. Next, we removed the spectra's 1/f component to better emphasize narrowband spectral peaks. To this end, we first interpolated the notch-filtered region (50, 100, 150, and 200 Hz, ± 5 Hz) of each spectrum (Modified Akima cubic Hermite algorithm). Then, we took the N2 spectrum of Cz and the channel-averaged N2 spectrum of the HPC, and fit each according to af^b using log-log least squares regression (89, 90). Fitting range was restricted to the 4–175 Hz range to avoid the often observed flattening of the spectrum below ~4 Hz (see Fig. 1AB insets) and the ~200 Hz interpolated data. Then, for each channel and stage, the N2 model fit was subtracted from the observed spectrum. We applied the N2 fit to all stages rather than using stage-specific fits to enable direct stage comparisons. Similarly, for HPC channels we subtracted the channel-averaged spectrum from each channel to allow channel comparisons. Adjusted spectra were resampled to log space and smoothed three times with a moving average window of length 5.

Time-frequency decomposition

In order to assess sleep oscillatory dynamics across several orders of magnitude, we decomposed the multichannel data with a family of complex Morlet wavelets. Each trial was extended with 5 s on either side to minimize edge artifacts. Wavelets were defined in terms of desired temporal resolution according to $e^{i2\pi t f} e^{-4\ln(2)t^2/h^2}$ (91), where i is the imaginary operator, t is time in seconds, f is frequency (50 logarithmically spaced frequencies between 0.5 and 200 Hz), and h is temporal resolution (full-width at half-maximum; FWHM) in seconds. We set h to be logarithmically spaced between 3 s (at 0.5 Hz) and 0.025 s (at 200 Hz), resulting in FWHM spectral resolutions of 0.3 and 35 Hz, respectively. Trial padding was trimmed from the convolution result and data were downsampled by a factor four to reduce the amount of data. We assessed PAC using a surrogate approach (see below). To make surrogate distributions independent of variable numbers and durations of trials, we first concatenated the convolution result of all trials of a given sleep stage, and then segmented them into 1 min fragments (discarding the final, incomplete segment). Raw PAC and surrogate distributions were then determined separately for each 1 min segment. Segment length was chosen to obtain stable PAC estimates that are not overly affected by cycle-by-cycle variations in coupling strength and/or phase.

Cross-frequency coupling

PAC was determined between all pairs of modulating frequency *f1* and modulated frequency *f2*, where *f2>2*f1*. We employed an adaptation of the mean vector length method (92) that adjusts for possible bias stemming from non-sinusoidal shapes of *f1* (93). Specifically, complex-valued debiased phase-amplitude coupling (dPAC) was calculated as:

$$\text{dPAC} = \frac{1}{n}\sum_{t=1}^{n}\left(amp_{f2}(t) * \left(e^{i\varphi_{f1}(t)} - B\right)\right)$$

where *i* is the imaginary operator, *t* is time, *amp*$_{f2}$*(t)* is the magnitude of the convolution result, or amplitude, of *f2*, φ_{f1} *(t)* is the phase of *f1*, and *B* is the mean phase bias:

$$B = \frac{1}{n}\Sigma_{t=1}^{n} e^{i\varphi_{f1}(t)}$$

Raw coupling strength (i.e., the degree to which the *f2* amplitude is non-uniformly distributed over *f1* phases) was defined as the magnitude (i.e., length) of the mean complex vector. For every 1 min segment, channel, and frequency pair, we constructed a surrogate distribution of coupling strengths by repeatedly (n = 100) time shifting the *f1* phase time series with respect to the *f2* amplitude time series by a random amount between 1 and 59 s, and recalculating the mean vector length for each iteration. Note that time shifting is a more conservative approach than fully scrambling time series, which may result in spurious effects (94). We then z-scored the observed coupling strength with respect to this null distribution of coupling

strength values. Thus, the z-scored measure (dPAC$_Z$) indicates how far, in terms of standard deviations, the observed coupling estimate is removed from the average coupling estimate under the null hypothesis of no coupling. All reported analyses of coupling strength are based on dPAC$_Z$, but for simplicity we refer to this measure as PAC. Coupling phase (i.e., the *f1* phase where *f2* amplitude is maximal) was defined as the phase angle of the mean complex vector.

Statistics

Statistical analyses were performed at three hierarchical levels of inference: the channel, the subject, and the group level. Specifically, power was analyzed at the subject and group (but not channel) levels, whereas PAC was analyzed at the channel and group (but not subject) levels. We used cluster-based permutation statistics (95) to determine the presence of PAC, and to compare power and PAC between sleep stages or brain regions. Briefly, a t-test was performed at each frequency bin (power) or frequency-frequency bin (PAC), and bins meeting a threshold ($P=0.1$) were allowed to form clusters based on spectral adjacency. T-values within each cluster were summed as a measure of effect size (cluster statistic). Next, condition labels were repeatedly shuffled and the largest cluster statistic at each iteration was stored. Number of iterations depended on the number of possible permutations: If the number of possible combinations was over 1,000, a random shuffle was performed for each of 1,000 iterations (i.e., Monte Carlo sampling), otherwise each unique combination was sampled exactly once (i.e., permutation sampling). Observed cluster statistics were then compared to the surrogate distribution of cluster statistics. Clusters were considered significant at $P<0.05$ for one-sided tests (presence of coupling), and at $P<0.025$ for two-sided tests (condition comparisons).

At the channel level (PAC only), the presence of coupling (e.g., Fig. 2A) was determined by comparing the distribution of dPAC$_Z$ values across segments to zero (paired t-test). Channel and stage differences (e.g., Fig. 2CG) were assessed using paired and unpaired t-tests, respectively. At the subject level (power only), trial-averaged spectra were compared between sleep stages across channels (e.g., Fig. 1DG, paired t-test). At the group level, power/dPAC$_Z$ values were first averaged across trials/segments, and, for HPC, relevant channels (i.e., anterior, middle, posterior, or all channels). The distribution of these values was compared to zero (e.g., Fig. 4A; paired t-tests), and contrasted between brain regions and stages (e.g., Fig. 1A, 4BC; paired t-tests).

Since our objective was to offer a comprehensive characterization of the oscillatory structure of human HPC, this study should be viewed as a hypothesis-generating (or exploratory), rather than a hypothesis-testing (or confirmatory) study. Consequently, while we employed rigorous permutation-based approaches for every analysis, we did not further adjust for multiple testing (e.g., no correction across all panels of Fig. 3AB).

Density-based clustering

To obtain a qualitative picture of how many individuals showed PAC for each frequency pair we employed the *density-based spatial clustering of applications with*

noise (DBSCAN) algorithm (63), an unsupervised clustering method that aggregates nearby points without requiring an *a priori* specification of the number of clusters. In detail, for every individual, sleep stage, and channel, we extracted the frequency-frequency coordinates where $dPAC_Z$ was at a local maximum. Note that local maxima were generally of larger magnitude for NREM than REM, allowing for the inclusion of weaker effects for REM. Local maxima carried forward to the clustering approach were selected based on both an adaptive and a fixed threshold approach. For the adaptive approach, maxima were rank ordered and thresholded according to: $prop_{inc}$ = 0.3 x N_{chan}, where $prop_{inc}$ is the proportion of local maxima that is included, and N_{chan} is the number of channels within a region. N_{chan} always equaled 1 at the scalp, but ranged from 1 to 8 for HPC. Hence, the adjustment for N_{chan} ensures we included approximately similar numbers of $dPAC_Z$ values (and corresponding frequency coordinates) for each subject, but also that each HPC channel was allowed to contribute its strongest effects. For the fixed approach, only local maxima with a $dPAC_Z$ score above 2 were included. Importantly, to allow large local maxima that were only present on a single channel to survive the clustering approach, frequency-frequency points were duplicated according to their $dPAC_Z$ value. Specifically, above-threshold $dPAC_Z$ scores were divided into 5 equally sized bins (ranging from the threshold value to the largest $dPAC_Z$ score), and each frequency-frequency point was multiplied n times according to the bin number containing its $dPAC_Z$ score. Then, for each individual, sleep stage, and region, resulting frequency coordinates were submitted to the DBSCAN algorithm. Frequency coordinates were assigned to the same cluster if they were a maximum distance of 1.5 frequency-frequency bins away, and clusters were required to contain a minimum of 2 points. Frequency coordinates not meeting these criteria were labeled as noise and discarded. We extracted each cluster's center of gravity as the $dPAC_Z$-weighted average of the frequency coordinates of that cluster's members, and assigned it to the closest discrete frequency-frequency bin. Examples of single-subject cluster assignment are presented in Supp. Fig. 4. A binary frequency-frequency image was generated for every individual, sleep stage, and region, with ones at cluster centers and zeros elsewhere. Binary images were smoothed with a 2D Gaussian filter of standard deviation 0.5, rounded upwards so as to consist of only zeros and ones, and summed across subjects to generate Fig. 5.

Competing interests

The authors declare that no financial or non-financial competing interests exist.

Funding information

This work was supported by the German Research Foundation (FE366/9-1 to JF) and a Wellcome Trust/Royal Society Sir Henry Dale Fellowship (107672/Z/15/Z to BPS).

References

1. Squire LR, Stark CEL, Clark RE (2004) The Medial Temporal Lobe. *Annu Rev Neurosci* 27(1):279–306.

2. Tulving E, Markowitsch HJ (1998) Episodic and declarative memory: Role of the hippocampus. *Hippocampus* 8(3):198–204.
3. Buzsáki G (1996) The Hippocampo-Neocortical Dialogue. *Cereb Cortex* 6(2):81–92.
4. Frankland PW, Bontempi B (2005) The organization of recent and remote memories. *Nat Rev Neurosci* 6(2):119–130.
5. Marr D (1971) Simple memory: a theory for archicortex. *Philos Trans R Soc Lond B Biol Sci* 262(841):23–81.
6. Zola-Morgan S, Squire L (1990) The primate hippocampal formation: evidence for a time-limited role in memory storage. *Science* 250(4978):288–290.
7. Ellenbogen JM, Hulbert JC, Stickgold R, Dinges DF, Thompson-Schill SL (2006) Interfering with Theories of Sleep and Memory: Sleep, Declarative Memory, and Associative Interference. *Curr Biol* 16(13):1290–1294.
8. Jenkins JG, Dallenbach KM (1924) Obliviscence during Sleep and Waking. *Am J Psychol* 35(4):605–612.
9. Walker MP, Brakefield T, Morgan A, Hobson JA, Stickgold R (2002) Practice with Sleep Makes Perfect: Sleep-Dependent Motor Skill Learning. *Neuron* 35:205–211.
10. Rasch B, Born J (2013) About Sleep's Role in Memory. *Physiol Rev* 93(2):681–766.
11. Stickgold R, Walker MP (2013) Sleep-dependent memory triage: evolving generalization through selective processing. *Nat Neurosci* 16(2):139–145.
12. Axmacher N, Elger CE, Fell J (2008) Ripples in the medial temporal lobe are relevant for human memory consolidation. *Brain* 131(7):1806–1817.
13. Cairney SA, Guttesen A á V, El Marj N, Staresina BP (2018) Memory Consolidation Is Linked to Spindle-Mediated Information Processing during Sleep. *Curr Biol* 28(6):948-954.e4.
14. Cox R, Hofman WF, de Boer M, Talamini LM (2014) Local sleep spindle modulations in relation to specific memory cues. *NeuroImage* 99:103–110.
15. Huber R, Ghilardi MF, Massimini M, Tononi G (2004) Local sleep and learning. *Nature* 430(6995):78–81.
16. Schönauer M, et al. (2017) Decoding material-specific memory reprocessing during sleep in humans. *Nat Commun* 8:15404.
17. Zhang H, Fell J, Axmacher N (2018) Electrophysiological mechanisms of human memory consolidation. *Nat Commun* 9(1). doi:10.1038/s41467-018-06553-y.
18. Andrillon T, et al. (2011) Sleep Spindles in Humans: Insights from Intracranial EEG and Unit Recordings. *J Neurosci* 31(49):17821–17834.
19. Frauscher B, et al. (2015) Interictal Hippocampal Spiking Influences the Occurrence of Hippocampal Sleep Spindles. *Sleep* 38(12):1927–1933.
20. Nir Y, et al. (2011) Regional Slow Waves and Spindles in Human Sleep. *Neuron* 70(1):153–169.
21. Staresina BP, et al. (2015) Hierarchical nesting of slow oscillations, spindles and ripples in the human hippocampus during sleep. *Nat Neurosci* 18(11):1679–1686.
22. Aru J, et al. (2015) Untangling cross-frequency coupling in neuroscience. *Curr Opin Neurobiol* 31:51–61.

23. Canolty RT, Knight RT (2010) The functional role of cross-frequency coupling. *Trends Cogn Sci* 14(11):506–515.
24. Cox R, van Driel J, de Boer M, Talamini LM (2014) Slow Oscillations during Sleep Coordinate Interregional Communication in Cortical Networks. *J Neurosci* 34(50):16890–16901.
25. Cox R, Mylonas DS, Manoach DS, Stickgold R (2018) Large-Scale Structure and Individual Fingerprints of Locally Coupled Sleep Oscillations. *Sleep*. doi:10.1093/sleep/zsy175.
26. Mak-McCully RA, et al. (2017) Coordination of cortical and thalamic activity during non-REM sleep in humans. *Nat Commun* 8:15499.
27. Mölle M, Bergmann TO, Marshall L, Born J (2011) Fast and Slow Spindles during the Sleep Slow Oscillation: Disparate Coalescence and Engagement in Memory Processing. *Sleep* 34(10):1411–1421.
28. Klinzing JG, et al. (2016) Spindle activity phase-locked to sleep slow oscillations. *NeuroImage* 134:607–616.
29. Clemens Z, et al. (2007) Temporal coupling of parahippocampal ripples, sleep spindles and slow oscillations in humans. *Brain* 130(11):2868–2878.
30. Clemens Z, et al. (2011) Fine-tuned coupling between human parahippocampal ripples and sleep spindles: Ripple-spindle events in human sleep. *Eur J Neurosci* 33(3):511–520.
31. Latchoumane C-FV, Ngo H-VV, Born J, Shin H-S (2017) Thalamic Spindles Promote Memory Formation during Sleep through Triple Phase-Locking of Cortical, Thalamic, and Hippocampal Rhythms. *Neuron* 95(2):424-435.e6.
32. Niethard N, Ngo H-VV, Ehrlich I, Born J (2018) Cortical circuit activity underlying sleep slow oscillations and spindles. *Proc Natl Acad Sci*:201805517.
33. Born J, Rasch B, Gais S (2006) Sleep to Remember. *The Neuroscientist* 12(5):410–424.
34. Steriade M, Nuñez A, Amzica F (1993) Intracellular analysis of relations between the slow (< 1 Hz) neocortical oscillation and other sleep rhythms of the electroencephalogram. *J Neurosci* 13(8):3266–3283.
35. Amzica F, Steriade M (1998) Electrophysiological correlates of sleep delta waves. *Electroencephalogr Clin Neurophysiol* 107(2):69–83.
36. Achermann P, Borbély AA (1997) Low-frequency (< 1 Hz) oscillations in the human sleep electroencephalogram. *Neuroscience* 81(1):213–222.
37. Benoit O, Daurat A, Prado J (2000) Slow (0.7–2 Hz) and fast (2–4 Hz) delta components are differently correlated to theta, alpha and beta frequency bands during NREM sleep. *Clin Neurophysiol* 111(12):2103–2106.
38. Buzsáki G (1986) Hippocampal Sharp Waves: Their Origin and Significance. *Brain Res* 398:242–252.
39. Oliva A, Fernández-Ruiz A, Fermino de Oliveira E, Buzsáki G (2018) Origin of Gamma Frequency Power during Hippocampal Sharp-Wave Ripples. *Cell Rep* 25(7):1693-1700.e4.
40. Gonzalez CE, et al. (2018) Theta bursts precede, and spindles follow, cortical and thalamic downstates in human NREM sleep. *J Neurosci* 38:9989 –10001.
41. Schreiner T, Doeller CF, Jensen O, Rasch B, Staudigl T (2018) Theta phase coordinated memory reactivation reoccurs in a slow-oscillatory rhythm during NREM sleep. *Cell Rep* 25:296–301.

42. Funk CM, Honjoh S, Rodriguez AV, Cirelli C, Tononi G (2016) Local Slow Waves in Superficial Layers of Primary Cortical Areas during REM Sleep. *Curr Biol* 26(3):396–403.
43. Uchida S, Maehara T, Hirai N, Okubo Y, Shimizu H (2001) Cortical oscillations in human medial temporal lobe during wakefulness and all-night sleep. *Brain Res* 891(1–2):7–19.
44. Bódizs R, et al. (2001) Rhythmic hippocampal slow oscillation characterizes REM sleep in humans: Hippocampal RSA in Human REM Sleep. *Hippocampus* 11(6):747–753.
45. Cantero JL, et al. (2003) Sleep-Dependent Theta Oscillations in the Human Hippocampus and Neocortex. *J Neurosci*:7.
46. Clemens Z, et al. (2009) Phase coupling between rhythmic slow activity and gamma characterizes mesiotemporal rapid-eye-movement sleep in humans. *Neuroscience* 163(1):388–396.
47. Bonnici HM, et al. (2012) Detecting Representations of Recent and Remote Autobiographical Memories in vmPFC and Hippocampus. *J Neurosci* 32(47):16982–16991.
48. Dandolo LC, Schwabe L (2018) Time-dependent memory transformation along the hippocampal anterior-posterior axis. *Nat Commun* 9(1):1205.
49. Ludowig E, et al. (2008) Intracranially Recorded Memory-related Potentials Reveal Higher Posterior than Anterior Hippocampal Involvement in Verbal Encoding and Retrieval. *J Cogn Neurosci* 20(5):841–851.
50. Poppenk J, Moscovitch M (2011) A Hippocampal Marker of Recollection Memory Ability among Healthy Young Adults: Contributions of Posterior and Anterior Segments. *Neuron* 72(6):931–937.
51. Vyazovskiy VV, et al. (2011) Local sleep in awake rats. *Nature* 472(7344):443–447.
52. Massimini M, Huber R, Ferrarelli F, Hill S, Tononi G (2004) The sleep slow oscillation as a traveling wave. *J Neurosci* 24(31):6862–6870.
53. De Gennaro L, Ferrara M, Vecchio F, Curcio G, Bertini M (2005) An electroencephalographic fingerprint of human sleep. *NeuroImage* 26(1):114–122.
54. Cox R, Schapiro AC, Manoach DS, Stickgold R (2017) Individual Differences in Frequency and Topography of Slow and Fast Sleep Spindles. *Front Hum Neurosci* 11:433.
55. Rusterholz T, et al. (2018) Nature and Nurture: Brain Region-Specific Inheritance of Sleep Neurophysiology in Adolescence. *J Neurosci* 38(43):9275–9285.
56. Todorova R, Zugaro M (2018) Hippocampal ripples as a mode of communication with cortical and subcortical areas. *Hippocampus*. doi:10.1002/hipo.22997.
57. Timofeev I (2011) Neuronal plasticity and thalamocortical sleep and waking oscillations. *Progress in Brain Research* (Elsevier), pp 121–144.
58. Mizuseki K, Miyawaki H (2017) Hippocampal information processing across sleep/wake cycles. *Neurosci Res* 118:30–47.
59. Zeitlhofer J, et al. (1997) Topographic distribution of sleep spindles in young healthy subjects. *J Sleep Res* 6(3):149–155.

60. Durán E, Oyanedel CN, Niethard N, Inostroza M, Born J (2018) Sleep stage dynamics in neocortex and hippocampus. *Sleep* 41(6). doi:10.1093/sleep/zsy060.
61. Emrick JJ, Gross BA, Riley BT, Poe GR (2016) Different Simultaneous Sleep States in the Hippocampus and Neocortex. *Sleep* 39(12):2201–2209.
62. Benjamini Y, Hochberg Y (1995) Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *J R Stat Soc Ser B Methodol* 57(1):289–300.
63. Ester M, Kriegel H-P, Sander J, Xu X (1996) A Density-Based Algorithm for Discovering Clusters in Large Spatial Databases with Noise. *Proceedings of 2nd International Conference on Knowledge Discovery and Data Mining*, p 6.
64. Scheffzük C, et al. (2011) Selective Coupling between Theta Phase and Neocortical Fast Gamma Oscillations during REM-Sleep in Mice. *PLoS ONE* 6(12):e28489.
65. Brankačk J, et al. (2012) Distinct features of fast oscillations in phasic and tonic rapid eye movement sleep: *Cross-frequency coupling in REM sleep. J Sleep Res* 21(6):630–633.
66. Muthukumaraswamy SD (2013) High-frequency brain activity and muscle artifacts in MEG/EEG: a review and recommendations. *Front Hum Neurosci* 7. doi:10.3389/fnhum.2013.00138.
67. Averkin RG, Szemenyei V, Bordé S, Tamás G (2016) Identified Cellular Correlates of Neocortical Ripple and High-Gamma Oscillations during Spindles of Natural Sleep. *Neuron* 92(4):916–928.
68. Mooij AH, Raijmann RCMA, Jansen FE, Braun KPJ, Zijlmans M (2017) Physiological Ripples (± 100 Hz) in Spike-Free Scalp EEGs of Children With and Without Epilepsy. *Brain Topogr* 30(6):739–746.
69. Mooij AH, et al. (2018) Ripples in scalp EEGs of children: co-occurrence with sleep-specific transients and occurrence across sleep stages. *Sleep* 41(11). doi:10.1093/sleep/zsy169.
70. Montplaisir J, Leduc L, Laverdière M, Walsh J, Saint-Hilaire JM (1981) Sleep Spindles in the Human Hippocampus: Normal or Epileptic Activity? *Sleep* 4(4):423–428.
71. Nakabayashi T, et al. (2001) Absence of sleep spindles in human medial and basal temporal lobes. *Psychiatry Clin Neurosci* 55(1):57–65.
72. Isomura Y, et al. (2006) Integration and Segregation of Activity in Entorhinal-Hippocampal Subregions by Neocortical Slow Oscillations. *Neuron* 52(5):871–882.
73. Fanselow MS, Dong H-W (2010) Are the Dorsal and Ventral Hippocampus Functionally Distinct Structures? *Neuron* 65(1):7–19.
74. Strange BA, Witter MP, Lein ES, Moser EI (2014) Functional organization of the hippocampal longitudinal axis. *Nat Rev Neurosci* 15(10):655–669.
75. Ujma PP, et al. (2015) A comparison of two sleep spindle detection methods based on all night averages: individually adjusted vs. fixed frequencies. *Front Hum Neurosci* 9. doi:10.3389/fnhum.2015.00052.
76. Cole SR, Voytek B (2017) Brain Oscillations and the Importance of Waveform Shape. *Trends Cogn Sci* 21(2):137–149.
77. Tort ABL, Komorowski R, Eichenbaum H, Kopell N (2010) Measuring Phase-Amplitude Coupling Between Neuronal Oscillations of Different Frequencies. *J Neurophysiol* 104(2):1195–1210.

78. Sirota A, Csicsvari J, Buhl D, Buzsáki G (2003) Communication between neocortex and hippocampus during sleep in rodents. *Proc Natl Acad Sci* 100(4):2065–2069.
79. Kanai R, Rees G (2011) The structural basis of inter-individual differences in human behaviour and cognition. *Nat Rev Neurosci* 12(4):231–242.
80. Seghier ML, Price CJ (2018) Interpreting and Utilising Intersubject Variability in Brain Function. *Trends Cogn Sci* 22(6):517–530.
81. Wagner T, Axmacher N, Lehnertz K, Elger CE, Fell J (2010) Sleep-dependent directional coupling between human neocortex and hippocampus. *Cortex* 46(2):256–263.
82. Jenkinson M, Beckmann CF, Behrens TEJ, Woolrich MW, Smith SM (2012) FSL. *NeuroImage* 62(2):782–790.
83. Smith SM (2002) Fast robust automated brain extraction. *Hum Brain Mapp* 17(3):143–155.
84. Zhang Y, Brady M, Smith S (2001) Segmentation of brain MR images through a hidden Markov random field model and the expectation-maximization algorithm. *IEEE Trans Med Imaging* 20(1):45–57.
85. Jenkinson M, Smith S (2001) A global optimisation method for robust affine registration of brain images. *Med Image Anal* 5(2):143–156.
86. Rechtschaffen A, Kales A (1968) *A Manual of Standardized Terminology, Techniques and Scoring System for Sleep Stages of Human Subjects*.
87. Silber MH, et al. (2007) The visual scoring of sleep in adults. *J Clin Sleep Med* 3(2):121–131.
88. Delorme A, Makeig S (2004) EEGLAB: an open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. *J Neurosci Methods* 134(1):9–21.
89. Miller KJ, Sorensen LB, Ojemann JG, den Nijs M (2009) Power-Law Scaling in the Brain Surface Electric Potential. *PLoS Comput Biol* 5(12):e1000609.
90. Vijayan S, Lepage KQ, Kopell NJ, Cash SS (2017) Frontal beta-theta network during REM sleep. *eLife* 6. doi:10.7554/eLife.18894.
91. Cohen MX (2018) A better way to define and describe Morlet wavelets for time-frequency analysis. *bioRxiv*:14.
92. Canolty RT, et al. (2006) High gamma power is phase-locked to theta oscillations in human neocortex. *Science* 313(September):1626–1628.
93. van Driel J, Cox R, Cohen MX (2015) Phase-clustering bias in phase–amplitude cross-frequency coupling and its removal. *J Neurosci Methods* 254:60–72.
94. Scheffer-Teixeira R, Tort AB (2016) On cross-frequency phase-phase coupling between theta and gamma oscillations in the hippocampus. *eLife* 5. doi:10.7554/eLife.20515.
95. Maris E, Oostenveld R (2007) Nonparametric statistical testing of EEG- and MEG-data. *J Neurosci Methods* 164(1):177–190.

Supplementary Figures and Tables

Supplementary Figure 1. Comparison of scalp and hippocampal spectra. Error shading: standard error of the mean across patients. Horizontal color bars above plots indicate significant (P<0.05) regional differences (cluster-based permutation test). Filled color reflects region with greater power. Note that while scalp and HPC signals have different amplitudes and are recorded using electrodes with different properties, the slope removal approach facilitates direct comparison between these sites.

Supplementary Figure 2. Individual scalp and hippocampal spectra of all patients. Colored lines indicate single-patient scalp (left) and HPC spectra averaged across available channels (right). Black line indicates group mean.

Supplementary Figure 3. Comparison of anterior, middle, and posterior hippocampal spectra. Single-patient HPC spectra averaged across available channels per region, followed by averaging across patients. Note the different sample sizes for different comparisons due to variable electrode coverage. Error shading: standard error of the mean across patients. No significant regional differences emerged (cluster-based permutation test).

Supplementary Figure 4. Examples of density-based clustering. Every dot indicates a suprathreshold (adaptive approach) local maximum in coupling strength submitted to the DBSCAN algorithm. Colored dots belong to a cluster, with different colors for different clusters. (Color coding different for each panel.) Small open dots indicate cluster centers. Black dots indicate points not assigned to cluster.

Supplementary Figure 5. Group-level phase-amplitude coupling for hippocampal regions. Mean coupling strength across patients for anterior, middle, and posterior HPC. White outlines indicate clusters of significantly higher than chance coupling (P<0.05, cluster-based permutation test). Note different sample sizes for each region due to variable electrode coverage.

Supplementary Figure 6. Stage differences in phase-amplitude coupling for hippocampal regions. Mean difference in coupling strength between sleep stages for anterior, middle, and posterior HPC. White outlines indicate significant positive clusters (P<0.05, cluster-based permutation test; no negative clusters present). Note different sample sizes for each region due to variable electrode coverage.

patient/channel	region	MNI x	y	z
p1	RIGHT			
TR07	P/wm	32	-31	-3
p2	LEFT			
TL05	A	-24	-12	-26
TL06	A	-25	-16	-22
TL07	A	-24	-19	-18
TL08	M	-24	-22	-16
TL09	M	-24	-26	-11
p3	RIGHT			
TRb02	A	27	-9	-27
TRb03	A	29	-11	-20
TRb04	A	29	-16	-17
TRb05	M	28	-20	-13
TRb06	M/wm	30	-25	-9
TRb07	P/wm	30	-29	-5
p4	LEFT			
TL02	A	-26	-14	-21
TL03	A	-26	-17	-18
TL04	M	-26	-20	-16
TL05	M	-26	-24	-13
TL06	P/wm	-27	-28	-10
TL07	P/wm	-28	-32	-5
TL08	P/wm	-29	-35	-1
p5	RIGHT			
TR01	A	26	-12	-24
TR02	A	27	-15	-21
TR03	A	27	-18	-18
TR04	A	27	-22	-14
TR05	M	27	-24	-11
TR06	M	27	-29	-9
TR07	M	27	-33	-4
TR08	P	29	-35	2
p6	LEFT			
TL06	A	-26	-17	-19
TL07	M	-27	-21	-16
TL08	M	-27	-25	-14
TL09	P	-28	-28	-12
TL10	P	-29	-24	-8

patient/channel	region	MNI x	y	z
p7	RIGHT			
TR04	A	22	-16	-22
TR05	A	22	-20	-21
TR06	M	22	-24	-19
TR07	M	24	-27	-14
TR08	M	25	-31	-9
TR09	P	28	-35	-5
TR10	P	29	-39	-2
p8	RIGHT			
TR03	A	28	-8	-21
TR04	A	28	-11	-19
TR05	M/lv	28	-15	-17
TR06	M/lv	29	-19	-15
TR07	M/wm	30	-23	-12
TR08	P/wm	31	-29	-8
p9*	LEFT			
TLa01	A	-27	-13	-25
TLa02	A	-31	-15	-24
p10	LEFT			
TL02	A/lv	-23	-13	-17
TL03	M/lv	-23	-15	-14
p11	LEFT			
TL03	A	-33	-19	-17
TL04	M	-33	-23	-16
TL05	M	-32	-26	-14
TL06	P	-32	-30	-11
TL07	P	-33	-34	-8
p12	RIGHT			
TR04	A	23	-5	-40
TR05	A	26	-17	-34
TR06	M	27	-22	-30
TR07	M	27	-25	-25
TR08	P	27	-29	-19
TR09	P	26	-31	-14
TR10	P	28	-35	-9

Supplementary Table 1. Locations of included hippocampal contacts. Electrodes were included when located wholly within HPC gray matter, on the HPC gray/white matter (wm) border, or the HPC gray matter/lateral ventricle (lv) border. Contacts were classified as anterior (A), middle (M) or posterior (P) based on individual anatomy. MNI coordinates after warping to MNI template are also provided. * indicates patient with lateral implantation.

frequency pair	N2 scalp			N2 HPC			N3 scalp			N3 HPC		
	f1	f2	N	f1	f2	N	f1	f2	N	f1	f2	N
SO–delta				1.0–1.2	4.5–5.1	4	0.6–0.7	4.0	4			
							1.0	3.5–4.5	3			
							1.3–1.7	3.5	3			
SO–theta	0.6	4.5–5.8	4				1.0	7.4	5			
	1.3	5.8	4				1.3	7.4–8.3	5			
SO–theta/slow spindle	0.8	8.3–9.4	4									
	0.6	8.3–9.4	3									
	1.5	8.3–10.6	3									
SO–fast spindle	1.0	13.6–15.3	5				**1.0**	**15.3**	**9**			
SO–beta/gamma	0.9–1.2	36.1	3	0.9	75.2–85.0	4				1.3	28.2–40.8	4
				0.9–1.0	156.7–177.0	3				0.8	75.2	4
										0.9–1.2	19.6	3
										1.3–1.5	156.6	3
delta–spindle/beta				1.5–1.7	19.6	3	1.9	17.3	3	2.8–3.1	19.6–25.0	5
										1.7	28.2–36.1	4
delta–beta/gamma	1.9	40.8	3	**3.1**	**75.2**	**12**				**2.8**	**75.2**	**11**
				3.1	**46.1**	**7**				**1.5**	**85.0**	**8**
				1.5	75.2	5				**2.5**	**46.1**	**7**
				2.8–3.1	108.5	5				2.2–2.8	36.1–46.1	6
				2.8–3.1	138.6	5				3.1–3.6	138.6	6
										2.5	122.6–138.6	5
										4.5	75.2	4
delta/theta–spindle				4.0–5.1	15.3–19.6	4				4.5	15.3–19.6	5
theta–gamma	8.3	58.9–66.5	3									
slow spindle/alpha–gamma							10.6–12.0	138.6	3			
spindle–ripple				13.6	75.2–85.0	3				12.1	85.0	3
beta–gamma	**36.1**	**156.6–177.0**	**7**				**22.1**	**108.5**	**7**			
	22.1–25.0	108.5	6				**32.0**	**156.6**	**7**			
	19.6	122.6	6				17.3–19.6	66.5	4			
	22.1	138.6	6									

Supplementary Table 2. Prevalence of NREM coupling following density-based clustering (adaptive approach). Indicated is how many individuals (N) expressed coupling for each combination of a modulating frequency range (f1) with a modulated frequency range (f2). Only effects present in at least 3 individuals (25%) are shown. Effects present in more than half of the sample (>6) are indicated in bold.