Infection of *Plasmodiophora brassicae* changes the fungal endophyte community of tumourous

stem mustard roots as revealed by high-throughput sequencing and culture-dependent methods

Abstract

Diverse fungal endophytes live in plants and are shaped by some abiotic and biotic stresses. Plant disease

as particular biotic stress possibly gives an impact on the communities of fungal endophytes. In this study,

clubroot disease caused by an obligate biotroph protist, *Plasmodiophora brassicae*, was considered to

analyze its influence on the fungal endophyte community using an internal transcribed spacer (ITS)

through high-throughput sequencing and culture-dependent methods. The results show that the diversity

of the endophyte community in the healthy roots was much higher than the clubroots. Ascomycota was

the dominant group of endophytes (including *Phoma*, *Mortierella*, *Penicillium*, etc.) in the healthy roots

while *P. brassicae* was the dominant taxon in the clubroots. Hierarchical clustering, principal component

analysis (PCA) and principal coordinates analysis (PCoA) indicated significant differences between the

endophyte communities in the healthy roots and clubroots. Linear discriminant analysis effect size (LefSe)

analysis showed that the dominant genera could be regarded as potential biomarkers. The endophyte

community in the healthy roots had a more complex network compared with the clubroots. Also, many

plant pathogenic *Fusarium* were isolated from the clubroots by the culture-dependent method. The

outcome of this study illustrates that *P. brassicae* infection may change the fungal endophyte community

associated with the roots of tumourous stem mustard and facilitates the entry of soil pathogen into the

roots.

Introduction

Fungal endophytes have a close relationship with the host plants and they live in their tissues [1,2]. They provide many ecological and physiological advantages to their hosts, such as growth promotion [3], resistance to plant pathogens and adaptability to various abiotic stresses such as temperature, pH, and osmotic pressure as well as biotic stresses [4-6]. Similarly, the fungal endophyte communities are also affected by abiotic and biotic stresses [7]. Plant disease as particular biotic stress results in significant changes in the physiology of the plant. These changes and pathogen itself may affect the diversity and composition of the fungal endophyte inhabiting the plant tissues and their interactions with their host plant [8-10].

Clubroot is a severe disease of cruciferous crops such as cabbage and cauliflower, which is caused by the biotrophic *Plasmodiophora brassicae* Woronin [11]. The infected root cells undergo abnormal cell division and enlargement resulting in the formation of spindle-like, spherical, knobby or club-shaped swellings [12]. The root galls significantly alter the morphology, development, and physiology of the diseased plants [13]. Moreover, *P. brassicae* consumes carbohydrates from the viable cells in gall, and make gall into a sink for nutrient substances [14,15].

Tumourous stem mustard (*Brassica juncea* var. *tumida*) is an economical and nutritionally important vegetable crop widely grown in Fuling County, a district in Chongqing Municipality, China. Clubroot is one of the most harmful diseases of tumourous stem mustard causing substantial economic damage [16]. The diseased tumourous stem mustard shows small plants and swollen roots and finally dies due to root rot during the late plant growth stage in the field, although *P. brassicae* is an obligate parasite. Moreover, many pathogenic fungi such as *Fusarium* sp. are isolated from the clubroots, which indicates that the community of fungal endophyte may change during the infection of *P. brassicae*. However, the specific

alteration in the community of fungal endophyte is still unclear.

In this context, the objectives of this study are, (1) to examine the diversity and composition of the

fungal endophyte community in the roots of tumourous stem mustard (2) to demonstrate the changes in

the community of fungal endophyte in the clubroots of tumourous stem mustard caused by *P. brassicae*

compared to healthy roots .

Materials and methods

Sample collection

P. brassicae-infected tumourous stem mustard roots were obtained from three fields (February 2, 2017)

in Fuling (29.21° N, 106.56° E) where clubroot disease was observed during the last 20 years. Roots

were collected from the plants at the harvest-stage. The roots were classified as either R (healthy roots,

no clubroot symptoms) or C (diseased roots, swollen clubroot symptoms). From one field, 30 plants

including 15 R and 15 C samples were randomly selected and formed as one group. By this, three groups

contained 90 plants from three fields and were referred to R1, C1, R2, C2, R3, and C3. The roots were

washed with tap water to remove the soil particles. The healthy roots with a diameter of 0.5 cm from

undiseased plants and clubroot galls with a diameter of 1 cm from diseased plants were cut off, and then

sterilized by 70 % (v/v) ethanol for 40 s, followed by 4 % (w/v) sodium hypochlorite for 60 s, and finally

washed three times using sterile distilled water. The surface sterilized healthy roots and galls peeled with

a sterilized razor were divided into three portions. One portion was used to extract genomic DNA by

cetyltrimethylammonium ammonium bromide (CTAB) DNA extraction method. The concentration and

purity of DNA were monitored on 1% agarose gel. One portion was made into paraffin section, dyed

with sarranine and observed under an optical microscope. Another portion was used to isolate endophytic

fungi by the culture-dependent method.

Isolation of endophytic fungi and its taxonomic identification

Three types of media, i.e. potato dextrose agar (BD Difco), rose bengal medium (BD Difco), and Czapek

medium (BD Difco) were used for the isolation of endophytic fungi. The surface sterilized healthy roots,

or clubroot samples were cut into tissue blocks of 5 mm x 5 mm, and were planted on a medium at 25

◦C in the dark. When mycelia appeared, they were transferred into another medium for purification. A

total of 156 fungal isolates were then identified based on the ITS sequence data. Mycelia of the fungal

isolates were ground with liquid nitrogen in a sterile mortar, and Genomic DNA from all the fungal

isolates was extracted using DNA extraction kit (TIANGEN Co. Ltd., Beijing, PR China). The internal

transcribed spacer (ITS) region was amplified with universal primers (ITS1 and ITS4) [17]. PCR mixture

contained 12 μL of 2× Taq PCR Mix (TIANGEN Co. Ltd., Beijing, PR China), 1 μL DNA template, 1

μL of each primer and 8 μL double distilled water. The PCR reaction conditions were as follows: initial

pre-heating at 94 ◦C for 3 min, 30 cycles of 94 ◦C for 30 s, 58 ◦C for 30 s, 72 ◦C for 30 s, and a final

extension at 72 ◦C for 10 min. PCR products were sequenced by 3730 sequencer (Majorbio Tech Co.

Ltd., Beijing, PR China). The ITS sequence was aligned in the NCBI database

(http://www.ncbi.nlm.nih.gov/) using the BLAST program. The highest hits (identity values higher than

97%) were regarded as the taxonomy of fungal isolates. The relative abundances were calculated

according to the taxonomy of fungal isolates.

High-throughput ITS Sequencing

ITS fragments were used as marker genes for high-throughput sequencing, and the primers of ITS

fragments target most fungal groups, as well as other eukaryotes. The ITS fragments were amplified by

PCR for barcoded pyrosequencing using the primers ITS1 and TS2 [18]. The employed PCR conditions

were: 95 °C for 2 min (one cycle), 95 °C for 30 s, 55 °C for 30 s, and 72 °C for 30 s (25 cycles), and

72 °C for 5 min (one cycle). The sequencing was performed using an Illumina MiSeq sequencer (Asbios

Technology Co., Ltd, China).

Processing of Bioinformatics and Data Analysis

For diversity analyses, the bioinformatics analysis pipeline was conducted on the free online platform,

Majorbio I–Sanger Cloud Platform (http://www.i-sanger.com). First, the raw sequences were processed

using the QIIME package (v1.8) [19]. The low-quality sequences, such as primer and barcode sequence

mismatch, and PCR-based or sequencing errors and chimeras, were removed. The taxonomy of each

operational taxonomic unit (OTU) representative sequence was performed using Unite (Release 7.2) and

ITS of *P. brassicae* under the threshold of 97% identity [20,21]. The Shannon index, Simpson index, and

rarefaction curves were calculated to evaluate the α-diversity. Relative abundance of endophyte was

calculated at phylum, genus and OTU levels. Analysis of variance (ANOVA) of the Shannon and

Simpson indices was obtained by SPSS 16.0 and the relative abundance was used to assess the differences

in the endophyte communities between the healthy roots and clubroots. For the evaluation of β-diversity,

hierarchical cluster dendrograms (Bray-Curtis distance dissimilarities) were constructed according to the

composition of OTU [22]. An unweighted UniFrac principal component analysis (PCA) and Principal

coordinates analysis (PCoA) were performed using R 3.1.1 statistical software [23].   The differentially

abundant genera between the healthy roots and clubroots were identified for finding the biomarker by

Discriminant Analysis Effect Size (LEfSe) software [24]. Co-occurrence network analysis was

conducted to reveal the relationship among the top 30 OTUs within the endophyte communities by

Networkx software [25]. Pearson's rank correlation coefficients between OTUs were calculated, and

only the correlations with r > 0.5 were selected to construct the network.

Results

Symptoms of Clubroot and Detection of *P. brassicae*

Healthy roots were not swollen, and the lateral roots appeared normal, whereas clubroots were swollen

and few lateral roots were observed (Fig. 1A). Abundant resting spores were found to be in the cells of

clubroots (Fig. 1B), but no resting spores were observed in the healthy roots (Fig. 1C).

Analysis of α-diversity

High-quality sequences of ITS were produced by the Miseq platform. The raw sequencing data for ITS

sequencing were deposited at the Sequence Read Archive (SRA, https://www.ncbi.nlm.nih.gov/sra)

under an accession number, SRP136514. The statistical formation of the sequences is shown in Table

S1. According to the taxonomy of the sequence and abundance (Table S2), the composition of the

endophyte communities was analyzed. Rarefaction curves analysis confirmed that the number of

observed OTUs increased asymptotically with an increase in the reads (Fig. 2A). Both a higher Shannon

index and Simpson index of the endophyte community were observed in the healthy roots than that in

the clubroots (Fig. 2B), suggesting a higher diversity of the endophyte community in the healthy roots.

Ascomycota was the dominant taxon (a relative abundance ranging from 76.3 to 78.5%) of the

endophyte community in the healthy roots followed by Zygomycota (RA, 15.5 to 19.9%), while

Cercozoa (RA, 92.0 to 94.3%) was the most dominant taxon of the endophyte community in the clubroots

followed by Ascomycota (RA, 5.6 to 7.6%) (Fig. 3A). The abundance of Basidiomycota and unclassified

endophytic fungi was higher in the healthy roots compared to clubroots.

At the genus level, a total of 40 and 27 genera were detected in the healthy roots and clubroots,

respectively. In the healthy roots, *Phoma* was the dominant genus with an RA of 19.15%, followed by

Mortierella (17.09%), *Penicillium* (11.49%), *Pseudallescheria* (10.75%), *Tetracladium* (9.98%), and

Fusarium (7.88%) (Fig. 3B). In the clubroots, *Plasmodiophora* was the dominant taxon (93.28 %)

followed by *Fusarium* (5.52 %) (Fig. 3B). Circos figure confirmed that *Plasmodiophora* was the most

abundant and was found only in the clubroots, whereas *Phoma*, *Mortierella*, and *Penicillium* were

predominant in the healthy roots (Fig. 3C).

A total of 66 and 40 OTUs were observed in the healthy roots and clubroots, respectively. Heatmap

showed that OTU268 was dominant in the endophyte community associated with clubroots, whereas

OTU106, OTU153, OTU192, OTU197, OTU114, OTU126, and OTU185 dominated the healthy roots

(Fig, S1).

Analysis of β-diversity

Hierarchical clustering analysis based on Bray-Curtis distance dissimilarities revealed that the

endophyte communities in the healthy roots and clubroots clustered in the two branches (Fig. 4A).

UniFrac-weighted PCA based on the composition of OTU showed variations between the healthy roots

and clubroots with the first two axes indicating 81.08 and 11.71 % of the total variation (Fig. 4B). PCoA

plot also clearly demonstrated similar results of PCA with the first two axes showing 99.34 and 0.62%

of the total variation (Fig. 4C). The endophyte community in the healthy roots was clustered on the right

side of the PCA and PCoA plot while the endophyte community in the clubroots was clustered on the

left side, indicating a clear separation between the endophyte community in R and C samples.

From the results of LEfSe, a significantly different taxa were found out between the two communities.

At the genus level, *Phoma*, *Mortierella*, *Penicillium*, unclassified Pleosporales, etc. were enriched in the

healthy root samples (Fig. 5A) and abundant *Plasmodiophora* in the clubroot samples possessed high

values of linear discriminant analysis (LDA) (Fig. 5B). These taxa with different abundance can be

regarded as potential biomarkers (LDA>3, P<0.05). Furthermore, 15 most abundant genera of the two

endophyte communities were compared by Student's t-test, where significant differences were observed

(Fig. S2). For instance, *Plasmodiophora* was significantly more abundant in the clubroots while all other

genera except *Fusarium* were significantly more abundant in the healthy roots.

Network Analysis

The endophyte community in the healthy roots had two centers with a complex network structure (28

nodes and 336 edges) (Fig. 6A) while the endophyte community in the clubroots had three centers with

a less complex network structure (18 nodes and 161edges) (Fig. 6B). Most of the nodes belong to

Ascomycota in both the healthy roots and clubroots networks and had the highest number of correlations

with other taxa. The node of the Cercozoa (i.e., *Plasmodiophora*) only appeared in the clubroot network

and had the highest percentage. When the relationship between *Plasmodiophora* and other fungi was

analyzed, it was found that 10 OTUs had a significant correlation with *Plasmodiophora* (3 OTUs had a

positive correlation, and 7 OTUs had negative correlation) (Table S3).

Culturable endophytic fungi

The same samples for Miseq sequencing were also used to isolate endophytic fungi by the culture-

dependent method. At the genus level, *Mortierella, Sarocladium, Phoma, Penicillium, Plectosphaerella,*

and *Fusarium* were obtained (Fig. 7A). *Fusarium* (a relative abundance of 81.4%) was the predominant

group in the clubroots, among which the following were present: *F. graminearum* (RA, 5.5 %), *F.*

oxysporum   (RA, 28.8 %), *F. solani*   (RA, 24.0 %),   *F. asiaticum*   (RA, 8.2 %), and *Fusarium* sp.

(RA 14.4 %). In the healthy roots, *Mortierella* was the main group with an RA of 36.2% followed

by *Phoma, Penicillium, Fusarium,* etc. (Fig. 7B).

Discussion

In this study, a high diversity of fungal endophyte in the healthy roots of tumourous stem mustard has

been found out using high-throughput sequencing. Our results agree with the early reports that the plants

can harbor a diversity of endophytic fungi in their healthy tissues, especially in the root system which is

considered to be the most suitable habitat for endophytic fungi [26,27]. Ascomycota is reported to be the

most common group of endophytic fungi [28], suggesting that they are suitable for the ecological niche

of plant tissue. Zhao also reported Ascomycota as the dominant group of endophytic fungi in the roots

of oilseed rape (*Brassica napus*) [29]. Also in this study, Ascomycota has been found to be the dominant

taxon in the healthy roots of tumourous stem mustard. In the phylum Ascomycota, *Phoma*, *Mortierella*,

Penicillium, and *Fusarium* were the common fungal genera in the healthy roots of tumourous stem

mustard. Arie et al. reported that *Phoma glomerata* produces epoxydon and inhibits clubroot disease,

demonstrating that *Phoma* in the healthy roots of tumourous stem mustard may perform a similar function

[30,31]. *Mortierella* was also detected in the roots of oilseed rape using high-throughput sequencing [29].

Melo et al. confirmed that the endophytic *M. alpina* in the moss Schistidium antarctici produces

antibiotics, antioxidant substances, and polyunsaturated fatty acids, which improve the environmental

suitability of the plant host [32]. Wani et al. also found that endophytic *M. alpina* promotes the

biosynthesis of Crocus apocarotenoid and enhances environmental stress tolerance [33]. These results

reflect that the endophytic *Mortierella* is a benefit to the plant, which indicates that *Mortierella* in

tumourous stem mustard is profitable to the host. Marinho et al. and Lin et al. both reported that

endophytic *Penicillium* produces active polyketides [34,35], reflecting that *Penicillium* in tumourous

stem mustard may produce similar substances.

Worldwide *Fusarium* is considered to be one of the most ubiquitous groups of fungi which can survive

in a wide range of plants, environments, and climates [36]. In this study, some OTUs identified as

Fusarium had a relatively high abundance in both the healthy roots and clubroots, reflecting the dominant

presence of *Fusarium* in tumorous stem mustard. Previous studies show that most of the *Fusarium*

species are nonpathogenic while only a small number are pathogenic [37]. Some OTU identified as

Fusarium in the healthy roots perhaps were nonpathogenic because of the absence of wilt symptoms.

OTU244 and OTU287 assigned as *F. solani*, a common soil-borne pathogen, were only and mainly found

in the clubroots and these populations may be pathogenic. Moreover, *F. solani* is the most fungal isolate

obtained by the culture-dependent method (Fig. 7A). Also, other pathogenic *Fusarium* such as *F.*

graminearum, *F. oxysporum,* and *F. asiaticum* also isolated from the clubroots rather than the healthy

roots, suggesting that *Plasmodiophora* infection many facilitate *Fusarium* to enter into the diseased roots.

The presence of a plenty of pathogenic *Fusarium* in the clubroots maybe the reason for the death of

tumourous stem mustard during the late plant growth stage in the field. In clubroots, *Plasmodiophora*

was the extremely abundant genus, although not belonging to fungi, amplified by PCR with ITS primers

for fungi and other eukaryotes. Zhao et al. also found that *Plasmodiophora* mainly enriched in the

clubroots of *Brassica napus* by high throughput sequencing using the same primers [29].

In general, the healthy plant tissues harbor a more diverse community of endophytic fungi than

diseased plant tissues [29,38]. Also, the endophyte community in the healthy roots of tumourous stem

mustard had significantly higher Shannon and Simpson indices compared with the clubroots. Also, a

higher number of genera and OTUs were obtained from the healthy roots than from the clubroots. The

endophyte communities in the healthy roots and clubroots also differed in the α- and β-diversity. The

marked difference in the composition of the endophyte community at the genus level between the healthy

roots and clubroots as revealed by LEfSe and Student's t-test showed that the dominant genera as

biomarkers caused differences in the two communities. These substantial discrimination in the dominant

genera was derived from the infection of *P. brassicae*. Pathogens may compete with the endophytic fungi

for space and nutrients in the same niche within the plant tissue [39]. We presume that a decrease in the

diversity of the fungal endophyte community in the clubroots of tumourous stem mustard has two causes.

First, the physiological changes and the formation of clubroots induced by *P. brassicae* may affect the

availability of nutrients for endophytes. Second, in clubroots, the resting spores of *P. brassicae* are

produced in the galls of diseased plants and may fill the gall cells. This may restrict the available space

for the endophytes. The phenomenon of a decrease in the diversity of endophytic fungi has been observed

in many diseased plant species [29,40,41].

In this study, some plant pathogenic fungi were detected from the clubroots, such as *F. solani*, but

not from the healthy roots of tumourous stem mustard. Similarly, Zhao et al. isolated many soil-borne

pathogenic fungi, including *Fusarium*, *Gibberella*, *Alternaria*, *Sclerotinia*, *Leptosphaeria,* and

Cylindrocarpon, from the oilseed rape roots infected with *P. brassicae* [29]. This multi-species

aggregation involving *P. brassicae* and other pathogenic fungi is similar to disease complexes containing

plant-parasitic nematodes and soil-borne pathogens [42]. The cuticle of clubroots is cracked, thereby

providing entry points for pathogens, even for opportunistic pathogens. Additionally, the cracked cuticle

aggravates the leaking of nutrients, such as amino acids and carbohydrates, into the rhizosphere soil

which may attract soil-borne pathogens and increase the risk of infection by these pathogens [43]. Root-

knot nematodes may form disease complexes with *Fusarium* species such as *F. oxysporum* and *F. solani*

[44]. In this study, the abundant presence of *F. solani* in the clubroots indicates that this pathogen and *P.*

brassicae may form a disease complex, which may make it more difficult to control the clubroot disease.

Further studies are necessary to clarify the potential existence of this disease complex. Besides, fungi

such as *C. cupreum*, *Phoma* sp. *Pochonia* sp. and *Myrmecridium schulzeri* were harvested and were

reported as biocontrol fungi [45-47], reflecting that these fungi may help the host to resist the infection

of *P. brassicae*.

Endophytic fungi in the plants must deal with many interactions and build a balanced network [48].

In this study, the endophyte community network in the healthy roots of tumourous stem mustard was

observed to be more complicated than in the clubroots, indicating a balanced network of interactions

among the fungal endophyte community in the healthy roots. In the clubroots, *P. brassicae* dominated

the endophyte community which was out of balance showing a weak network of interactions. In general,

endophytes are more diverse and abundant than the pathogens in a healthy plant microbiome, and *vice*

versa in a diseased plant microbiome [49].

Conclusions

In conclusion, the obtained data from this study showed that the fungal endophyte community in the

clubroots are markedly different from the healthy roots in terms of alpha and beta diversity, suggesting

that the infection of *P. brassicae* changes the fungal endophyte community in tumourous stem mustard

roots. Future work should involve identifying the pathogenicity of *F. oxysporum* and *F. solani* on

tumourous stem mustard. Moreover, some endophytic fungi with biocontrol activity against *P. brassicae*

could also be considered in the evaluation.

Figure legends

Fig 1. **Healthy roots and clubroots of tumourous stem mustard caused by the infection of *P.**

*brassicae*** (A) roots of a healthy plant (left) and a diseased plant (right) (B) resting spores (pots) in the

root cells of cluboorts coloured with fast green (C) the cell of healthy roots.

Fig 2. **Rarefaction curves (A) and diversity indices (B) of the endophyte communities associated with the healthy roots and clubroots of tumourous stem mustard infected with *P. brassicae*.** R, healthy roots. C, clubroots. ** differences at 0.01 level.

Fig 3. **Relative abundance (%) of the endophyte at phylum (a), genus (B) and Circos figure at the genus level (C) in the healthy roots and clubroots of tumourous stem mustard infected with *P. brassicae*.** R, healthy roots. C, clubroots.

Fig 4. **Hierarchical clustering analysis (a) UniFrac-weighted PCA (B) and PCoA (C) of the endophyte communities associated with the healthy roots and clubroots of tumourous stem mustard infected with *P. brassicae*.** R, healthy roots. C, clubroots.

Fig 5. **LefSe analysis (A) The cladogram diagram shows the taxas with marked differences in the two endophyte communities**. Red and green indicate different groups, with the classification of taxas at the level of phylum, class, order, family, and genus shown from inside to the outside. The red and green nodes in the phylogenetic tree represent taxas that play an important role in the two endophyte communities, respectively. Yellow nodes represent taxas with no significant difference. (B) Species with the significant difference that have an LDA score higher than the estimated value; the default score is 3.0. The length of the histogram represents LDA score; i.e., the degree of influence of taxas with a significant difference between different groups. R, healthy roots. C, clubroots.

Fig 6. **Co-occurrence network analysis of the two endophyte communities within the healthy roots**

and clubroots of tumourous stem mustard infected with *P. brassicae* (A) Healthy roots (B) Clubroots.

Each node represents taxa affiliated at the OTU level, and the size of the nodes represent an average

abundance of OTU. The lines represent the connections between each OTU. A red line indicates a

positive correlation. whereas a green line indicates a negative correlation.

Fig 7. The endophytic fungi isolated from the clubroots (A) and healthy roots (B) by the culture-

dependent method.

Supporting information

Table S1 Sequence information of each sample. R, healthy roots. C, clubroots.

Table S2 Taxonomy and distribution of the OTUs. Taxonomy at Phylum, Class, Order, Family,

Genus, Species, and OTU level. R, Healthy roots. C, Clubroots. The number in the table cell is the

number of sequences of each OTU.

Table S3 Pearson's correlation relationships between *Plasmodiophora* and OTUs showed in the

network.

Fig S1Heatmap of the 50 most abundant OTU from R and C samples. R, healthy roots; C, clubroots.

Fig S2 Student's t-test bar plot of the endophyte communities at the genus level in the healthy roots

and clubroots of tumourous stem mustard infected with *P. brassicae*. $p<0.05^*$, $p <0.01^{**}$,

$p<0.001^{***}$. R, healthy roots. C, clubroots.

Figure1

Figure2

Figure3

Figure4

Figure5

a

b

Basidiomycota
Ascomycota
Zygomycota
Cercozoa
Unclasified fungi

Figure6

a

Legend a (left)	
■ Mortierella	■ Sarocladium	■ Phoma
■ Penicillium	■ Plectosphaerella	■ Fusarium

81.4%
8.3%
2.1%
6.2%
1.4%
1.4%

5.5 %
14.4 %
8.2 %
24.0 %
28.8 %

■ Fusarium graminearum
■ Fusarium oxysporum
■ Fusarium solani
■ Fusarium asiaticum
■ Fusarium sp.

b

1.3%
7.9%
12.5%
13.8%
21.1%
3.9%
3.3%
36.2%

■ Penicillium   ■ Fusarium   ■ Mortierella   ■ Pochonia

■ Chaetomium   ■ Phoma   ■ Myrmecridium   ■ Pleosporales

Figure7