March 24, 2021

The Degree of Polymerization and Sulfation Patterns in Heparan Sulfate are

Critical Determinants of Cytomegalovirus Entry into Host Cells

Mohammad H. Hasan[1#], Dipanwita Mitra[1#], Rinkuben C. Parmar[1], Lauren A. Fassero, Quntao

Liang[2,6], Hong Qiu[3], Vaibhav Tiwari[4], Joshua S. Sharp[2], Lianchun Wang[5] and Ritesh Tandon[1]*

[1]Department of Microbiology and Immunology, University of Mississippi Medical Center, 2500

North State Street, Jackson, MS 39216, USA. [2]Department of Biomolecular Sciences, School of

Pharmacy, University of Mississippi, Oxford, MS 38677, USA. [3]Complex Carbohydrate

Research Center, University of Georgia, Athens, GA 30602. [4]Department of Microbiology and

Immunology, Midwestern University, Downers Grove, IL, 60515, USA. [5]Department of Molecular

Pharmacology and Physiology, University of South Florida, Tampa, FL 33612, USA. [6]College of

Biological Science and Engineering, University of Fuzhou, Fujian, 350108, China.

#Equal contributions

*Corresponding Author: Ritesh Tandon

Phone: 601-984-1705, Fax: 601-984-1708, Email: rtandon@umc.edu

Keywords: CMV, herpesviruses, heparan sulfate, sulfation

Running title: HS as CMV attachment receptors

ABSTRACT

Several enveloped viruses, including herpesviruses attach to host cells by initially interacting with cell surface heparan sulfate (HS) proteoglycans followed by specific coreceptor engagement which culminates in virus-host membrane fusion and virus entry. Interfering with HS-herpesvirus interactions has long been known to result in significant reduction in virus infectivity indicating that HS play important roles in initiating virus entry. In this study, we provide a series of evidence to prove that specific sulfations as well as the degree of polymerization (*dp*) of HS govern human cytomegalovirus (CMV) binding and infection. First, purified CMV extracellular virions preferentially bind to sulfated longer chain HS on a glycoarray compared to a variety of unsulfated glycosaminoglycans including unsulfated shorter chain HS. Second, the fraction of glycosaminoglycans (GAG) displaying higher *dp* and sulfation has a larger impact on CMV titers compared to other fractions. Third, cell lines deficient in specific glucosaminyl sulfotransferases produce significantly reduced CMV titers compared to wild-type cells and virus entry is compromised in these mutant cells. Finally, cells pretreated with a peptide that specifically binds sulfated-HS produce significantly reduced virus titers compared to the control peptide treated cells. Taken together, these results highlight the significance of HS chain length and sulfation patterns in CMV attachment and infectivity.

IMPORTANCE

Heparan sulfate (HS) is a linear polysaccharide found in all animal tissues. It binds to a variety of protein ligands, including cytokines, chemokines, growth factors and morphogens and regulates a wide range of biological activities, including developmental processes, angiogenesis, blood coagulation, and tumor metastasis. The molecular diversity in HS chains generates unique binding sites for specific ligands and can offer preferential binding for a specific virus over other viruses or cellular ligands. In the current study human cytomegalovirus (CMV) was found to bind preferentially to uniquely sulfated and polymerized HS. The HS mimics designed with these properties inhibited CMV infection. The results were corroborated by parallel studies in mutant mouse cells as well as using peptide inhibition. Combined together, the data suggests that CMV preferentially attaches to uniquely modified HS and thus this virus-host interaction is amenable to targeting by specifically designed HS mimics or peptides.

INTRODUCTION

The heparan sulfate (HS) proteoglycans are present on most cell types and function as cellular attachment receptors for medically important viruses, including human immunodeficiency virus (HIV), hepatitis-C virus (HCV), human papillomavirus (HPV), Dengue virus (DENV) and the recently emerged SARS-CoV-2 (1, 2) (3-6). In addition, virtually all human herpesviruses, with the possible exception of Epstein Barr virus, use HS as an initial co-receptor for entry (7). The interaction between cell surface HS and virus envelope is the initial event in the complex process of virus entry. A successful virus entry involves downstream co-receptor interactions ultimately leading to fusion between the virus envelope and the cell membrane (8).

Herpesviruses and several other enveloped viruses enter the host cells using two distinct pathways: 1) A pH-independent pathway which involves the fusion of the virus envelope with the plasma membrane; and 2) A pH-dependent pathway that involves endocytosis of the virus particle (9). In cells, where binding of virus to cell surface receptors induces endocytosis, the usual consequence is the acidification of the endosome, which ultimately triggers fusion between the virus envelope and endosomal membrane (7). Interestingly, human cytomegalovirus (HCMV) entry follows direct fusion at the cell surface in fibroblasts, while entry into other relevant cell types, such as endothelial cells, follows an endocytic route (10, 11). Different virus glycoprotein complexes are involved in each case; however, HS functions as the primary attachment receptor. Moreover, the presence of HS receptors are well documented in endosomal membranes and HS receptors likely play roles in intracellular virus trafficking (12-15).

The herpesvirus envelope is a lipid bilayer derived from host cell membranes in which

most cellular proteins have been displaced by viral membrane proteins. For HCMV, at

least twenty three different viral glycoproteins have been found to be associated with

purified virion preparations (16). For most herpesviruses, the conserved glycoprotein B

(gB) is required for virus entry and binds to cell surface molecules, including HS, which

is present not only as a constituent of cell surface proteoglycans but also as a

component of the extracellular matrix and basement membranes in organized tissues

(7, 17). HCMV gB binds to HS resulting in virus attachment (18) similar to its

counterparts in herpes simplex virus (HSV)-1 (17, 19) and varicella-zoster virus (VZV)

(20). Subsequently, HCMV gB binds to cellular protein receptors such as EGFR (21),

PDGFα (22), and integrins (23, 24), which culminates in virus entry. Treatment of cells

with soluble form of gB inhibits HCMV entry (25). HCMV binding and infection are

reduced by soluble heparin and HS, as well as in cells treated with heparinases or those

unable to produce HS (26). A better structural understanding of these inhibitions will

pave the way to design effective antivirals that are highly specific as well as more

effective.

The synthesis of HS is a complex process involving multiple specialized enzymes

and is initiated from a tetrasaccharide (GlcA-Gal-Gal-Xyl) that is attached to the core

protein (Fig. 1A). HS polymerase is responsible for building the polysaccharide

backbone with a repeating unit of -GlcA-GlcNAc- (Fig. 1B). The backbone is then

modified by N-deacetylase/N-sulfotransferase (NDST) responsible for N-deacetylation

and N-sulfation of selected glucosamine residues, C5-epimerase responsible for

epimerization of selected glucuronic moieties to iduronic acid, 2-*O*-sulfotransferase

(Hs2st; 2-*O*-ST) responsible for 2-*O*-sulfation of selected iduronic acid residues, 6-*O*-

sulfotransferase (H6st; 6-*O*-ST) for 6-*O*-sulfation and finally (but rarely) 3-*O*-

sulfotransferases (Hs3st; 3-*O*-ST) responsible for 3-*O*-sulfation (27, 28). The substrate

specificities of these biosynthetic enzymes dictate the structures of HS products,

including sulfation levels, the contents of L-iduronic acid (IdoA) units and the size of the

polysaccharides (27). The location of the sulfo groups and IdoA in turn play a crucial

role in determining the binding and functions of HS.

In the current study, we investigated the impact of specific sulfations as well as

degree of polymerization in terms of numbers of monosaccharide units (*dp*) in HS chain

on both human and mouse CMV infection and binding. Purified CMV extracellular

virions preferentially bound strongly to the longer sulfated HS chains but not to the

shorter unsulfated HS chains on a glycoarray. Glycosaminoglycans of different *dp* were

derivatized from enoxaparin (a low molecular weight heparin) and tested for their ability

to inhibit CMV infection in cell culture. The results show that longer glycan chains are

more efficient at reducing CMV titers in cells compared to shorter chain glycans. Also,

the cell lines defective in expression of various sulfotransferases showed significantly

reduced CMV entry and replication. Finally, cells pre-treated with peptides that bind HS

and 3-*O*-S HS (29-32) significantly reduced HCMV titers compared to a control peptide.

Overall, these results indicate that CMV binding to cell surface glycans is dependent on

branch length and sulfation pattern of HS.

MATERIALS AND METHODS

Preparation of glycosaminoglycans (GAGs) oligosaccharides. Glycosaminoglycans

of different *dp* were fractionated from enoxaparin (a low molecular weight heparin) by

Bio-Gel P-10 chromatography as previously described (33). Briefly, 15 mg/mL

enoxaparin sodium derived from porcine intestinal mucosa (Sanofi-Aventis U.S.,

Bridgewater, NJ) was applied to a Bio-Gel P-10 column (2.5X120 cm, Bio-Rad,

Hercules CA) and eluted with 0.2 M NH_4HCO3 at a flow rate of 14 ml/h. Elution of

oligosaccharides was monitored by absorbance at 232 nm. NH_4HCO3 was removed by

heating in oven at 50 °C for 24 h.

Preparation of the 6-*O* desulfated Arixtra with MTSTFA. A detailed procedure on the

preparation of 6-*O* desulfated Arixtra was published previously (34). Briefly, 4 mg of

Arixtra was added to 10 volumes (w/w) of N-Methy-N-(trimethylsilyl)-trifluoroacetamide

(MTSTFA, Sigma, ≥98.5%) and 100 volumes (v/w) of pyridine. The mixture was heated

at 100 ℃ for 30 min, then quickly cooled in an ice-bath, followed by extensive dialysis

and freeze-drying. The sample was resuspended in 50% acetonitrile/water at a

concentration of 30 µM for later LC-MS/MS analysis.

LC-MS/MS analysis. The 6-*O*-desulfated Arixtra (30 µM) was analyzed on a Thermo

Orbitrap Fusion Tribrid (Thermo Fisher Scientific) coupled with an Ultimate 3000 Nano

LC system (Dionex) using direct infusion. The flow rate was set to 1 µl/min. Mobile-

phase was 50% acetonitrile. Nanoelectrospray voltage was set to 2.0 kV in negative ion

mode. Full MS scan range was set to 200-2000 m/z at a resolution of 60,000, RF lens

was 6%, and the automatic gain control (AGC) target was set to 2.0 × 105. For the

MS/MS scans, the resolution was set to 50,000, the precursor isolation width was 3 m/z

units, and ions were fragmented by collision-induced dissociation (CID) at a normalized

collision energy of 80%.

Cells. Mouse embryonic fibroblasts (MEF) and human foreskin fibroblasts (HFF) were

cultured in Dulbecco's modified Eagle's medium (DMEM, Cellgro, Manassas, VA)

containing 4.5 g/ml glucose, 10% fetal bovine serum (SAFC, Lenexa, KS), 1 mM

sodium pyruvate, 2 mM L-glutamine, and 100 U/ml penicillin-streptomycin (Cellgro,

Manassas, VA) at 37 °C with 5% CO2. Mouse lung endothelial cells (WT, H3st1$^{-/-}$,

H3st4$^{-/-}$, H3st1/4-double-knockout, H6st1$^{-/-}$, H6st2$^{-/-}$, and H6st1/2 double-knockout) were

obtained from Wang laboratory at University of South Florida and maintained as

described earlier (35).

Virus. MCMV (strain K181) was grown in MEF cells, while HCMV (Towne strain) was

grown on HFF cells. Virus stock was prepared in 3X autoclaved milk, sonicated 3 times

and stored at −80 °C. 3X autoclaved milk is prepared from Carnation (Nestle) instant

nonfat dry milk powder. 10% milk was prepared in nano pure water, pH was adjusted to

7.0 and was autoclaved for 3 times. During infection, media was removed from the wells

of cell culture plates and appropriately diluted virus stock was absorbed onto the cells in

DMEM without serum. Cells were incubated for 1 hour with gentle shaking every 10

mins followed by washing 3X with PBS. Fresh complete medium was added and cells

were incubated until the end point. For extracellular virus (ECV) purification, HFF were

seeded in roller bottles, grown to confluency and infected with HCMV (Towne strain) at

MOI of 0.01. Two days after 100% cytopathic effect was observed, infected cell medium

was collected and centrifuged at low speed to pellet cellular debris, and the supernatant

was transferred to new tubes and centrifuged at 20,000 g for 1 hour to pellet the ECV.

This ECV pellet was re-suspended in phosphate buffer, sonicated to eliminate any

aggregates, loaded over 15-50% continuous sucrose gradients and centrifuged in a

SW-41 rotor at 39,000 RPM for 20 min. ECV bands were visualized in incandescent

light and harvested by puncturing the sides of the centrifuge tubes. These bands were

washed once with phosphate buffer, spun again and the final pellet resuspended in low

salt phosphate buffer. An aliquot of the sample was used for assessment of initial

quality of ECV by negative staining and transmission electron microscopy. Purified ECV

were shipped on ice to Z biotech (Aurora, CO) for glycoarray binding analysis.

Cell viability assay. HFF cells plated in 12 well tissue culture plates were grown to

confluency and pretreated for 1h with 10 μM concentration of candidate HS and then

infected with HCMV (Towne strain) at a multiplicity of infection (MOI) of 3.0 or mock-

infected in the presence of candidate HS . Five hundred μl of fresh complete medium

containing HS was added to the wells on day 3 and day 6. At the designated time

points, media was removed and cells were harvested by trypsinization. Cell viability was

determined using trypan blue exclusion on TC20 automated cell counter (BioRad

Laboratories, Hercules, CA) following manufacturer's protocol.

Virus titers. Infected or mock-infected samples were harvested within the medium at

the designated end points and stored at –80 °C before titration. In some experiments,

media and cells were separated by low-speed (< 1000 × g) centrifugation and viral

loads in supernatant and cells were quantified by titering on wild-type cells. Titers were

performed as described earlier (36) with some modifications. In brief, monolayers of

fibroblasts grown in 12 well plates and serial dilutions of sonicated samples were

absorbed onto them for 1 h, followed by 3X washing with PBS. Carboxymethylcellulose

(CMC) (Catalog No. 217274, EMD Millipore Corp., Billerica, MA) overlay with complete

DMEM media (1-part autoclaved CMC and 3 parts media) was added and cells were

incubated for 5 days. At the end point, overlay was removed and cells were washed 2X

with PBS. Infected monolayers were fixed in 100% methanol for 7 min, washed once

with PBS and stained with 1% crystal violet (Catalog No. C581-25, Fisher Chemicals,

Fair Lawn, NJ) for 15 min. Plates were finally washed with tap water, air dried and

plaques with clear zone were quantified.

Immunoblots. Mouse lung endothelial cells (WT, H3st1$^{-/-}$, H3st4$^{-/-}$, H3st1/4-double-

knockout, H6st1$^{-/-}$, H6st2$^{-/-}$, and H6st1/2 double-knockout) were infected with MCMV

(K181 strain) and an MOI of 3.0 and the whole cell lysates were harvested at 2 hours

post infection for analysis. The blot was probed with anti -IE1 antibody (catalog no. HR-

MCMV-12, Center for Proteomics, University of Rijeka, Croatia) and HRP-conjugated

goat anti mouse antibody (PI131444, Invitrogen) was used as the secondary antibody.

Glycoarrays. A dilution series of purified HCMV virions were incubated on two different

custom glycoarrays (Table 1 and Table 2, Z-Biotech) using established protocols (37)

and the arrays were analyzed to assess specific virus binding. Briefly, 10^5 to 10^8 pfu/ml

of purified virions were incubated for an hour on glycoarrays containing six replicates of

each glycosaminoglycan. After incubation, staining with primary antibody (mouse anti

gB (clone 2F12, Virusys Inc, Taneytown, MD) was done at 100 µg/ml and secondary

antibody (Goat anti mouse IgG AlexaFlour555) was done at 1µg/ml. Maximum strength

fluorescent signal was obtained for 10^8 pfu/ml concentration of the virus, therefore, only

this concentration is represented in the final data obtained for plotting the graphs.

Treatment of HFF cells with anti-HS and anti-3-*O*-S HS peptides. HFF cells were

pre-treated with the phage display derived peptides (1 mg/ml) generated against wild-

type HS (LRSRTKIIRIRH), and 3-*O*-S HS (MPRRRRIRRRQK) (29) that bind specifically

to HS and 3-*O*-S HS respectively, or left mock treated for 4 hours before the cells were

infected with β-galactosidase expressing CMV (ATCC) for 9 days. β-Galactosidase

assay was performed using X-gal (Sigma). The effect of entry- blocking activity of

peptide was examined by counting number of virus foci.

Statistics: Student's t-tests were conducted in Graphpad Prism comparing the means

of different groups (GraphPad Prism version 8.0.0, GraphPad Software, San Diego,

California USA, www.graphpad.com). Standard error of mean was plotted as error bars.

A p value of <0.05 was considered significant.

RESULTS

Purified HCMV extracellular virions preferentially bind to sulfated glycosaminoglycans with increased degree of polymerization.

First, we sought to establish the category of GAG that preferentially binds to purified HCMV virions. HCMV extracellular virions were purified as described above and incubated with custom glycoarrays containing increasing molecular weight species of hyaluronic acid, heparin, chondroitin sulfate, and dermatan sulfate (Table 1). HCMV binding to non-sulfated hyaluronic acid (HA10 to HA20 and HA93 polymer) was negligent but significant binding to all heparin species was detected with a trend of increased binding to heparins as their *dp* increased (Fig. 2). HCMV also bound to large size chondroitin sulfate D (GAG28, *dp20*), and dermatan sulfate oligosaccharides (GAG32- GAG34, *dp16-dp20*) but not to chondroitin sulfate AC (GAG17- GAG22). It is important to note that while the chondroitin sulfate A (CS-A) is sulfated at C4 of the GalNAc, and the chondroitin sulfate C (CS-C) is sulfated at the C6 of the GalNAc only, the chondroitin sulfate D is sulfated at C2 of the glucuronic acid as well as the C6 of the GalNAc sugar and hence has double the amount of sulfation compared to CS-A and CS-C. Dermatan sulfate (DS), formerly referred to as CS-B, is formed from the polymer backbone of chondroitin sulfate by the action of chondroitin-glucuronate C5 epimerase, which epimerizes individual d-glucuronic acid residues to l-iduronic acid. The binding affinity to DS was also size-dependent increasing from GAG32- GAG34 (*dp16- dp20*). Heparin (*dp30*) was the best HCMV binder in this assay. The virus preparation does not contain streptavidin label and thus positive control 1 (PC1) serves as a negative rather than a positive control.

On a second HS specific array (Table 2), HCMV showed strong binding to the

HS with longer monosaccharide chains (HS007 to HS024) and minimal binding to

unsulfated glycans (HS001-HS006) (Fig. 3). The maximum binding was observed for

HS014, HS015 and HS016, which are all 6-O-S 9-mers with moderate amount of

sulfation (1.3-1.8 sulfate group per disaccharide). Also, significant amount of binding

was observed for 2-O-S (HS17-HS19), 6-O-S/2-O-S (HS20-22) and 2-O-S/6-O-S/3-O-S

(HS23-24) HS that had high amount of sulfation (1.3-2.7 sulfate group per disaccharide)

and 6-8 disaccharides per chain. Overall the data from these experiments indicate that

the dp of HS as well as sulfation is important for HCMV binding.

The degree of polymerization of GAG chains impacts CMV infectivity.

Glycosaminoglycans of different dp were fractionated from enoxaparin (a low molecular

weight heparin). All of these GAGs are based on a HS backbone and differ in either dp

or degree/place of sulfation or both (Fig. 4, S1). These GAGs, along with heparin and

Arixtra (fondaparinux sodium), were first screened in a GFP-based preliminary virus

focus reduction assay using GFP tagged HCMV (Towne strain). The viral GFP

expression was most efficiently reduced by heparin salt (PIHSS; Heparin sodium salt

from porcine intestinal mucosa) whereas Arixtra, 6-O-desulfated Arixtra and enoxaparin

had little to no impact on GFP expression (Fig. 4). In general, enoxaparin derived GAGs

with higher dp were more efficient in reducing viral GFP compared to low dp derivatives.

To follow up on this primary GFP based screening, we performed viral titer assay using

HCMV (Towne strain) that measures total virus yields at 5 days post-infection. Most

reduction in viral titers was observed for heparin (PIHSS) followed by enoxaparin

derivative with >20 *dp* (Fig. 5A). Plotting of viral titer reduction as a function of *dp*

revealed a general trend where higher *dp* derivatives lead to higher reduction in viral

titers (Fig. 5B). Thus, this experiment indicated that longer HS chains are more efficient

at reducing HCMV titers in cells. To investigate whether this inhibitory effect was due to

an increase in the number of HCMV binding sites per chain of longer chain GAGs

towards virus particles, the experiments were repeated at 0.05 g/L concentrations of

GAGs instead of the previously used molar equivalent concentrations (Fig. 6A). As

micromolar concentration (10 µM) of GAGs is based on number of molecules provided,

GAGs consisting of longer chain will have more potential binding sites for virus than

those of shorter chains. The other concentration (0.05 g/L) is based on weight; thus this

concentration normalizes the number of potential virus binding sites for GAGs

consisting of both long and short chains. Interestingly, similar trend of inhibitory results

leaning towards efficacy of higher *dp* against HCMV infection were obtained at 0.05 g/L

indicating that this effect is not merely due to a higher number of potential independent

binding sites in the longer GAG chains and instead involves a difference in the

molecular interaction between HCMV and the longer GAG chains. A line graph for each

concentration of GAGs was generated that demonstrates the relationship of viral titer

and degree of polymerization (Fig. 6B). Although GAG treatment is not known to induce

cell death, to rule out that these effects on virus titers could be attributed to the health of

cells, we performed cell viability assays in both uninfected and infected settings. Cell

viability was not affected at the treated concentrations of any of our test GAGs (Fig. 7A).

Moreover, heparin (PIHSS) and enoxaparin derivatives (*dp* 12 or greater) efficiently

protected cells from virus induced lytic death (Fig. 7B). These results corroborate the

results of our glycoarray experiments that showed that GAG with higher *dp* have higher

CMV binding compared to GAG with lower *dp* (Fig. 3).

Cell lines defective in expression of specific sulfation enzymes have reduced CMV titers

and reduced virus entry.

Due to species specificity of HCMV, animal models are frequently used to study CMV

pathogenesis (38, 39). Studies of murine CMV (MCMV) infections of mice have served

a major role as a model of CMV biology and pathogenesis (40). Lung endothelial cell

lines from adult mice were mutated for specific sulfotransferase enzymes by a CRISPR-

Cas9 based gene editing system (35, 41, 42). Since previous studies showed that 3-*O*-

S HS is important for HCMV entry in human iris stromal cells (43), we analyzed virus

replication in Hs3st1 and Hs3st4 (Glucosaminyl 3-*O*-sulfotransferase 1 and 4,

respectively) knockout cell lines as well as the Hs3st1/4 double knockout cell line. At

high (5.0) as well as low (0.01) multiplicity of infection (MOI), MCMV growth was

significantly reduced in the single Hs3st1 and Hs3st4 knockouts as well as in the double

Hs3st1/4 knockouts, indicating that 3-*O*-sulfation of HS is important for HCMV infection

(Fig. 8). Further, we probed whether virus entry is impacted in the cells knocked out for

different combinations of these sulfotransferases. Expression of viral immediate early

protein (IE1) has been used as a surrogate for virus entry since it's one of the earliest

events after a successful virus entry (44-46). Results of an immunoblot probing for IE1

show that virus entry is significantly impacted in H3st1$^{-/-}$, H3st4$^{-/-}$, H3st1/4-double-

knockout, H6st1$^{-/-}$, H6st2$^{-/-}$, and H6st1/2 double-knockout cells, compared to the wild-

type cells (Fig. 9A and B). The H6st1$^{-/-}$, H6st2$^{-/-}$ and H3st4$^{-/-}$ mutants showed the most

impact on MCMV entry.

Peptides generated against HS and 3-*O*-S HS block HCMV infection.

In order to examine the effect of sulfated HS on HCMV infection, we utilized phage

display derived anti-HS and anti-3-*O*-S HS peptide (29). These peptides bind

specifically to cellular HS and 3-*O*-S HS respectively, and have been shown to block

binding of HSV-1. The HFF cells were pre-treated with either the anti-HS peptide or the

anti-3-*O*-S HS peptide and the mock-treated cells were used as control. As indicated in

Fig. 10, the anti-3-*O*-S HS peptide treatment resulted in a significant reduction of HCMV

foci in HFF cells compared to an anti-HS peptide or the mock-treated cells. The fact that

anti-3-*O*-S HS peptide was much more effective at reducing HCMV foci compared to an

anti-HS peptide indicates that this effect is governed by HS structure and not only by the

peptide charge.

DISCUSSION

In this study, we utilized multiple approaches, including glycoarray binding analysis, HS

mimics, HS mutant cell lines, and anti-HS/3-*O*-S HS peptides to demonstrate that

specifically sulfated HS with higher degree of polymerization affect CMV infection and

binding. The results significantly advance the age-old knowledge of HS binding to

herpesviruses by illustrating the importance of HS structural modifications in CMV

binding and infection. We first screened several sulfated or unsulfated GAGs with

complex sugar structure to investigate which GAGs are more efficient at binding to HCMV virions. This glycoarray analysis indicated that HCMV bound heparins with strong affinity and showed increased affinity for longer chain length heparins (Fig. 2). To further investigate this binding, we utilized another glycoarray consisting of HS of varied polymerization and sulfation levels. The results from this glycoarray indicated that HCMV binds strongly with HS having both longer monosaccharide chain and a moderate level of sulfation (Fig. 3). Thus, sulfated HS with more complex branches and sulfation patterns preferentially bind to HCMV. Next, we fractioned HS by length (*dp* 2-20) from enoxaparin and tested their ability to inhibit HCMV growth in cell culture by competing with HCMV binding. The GFP tagged HCMV was used and the number of GFP+ foci was quantified in the presence of increasing HS chain length. Amounts of viral GFP was more effectively reduced when cells were pretreated and maintained with GAGs having a higher *dp* (Fig. 4). This assay served as a surrogate for a virus entry assay since the GFP is independently expressed from an early promoter in the virus genome (47). For a deeper understanding of this reduction, we performed a similar experiment where HCMV Towne strain was used and viral load was quantified at 5 days post-infection. Significant reduction in virus titers was observed in samples treated with higher *dp* of GAG but not with lower *dp* corroborating the results from glycoarray experiments that chain length of GAG is an important factor in determining HCMV binding. Also, this effect was not due to a simple increase in the number of potential HCMV binding sites per mole of GAG, as evidenced by similar trend of inhibition obtained when treating cells with equivalent µM or g/L concentrations of GAGs. Treatment of cells with these GAGs did not affect cell viability for the duration of

treatment (Fig. 7A) confirming that the observed reduction in virus titer was not due to

the cell death. Moreover, cells pretreated and maintained with GAGs of longer *dp*

resisted infection induced cell death at late time post-infection (Fig. 7B). We also tested

the impact of specific HS sulfation mutants on MCMV infection. As 3-*O* sulfation has

been reported to be critical for herpesvirus entry (43, 48), we tested MCMV growth in

Hs3st1, Hs3st4 and dual Hs3st1/4 knockout cells. For both high and low MOI, virus titer

was significantly reduced in Hs3st1, Hs3st4 and dual Hs3st1/4 knockout cells (Fig. 8).

Although the differences were statistically significant, we did not see a robust inhibition

of virus titers in these assays and also the double knockout had somewhat less

reduction overall. This could largely be explained by the possible presence of additional

isoforms of Hs3st1 and Hs3st4 in these cells (49). To directly assess the impact of HS

sulfation on virus entry, we used several mutant cell lines deficient in HS sulfation

enzymes. All of these cells were defective in virus entry as assessed by IE1 protein

expression (Fig. 9.). Additional data from anti-3-*O*-S HS peptide confirmed the

significance of sulfation in HCMV infectivity. The GAG experiments used a fibroblast cell

culture system and a fibroblast tropic strain of HCMV (Towne), whereas MCMV

experiments used lung epithelial cells, thus providing the experimental data from

multiple cell types and two different viruses. While the GFP based assays provide a

surrogate for virus entry assays, the real impact of virus entry inhibitors would be a

reduction in viral titers at the end of infection since an entry inhibitor that only delays

virus entry would be of little translational value. Thus virus yield and titers were used a

measure of effectiveness of GAG inhibitors.

The enzymatic modification of HS chains is known to generate unique binding sites for

viral ligands. For example, 3-*O*-sulfation modification in HS chain generates fusion

receptor for HSV glycoprotein D (gD) promoting viral entry and spread (50). The 3-*O*-S

HS is a product of enzymatic modification at C3 position of glucosamine residue, which

is relatively rare in comparison to other HS modifications (Fig. 1B). Expression of Hs3st

can make normally resistant Chinese hamster ovary (CHO-K1) cells susceptible to

HSV-1 infection (51). Studies in clinically relevant primary human corneal fibroblasts

have also shown 3-*O*-S HS as a primary attachment receptor for HSV entry (48).

Interestingly, both HSV-1 and HSV-2 use HS as an attachment receptor but HSV-1

binds to distinct modification sites on HS that HSV-2 is unable to, which could explain

some of the differences in cell tropism exhibited by these two viruses (52). For example,

while N-sulfation and carboxyl groups are required for both HSV-1 and HSV-2 binding,

only HSV-1 is able to bind the specific modification sites generated by 2-*O*, 6-*O,* and 3-

O-sulfations (53). The *O*-desulfated heparins have little or no inhibitory effect on HSV-1

infection but inhibit HSV-2 infection. This susceptibility to *O*-desulfated heparins can be

transferred to HSV-1 by recombinant transfer of the gene for glycoprotein C (gC-2) from

HSV-2 (53). We reported earlier that 3-*O*-S HS are important for HCMV entry in human

iris stromal (HIS) cells (43). The expression of Hs3st in HIS cells promoted HCMV

internalization, while pretreatment of HIS cells with heparinase enzyme or treatment

with anti-3-*O*-S HS (G2) peptide significantly reduced HCMV plaques/foci formation. In

addition, co-culture of the HCMV-infected HIS cells with CHO-K1 cells expressing 3-*O*-S

HS significantly enhanced cell fusion. A similar trend of enhanced fusion was observed

with cells expressing HCMV glycoproteins (gB, gO, and gH-gL) co-cultured with 3-*O*-S

HS cells. These results highlight the role of 3-*O*-S HS during HCMV entry.

Owing to their inherent structural features, certain sulfated glycans can exert

therapeutic effects against infections caused by pathogenic microorganisms. A study by

Pomin *et al.*, showed that administering sulfated glycans can disrupt the pathogen

protein-host glycosaminoglycan (GAG) complex formation causing impairment of

microbial binding onto host cells (54). Similarly, sulfated GAG, glycosphingolipids and

lectins have been shown to inhibit DENV entry (55). Heparan sulfate mimics, such as

suramin, pentosan polysulfate, and PI-88, SPGG (56, 57) have been reported to be

effective against multiple viruses including herpesviruses (4, 58, 59). The inhibitory

activity of HS mimics, including these compounds, is believed to be due to their

association with GAG binding sites of the putative receptor-binding domain on the viral

protein (4, 60). Thus, HS mimics can inhibit virus adsorption and entry.

Overall, the data from these studies indicate that *dp* of GAGs as well as specific

sulfation patterns govern HCMV infection of cells. These studies show the promise of

highly polymerized sulfated-HS as effective anti-CMV agents. Future studies will be

aimed at confirming the CMV glycoproteins that specifically bind to HS on cell surface

and their possible structural illustrations.

Acknowledgments

The research was supported by NASA (Award #80NSSC19K1603, PI: Tandon) and NIH

(Award R21HL131553, PI: L.W.). QL, JSS and LW acknowledge funding from the

Author Contributions

438   RT, JSS, and LW designed the experiments; MHH, DM, LAF, RBP, QL, JSS, VT and

439   RT performed the experiments and analyzed the data. RT and MHH wrote and edited

440   the manuscript.

REFERENCES

1. Clausen TM, Sandoval DR, Spliid CB, Pihl J, Perrett HR, Painter CD, Narayanan A, Majowicz SA, Kwong EM, McVicar RN, Thacker BE, Glass CA, Yang Z, Torres JL, Golden GJ, Bartels PL, Porell RN, Garretson AF, Laubach L, Feldman J, Yin X, Pu Y, Hauser BM, Caradonna TM, Kellman BP, Martino C, Gordts P, Chanda SK, Schmidt AG, Godula K, Leibel SL, Jose J, Corbett KD, Ward AB, Carlin AF, Esko JD. 2020. SARS-CoV-2 Infection Depends on Cellular Heparan Sulfate and ACE2. Cell 183:1043-1057 e15.

2. Tandon R, Sharp JS, Zhang F, Pomin VH, Ashpole NM, Mitra D, McCandless MG, Jin W, Liu H, Sharma P, Linhardt RJ. 2021. Effective Inhibition of SARS-CoV-2 Entry by Heparin and Enoxaparin Derivatives. J Virol 95.

3. Barth H, Schäfer C, Adah MI, Zhang F, Linhardt RJ, Toyoda H, Kinoshita-Toyoda A, Toida T, van Kuppevelt TH, Depla E. 2003. Cellular binding of hepatitis C virus envelope glycoprotein E2 requires cell surface heparan sulfate. Journal of Biological Chemistry 278:41003-41012.

4. Chen Y, Maguire T, Hileman RE, Fromm JR, Esko JD, Linhardt RJ, Marks RM. 1997. Dengue virus infectivity depends on envelope protein binding to target cell heparan sulfate. Nature medicine 3:866.

5. Giroglou T, Florin L, Schäfer F, Streeck RE, Sapp M. 2001. Human papillomavirus infection requires cell surface heparan sulfate. Journal of virology 75:1565-1570.

6.   Tyagi M, Rusnati M, Presta M, Giacca M. 2001. Internalization of HIV-1 tat

requires cell surface heparan sulfate proteoglycans. Journal of Biological

Chemistry 276:3254-3261.

7.   Shukla D, Spear PG. 2001. Herpesviruses and heparan sulfate: an intimate

relationship in aid of viral entry. The Journal of clinical investigation 108:503-510.

8.   Shukla D, Liu J, Blaiklock P, Shworak NW, Bai X, Esko JD, Cohen GH,

Eisenberg RJ, Rosenberg RD, Spear PG. 1999. A novel role for 3-O-sulfated

heparan sulfate in herpes simplex virus 1 entry. Cell 99:13-22.

9.   Blanchard E, Belouzard S, Goueslain L, Wakita T, Dubuisson J, Wychowski C,

Rouillé Y. 2006. Hepatitis C virus entry depends on clathrin-mediated

endocytosis. Journal of virology 80:6964-6972.

10.   Compton T, Feire A. 2007. Early events in human cytomegalovirus infection,

Human herpesviruses: biology, therapy, and immunoprophylaxis. Cambridge

University Press.

11.   Ryckman BJ, Jarvis MA, Drummond DD, Nelson JA, Johnson DC. 2006. Human

cytomegalovirus entry into epithelial and endothelial cells depends on genes

UL128 to UL150 and occurs by endocytosis and low-pH fusion. Journal of

virology 80:710-722.

12.   Podyma-Inoue KA, Moriwaki T, Rajapakshe AR, Terasawa K, Hara-Yokoyama

M. 2016. Characterization of heparan sulfate proteoglycan-positive recycling

endosomes isolated from glioma cells. Cancer Genomics-Proteomics 13:443-

452.

13. Park H, Kim M, Kim H-J, Lee Y, Seo Y, Pham CD, Lee J, Byun SJ, Kwon M-H.

2017. Heparan sulfate proteoglycans (HSPGs) and chondroitin sulfate

proteoglycans (CSPGs) function as endocytic receptors for an internalizing anti-

nucleic acid antibody. Scientific reports 7:14373.

14. Christianson HC, Belting M. 2014. Heparan sulfate proteoglycan as a cell-surface

endocytosis receptor. Matrix Biology 35:51-55.

15. Sarrazin S, Lamanna WC, Esko JD. 2011. Heparan sulfate proteoglycans. Cold

Spring Harbor perspectives in biology 3:a004952.

16. Varnum SM, Streblow DN, Monroe ME, Smith P, Auberry KJ, Paša-Tolić L,

Wang D, Camp DG, Rodland K, Wiley S. 2004. Identification of proteins in

human cytomegalovirus (HCMV) particles: the HCMV proteome. Journal of

virology 78:10960-10966.

17. Spear PG, Shieh M-T, Herold BC, WuDunn D, Koshy TI. 1992. Heparan sulfate

glycosaminoglycans as primary cell surface receptors for herpes simplex virus, p

341-353, Heparin and related polysaccharides. Springer.

18. Compton T, Nowlin DM, Cooper NR. 1993. Initiation of human cytomegalovirus

infection requires initial interaction with cell surface heparan sulfate. Virology

193:834-841.

19. Laquerre S, Argnani R, Anderson DB, Zucchini S, Manservigi R, Glorioso JC.

1998. Heparan sulfate proteoglycan binding by herpes simplex virus type 1

glycoproteins B and C, which differ in their contributions to virus attachment,

penetration, and cell-to-cell spread. Journal of virology 72:6119-6130.

20.   Jacquet A, Haumont M, Chellun D, Massaer M, Tufaro F, Bollen A, Jacobs P.

507   1998. The varicella zoster virus glycoprotein B (gB) plays a role in virus binding

508   to cell surface heparan sulfate proteoglycans. Virus research 53:197-207.

509   21.   Wang X, Huong SM, Chiu ML, Raab-Traub N, Huang ES. 2003. Epidermal

510   growth factor receptor is a cellular receptor for human cytomegalovirus. Nature

511   424:456-61.

512   22.   Soroceanu L, Akhavan A, Cobbs CS. 2008. Platelet-derived growth factor-alpha

513   receptor activation is required for human cytomegalovirus infection. Nature

514   455:391-5.

515   23.   Feire AL, Roy RM, Manley K, Compton T. 2010. The glycoprotein B disintegrin-

516   like domain binds beta 1 integrin to mediate cytomegalovirus entry. J Virol

517   84:10026-37.

518   24.   Feire AL, Koss H, Compton T. 2004. Cellular integrins function as entry receptors

519   for human cytomegalovirus via a highly conserved disintegrin-like domain. Proc

520   Natl Acad Sci U S A 101:15470-5.

521   25.   Boyle KA, Compton T. 1998. Receptor-binding properties of a soluble form of

522   human cytomegalovirus glycoprotein B. Journal of virology 72:1826-1833.

523   26.   Song BH, Lee GC, Moon MS, Cho YH, Lee CH. 2001. Human cytomegalovirus

524   binding to heparan sulfate proteoglycans on the cell surface and/or entry

525   stimulates the expression of human leukocyte antigen class I. Journal of General

526   Virology 82:2405-2413.

527   27.   Esko JD, Selleck SB. 2002. Order out of chaos: assembly of ligand binding sites

528   in heparan sulfate. Annual review of biochemistry 71:435-471.

529   28.   Multhaupt HA, Couchman JR. 2012. Heparan sulfate biosynthesis: methods for

530   investigation of the heparanosome. Journal of Histochemistry & Cytochemistry

531   60:908-915.

532   29.   Tiwari V, Liu J, Valyi-Nagy T, Shukla D. 2011. Anti-heparan sulfate peptides that

533   block herpes simplex virus infection in vivo. J Biol Chem 286:25406-15.

534   30.   Baldwin J, Maus E, Zanotti B, Volin MV, Tandon R, Shukla D, Tiwari V. 2015. A

535   role for 3-O-sulfated heparan sulfate in promoting human cytomegalovirus

536   infection in human iris cells. J Virol 89:5185-92.

537   31.   Tiwari V, Tarbutton MS, Shukla D. 2015. Diversity of heparan sulfate and HSV

538   entry: basic understanding and treatment strategies. Molecules 20:2707-27.

539   32.   Ali MM, Karasneh GA, Jarding MJ, Tiwari V, Shukla D. 2012. A 3-O-sulfated

540   heparan sulfate binding peptide preferentially targets herpes simplex virus 2-

541   infected cells. J Virol 86:6434-43.

542   33.   Wei Z, Lyon M, Gallagher JT. 2005. Distinct substrate specificities of bacterial

543   heparinases against N-unsubstituted glucosamine residues in heparan sulfate. J

544   Biol Chem 280:15742-8.

545   34.   Kariya Y, Kyogashima M, Suzuki K, Isomura T, Sakamoto T, Horie K, Ishihara M,

546   Takano R, Kamei K, Hara S. 2000. Preparation of completely 6-O-desulfated

547   heparin and its ability to enhance activity of basic fibroblast growth factor. J Biol

548   Chem 275:25949-58.

549   35.   Qiu H, Shi S, Yue J, Xin M, Nairn AV, Lin L, Liu X, Li G, Archer-Hartmann SA,

550   Dela Rosa M, Galizzi M, Wang S, Zhang F, Azadi P, van Kuppevelt TH, Cardoso

551   WV, Kimata K, Ai X, Moremen KW, Esko JD, Linhardt RJ, Wang L. 2018. A

mutant-cell library for systematic analysis of heparan sulfate structure-function

relationships. Nat Methods 15:889-899.

36. Zurbach KA, Moghbeli T, Snyder CM. 2014. Resolving the titer of murine

cytomegalovirus by plaque assay using the M2-10B4 cell line and a low viscosity

overlay. Virol J 11:71.

37. Alam SM, Aussedat B, Vohra Y, Meyerhoff RR, Cale EM, Walkowicz WE,

Radakovich NA, Anasti K, Armand L, Parks R, Sutherland L, Scearce R, Joyce

MG, Pancera M, Druz A, Georgiev IS, Von Holle T, Eaton A, Fox C, Reed SG,

Louder M, Bailer RT, Morris L, Abdool-Karim SS, Cohen M, Liao HX, Montefiori

DC, Park PK, Fernandez-Tejada A, Wiehe K, Santra S, Kepler TB, Saunders KO,

Sodroski J, Kwong PD, Mascola JR, Bonsignori M, Moody MA, Danishefsky S,

Haynes BF. 2017. Mimicry of an HIV broadly neutralizing antibody epitope with a

synthetic glycopeptide. Sci Transl Med 9.

38. Reddehase MJ, Lemmermann NAW. 2018. Mouse Model of Cytomegalovirus

Disease and Immunotherapy in the Immunocompromised Host: Predictions for

Medical Translation that Survived the "Test of Time". Viruses 10.

39. Brune W, Hengel H, Koszinowski UH. 2001. A mouse model for cytomegalovirus

infection. Curr Protoc Immunol Chapter 19:Unit 19 7.

40. Cekinovic D, Lisnic VJ, Jonjic S. 2014. Rodent models of congenital

cytomegalovirus infection. Methods Mol Biol 1119:289-310.

41. Sauer B. 1987. Functional expression of the cre-lox site-specific recombination

system in the yeast Saccharomyces cerevisiae. Mol Cell Biol 7:2087-96.

42.   Zhang F, Wen Y, Guo X. 2014. CRISPR/Cas9 for genome editing: progress,
575   implications and challenges. Hum Mol Genet 23:R40-6.

576   43.   Baldwin J, Maus E, Zanotti B, Volin MV, Tandon R, Shukla D, Tiwari V. 2015. A
577   role for 3-O-sulfated heparan sulfate in promoting human cytomegalovirus
578   infection in human iris cells. Journal of virology 89:5185-5192.

579   44.   Oduro JD, Uecker R, Hagemeier C, Wiebusch L. 2012. Inhibition of human
580   cytomegalovirus immediate-early gene expression by cyclin A2-dependent
581   kinase activity. J Virol 86:9369-83.

582   45.   Adamson CS, Nevels MM. 2020. Bright and Early: Inhibiting Human
583   Cytomegalovirus by Targeting Major Immediate-Early Gene Expression or
584   Protein Function. Viruses 12.

585   46.   Archer MA, Brechtel TM, Davis LE, Parmar RC, Hasan MH, Tandon R. 2017.
586   Inhibition of endocytic pathways impacts cytomegalovirus maturation. Sci Rep
587   7:46069.

588   47.   Dunn W, Chou C, Li H, Hai R, Patterson D, Stolc V, Zhu H, Liu F. 2003.
589   Functional profiling of a human cytomegalovirus genome. Proc Natl Acad Sci U S
590   A 100:14223-8.

591   48.   Tiwari V, Clement C, Xu D, Valyi-Nagy T, Yue BY, Liu J, Shukla D. 2006. Role
592   for 3-O-sulfated heparan sulfate as the receptor for herpes simplex virus type 1
593   entry into primary human corneal fibroblasts. Journal of virology 80:8970-8980.

594   49.   Qiu H, Shi S, Yue J, Xin M, Nairn AV, Lin L, Liu X, Li G, Archer-Hartmann SA,
595   Dela Rosa M, Galizzi M, Wang S, Zhang F, Azadi P, van Kuppevelt TH, Cardoso
596   WV, Kimata K, Ai X, Moremen KW, Esko JD, Linhardt RJ, Wang L. 2018. A

mutant-cell library for systematic analysis of heparan sulfate structure–function

relationships. Nature Methods doi:10.1038/s41592-018-0189-6.

50.   Tiwari V, Tarbutton M, Shukla D. 2015. Diversity of heparan sulfate and HSV

entry: basic understanding and treatment strategies. Molecules 20:2707-2727.

51.   Tiwari V, O'Donnell C, Copeland RJ, Scarlett T, Liu J, Shukla D. 2007. Soluble 3-

O-sulfated heparan sulfate can trigger herpes simplex virus type 1 entry into

resistant Chinese hamster ovary (CHO-K1) cells. Journal of General Virology

88:1075-1079.

52.   O'Donnell CD, Kovacs M, Akhtar J, Valyi-Nagy T, Shukla D. 2010. Expanding the

role of 3-O sulfated heparan sulfate in herpes simplex virus type-1 entry. Virology

397:389-398.

53.   Herold BC, Gerber SI, Belval BJ, Siston AM, Shulman N. 1996. Differences in the

susceptibility of herpes simplex virus types 1 and 2 to modified heparin

compounds suggest serotype differences in viral entry. Journal of virology

70:3461-3469.

54.   Pomin VH. 2017. Antimicrobial Sulfated Glycans: Structure and Function. Curr

Top Med Chem 17:319-330.

55.   Hidari KI, Abe T, Suzuki T. 2013. Carbohydrate-related inhibitors of dengue virus

entry. Viruses 5:605-18.

56.   Gangji RN, Sankaranarayanan NV, Elste J, Al-Horani RA, Afosah DK, Joshi R,

Tiwari V, Desai UR. 2018. Inhibition of Herpes Simplex Virus-1 Entry into Human

Cells by Nonsaccharide Glycosaminoglycan Mimetics. ACS Med Chem Lett

9:797-802.

57.   Majmudar H, Hao M, Sankaranarayanan NV, Zanotti B, Volin MV, Desai UR,

Tiwari V. 2019. A synthetic glycosaminoglycan mimetic blocks HSV-1 infection in

human iris stromal cells. Antiviral Res 161:154-162.

58.   Lee E, Pavy M, Young N, Freeman C, Lobigs M. 2006. Antiviral effect of the

heparan sulfate mimetic, PI-88, against dengue and encephalitic flaviviruses.

Antiviral Res 69:31-8.

59.   Marks RM, Lu H, Sundaresan R, Toida T, Suzuki A, Imanari T, Hernaiz MJ,

Linhardt RJ. 2001. Probing the interaction of dengue virus envelope protein with

heparin: assessment of glycosaminoglycan-derived inhibitors. J Med Chem

44:2178-87.

60.   Modis Y, Ogata S, Clements D, Harrison SC. 2003. A ligand-binding pocket in

the dengue virus envelope glycoprotein. Proc Natl Acad Sci U S A 100:6986-91.

61.   Poulain FE, Yost HJ. 2015. Heparan sulfate proteoglycans: a sugar code for

vertebrate development? Development 142:3456-67.

FIGURE LEGENDS

Figure 1. Structural features of heparan sulfate. (A) HS is a linear polysaccharide composed of repeating uronic acid [D-glucuronic acid (GlcA) or L-iduronic acid (IdoA)] and D-glucosamine (GlcN) disaccharide subunits. Synthesized chain of HS represents assembly of the tetrasaccharide linker region (GlcA-Gal-Gal- Xyl) at reducing end on serine residues of the protein core followed by the addition of alternating GlcA and GlcNAc residues. The chain extension is also accompanied by a series of modifications, which include 6-*O*, 3-*O* sulfations on GlcN and the 2-*O* sulfation on GlcA. The arrow shows the 3-*O* position of the GlcN where sulfation is important for herpesvirus binding (15, 50).

(B) Heparan sulfate chains are initially synthesized as repeating disaccharide units of N-acetylated glucosamine and glucuronic acid. HS can then be modified by a series of enzymatic reactions, including N-deacetylation and N-sulfation of N-acetylated glucosamine converting it to N-sulfo-glucosamine, C5 epimerization of glucuronic acid to iduronic acid, and O-sulfation at the 2-OH, 6-OH, and 3-OH positions. Among sulfations, first is 2-*O*-sulfation of iduronic acid and glucuronic acid, followed by 6-*O*-sulfation of N-acetylated glucosamine and N-sulfo-glucosamine units, and finally 3-*O*-sulfation of glucosamine residues (52, 61).

Fig. 1A.

Fig. 1B

Figure 2. Binding of purified extracellular CMV virions on a custom designed glycosaminoglycan glycoarray. Relative fluorescence units (RFU), which are directly proportional to the amount of virus binding, are plotted on the Y-axis in the graph. Ligand descriptions and chain structures are provided in Table 1. Six replicates for each GAG were used in the assay. NC: Negative control (print buffer), PC1: positive control (Biotinylated Glycan), PC2: human IgG (0.1 mg/ml), PC3: mouse IgG (0.1 mg/ml), PC4: rabbit IgG (0.1 mg/ml), *dp*: degree of polymerization, triangles at the bottom of the graph represent an increasing degree of polymerization of GAGs from left to right. The virus preparation does not contain streptavidin label and thus PC1 serves as a negative rather than a positive control.

Figure 3. Binding of purified extracellular CMV on a custom designed heparan sulfate glycoarray. Relative fluorescence units (RFU), which are directly proportional to the amount of virus binding, are plotted on the Y-axis in the graph. Ligand descriptions and chain structures are provided in Table 2. Six replicates for each ligand were used. NC: negative control (print buffer) PC1: positive control (biotinylated glycan), PC2: human IgG (0.1 mg/ml), PC3: mouse IgG (0.1 mg/ml), PC4: rabbit IgG (0.1 mg/ml). The virus preparation does not contain streptavidin label and thus PC1 serves as a negative rather than a positive control.

Figure 4. Inhibition of HCMV growth by glycosaminoglycan derivatives. Primary human foreskin fibroblasts (HFF) grown in 96 well plate were pretreated for one hour with 10 µM of 1) 6-*O*-desulfated Arixtra, 2) Unmodified Arixtra, 3) Heparin sodium salt from porcine intestinal mucosa (PIHSS), 4) Enoxaparin, or series of heparin oligosaccharide from enoxaparin: 5) *dp2*, 6) *dp4*, 7) *dp6*, 8) *dp8*, 9) *dp10*, 10) *dp12*, 11) *dp14*, 12) *dp16*, 13) *dp18*, 14) dp20 15) > *dp20* or control (dH2O). Cells were infected with GFP tagged HCMV (Towne strain) virus at an MOI of 3.0 in the presence of the test glycosaminoglycans. At 5 days post-infection, cells were fixed and number of foci (GFP) was counted under an epifluorescent microscope. Percent of viral GFP was calculated compared to virus only infected control (100% GFP expression). Results are representative of three independent replicates. Standard error of mean was plotted as error bars.

Figure 5. Effect of glycosaminoglycan derivatives at 10 µM concentration on HCMV growth. (A) Primary human foreskin fibroblasts (HFF) were pretreated for one hour with 10 µM of 1) 6-*O*-desulfated Arixtra, 2) Regular Arixtra, 3) Heparin sodium salt from porcine intestinal mucosa (PIHSS), 4) Enoxaparin, or series of heparin oligosaccharide from enoxaparin: 5) *dp2*, 6) *dp4*, 7) *dp6,* 8) *dp8*, 9) *dp10*, 10) *dp12*, 11) *dp14*, 12) *dp16*, 13) *dp18*, 14) *dp20* 15) > *dp20* or control (dH2O). Cells were infected with HCMV (Towne strain) virus at an MOI of 3.0 in the presence of test glycosaminoglycans. Cells and media were harvested at 5 days post-infection and titered for HCMV plaque forming units (pfu) on fresh fibroblasts in tissue culture dishes. Individual samples (3 replicates each) were quantified and displayed as total pfu/ml on Y-axis. (B) Virus titer is plotted (Y-axis) against degree of polymerization (X-axis). Data points ahead of the broken line is for a mixture of GAGs (*dp*>20). Results are representative of three independent replicates. Standard error of mean was plotted as error bars.

Figure 6. Effect of glycosaminoglycan derivatives at 0.05 g/L concentration on HCMV growth. (A) Primary human foreskin fibroblasts (HFF) were pretreated for one hour with 0.05 g/L of 1) 6-*O*-desulfated Arixtra, 2) Regular Arixtra, 3) Heparin sodium salt from porcine intestinal mucosa (PIHSS), 4) Enoxaparin, or series of heparin oligosaccharide from enoxaparin: 5) *dp2*, 6) *dp4*, 7) *dp6*, 8) *dp8*, 9) *dp10*, 10) *dp12*, 11) *dp14*, 12) *dp16*, 13) *dp18*, 14) *dp20* 15) > *dp20* or control (dH2O). Cells were infected with HCMV (Towne strain) virus at an MOI of 3.0 in the presence of test glycosaminoglycans. Cells and media were harvested at 5 days post-infection and titered for HCMV plaque forming units (pfu) on fresh fibroblasts in tissue culture dishes. Individual samples (3 replicates each) were quantified and displayed as total pfu/ml on Y-axis. (B) Virus titer is plotted (Y-axis) against degree of polymerization (X-axis). Data points ahead of the broken line is for a mixture of GAGs (*dp*>20). Results are representative of three independent replicates. Standard error of mean was plotted as error bars.

Figure 7. Effect of GAG treatment on cell viability of HFF cells. Primary HFF were pretreated for one hour with 10 µM of 1) 6-*O*-desulfated Arixtra, 2) Regular Arixtra, 3) Heparin sodium salt from porcine intestinal mucosa (PIHSS), 4) Enoxaparin, or series of heparin oligosaccharide from enoxaparin: 5) *dp2*, 6) *dp4*, 7) *dp6*, 8) *dp8*, 9) *dp10*, 10) *dp12*, 11) *dp14*, 12) *dp16*, 13) *dp18*, 14) *dp20*, 15) > *dp20* or control (dH2O). Cells were either mock infected (A) or infected with HCMV (Towne strain) virus at an MOI of 3.0 (B) in the presence of test glycosaminoglycans. Cells were harvested at 5 days post-infection and cell viability was assessed using Trypan Blue exclusion assay. Results are representative of three independent replicates. Standard error of mean was plotted as error bars.

A.

B.

Figure 8. Mouse CMV replication in sulfotransferase knockout cell lines. Cells were grown to 90% confluency and infected with wild-type MCMV (strain K181) at low (0.01, A, B) and high (3.0, C, D) MOI. Cells and the medium were harvested at 3- and 5-days post-infection, sonicated to release the virus and diluted for plating on to wild-type MEF in tissue culture dishes in order to enumerate total MCMV pfu/ml. Results are representative of three independent replicates. A student's t-test was conducted in Graphpad Prism comparing the means of different groups. A p-value <0.05 was considered significant. Standard error of mean was plotted as error bars. An asterisk (*) indicates significant inhibition compared to wild-type. Hs3st1 and Hs3st4: Glucosaminyl 3-*O*-sulfotransferase 1 and 4, respectively. WT: wild type; KO: knockout.

MOI 0.01

MOI 3.0

Figure 9. Mouse CMV entry in sulfotransferase knockout cell lines. A. Cells were grown to 90% confluency and infected with wild-type MCMV (strain K181) at an MOI of 3.0. Cells were harvested at 2 hours post-infection, and whole cell lysates were loaded on a polyacrylamide gel for blotting. The blots were probed with anti IE1 antibody. Beta-actin was used as a loading control. B. Bands from two independent experiments were quantified by densitometry and means were plotted. Standard error of mean was plotted as error bars. Hs6st1 and Hs6st4: Glucosaminyl 6-O-sulfotransferase 1 and 4, respectively. Hs3st1 and Hs3st4: Glucosaminyl 3-O-sulfotransferase 1 and 4, respectively. WT: wild type; dko: double knockout.

A.

Uninfected control H$_3$$_s$t1/4dko H$_3$$_s$t4ko H$_3$$_s$t1ko H$_6$$_s$t1/2dko H$_6$$_s$t2-/- H$_6$$_s$t1-/- H$_6$$_s$t WT

IE1

Actin

B.

Figure 10. Effect of anti-3-O-S HS peptide on CMV infectivity in HFF cells. HFF

cells were pre-treated with wild-type HS peptide, anti 3-OS HS peptide for 4 hrs. The

mock treated cells were used as a positive control. The cells were infected with β-

galactosidase expressing CMV for 9 days. β-Galactosidase assay was performed using

X-gal (Sigma). The effect of entry-blocking activity of peptide was examined by counting

number of foci. Results are representative of three independent experiments. A

student's t-test was conducted in Graphpad Prism comparing the means of each group.

A p value <0.05 was considered significant. Standard error of mean was plotted as error

bars. Both anti-HS and anti 3-O-S HS showed significant inhibition compared to mock.