Functional genomics of a symbiotic community: shared traits in the olive fruit fly gut microbiota

Frances Blow[1,2], Anastasia Gioti[3], Ian B. Goodhead[1,4], Maria Kalyva[3,5] Anastasia Kampouraki[6,7], John Vontas[6,7], Alistair C. Darby[1*]

1. Institute of Integrative Biology, University of Liverpool, Liverpool, Merseyside, UK
2. Department of Entomology, Cornell University, Ithaca, NY 14853, USA
3. Perrotis College, American Farm School, Thessaloniki, Greece
4. School of Environment and Life Sciences, Peel Building, University of Salford, M5 4WT, UK
5. Department of Biology, University of Crete, Voutes Campus, Heraklion, Crete, Greece
6. Institute of Molecular Biology & Biotechnology, Foundation for Research & Technology Hellas, Heraklion, Crete, Greece
7. Pesticide Science, Agricultural University of Athens, Iera Odos 75, 11855, Athens, Greece

*Author for Correspondence:
Alistair C. Darby, Institute of Integrative Biology, University of Liverpool, Liverpool, Merseyside, UK email: acdarby@liverpool.ac.uk

Abstract

The olive fruit fly *Bactrocera oleae* is a major pest of olives worldwide and houses a specialized gut microbiota dominated by the obligate symbiont "*Candidatus* Erwinia dacicola*". Ca.* E. dacicola is required for juvenile development in ripening olives and is thought to supplement dietary nitrogen in adults, with only indirect evidence for this hypothesis so far. Here, we sought to functionally investigate the symbiosis and its contribution to insect fitness. Inspection of three available *Ca.* E. dacicola genomes and one transcriptome provided evidence that the symbiont can hydrolyze urea for use as a nitrogen source, most probably by the adult insect: A urease operon was identified, not expressed at the juvenile stage, for which we provide evidence for a likely acquisition by horizontal gene transfer. Moreover, to explore the ecology of the *B. oleae* gut, we examined the composition of bacterial communities associated with Cretan olive fruit fly populations. We identified and reconstructed the genome of a novel component of the gut microbiota, *Tatumella* sp. TA1, which is stably associated with Mediterranean olive fruit fly populations. We further detected, in the genomes of both obligate and facultative members of the gut microbiota, components of extracellular surface structures, which may mediate interactions with the host and between microbes, including the exchange of genetic material. Our findings suggest that urea hydrolysis is a conserved functional trait for microbes stably inhabiting the *B. oleae* gut, and that genetic exchanges between microbes may have shaped the symbiosis between *B. oleae* and *Ca.* E. dacicola.

Key words: Symbiosis, horizontal gene transfer, urease, adhesion, *Candidatus* Erwinia dacicola, *Tatumella* sp. TA1

Introduction

Many insects house gut microbial communities that perform essential functions related to their diet or lifestyle (Dillon and Dillon 2004). Insect species with a dependence on a specialized gut microbial community for their fitness often have specialized alimentary structures to house microbes, and these microbes are also often vertically transmitted between mother and offspring to ensure the inoculation of subsequent generations (Salem et al. 2015). The gut microbiota of wild populations of the olive fruit fly *Bactrocera oleae* (Tephritidae) is numerically dominated by a single Gammaproteobacterium, "*Candidatus* Erwinia dacicola" (Capuzzo et al. 2005; Estes et al. 2009; Kounatidis et al. 2009; Estes et al. 2012; Ben-Yosef et al. 2015). The symbiont *Ca.* E. dacicola resides in the digestive tract throughout the insect's lifecycle, in larvae within the midgut caeca and in adults in a specialized foregut diverticulum structure called the esophageal bulb, and is maternally transmitted between generations by egg smearing (Estes et al. 2009). *Ca.* E. dacicola has not been detected outside of the insect by environmental sampling, while efforts to culture it *in vitro* have so far been unsuccessful (Capuzzo et al. 2005). From the above, the bacterium qualifies as a specific obligate symbiont of *B. oleae*, but also a challenge to study experimentally. In this context, sequence data-based methods are of high value for the generation of hypotheses on the different roles of *Ca.* E. dacicola in host fitness.

In common with many holometabolous insects, the adult and juvenile stages of *B. oleae* feed on different food sources (Tzanakakis 2003), therefore the *Ca.* E. dacicola symbiosis is hypothesized to perform alternative functions during these developmental stages. In larvae, the symbiosis has been hypothesized to allow the insect to use ripening fruit by detoxifying defensive plant phenolic compounds, such as oleuropein (Soler-Rivas et al. 2000; Ben-Yosef et al. 2015), and sequestering amino acids for protein synthesis (Ben-Yosef et al. 2015). During adulthood, the native gut microbiota is thought to provision nitrogen to the host: females harboring their native microbiota have a higher reproductive output on diets lacking essential amino acids (EAAs), and diets containing only nitrogen sources that are metabolically intractable to *B. oleae*, such as urea (Ben-Yosef et al. 2010; Ben-Yosef et al. 2014). It is common for many insects to house microbes that increase the quantity or quality of dietary nitrogen by performing novel metabolic functions (Douglas 2009). For example, the intracellular symbiont *Blochmannia* within the bacteriocytes of *Camponotus* ants encodes urease that metabolizes dietary urea to ammonia from which it synthesizes essential and non-essential amino acids, subsequently transported to the hemolymph for host consumption (Feldhaar et al. 2007). A similar process is proposed to occur in *B. oleae* adults, which consume bird droppings containing ammonia and urea as part of their

omnivorous diet (Ben-Yosef et al. 2014), but do not encode endogenous ureases (*B. oleae*

genome accession number: GCF_001188975.1, Djambazian et al. 2018). However, the

specific pathways and related enzymes involved in the metabolism and uptake of

nitrogenous substrates remain to be elucidated in *B. olea*e.

Previous experimental approaches aiming to evaluate symbiotic function have tested the

native microbial community of the olive fruit fly, which is dominated by, but not restricted to

Ca. E. dacicola (Estes et al. 2009; Ben-Yosef et al. 2015; Blow, Vontas, et al. 2016; Blow et

al. 2017). In this study, we took advantage of recently available –omic resources for the

symbiont (Blow, Gioti et al. 2016, Pavlidi et al. 2017, Estes et al, 2018) and further

generated new sequence data from single-culture and metagenomic samples of the *B. oleae*

gut community to 1) investigate whether *Ca.* E. dacicola encodes the full gene repertoire

required to provide dietary nitrogen to *B. oleae* and 2) identify other members of the *B. oleae*

gut microbiota potentially capable of performing these functions. For this purpose, we

employed comparative genomic and phylogenetic analyses, considering the function of the

gut community as a whole in the context of symbiosis.

Results and discussion

A novel member of the gut microbiota associated with Mediterranean populations of

B. oleae

To explore the ecology of the *B. oleae* gut, we examined its bacterial community

composition by 16S rRNA gene amplicon sequencing of samples from olive fruit fly

populations collected in Crete. This approach, combined with culturing of selected isolates,

allowed identifying and isolating in to axenic culture a culture-viable member of the gut

microbiota that has not been characterized before, referred to here as *Tatumella* sp. TA1.

Tatumella sp. TA1 was initially identified in wild Cretan populations of adult *B. oleae,* and

reanalysis of 16S rRNA gene data from a previous study demonstrated that *Tatumella* sp.

TA1 is also present in wild populations from Israel (Ben-Yosef et al. 2015). Notably,

Tatumella sp. TA1 was not detected in any of the studies of bacterial communities

associated with US populations of *B. oleae* (Estes et al. 2009; Estes et al. 2012). These

results suggest that TA1 is a facultative member of the gut community and that it is restricted

to Mediterranean populations of the olive fruit fly. We detected *Tatumella* sp. TA1 in 88.9-90

% of individuals from populations in Israel, in both adults and larvae, and in 26.9 % of

individuals from populations in Crete (Table 1). We did not detect *Enterobacter* sp. OLF,

another facultative member of the gut community identified in US populations of *B. oleae*

(Estes et al. 2009), in either of the Mediterranean populations studied, suggesting that the

two taxa are geographically restricted.

We used PacBio RS II sequencing to generate a draft genome sequence for *Tatumella* sp.

TA1. The assembly comprised a chromosomal sequence of 3,389,139 bp and a circular

plasmid sequence of 49,211 bp (total genome size 3.4 Mb) encoding a total of 3,309 protein

coding genes, 72 tRNAs, and 22 rRNAs. Both the chromosome and plasmid sequences

have been circularized. The GC content of the genome was 48.6 %, and both this and the

genome size were comparable to those of previously-published genomes of *Tatumella*

species, many of which are host-associated (Hollis et al. 1981; Marín-Cevada et al. 2010;

Chandler et al. 2014). The assembly is predicted to be 100 % complete following the method

used in (Rinke et al. 2013), which detected 138/138 single-copy marker genes.

Phylogenetic placement of obligate and facultative members of the *B. oleae* gut

microbiota

In agreement with a previously-published comprehensive phylogeny (Palmer et al. 2017),

ortholog analysis of 46 bacterial taxa belonging to the genera *Erwinia*, *Tatumella* and

Pantoea (full list in Table S1A) supports the phylogenetic placement of *Tatumella* sp. TA1

within the *Tatumella* genus (Figure 1). Phylogenetic analyses confirmed that *Tatumella* sp.

TA1 and *Enterobacter* sp. OLF are distinct members of the order *Enterobacterales*,

indicating that they are unique components of the *B. oleae* gut microbiota. In addition, the

phylogeny showed that, despite an evolutionary association with *B. oleae* and vertical

transmission, which are expected to change genome characteristics through altered

selection pressures and drift (McCutcheon and Moran 2011), *Ca.* E. dacicola clusters within

the *Erwinia* genus (Figure 1). The closest phylogenetic relatives of *Tatumella* sp. TA1 and

Ca. E. dacicola are the thrip symbionts BFo2 and BFo1, respectively (Facey et al. 2015).

The clustering of these taxa may represent comparable lifestyles. In a result analogous to

diet experiments in olive flies (Ben-Yosef et al. 2010; Ben-Yosef et al. 2014), the fitness

benefits of the WFT gut-lumen symbiont BFo1 were condition-dependent and only apparent

on diets lacking a balanced source of amino acids, when gut bacteria are presumed to

synthesize and provision adult WFT with amino acids required for protein synthesis (de Vries

et al. 2004). BFo1 and BFo2 also inhabit the gut lumen and are vertically transmitted

between generations by an extracellular route (de Vries et al. 2001). Further functional

studies may elucidate whether specific characteristics shared by these taxa predispose them

to this lifestyle.

Functional redundancy of nitrogen assimilation genes in members of the *B. oleae* gut microbiota

We investigated the hypothesis that members of the *B. oleae* gut microbiota supplement the olive fruit fly diet with nitrogen by focusing on both obligate and facultative bacterial members. The annotated genomes of the obligate *Ca.* E. dacicola (Blow, Gioti et al. 2016) and facultative *Tatumella* sp. TA1 (this study) and *Enterobacter* sp. OLF (Estes et al. 2018) all encode enzymes that metabolize urea to ammonia and carbon dioxide (Figure 2, Table S2A): *Ca.* E. dacicola and *Enterobacter* sp. OLF encode ureases (EC 3.5.1.5), and *Tatumella* sp. TA1 encodes urea carboxylase (EC 6.3.4.6) and allophanate hydrolase (EC 3.5.1.54). None of these enzymes are encoded in the genome of *B. oleae* (Table S2A). A BlastP and TblastN search of the *Ca.* E. dacicola transcriptome data from larvae developing in green and black olives (Pavlidi et al. 2017) did not identify any of the urease genes, indicating that urease may not be expressed at the larval stage in the obligate symbiont. This result is expected, since at this stage there is no dietary source of urea. Therefore, a hypothesis to further explore is that urease expression in *Ca.* E. dacicola is inducible by host or dietary cues present during *B. oleae* adulthood. Similarly, the closest homologs to the *Ca.* E. dacicola urease gene (see next section) come from two plant pathogens, *Gibbsiella quercinecans* and *Brenneria goodwinii,* previously characterized as negative for urease function (Brady et al. 2010; Denman et al. 2012), which indicates that urease function may be inducible under specific conditions.

In addition to urea hydrolysis, *Ca.* E. dacicola, *Tatumella* sp. TA1 and *Enterobacter* sp. OLF all encode biosynthetic operons for non-essential and essential amino acids (EAA), with further evidence for expression during larval feeding in *Ca.* E. dacicola (Table S2B, sequences available in Table S3). Glutamine synthetase (EC 6.3.1.2), is also present in all three taxa and expressed in the *Ca.* E. dacicola transcriptome data. This enzyme, also encoded by the host genome (RefSeq accession GCF_001188975.1), incorporates nitrogen from ammonia into the non-essential amino acid glutamine (Figure 2), which can be channeled into downstream metabolic processes including EAA biosynthesis (Feldhaar et al. 2007; Sabree et al. 2009).

We also investigated the hypothesis that host nitrogenous waste, which would otherwise be excreted, is the source of nitrogen assimilated to non-essential amino acids by the gut microbiota. Urea could come from recycling of host waste uric acid (Ben-Yosef et al. 2014), as has been observed in other insect symbionts that upgrade the nitrogen content of the host diet (Potrikus and Breznak 1981; Sasaki et al. 1996; Kashima et al. 2006; Feldhaar et al. 2007; Sabree et al. 2009). We found no genomic evidence that urea can be produced via uricolysis in either the host or any member of the so-far identified and sequenced microbiota: BlastP searches against shotgun sequencing data of olive fruit fly gut samples and

corresponding metagenomic assemblies of the gut microbial community (NCBI BioProject
accession PRJNA326914) failed to identify the required enzyme allantoicase (EC 3.5.3.4;
Figure S1, Table S2A). However, *B. oleae* does encode an arginase (EC 3.5.3.1), which can
hydrolyze the essential amino acid arginine to urea and ornithine and could provide urease-
degrading bacteria with a supply of urea (Figure 2). One hypothesis would be that *B. oleae*
might enrich its gut microbiota with ureolytic bacteria such as *Ca.* E. dacicola through
inducible delivery of urea produced by arginase activity. Signal peptides were not detected
when individual urease proteins or the full operon from *Ca.* E. dacicola were analyzed with
SignalP version 4.0 (Petersen et al. 2011), indicating that, if expressed, the urease encoded
by *Ca.* E. dacicola is intracellular. This suggests that specific uptake of urea and excretion of
the products of its hydrolysis must either be coordinated with the host, or that microbial cells
are lysed after the hydrolysis of urea, in order for the host to gain the observed nutritional
benefit when urea is added to the diet (Ben-Yosef et al. 2010; 2014). A key component of
future studies should be to track the fate of metabolites resulting from urea hydrolysis:
whether they are retained by microbes for endogenous metabolism, or whether they are
trafficked back to the host as e.g. ammonia or glutamine.
205

Evidence for horizontal gene transfer of the urease operon in the obligate symbiont
***Ca.* E. dacicola**
The *Ca.* E. dacicola urease enzyme is encoded, as in all ureolytic bacteria, by a cluster of
eight adjacent genes arranged in an operon and denoted as *ureDABCEJFG*, with *ureABC*
encoding structural proteins of the enzymatic complex, *ureDEFG* accessory proteins, and
ureJ the transcriptional regulator. The urease operon appears incomplete at its 3' end in the
ErwSC assembly (Blow, Gioti et al. 2016), missing *ureG*. However, *ureG* is present in the
Ca. E. dacicola genome, as confirmed by BlastP queries against the initial set of
metagenomic reads obtained from shotgun sequencing (NCBI BioProject accession
PRJNA326914). The urease operon is also present in two additional *Ca.* E. dacicola
assemblies (IL and Oroville) reconstructed with different methods and from different olive fly
populations (Table S1A), and is complete (5,962 bp total length) in these assemblies,
including *ureG*. The three 'versions' of the operon from different assemblies are 99% similar
at the nucleotide level for the genes present. One explanation for the absence of *ureG* in the
ErwSC assembly is over-trimming, aiming to minimize the risk of including non-*Ca.* E.
dacicola scaffolds, since metagenomic sequencing data included DNA from several species,
such as *Tatumella* sp. TA1. The differences between the reduced ErwSC and the more
complete Oroville assembly are outlined in detail in (Estes, Hearn, Agrawal, et al. 2018). In
any case, the presence and similarity of the operon in all three *Ca* E. dacicola assemblies
despite their differences and despite the presence of different facultative taxa in the gut

microbiome of *B. oleae* from distinct geographic populations, indicate that the urease operon

is highly conserved in *Ca* E. dacicola.

To the best of our knowledge, *Ca.* E. dacicola is the only species of the genus *Erwinia* that

encodes urease genes. Examination of the phylogeny of *ureC*, the longest structural gene of

the operon, provided evidence for the acquisition of the urease operon in *Ca.* E. dacicola by

horizontal gene transfer (HGT): The *Ca.* E. dacicola *ureC* history (Figure 3A), is different

from that of the species history, as shown by the comparison to a 'reference' tree,

reconstructed from 8 neutral single-copy orthologs (Figure 3B). The reference tree

accurately depicts the known phylogenetic relationships between *Ca.* E. dacicola and other

Proteobacteria, in agreement with the recently published comprehensive phylogeny (Palmer

et al. 2017) and the species tree presented in this study (Figure 1). By contrast, the *Ca.* E.

dacicola *ureC* protein groups together with proteins from more distant Gammaproteobacteria

of the *Enterobacterales* order (*G. quercinecans*, *B. goodwinii*), and members of the distantly

related Betaproteobacteria (Figure 3). Hacker and Kaper (2000) defined genomic islands

(GEIs) in bacteria as small (<10 Kbp), syntenic blocks of genes acquired by HGT. It has

been observed that GEIs share some common characteristics, such as the encoding of

genes offering a selective advantage to the host, flanking by mobile genetic elements

(MGEs) such as transposases, insertion close to tRNA genes, and a GC content different

from that of the rest of the genome (Juhas et al. 2009). The ureolytic capacity conferred to

Ca. E. dacicola by the urease operon is potentially an example of acquired selective

advantage, since it increases the availability of nutrients to the symbiont itself or to the host,

upon which it is dependent. Thus, it is tempting to argue that the urease operon represents a

GEI in *Ca.* E. dacicola. Similarly to other GEIs, it has a distinct (lower) GC content compared

to the rest of the genome (mean GC_{urease}=0.502, mean GC_{genome}=0.535), and this difference

is statistically significant (p = 5.085e-06). Moreover, the distribution of GC values sampled

from the urease operon is significantly different from the whole-genome distribution (p

=1.031E-05). Unfortunately, proximity of the operon to MGEs and tRNAs could not be

confirmed in *Ca.* E. dacicola, since its genomic architecture remains unresolved: The operon

appears in a separate scaffold in all three available assemblies (e.g. IL assembly accession

number: LJAM02000124.1), none of which have been further extended and ordered with the

use of long insert-size libraries. MGEs are frequently associated with fragmented

assemblies, and in the ErwSC assembly, we detected a transposase fragment upstream of

the first gene of the operon, *ureD*, in the scaffold containing the urease. However, mate-pair

library sequencing data from Cretan olive fruit fly guts (accession number: SRX1896451) did

not support the presence of the transposase upstream of *ureD*, and did not allow

identification of the operon's flanking scaffolds.

The present data do not allow us to identify the date of the acquisition of the urease operon,

nor its taxonomic origin. The observation that the GC content of the operon is distinct from

that of its host argues in favor of a relatively recent acquisition, as previously proposed for a

different *Ca.* E. dacicola GEI (Estes et al. 2018), but one cannot conclude without dating the

symbiosis. Similarly, regarding the donor, the low bootstrap support for the clades that link

the groups comprising *Ca.* E. dacicola, *G. quercinecans* and *Brenneria* to the rest of the

clades indicates that the original donor of the urease cluster might be extinct or not yet

sequenced. One possibility is that it is a free-living bacterium encountered by *Ca.* E. dacicola

before its transition to obligacy with *B. oleae* or a yet-uncharacterized gut bacterium. In line

with the latter, there is abundant evidence that members of gut microbial communities in

vertebrates and invertebrates undergo HGT, and that this can subsequently influence

functional traits (Petridis et al. 2006; Smillie et al. 2011; Stecher et al. 2012; Baker et al.

2018). Linked to HGT events, mobile genetic elements (MGEs), including transposases,

repetitive DNA regions and phages, are abundant in all three *Ca.* E. dacicola assemblies:

MGEs represent an average 34 % of the obligate symbiont's genome, in contrast to the

corresponding MGE content in *Tatumella* sp. TA1 and *Enterobacter* sp. OLF (Table S4).

However, these percentages should not to be taken at absolute value due to the inherent

difficulties in computational annotation of MGEs, especially from metagenomic samples

(Jørgensen et al. 2014), while they might represent an overestimate due to the draft - and

often fragmented - nature of the assemblies. Still, it is notable that high transposase content

has been often observed in bacteria that recently adopted a symbiotic lifestyle (Gil et al.

2008; Ran et al. 2010), while MGEs were previously shown to be transcriptionally active in

B. oleae larvae feeding on green and black olives (Pavlidi et al. 2017). These findings overall

support the idea that genetic exchange between *Ca.* E. dacicola and members of the *B.*

oleae gut microbial community may occur.

Genes encoding extracellular surface structure components in the *B. oleae* gut

microbiota

Exchanges of genetic material between bacteria are facilitated by extracellular surface

structures (Thomas and Nielsen 2005), components of which are encoded in the genomes

of the *B. oleae* gut microbiota: Components of the IncF transfer system of conjugative

plasmids, encoding F-like pili, are present in the two most complete *Ca.* E. dacicola

assemblies, IL and Oroville (Table S5). In addition, the chromosome of *Tatumella* sp. TA1

encodes a Trb operon for conjugative transfer, potentiating the exchange of genetic material

(Table S5). Notably, plasmid pTA1 from *Tatumella* sp. TA1, identified in the present study

(GenBank accession number: CP033728), is present in the *Ca.* E. dacicola Oroville

assembly, which was generated from *B. oleae* populations where *Tatumella* sp. TA1 has not

yet been detected. This result, most likely not due to 'contamination' as confirmed by the

absence of 16S rRNA gene sequences from taxa other than *Ca.* E. dacicola in the Oroville

assembly (data not shown), indicates that transfers of genetic material among members of

the *B. oleae* gut microbiota occur.

Some extracellular surface structures that mediate the exchange of genetic material

between bacteria, such as type IV secretion systems (T4SS), also facilitate adhesion and

molecular exchange with eukaryotic cells (Alvarez-Martinez and Christie 2009). T4SS are

diverse; VirB-like T4SS encode short and rigid pili, whereas the F-pili T4SS for conjugative

transfer tend to be long and flexible (Clarke et al. 2008). Depending on their structure, pili

can deliver a range of effector molecules including proteins and nucleic acids to both

prokaryotic and eukaryotic cells, and can also be involved in the transition between motility

and sessility, for example in the process of biofilm formation (Christie and Vogel 2000).

T4SS mediate the delivery of effector proteins directly to the cytoplasm of eukaryotic cells

during the process of infection in pathogenic bacteria (Cascales and Christie 2003), and root

nodule colonization in symbiotic *Rhizobia* (Deakin and Broughton 2009). Effector proteins

can induce regulatory changes in the eukaryotic cell, triggering modification of cellular

conditions or conditions in the environment to facilitate growth or invasion (Burns 2003). A

conserved VirB-like T4SS operon encoding the essential genes for pilus formation and

substrate trafficking (VirB1-11 and VirD4, (Wallden et al. 2010)) was detected in all three *Ca.*

E. dacicola assemblies, but was not detected in either *Enterobacter* sp. OLF or *Tatumella*

sp. TA1 (Table S5). Evidence for expression of the *Ca.* E. dacicola VirB-like T4SS operon

during *B. oleae* juvenile development in ripening olives (Pavlidi et al. 2017) and for the

presence of pili-like structures in the esophageal bulb (Poinar et al. 1975) suggest that the

T4SS system pili may also play a - yet unknown- role in the *B. oleae* - *Ca.* E. dacicola

symbiosis, for example by the establishment of *Ca.* E. dacicola biofilms (Estes et al. 2009).

One potential function to explore in future studies might be the coordination of gut lumen

colonization and subsequent resource exchange between the host and symbiont.

Conclusions

In this study, we aimed to gain understanding on the advantages of symbiosis for the olive

fruit fly *B. oleae* by integrating various sources of genetic information on its microbial gut

community. This led to the identification, by 16S rRNA gene sequencing and culture-

dependent methods, of a novel facultative member of the gut microbiota, *Tatumella* sp. TA1,

which exclusively associates with Mediterranean populations of *B. oleae* throughout the

lifecycle. *Tatumella* sp. TA1 is phylogenetically distinct from the US-restricted facultative

symbiont *Enterobacter* sp. OLF, highlighting the population-dependent nature of gut

microbiota composition in the olive fruit fly. Comparative genomics indicated that the obligate

symbiont *Ca.* E. dacicola, as well as *Tatumella* sp. TA1 and *Enterobacter* sp. OLF, which are

all stable components of the *B. oleae* microbiota throughout the olive fruit fly lifecycle,

encode genes that allow the use of urea as a nitrogen source. The hydrolysis of urea to

ammonia is encoded by genes with distinct phylogenetic origins in each organism: HGT of a

urease operon in *Ca.* E. dacicola, an endogenous urease operon in *Enterobacter* sp. OLF,

and the presence of alternative enzymatic machinery (urea carboxylase and allophanate

hydrolase) in *Tatumella* sp. TA1. These findings provide a potential mechanistic basis for

previous experimental evidence of gut microbiota-mediated dietary nitrogen provisioning

during adulthood, but emphasize the need for experimental validation of metabolic cross-

feeding between the host and obligate symbiont. A hypothesis in this direction, warranting

focus in future studies, is that a VirB-like T4SS encoded by *Ca.* E. dacicola, but missing from

the genomes of *Enterobacter* sp. OLF and *Tatumella* sp. TA1, may facilitate specific

interactions with the host at the symbiotic interface. Besides T4SS, our study further

highlighted extracellular surface structures, encoded in the genomes of the obligate and the

facultative symbionts, as mediators of DNA transfer: Detection of a plasmid shared by

geographically-distinct *Ca.* E. dacicola and *Tatumella* sp. TA1, along with the horizontal

acquisition of genes important to symbiosis function (urease in this study, genes related to

amino-acid degradation in (Estes, Hearn, Agrawal, et al. 2018)) and the abundance of

mobile genetic elements in the obligate symbiont genome, indicate that previous and on-

going genetic exchanges between gut community members are important determinants of

symbiotic interactions with *B. oleae*.

Materials and Methods

Insect material

B. oleae adults were obtained by collecting infested olives from trees in the grounds of the

University of Crete (Heraklion, Greece) in October and November of 2014. Infested olives

were suspended over sterile sand with a wire mesh, and third-instar larvae were allowed to

emerge from the fruit. Flies pupated in the sand and were placed into Petri dishes in 10 cm^3

plastic cages prior to emergence. Flies were maintained at 25 ^{o}C and 60 % relative humidity

and were supplied with artificial diet (19 % hydrolyzed yeast, 75 % icing sugar, 6 % egg

yolk). Each cage was provided with Milli-Q water in a clean plastic container and wax cones

for oviposition.

Preparation and sequencing of 16S rRNA gene amplicon libraries

DNA was extracted from a total of 52 whole *B. oleae* adults using the Qiagen DNeasy Blood

and Tissue kit for Gram-positive bacteria (Qiagen, UK) following the manufacturer's

instructions. An additional bead-beating step using 3 mm carbide beads (Qiagen, UK) in a

Qiagen tissue lyzer (Qiagen, UK) at 25 Hz for 30 seconds was employed. In order to assess

the diversity and composition of the bacterial communities, bacterial 16S rRNA V4 regions

were amplified by PCR using the universal primers F515 (5'-GTGCCAGCMGCCGCGGTAA-

3') and R806 (5'- GGACTACHVGGGTWTCTAAT-3') (Caporaso et al. 2011). Samples were

dual-indexed for sequencing following the method in D'Amore et al. (2016). PCR reactions

were performed in a total volume of 20 µl, containing 5 ng of template DNA, 10 µl NEBNext

2x High-Fidelity Master Mix (New England Biolabs), 0.3 µM of each primer, and 3.4 µl PCR-

clean water. Thermal cycling conditions were 98 °C for 2 minutes, 10 cycles of 98 °C for 20

seconds, 60 °C for 15 seconds, and 70 °C for 30 seconds, with a final extension at 72 °C for

5 minutes. PCR products were purified with Agencourt AMPure XP beads (Beckman Coulter

Genomics) and used as template for the second PCR reaction. Purified first-round PCR

products were combined with 10 µl NEBNext 2x High-Fidelity Master Mix (New England

Biolabs), and 0.3 µM of each barcoding primer containing adapters and indexes to a total

volume of 20 µl. Thermal cycling conditions were 98 °C for 2 minutes, 15 cycles of 98 °C for

20 seconds, 55 °C for 15 seconds and 72 °C for 40 seconds, with a final extension at 72 °C

for 60 seconds. PCR products were purified with Agencourt AMPure XP beads (Beckman

Coulter Genomics) and quantified with the Qubit dsDNA High-Sensitivity assay (Life

Technologies), and an Agilent Bioanalyzer High-Sensitivity DNA chip (Agilent). Samples

were pooled at equimolar concentrations and size-selected in a range of 350-450 bp by

Pippin-Prep (Sage Science). Sequencing was performed at the University of Liverpool

Centre for Genomic Research on an Illumina MiSeq platform with V2 chemistry, generating

250 bp paired-end reads. All raw sequencing reads were deposited at NCBI under the

BioProject accession PRJNA321174.

Computational analyses of 16S rRNA sequencing data

Raw sequencing reads were de-multiplexed and converted to FASTQ format using CASAVA

version 1.8 (Illumina 2011). Cutadapt version 1.2.1 (Martin 2011) was used to trim Illumina

adapter sequences from FASTQ files. Reads were trimmed if 3 bp or more of the 3' end of a

read matched the adapter sequence. Sickle version 1.200 (Joshi & Fass 2011) was used to

trim reads based on quality: any reads with a window quality score of less than 20, or were

less than 10 bp long after trimming, were discarded. BayesHammer was used to correct

reads based on quality (Nikolenko et al. 2013). Paired-end reads were merged with a

minimum overlap of 50 bp using PandaSeq (Masella et al. 2012). All subsequent analyses

were conducted in QIIME version 1.8.0 (Caporaso, Kuczynski, et al. 2010). Sequences were

clustered into Operational Taxonomic Units (OTUs) by *de novo* OTU picking with USEARCH

(Edgar 2010). Chimeras were detected and omitted with UCHIME (Edgar et al. 2011) and

the QIIME-compatible version of the SILVA 111 release database (Quast et al. 2013). The

most abundant sequence was chosen as the representative for each OTU, and taxonomy

was assigned to representative sequences by BLAST (Altschul et al. 1990) against the

SILVA 111 release database, which was supplemented with several reference sequences

for *Ca.* E. dacicola. OTUs were filtered from the dataset if they matched the SILVA 111

database for chloroplasts or mitochondria. OTU representative sequences were aligned

against the Greengenes core reference alignment (DeSantis et al. 2006) using PyNAST

(Caporaso, Bittinger, et al. 2010). Previously published 16S rRNA gene amplicons from flies

collected in Israel (Ben-Yosef et al. 2015) were employed for comparison with the above

data. Since data generation methods for the samples from Israel varied slightly, all data

analysis methods were standardized from read-error correction onwards in this study.

Tatumella sp. TA1 prevalence was calculated as the proportion of individuals where

Tatumella sp. TA1 16S rRNA was > 0.01% of the total 16S rRNA gene copies from that

sample, and relative abundance was the proportion of the total 16S rRNA gene copies.

Isolation, culture and identification of *Tatumella* sp. TA1 from *B. oleae*

Tatumella sp. TA1 was identified from a two-day old male fly following the below isolation

and culture procedures, all conducted under sterile conditions using aseptic technique or in

a laminar flow cabinet. Individual two-day old adult *B. oleae* collected in Crete were surface-

sterilized with 70 % ethanol and rinsed twice in distilled water prior to homogenization with a

plastic pestle in 20 µl nuclease-free water. 10 µl of the homogenate was spotted on to

Columbia agar (Oxoid, UK) supplemented with 5 % defibrinated horse blood (TCS

Biosciences, UK), and plates were incubated at 25 $^{\circ}$C for 48 h. Single colonies were picked

and streaked onto Brain Heart Infusion (BHI) agar (Oxoid, UK) to establish pure cultures.

Pure liquid cultures were generated by inoculating single colonies from BHI agar plates in to

BHI liquid medium and incubating them at 25 $^{\circ}$C for 48 h. Liquid cultures were cryopreserved

following the addition of a cryoprotectant (20 % [v/v] glycerol final concentration) and storage

at -80 $^{\circ}$C. In order to identify isolates, single colonies from BHI agar plates were picked and

inoculated into 10 µl nuclease-free water in a PCR tube. Tubes were incubated at 95 $^{\circ}$C for

5 minutes to lyse cells and isolate DNA. A 1500 bp region of the 16S rRNA gene was

amplified with universal primers 8F (5'–AGAGTTTGATCMTGGCTCAG–3') and 1492R

(5'-CCCCTACGGTTACCTTGTTACGAC–3'). Reactions were performed in a total volume of

25 µl containing 12.5 µl 2 x MyTaq Red (Bioline, UK), 0.5 µl of each 10 µM primer stock, 1 µl

template DNA, and 10.5 µl nuclease-free water. Thermal cycling conditions were 95 $^{\circ}$C for 5

minutes, 30 cycles of 95 $^{\circ}$C for 30 seconds, 56 $^{\circ}$C for 45 seconds, 72 $^{\circ}$C for 90 seconds, and

a final extension at 72 $^{\circ}$C for 7 minutes. PCR products were Sanger sequenced with forward

primer 8F by GATC (GATC, Cologne) and resulting sequences were subjected to BLAST

analysis (Altschul et al. 1990) against the GenBank database (http://www.ncbi.nlm.nih.gov/)

to allow taxonomic identification by similarity.

Library preparation and sequencing of the *Tatumella* sp. TA1 genome

DNA from *Tatumella* sp. TA1 was extracted as follows: Cryopreserved isolates were revived

by streaking on to BHI agar and incubated at 25 $^{\circ}$C for 72h. Single colonies were inoculated

into BHI broth and incubated at 25 $^{\circ}$C until cultures reached an OD600 of 0.3. Cultures were

pelleted by centrifugation at 6,000 x g for 6 minutes. The supernatant was removed, and

cells were resuspended in DNA elution buffer at a concentration of 1 x 10^5 Colony Forming

Units (CFUs) mL^{-1}. DNA was extracted using the Zymo Quick DNA Universal Kit (Zymo, UK)

following the manufacturer's instructions for biological fluids and cells with the following

amendments to the protocol: samples were incubated with proteinase K at 55 $^{\circ}$C for 30

minutes rather than 10 minutes. DNA was purified with Ampure beads (Agencourt) at a 1:1

ratio, and stored at 4 $^{\circ}$C until library preparation. DNA was sheared to 10 kb using Covaris

G-tubes following the manufacturer's guidelines, and library preparation was performed with

the SMRTbell library preparation kit (Pacific Biosciences) following the manufacturer's

instructions. The Qubit dsDNA HS assay (Life Technologies, UK) was used to quantify the

library, and the average fragment size was determined using the Agilent Bioanalyzer HS

assay (Agilent). Size selection was performed with the Blue Pippin Prep (Sage Science)

using a 0.75 % agarose cassette and the S1 marker. The final SMRT bell was purified with

Agencourt AMPure XP beads (Beckman Coulter Genomics) and quantified with the Qubit

dsDNA High-Sensitivity assay (Life Technologies), and an Agilent Bioanalyzer High-

Sensitivity DNA chip (Agilent). The SMRTbell library was annealed to sequencing primers at

values predetermined by the Binding Calculator (Pacific Biosciences), and sequencing was

performed on two SMRT cells using 360-minute movie times. Pacific Biosciences

sequencing and library preparation of the *Tatumella* sp. TA1 isolate was performed at the

University of Liverpool Centre for Genomic Research on a Pacific Biosciences RS II

sequencer.

Genome assembly and annotation

The *Tatumella* sp. TA1 draft genome was assembled from 365,445 sub-reads with a mean

length of 6,926 bp using the Hierarchical Genome Assembly Process (HGAP) workflow

(Chin et al. 2013). It is available at NCBI under the accession numbers CP033727-

CP033728. The HGAP pipeline comprised pre-assembly error correction of sub-reads based

on read length and quality, assembly with Celera, and assembly polishing with Quiver. The

assembled genome was annotated with PROKKA version 1.5.2 (Seemann 2014). To further

investigate the extracellular membrane structures and mobile elements encoded in the

Enterobacter sp. OLF, *Tatumella* sp. TA1 and three *Ca.* E. dacicola draft assemblies,

genomes were re-annotated with RAST (Overbeek et al. 2014). These annotations are

available in Table S3.

Phylogenetic analyses

For the species tree, allowing phylogenetic placement of *Tatumella* sp. TA1 and Ca. *E.*

dacicola, taxa were selected based on (Palmer et al. 2017), which presents a robust

phylogeny of the *Erwinia*, *Pantoea* and *Tatumella* genera using genome-wide orthologs.

Where Palmer et al. (2017) included several conspecific taxa, we chose the most

biologically-relevant of the identical taxa. The Western Flower Thrip (WFT)-associated

bacteria BFo1 and BFo2 were included in order to validate phylogenetic similarities with *Ca.*

E. dacicola and *Tatumella* TA1 based on 16S rRNA sequences. A full list of the 46 genomes

employed for this analysis can be found in Table S1A, including the 6 outgroup taxa and the

3 available *Ca.* E. dacicola draft genome assemblies (ErwSC, IL and Oroville). Genomes

were annotated with PROKKA version 1.5.2 using the default settings, and 289 single-copy

orthologs present in all taxa were identified using OrthoMCL version 1.4 with default

parameters (Li et al. 2003). The 'reference' tree for the urease operon analysis was

estimated from 8 genes randomly selected from the set of 289 single-copy orthologs, with a

total of 5,572 informative sites; we assumed neutral evolution status for these genes based

on their predicted housekeeping functions (Table S1A). For the urease subunit alpha (*ureC*)

gene tree, homologs were retrieved based on BlastP queries against the nr database at

NCBI (restricted to taxid=Bacteria). We included as many taxa common to the reference and

species tree as possible, to ensure meaningful comparisons. Additional homologs were

retrieved by targeted searches in RefSeq (full list of species names and accession numbers

in Table S1B). For all trees, amino acid sequences were aligned with MUSCLE version

3.8.31 (Edgar 2004), and informative sites were selected with Gblocks version 091b

(Castresana 2000). Maximum-Likelihood trees were estimated using IQ-TREE version 1.5.5

after automated model selection (Nguyen et al. 2015), with 300 random trees as burn-in.

Node support was calculated using 1000 ultra-fast bootstraps (Minh et al. 2013). The multi-

locus coalescent-based species and reference trees were estimated from individual ortholog

trees using Astral III, with node support calculated using 1000 bootstraps (Zhang et al.

2018). Trees were rooted and visualized using FigTree v.1.4.0

(http://tree.bio.ed.ac.uk/software/figtree/).

Statistical analyses

All statistical analyses were performed in R version 3.3.3 (R Core Team 2017). Comparisons

of GC content mean ranks, means and distributions were performed with Mann-Whitney,

Welch's t-test (allowing for comparison of unequal variance-samples) and Kolmogorov-

Smirnov tests. For these comparisons, the GC content of the urease operon and the whole

genome of *Ca.* E. dacicola (ErwSC assembly) were calculated using overlapping 50 bp

windows.

Data availability

16S rRNA amplicon sequencing data are available as part of NCBI BioProject

PRJNA321174. The *Tatumella* sp. TA1 genome is available at NCBI under accession

numbers CP033727-CP033728. Alignments for Figs 1 and 3 are available upon request

from the corresponding authors.

Acknowledgements

This work was supported by a BBSRC Industrial CASE Award BB/K501773/1 to ACD, awarded by the Biosciences Knowledge Transfer Network to the Food, Industrial Bioscience and Plants and crops Sectors. This work was further supported by Greek national funds through the Public Investments Program (PIP) of the General Secretariat for Research & Technology (GSRT), under the Emblematic Action "The Olive Road" (project code: 2018ΣE01300000), awarded to JV. The authors are grateful to Greg Blow and Michael Capper for reading drafts of the manuscript, and to Paschalia Kapli and Pavlos Pavlidis for discussions and technical advice on horizontal gene transfer and phylogenetic trees.

Author Contributions

F.B., A.G., J.V. and A.C.D designed the study. F.B. and A.K. collected the samples. F.B. and I.B.G. prepared the sequencing libraries. F.B., A.G., I.B.G. M.K. and A.C.D. performed data analysis. F.B., A.G. and A.C.D. wrote the manuscript, and all authors read and made comments on the manuscript prior to submission.

Figure and Table legends

Table 1. Prevalence of *Tatumella* sp. TA1 in Mediterranean populations of *B. oleae*.

Fig. 1. Maximum-Likelihood tree estimated from amino acid alignments of 289 single-copy orthologs retrieved from whole genome assemblies of 39 *Erwinia, Pantoea* and *Tatumella* and 7 outgroup taxa. Full names and accession numbers of the genes used are available in Table S1A. Numbers on nodes correspond to bootstrap support (only values <100 are shown).

Fig. 2. Proposed pathways for urea hydrolysis and recycling of nitrogen into host and microbial metabolic pathways via ammonia, glutamine and glutamate. The genes encoding the illustrated enzymes were detected in genome assemblies of *Ca.* E. dacicola (Blow, Gioti, et al. 2016), *Tatumella* sp. TA1 (this study), both annotated with PROKKA, and *Enterobacter* sp. OLF annotated as described in (Estes, Hearn, Nadendla, et al. 2018). GS (Glutamine synthetase E.C. 6.3.1.2); CPS (Carbamoyl-phosphate synthase E.C. 6.3.4.16; E.C. 6.3.5.5); NAGS (N-acetylglutamate synthase E.C. 2.3.1.1); AGK (Acetylglutamate kinase E.C. 2.7.2.8); NAGPR (N-acetyl-gamma-glutamyl-phosphate reductase E.C. 1.2.1.38); AOAT (Acetylornithine/succinyldiaminopimelate aminotransferase E.C. 2.6.1.11); AODA (Acetylornithine deacetylase E.C. 3.5.1.16); OCT (Ornithine carbamoyltransferase E.C. 2.1.3.3); ASS (Argininosuccinate synthase E.C. 6.3.4.5); ASL (Argininosuccinate lyase E.C. 4.3.2.1); ARG (Arginase E.C. 3.5.3.1); UC (Urea carboxylase E.C. 6.3.4.6); AH (Allophanate hydrolase E.C. 3.5.1.54); URE (Urease E.C. 3.5.1.5).

Fig. 3. Maximum-Likelihood trees estimated from amino acid alignments of (*A*) the urease subunit alpha (*ureC*) gene and (*B*) 8 single-copy orthologs, representing a 'neutral reference'. Rooting of both trees was performed with the most phylogenetically remote taxon, the Alphaproteobacterium *Sinorhizobium meliloti*. Full names and accession numbers for the genes (*A*) and genomes (*B*) used are available in Table S1B and Table S1A, respectively. Numbers on nodes correspond to bootstrap support.

Supplementary material

Fig. S1. Hypothesized microbial nitrogen recycling pathways from host waste products. UO (Uricase E.C. 1.7.3.3); ALN (Allantoinase E.C. 3.5.2.5); ALC (Allantoicase E.C. 3.5.3.4); UL (Ureidoglycolate lyase E.C. 4.3.2.3); URE (Urease 3.5.1.5); GS (Glutamine synthetase E.C. 6.3.1.2).

Table S1. GenBank accession numbers and metadata for sequences used in phylogenetic analyses. (*A*) Whole genome sequences used for ortholog picking and phylogenetic reconstruction of the species and reference trees (*B*) Amino acid sequences used for the *ureC* gene tree.

Table S2. Presence and expression of genes related to nitrogen assimilation in *Ca.* E. dacicola and *Tatumella* sp. TA1. *(A)* Urea hydrolysis and nitrogen recycling pathways *(B)* Amino acid biosynthetic pathways.

Table S3. Functional annotations and corresponding sequences for the genomes of *Ca.* E. dacicola (three currently available assemblies), *Tatumella* sp. TA1 and *Enterobacter* sp. OLF, as well as for the transcriptome of *Ca.* E. dacicola (ribosomal RNAs excluded). The annotations come from RAST subsystems.

Table S4. Abundance of different categories of mobile genetic elements in the *Ca.* E. dacicola (three currently available assemblies), *Tatumella* sp. TA1 and *Enterobacter* sp. OLF genomes, as annotated by RAST subsystems.

Table S5. Functional annotation of *Ca.* E. dacicola, *Tatumella* sp. TA1 and *Enterobacter* sp. OLF genomes for extracellular surface structure components, by assignment of genes to the KEGG category "Membrane Transport" using RAST subsystems.

Table 1. Prevalence of *Tatumella* sp. TA1 in Mediterranean populations of *B. oleae*.

Population	Lifestage	No. individuals	Prevalence	% 16S rRNA gene copies	
				Mean	Max.
Crete	Adults	52	26.9%	0.3%	7.4%
Israel	Adults	9	88.9%	1%	3.5%
	Third-instar larvae	10	90%	3.5%	14.5%

References

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. J. Mol. Biol. 215:403–410.

Alvarez-Martinez CE, Christie PJ. 2009. Biological diversity of prokaryotic type IV secretion systems. Microbiol. Mol. Biol. Rev. 73:775–808.

Baker KS, Dallman TJ, Field N, Childs T, Mitchell H, Day M, Weill F-X, Lefèvre S, Tourdjman M, Hughes G, et al. 2018. Horizontal antimicrobial resistance transfer drives epidemics of multiple Shigella species. Nat. Commun. 9:1462.

Ben-Yosef M, Aharon Y, Jurkevitch E, Yuval B. 2010. Give us the tools and we will do the job: symbiotic bacteria affect olive fly fitness in a diet-dependent fashion. Proc. Biol. Sci. 277:1545–1552.

Ben-Yosef M, Pasternak Z, Jurkevitch E, Yuval B. 2014. Symbiotic bacteria enable olive flies (Bactrocera oleae) to exploit intractable sources of nitrogen. J. Evol. Biol. 27:2695–2705.

Ben-Yosef M, Pasternak Z, Jurkevitch E, Yuval B. 2015. Symbiotic bacteria enable olive fly larvae to overcome host defences. Royal Society Open Science 2:150170.

Blow F, Gioti A, Starns D, Ben-Yosef M, Pasternak Z, Jurkevitch E, Vontas J, Darby AC. 2016. Draft Genome Sequence of the Bactrocera oleae Symbiont "Candidatus Erwinia dacicola." Genome Announc. [Internet] 4. Available from: http://dx.doi.org/10.1128/genomeA.00896-16

Blow F, Vontas J, Darby AC. 2016. Draft Genome Sequence of Stenotrophomonas maltophilia SBo1 Isolated from Bactrocera oleae. Genome Announc. [Internet] 4. Available from: http://dx.doi.org/10.1128/genomeA.00905-16

Blow F, Vontas J, Darby AC. 2017. Draft Genome Sequence of Chryseobacterium Strain CBo1 Isolated from Bactrocera oleae. Genome Announc. [Internet] 5. Available from: http://dx.doi.org/10.1128/genomeA.00177-17

Burns DL. 2003. Type IV transporters of pathogenic bacteria. Curr. Opin. Microbiol. 6:29–34.

Caporaso JG, Bittinger K, Bushman FD, DeSantis TZ, Andersen GL, Knight R. 2010. PyNAST: a flexible tool for aligning sequences to a template alignment. Bioinformatics 26:266–267.

Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, Fierer N, Peña AG, Goodrich JK, Gordon JI, et al. 2010. QIIME allows analysis of high-throughput community sequencing data. Nat. Methods 7:335–336.

Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Lozupone CA, Turnbaugh PJ, Fierer N, Knight R. 2011. Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample. Proc. Natl. Acad. Sci. U. S. A. 108 Suppl 1:4516–4522.

Capuzzo C, Firrao G, Mazzon L, Squartini A, Girolami V. 2005. "Candidatus Erwinia dacicola", a coevolved symbiotic bacterium of the olive fly Bactrocera oleae (Gmelin). Int. J. Syst. Evol. Microbiol. 55:1641–1647.

Cascales E, Christie PJ. 2003. The versatile bacterial type IV secretion systems. Nat. Rev. Microbiol. 1:137–149.

Castresana J. 2000. Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. Mol. Biol. Evol. 17:540–552.

Chandler JA, James PM, Jospin G, Lang JM. 2014. The bacterial communities of Drosophila suzukii collected from undamaged cherries. PeerJ 2:e474.

Chin C-S, Alexander DH, Marks P, Klammer AA, Drake J, Heiner C, Clum A, Copeland A, Huddleston
J, Eichler EE, et al. 2013. Nonhybrid, finished microbial genome assemblies from long-read
SMRT sequencing data. Nat. Methods 10:563–569.

Christie PJ, Vogel JP. 2000. Bacterial type IV secretion: conjugation systems adapted to deliver
effector molecules to host cells. Trends Microbiol. 8:354–360.

Clarke M, Maddera L, Harris RL, Silverman PM. 2008. F-pili dynamics by live-cell imaging. Proc. Natl.
Acad. Sci. U. S. A. 105:17978–17981.

Deakin WJ, Broughton WJ. 2009. Symbiotic use of pathogenic strategies: rhizobial protein secretion
systems. Nat. Rev. Microbiol. 7:312–320.

DeSantis TZ, Hugenholtz P, Larsen N, Rojas M, Brodie EL, Keller K, Huber T, Dalevi D, Hu P,
Andersen GL. 2006. Greengenes, a chimera-checked 16S rRNA gene database and workbench
compatible with ARB. Appl. Environ. Microbiol. 72:5069–5072.

Dillon RJ, Dillon VM. 2004. The gut bacteria of insects: nonpathogenic interactions. Annu. Rev.
Entomol. 49:71–92.

Djambazian H, Bayega A, Tsoumani KT, Sagri E, Gregoriou M-E, Giorda K, Tsiamis G, Bourtzis K,
Oikonomopoulos S, Dewar K, et al., unpublished data,
https://www.biorxiv.org/content/biorxiv/early/2018/12/26/505040, last accessed March 27, 2019.

Douglas AE. 2009. The microbial dimension in insect nutritional ecology. Funct. Ecol. 23:38–47.

Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput.
Nucleic Acids Res. 32:1792–1797.

Edgar RC. 2010. Search and clustering orders of magnitude faster than BLAST. Bioinformatics
26:2460–2461.

Edgar RC, Haas BJ, Clemente JC, Quince C, Knight R. 2011. UCHIME improves sensitivity and
speed of chimera detection. Bioinformatics 27:2194–2200.

Estes AM, Hearn DJ, Agrawal S, Pierson EA, Dunning Hotopp JC. 2018. Comparative genomics of
the Erwinia and Enterobacter olive fly endosymbionts. Sci. Rep. 8:15936.

Estes AM, Hearn DJ, Bronstein JL, Pierson EA. 2009. The Olive Fly Endosymbiont, "Candidatus
Erwinia dacicola," Switches from an Intracellular Existence to an Extracellular Existence during
Host Insect Development. Appl. Environ. Microbiol. 75:7097–7106.

Estes AM, Hearn DJ, Burrack HJ, Rempoulakis P, Pierson EA. 2012. Prevalence of Candidatus
Erwinia dacicola in wild and laboratory olive fruit fly populations and across developmental
stages. Environ. Entomol. 41:265–274.

Estes AM, Hearn DJ, Nadendla S, Pierson EA, Dunning Hotopp JC. 2018. Draft Genome Sequence
of Enterobacter sp. Strain OLF, a Colonizer of Olive Flies. Microbiol Resour Announc [Internet] 7.
Available from: http://dx.doi.org/10.1128/MRA.01068-18

Facey PD, Méric G, Hitchings MD, Pachebat JA, Hegarty MJ, Chen X, Morgan LVA, Hoeppner JE,
Whitten MMA, Kirk WDJ, et al. 2015. Draft Genomes, Phylogenetic Reconstruction, and
Comparative Genomics of Two Novel Cohabiting Bacterial Symbionts Isolated from Frankliniella
occidentalis. Genome Biol. Evol. 7:2188–2202.

Feldhaar H, Straka J, Krischke M, Berthold K, Stoll S, Mueller MJ, Gross R. 2007. Nutritional
upgrading for omnivorous carpenter ants by the endosymbiont Blochmannia. BMC Biol. 5:48.

Gil R, Belda E, Gosalbes MJ, Delaye L, Vallier A, Vincent-Monégat C, Heddi A, Silva FJ, Moya A,
Latorre A. 2008. Massive presence of insertion sequences in the genome of SOPE, the primary
endosymbiont of the rice weevil Sitophilus oryzae. Int. Microbiol. 11:41–48.

Hollis DG, Hickman FW, Fanning GR, Farmer JJ 3rd, Weaver RE, Brenner DJ. 1981. Tatumella
ptyseos gen. nov., sp. nov., a member of the family Enterobacteriaceae found in clinical
specimens. J. Clin. Microbiol. 14:79–88.

Jørgensen TS, Kiil AS, Hansen MA, Sørensen SJ, Hansen LH. 2014. Current strategies for mobilome
research. Front. Microbiol. 5:750.

Joshi N, Fass JN. 2011. Sickle-A windowed adaptive trimming tool for FASTQ files using quality.
Online publication. https://github. com/najoshi/sickle

Juhas M, van der Meer JR, Gaillard M, Harding RM, Hood DW, Crook DW. 2009. Genomic islands:
tools of bacterial horizontal gene transfer and evolution. FEMS Microbiol. Rev. 33:376–393.

Kashima T, Nakamura T, Tojo S. 2006. Uric acid recycling in the shield bug, Parastrachia japonensis
(Hemiptera: Parastrachiidae), during diapause. J. Insect Physiol. 52:816–825.

Kounatidis I, Crotti E, Sapountzis P, Sacchi L, Rizzi A, Chouaia B, Bandi C, Alma A, Daffonchio D,
Mavragani-Tsipidou P, et al. 2009. Acetobacter tropicalis Is a Major Symbiont of the Olive Fruit
Fly (Bactrocera oleae). Appl. Environ. Microbiol. 75:3281–3288.

Li L, Stoeckert CJ Jr, Roos DS. 2003. OrthoMCL: identification of ortholog groups for eukaryotic
genomes. Genome Res. 13:2178–2189.

Marín-Cevada V, Caballero-Mellado J, Bustillos-Cristales R, Muñoz-Rojas J, Mascarúa-Esparza MA,
Castañeda-Lucio M, López-Reyes L, Martínez-Aguilar L, Fuentes-Ramírez LE. 2010. Tatumella
ptyseos, an unrevealed causative agent of pink disease in pineapple. Journal of phytopathology
158:93–99.

Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet. journal 17:-10.

Masella AP, Bartram AK, Truszkowski JM, Brown DG, Neufeld JD. 2012. PANDAseq: paired-end
assembler for illumina sequences. BMC Bioinformatics 13:31.

Minh BQ, Nguyen MAT, von Haeseler A. 2013. Ultrafast approximation for phylogenetic bootstrap.
Mol. Biol. Evol. 30:1188–1195.

Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. 2015. IQ-TREE: a fast and effective stochastic
algorithm for estimating maximum-likelihood phylogenies. Mol. Biol. Evol. 32:268–274.

Nikolenko SI, Korobeynikov AI, Alekseyev MA. 2013. BayesHammer: Bayesian clustering for error
correction in single-cell sequencing. BMC Genomics 14 Suppl 1:S7.

Overbeek R, Olson R, Pusch GD, Olsen GJ, Davis JJ, Disz T, Edwards RA, Gerdes S, Parrello B,
Shukla M, et al. 2014. The SEED and the Rapid Annotation of microbial genomes using
Subsystems Technology (RAST). Nucleic Acids Res. 42:D206–D214.

Palmer M, Steenkamp ET, Coetzee MPA, Chan W-Y, van Zyl E, De Maayer P, Coutinho TA, Blom J,
Smits THM, Duffy B, et al. 2017. Phylogenomic resolution of the bacterial genus Pantoea and its
relationship with Erwinia and Tatumella. Antonie Van Leeuwenhoek 110:1287–1309.

Pavlidi N, Gioti A, Wybouw N, Dermauw W, Ben-Yosef M, Yuval B, Jurkevich E, Kampouraki A, Van
Leeuwen T, Vontas J. 2017. Transcriptomic responses of the olive fruit fly Bactrocera oleae and
its symbiont Candidatus Erwinia dacicola to olive feeding. Sci. Rep. 7:42633.

Petersen TN, Brunak S, von Heijne G, Nielsen H. 2011. SignalP 4.0: discriminating signal peptides
from transmembrane regions. Nat. Methods 8:785–786.

Petridis M, Bagdasarian M, Waldor MK, Walker E. 2006. Horizontal transfer of Shiga toxin and
antibiotic resistance genes among Escherichia coli strains in house fly (Diptera: Muscidae) gut. J.
Med. Entomol. 43:288–295.

Poinar GO Jr, Hess RT, Tsitsipis JA. 1975. Ultrastructure of the Bacterial Symbiotes in the
Pharyngeal Diverticulum of Dacus oleae (Gmelin) (Trypetidae; Diptera). Acta Zool. 56:77–84.

Potrikus CJ, Breznak JA. 1981. Gut bacteria recycle uric acid nitrogen in termites: A strategy for
nutrient conservation. Proc. Natl. Acad. Sci. U. S. A. 78:4601–4605.

Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glöckner FO. 2013. The
SILVA ribosomal RNA gene database project: improved data processing and web-based tools.
Nucleic Acids Res. 41:D590–D596.

R Core Team (2017). R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Ran L, Larsson J, Vigil-Stenman T, Nylander JAA, Ininbergs K, Zheng W-W, Lapidus A, Lowry S,
Haselkorn R, Bergman B. 2010. Genome Erosion in a Nitrogen-Fixing Vertically Transmitted
Endosymbiotic Multicellular Cyanobacterium. PLoS ONE [Internet] 5:e11486. Available from:
http://dx.doi.org/10.1371/journal.pone.0011486

Rinke C, Schwientek P, Sczyrba A, Ivanova NN, Anderson IJ, Cheng J-F, Darling A, Malfatti S, Swan
BK, Gies EA, et al. 2013. Insights into the phylogeny and coding potential of microbial dark
matter. Nature 499:431–437.

Sabree ZL, Kambhampati S, Moran NA. 2009. Nitrogen recycling and nutritional provisioning by
Blattabacterium, the cockroach endosymbiont. Proc. Natl. Acad. Sci. U. S. A. 106:19521–19526.

Salem H, Florez L, Gerardo N, Kaltenpoth M. 2015. An out-of-body experience: the extracellular
dimension for the transmission of mutualistic bacteria in insects. Proc. Biol. Sci. 282:20142957.

Sasaki T, Kawamura M, Ishikawa H. 1996. Nitrogen recycling in the brown planthopper, Nilaparvata
lugens: Involvement of yeast-like endosymbionts in uric acid metabolism. J. Insect Physiol.
42:125–129.

Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. Bioinformatics 30:2068–2069.

Smillie CS, Smith MB, Friedman J, Cordero OX, David LA, Alm EJ. 2011. Ecology drives a global
network of gene exchange connecting the human microbiome. Nature 480:241–244.

Soler-Rivas C, Espin JC, Wichers HJ. 2000. Oleuropein and related compounds. J. Sci. Food Agric.
80:1013–1023.

Stecher B, Denzler R, Maier L, Bernet F, Sanders MJ, Pickard DJ, Barthel M, Westendorf AM,
Krogfelt KA, Walker AW, et al. 2012. Gut inflammation can boost horizontal gene transfer
between pathogenic and commensal Enterobacteriaceae. Proc. Natl. Acad. Sci. U. S. A.
109:1269–1274.

Thomas CM, Nielsen KM. 2005. Mechanisms of, and barriers to, horizontal gene transfer between
bacteria. Nat. Rev. Microbiol. 3:711–721.

Tzanakakis M. 2003. Seasonal development and dormancy of insects and mites feeding on olive: a
review. Neth. J. Zool. 52:87–224.

de Vries EJ, Jacobs G, Breeuwer JA. 2001. Growth and transmission of gut bacteria in the Western
flower thrips, Frankliniella occidentalis. J. Invertebr. Pathol. 77:129–137.

de Vries EJ, Jacobs G, Sabelis MW, Menken SBJ, Breeuwer JAJ. 2004. Diet-dependent effects of gut
bacteria on their insect host: the symbiosis of Erwinia sp. and western flower thrips. Proc. Biol.
Sci. 271:2171–2178.

Wallden K, Rivera-Calzada A, Waksman G. 2010. Microreview: Type IV secretion systems: versatility
and diversity in function. Cell. Microbiol. 12:1203–1212.

Zhang C, Rabiee M, Sayyari E, Mirarab S. 2018. ASTRAL-III: polynomial time species tree
reconstruction from partially resolved gene trees. BMC Bioinformatics 19:153.

BFo2
Tatumella sp. TA1
Tatumella morbirosei
Tatumella saanichensis
Tatumella sp. UCD
Pantoea wallisii
Pantoea rodasii
Pantoea rwandensis
Pantoea stewartii
Pantoea allii
Pantoea anthophila
Pantoea conspicua
Pantoea vagans
Pantoea agglomerans
Pantoea gaviniae
Pantoea calida
Erwinia billingiae
Pantoea sp. IMH
Erwinia tracheiphila
Erwinia sp. 9145
Erwinia mallotivora
Erwinia tasmaniensis
Erwinia amylovora
Erwinia pyrifoliae
BFo1
Ca. Erwinia dacicola
Klebsiella michiganensis
Klebsiella pneumoniae
Lelliottia amnigena
Leclercia adecarboxylata
Enterobacter sp. OLF
Enterobacter cloacae
Gibbsiella quercinecans
Brenneria goodwinii
Pseudomonas sagittaria
Pseudomonas monteilii
Lautropia mirabilis
Lampropedia cohaerens
Sinorhizobium meliloti

97.4
99.5
47.6
37.6
97.9
72.3
73.5
49
89.6
99
99.8

2.0