Global phylogeographical patterns in the spread of foot-and-mouth disease virus

Running title: Phylogeography of foot-and-mouth disease virus

Manuel Jara[1], Alba Frias-De-Diego[1], Simon Dellicour[2,3], Guy Baele[3], and Gustavo Machado[1,*]

[1]Department of Population Health and Pathobiology, College of Veterinary Medicine, North

Carolina State University, Raleigh, NC, USA

[2]Spatial Epidemiology Lab (SpELL), Université Libre de Bruxelles, CP160/12 50, av. FD

Roosevelt, 1050 Bruxelles, Belgium

[3]Department of Microbiology and Immunology, Laboratory for Clinical and Epidemiological

Virology, Rega Institute, KU Leuven, Leuven, Belgium

[*]Corresponding author: gmachad@ncsu.edu

Abstract

The economic impacts of Foot-and-mouth disease virus (FMDV) have prompted most countries

worldwide to embark on local, regional and national control and eradication programs. In

conjunction with the global efforts to reduce further virus dissemination into free-zones and

contain or eradicate circulation in endemic regions, global markets and animal trade have

increased linearly in past decades facilitating virus spread to other countries. FMDV proved its

potential to propagate across borders on numerous occasions, but yet details regarding how

exactly the spread between countries occurred at global scale remain missing. To elucidate the

FMDV global spread characteristics, we studied its spatiotemporal dynamics using Bayesian

macroevolutionary analyses for serotypes O, A, Asia1, SAT1, SAT2, and SAT3, with a dataset

that comprises 58 years of phylogenetic and epidemiological information. We reconstructed the

evolutionary routes and historical spread of FMDV at global scale level. Our results highlight the

main differences in phylogeographic patterns, dispersal rates, and the role of host species in the

spread and maintenance of virus circulation. Contrary to previous studies, our results showed

that four FMDV serotypes were monophyletic (O, A, Asia1 and C), while all SATs serotypes did

not evidence a defined common ancestor. We also observed that FMDV diffusion was

characterized by a great variability on the spatiotemporal dynamics exhibited by serotypes A

and O, reflected by the wide variety of colonized countries. On the other hand, we observed that

serotypes with local spread have been historically constrained into their endemic areas (Asia 1

and SATs), which in the case of SAT serotypes, may be related to a lower variety of hosts. This

study provides significant insights into the spatiotemporal dynamics of the global circulation of

FMDV serotypes, by characterizing the viral routes of spread at serotype level, allowing to use

the past knowledge to improve future decisions for a more efficient control and eradication of

this disease.

Keywords: Disease dispersion, transboundary emerging diseases, risk factors, evolution

INTRODUCTION

The rapid growth of the global population along with the current demand for animal protein and

the increasing animal trade have positively impacted the spread of a broad range of

transboundary diseases. Clear examples of this phenomena are the recent emergence of

African Swine Fever in China, Avian Influenza or Contagious Bovine Pleuropneumonia among

others (Daszak et al., 2001; Kouba, 2003; Smith et al., 2007; Kilpatrick, 2011; Wang et al.,

2018). Understanding the tempo and mode of disease evolution allows to estimate the impact of

those and other factors on the evolutionary patterns exhibited by these diseases, which cannot

be analyzed by classic epidemiological studies (Read et al., 1999; Trevathan, 1999; Stearns

and Ebert, 2001).

Evolutionary ecology has shown to be an accurate and highly impacting approach to

understand the spatiotemporal dynamics of infectious diseases, with the capacity to explain

disease spread, virulence and invasion potential (Galvani, 2003; Mideo et al., 2008; Magiorkinis

et al., 2009; Weaver and Forrester, 2015; Lion and Gandon, 2016; Dellicour et al., 2017; Forni

et al., 2018), as well as the relationships with its underlying environment as a potential risk

factor (Stephens et al., 2016; Dellicour et al., 2017; Jacquot et al., 2017). In the case of

transboundary diseases, evolutionary ecology has proven to be a useful approach, providing

accurate knowledge for the control of pathogens worldwide (Auguste et al., 2015; Pigeault et al.,

2015; Rogalski et al., 2017; Fountain-Jones et al., 2018).

Foot-and-mouth disease virus (FMDV) is the most influential transboundary animal

disease (TAD) with historical worldwide circulation reported in domestic and wildlife reservoirs

(Grubman and Baxt, 2004; Brito et al., 2017). FMDV is a highly contagious disease caused by a

small single-strained RNA virus of the genus *Aphthovirus*, member of the family Picornaviridae

(Belsham, 1993) and classified into seven different serotypes; O, A, C, Asia 1 and Southern

African Territories (SATs) 1, 2 and 3 (Sobrino et al., 2001; Brown, 2003; OIE, 2009) which affect

severely the productivity of domesticated livestock, causing great economic losses (Abdul-

Hamid et al., 2011; Kandeil et al., 2013; Ganji et al., 2018). The United States Department of

Agriculture has recently updated calculations suggesting that a possible introduction of FMDV

could result in losses between \$15 to \$100 billion (Perez et al., 2004a; Knight-Jones and

Rushton, 2013; Rushton and Knight-Jones, 2015; USDA-APHIS, 2017). One of the main

reasons for this great impact is the wide variety of hosts known for FMDV (i.e., cattle, buffalo,

swine, sheep, and deer), from which a remarkable percentage is part of the food industry.

74   FMDV is known to be transmitted locally and globally, often associated with infected

75   animal products and human and animal movements (IUCN/SSC, 2018). Local transmission, on

76   the other hand, is based on airborne spread, animal direct contact with infected individuals or

77   carcasses and translocation of contaminated staff, equipment, and machinery (Ferguson et al.,

78   2001; Orsel et al., 2009).

79   Phylogenetic relationships among FMDV serotypes have been widely studied, mostly

based on individual genes (Fares and Tully, 2008; Lewis-Rogers et al., 2008; Schumann et al.,

2008; Priyadarshini et al., 2009), but also, in a lower level, on whole genome sequences (WGS)

(Carrillo et al., 2005; Cooke and Westover, 2008; Yoon et al., 2011). Nevertheless, most of the

studies mentioned above need to be updated due to the small available number of sequences

<200, and their inaccurate evolutionary analyses, which in most of the cases (excluding Yoon et

al (2011)) they did not consider the Bayesian phylogenetic methods (with the robustness

provided by bootstrap or Markov Chain Monte Carlo statistics), making the results less reliable.

87   Likewise, along with its evolution, several studies have explored the spatiotemporal

evolutionary dynamics of FMDV in different parts of the world, mainly focusing on the diffusion

patterns across its endemic regions: Asia and Africa (Abdul-Hamid et al., 2011; Di Nardo et al.,

2011; de Carvalho et al., 2013; Brito et al., 2017; Bachanek-Bankowska et al., 2018). However,

studies considering all serotypes are only available within small geographic regions, and global

studies only assessed some of the viral serotypes (Di Nardo et al., 2011; Alkhamis et al., 2018;

Brito et al., 2018).

94   In this study, we addressed the spatiotemporal dynamics of FMDV by using Bayesian

macroevolutionary analyses, including a comprehensive genetic, geographical and temporal

dataset regarding past and current FMDV outbreaks. The main motivation of this work was to

shed light on the complex evolutionary history of FMDV via the i) reconstruction of the

spatiotemporal phylogeographic history of FMDV at serotype level and ii) determine and

compare the global spatial spread of each FMDV serotype.

MATERIALS AND METHODS

Data collection and curation

We built a comprehensive genetic database comprising 543 publicly available whole genome

sequences from six FMDV serotypes (A, O, Asia1, SAT 1, SAT 2 and SAT 3), with sampling

dates ranging from 1959 to 2017 (GenBank ID numbers in Supplementary material Table S1).

Serotype C was not included in this study due to data unavailability (only 3 sequences

available). Our dataset gathers information from 43 countries and 4 continents obtained from

the Virus Pathogen Resource database, available at https://www.viprbrc.org (See Table S1).

To determine accurate phylogenetic relationship among FMDV reports, we combined the

available genetic information along with collection date, host species (i.e., cattle, ankole cow,

buffalo, swine and sheep) and location (discrete information at country level) as metadata

information. Any sample lacking one of these three characteristics was discarded.

Discrete phylogeographical analysis

Sequences were aligned using Mega X, available at www.megasoftware.net/ (Kumar et al.,

2008). To determine whether there was a sufficient temporal molecular evolutionary signal of

the FMDV sequences used for each serotype phylogeny, we used TempEst v1.5 (Rambaut et

al., 2016). To calculate the *P*-values associated with the phylogenetic signal analysis, we used

the approach described by (Murray et al., 2016) based on 1,000 random permutations of the

sequence sampling dates (Navascués et al., 2010). The relationship found between root-to-tip

divergence and sampling dates (years) supported the use of molecular clock analysis in this

study. Root-to-tip regression results for each serotype are reported in Supplementary Table S2,

all the results supported a significant temporal signal (*P*-value<0.05).

Phylogeographic history of FMDV dispersal was recovered from the obtained

spatiotemporal phylogenies for each serotype, where each tip represents the geographic

centroid where the sequence was extracted. Phylogenetic trees were generated by a discrete

phylogeography estimation by Bayesian inference through Markov chain Monte Carlo (MCMC),

implemented in BEAST v2.5.0 (Bouckaert et al., 2014). We partitioned the coding genes into

first+second and third codon positions and applied a separate Hasegawa-Kishino-Yano

(HKY+G; (Hasegawa et al., 1985)) substitution model with gamma-distributed rate

heterogeneity among sites to each partition (Shapiro et al., 2006). To account for evolutionary

rate variation among lineages, we used an uncorrelated lognormal relaxed molecular clock

(Drummond et al., 2006) using a coalescent constant population model. For each serotype, we

applied an uncorrelated relaxed clock model with an underlying lognormal distribution. All

analyses were developed for 200 million generations, sampling every 10,000th generation and

removing 10% as chain burn-in. All the Markov chain Monte Carlo analyses for each serotype

were investigated using Tracer software v1.6 (Drummond et al., 2006) to ensure adequate

effective sample sizes (ESS) (above 200) which were obtained for all parameters. Final trees

were summarized and visualized via Tree Annotator v. 2.3.0 and FigTree 1.4.3 respectively

(included in BEAST v2.5.0) (Rambaut and Drummond, 2016; Rambaut, 2017). Finally, to

visualize the spatiotemporal diffusion of each serotype we used Spatial Phylogenetic

Reconstruction of Evolutionary Dynamics using Data-Driven Documents (D3) SPREAD3

software (Bielejec et al., 2016).

RESULTS

The number of sequences per country varied from 1 to 100 samples. United Kingdom,

Argentina, India, and Japan, that have experienced the most severe impacts during this century,

were the countries with the highest number available genomes (Figure 1, Supplementary Table

S3). Likewise, several countries have been historically affected by more than one serotype,

particularly in Asia and Africa, where we observed that Uganda presented the highest virus

diversity, including serotypes O, A, and all SATs (Supplementary Table S3, Supplementary

Figure S1).

FIGURE 1. Spatial distribution of Foot-and-mouth disease virus. Blue range represents the

number of sequences available at the country level.

Spatiotemporal dynamics of FMDV

The spatiotemporal dynamics obtained for all FMDV serotypes is described in Figure 2, while a

more detailed explanation that shows the historical spread of each serotype in the phylogenetic

and geographic space is represented in Figures 3 to 8.

Phylogeographic analyses highlighted great asymmetries in the tempo and mode of

each serotype evolution (Figure 2). SAT1 appeared to be the basal clade of the entire group,

originating SAT2, SAT3 and serotype A, which later diversified into serotype Asia1, O and C.

Our analysis suggested O seemed to be the most recent, prolific and widespread lineage, with

the higher number of sequences available worldwide. Serotype A and Asia 1 appeared second

and third in the number of available sequences, followed by all SATs serotypes. The less

represented lineage was Serotype C, for which only 3 available sequences were available.

Maximum clade credibility phylogeny showed the monophyly of serotypes O, A, Asia 1 and C

(each serotype shared a common ancestor), while SATs serotypes appeared to have multiple

origins (Figure 2).

The phylogeographic analysis of FMDV was only performed at serotype level, due to the

low effective sample size value obtained when we analyzed FMDV as a whole (ESS<100),

which can generate a poor estimation of the posterior distributions.

FIGURE 2. Epidemiological statistics estimated for Foot-and-mouth disease. A Maximum clade

credibility phylogeny of the global FMDV epidemic, colored according to all serotypes (A, O, C,

Asia1, SAT1, SAT2, and SAT3).

Spatiotemporal diffusion among serotypes

Serotype O

We analyzed 360 sequences from serotype O, which comprise 66% of the global FMDV tree

(Figure 2). This serotype also presents the widest distribution of all, with records from 29

countries. Based on our analysis, its most likely center of origin was China from which it spread

in all directions. Viral spread occurred both locally (30% of the cases) to the surrounding

countries (the countries that are part of its international borders) and across long distances

(70%) to further countries (i.e., United Kingdom, Greece, Uganda and Malaysia), spreading

globally and colonizing most of Asia, Northeastern Africa and South America in less than 50

years (Figures 3a and 3b) (see Supplementary Video S1 for detailed footage). Phylogenetic

reconstruction identified clusters formed by different sublineages, where the most

phylogenetically distant were the group Asia, Africa and UK (outbreak in 2001) from the group

conformed by the UK (outbreak in 1967-68 + 2007) and Ecuador. The biggest centers of

dispersal events (geographic spread accompanied by diversification) of this serotype were

China, Malaysia, Turkey, South Korea, and Uganda, which in the case of China, South Korea,

and Turkey were among the countries with the highest number of sequences (see Figure 1b).

Serotype O also has the highest host diversity, where the most representative were cattle (71%

of the sequences) and swine (20%). Cattle were not only the most important host for this

serotype but also the most likely initial host of the ancestral lineages. On the other hand, sheep,

buffalo and ankole cow represented less than 10% of the total diversity (Figure 3a).

Spatiotemporal diffusion map shows the serotype O global distribution with great evidence of

long-range movements across countries and continents (Figure 3b).

FIGURE 3. Epidemiological statistics estimated for Foot-and-mouth disease (Serotype O)

epidemic. (a) Maximum clade credibility phylogeny of the FMDV epidemic colored according to

the countries where the disease has been registered. Dots at the end of the branches represent

the host species (buffalo, cattle, ankole cow, sheep, and swine). The probabilities of ancestral

states (inferred from the Bayesian MCMC discrete traits analysis) are shown in pie charts at

each node, where charts above branches represent locations and charts below branches most

likely hosts. (b) Reconstructed spatiotemporal diffusion of FMD serotype O spread, where

diameters of the colored circles are proportional to the square root of the number of MCC

branches maintaining a particular location state at each time period. The color of the branches

represents the age of the internal nodes, where darker red colors represent older spread

events.

Serotype A

Phylogeographic relationships obtained for serotype A indicated India as the most likely center

of origin. Besides, India was represented as the source of the highest number of dispersal

events, not only for the spread across Asia and Africa but also to South America, where we

observed a high number of diversification events in a short period of time (mainly associated

with the Argentina outbreak in 2001). Several phylogenetic groups were observed inside this

serotype, and the most distinguishable were the ancestral African group conformed by (Egypt,

Chad, and Zambia) and the most recent group of conformed by the rest of the sublineages. We

also observed that long-distance dispersal events were the most representative of the

spatiotemporal dynamics of this serotype, accounting for 73% of the total spread. Another

characteristic of this event was their multi-directionality, coming back to the point of origin in

numerous occasions. The most recent geographic dispersal events (from 2010) also suggested

India as one of the most important centers of origin for the spread of the virus and its

diversification, followed by Pakistan and Malaysia. The hosts associated with serotype A were;

cattle (93%), buffalo (5%), ankole cow and swine, which only represented a 2%. Viral circulation

was found to be widely spread in cattle population, while infected buffalos were only registered

in India, Vietnam, and Pakistan (Figures 4a and 4b; Supplementary Video S2).

FIGURE 4. Epidemiological statistics estimated for Foot-and-mouth disease (Serotype A)

epidemic. (a) Maximum clade credibility phylogeny of the FMDV epidemic colored according to

the countries where the disease has been registered. Dots at the end of the branches represent

the host species (buffalo, cattle, sheep, and swine). The probabilities of ancestral states

(inferred from the Bayesian MCMC discrete traits analysis) are shown in pie charts at each

node, where charts above branches represent locations and charts below branches most likely

hosts. (b) Reconstructed spatiotemporal diffusion of FMD serotype A spread, where diameters

of the colored circles are proportional to the square root of the number of MCC branches

maintaining a particular location state at each time period. The color of the branches represents

the age of the internal nodes, where darker red colors represent older spread events.

Serotype Asia 1

In the case of serotype A, phylogeographic analyses also indicated India as the most likely

origin, from which it diverged in all directions. Two main sublineages were identified, one of

them related to India, Bangladesh and the western side (i.e., Pakistan, Afghanistan, and

Turkey), while the other group gathered the dispersion through the southeastern side, including

India, China, Malaysia, and Vietnam. These dispersal events occurred mainly through local

movements, representing 56% of the spatiotemporal dynamics of this serotype (Figures 5a and

5b) (for detailed footage see Supplementary Video S3). The most important centers of

diversification for this serotype were India, China, and Malaysia (Figure 5a). The main hosts

associated to the dispersion of this serotype were cattle (representing 66% of the sequences),

followed by swine (14%, exclusively in China), buffalo (6%, it was observed in India, Vietnam,

and China) and sheep (~3%, also observed only in China). Its historical phylogeography also

revealed a relatively constant dispersion to new places marked by a slowdown between 1930

and 1960 showing a local dispersion during that time for some lineages. However, we observed

a clear increase in the diversification events related to the arrival of this serotype to Vietnam

from Pakistan and its spread back to Turkey, which also corresponds to the biggest dispersion

event occurring for this specific serotype.

FIGURE 5. Epidemiological statistics estimated for Foot-and-mouth disease (Serotype Asia 1)

epidemic. (a) Maximum clade credibility phylogeny of the FMDV epidemic, colored according to

the countries where the disease has been registered. Dots at the end of the branches represent

the host species (buffalo, cattle, sheep, and swine). The probabilities of ancestral states

(inferred from the Bayesian MCMC discrete traits analysis) are shown in pie charts at each

node, where charts above branches represent locations and charts below branches most

probable hosts. (b) Reconstructed spatiotemporal diffusion of FMD serotype Asia1 spread,

where diameters of the colored circles are proportional to the square root of the number of MCC

branches, maintaining a particular location state at each time period. The color of the branches

represents the age of the internal nodes, where darker red colors represent older spread

events.

Serotype SAT1

The phylogeographic patterns of SAT1 exposed Uganda as the most likely country of origin,

from where it diverged to Namibia, Nigeria, and Chad (Figures 6a and 6b). Phylogenetic

relationship identified three main sublineages, one of them represented by the ancestor Uganda

and Chad, other group by the countries that are part of the southern spread (i.e., Botswana,

Mozambique, Namibia, South Africa and Tanzania), and the last group represented by Nigeria,

which is also the sublineage with the most recent appearance. Contrary to the previous

serotypes, SAT1 did not present a clear source of dispersal events, although its spread was

mostly concentrated on the Eastern part of Africa (Figure 6b). Despite SATs serotypes are only

distributed across Africa, in the case of SAT1, the highest proportion of its dispersal events

occurred across long distance countries (63% of the cases. See Supplementary Video S4 for

detailed footage). The hosts associated with the dispersion of this serotype were cattle and

buffalo (Figure 6a). Cattle, with 65% of the reports, had a higher representation, mainly in the

north and central Africa, while buffaloes, which represented 35% of the reports, appeared as the

most common host in the southern countries (i.e., Namibia, Botswana, and South Africa).

Intriguingly, the phylogenetic analysis showed cattle as the host of the ancestral lineages of

SAT1, which would later adapt to infect buffalo in some regions (Figures 6a and 6b).

FIGURE 6. Epidemiological statistics estimated for Foot-and-mouth disease (Serotype SAT1)

epidemic. (a) Maximum clade credibility phylogeny of the FMDV epidemic, colored according to

the countries where the disease has been registered. Dots at the end of the branches represent

the host species (buffalo, cattle, sheep, and swine). The probabilities of ancestral states

(inferred from the Bayesian MCMC discrete traits analysis) are shown in pie charts at each

node, where charts above branches represent locations and charts below branches most

probable hosts. (b) Reconstructed spatiotemporal diffusion of FMD serotype SAT1 spread,

where diameters of the colored circles are proportional to the square root of the number of MCC

branches, maintaining a particular location state at each time period. The color of the branches

represents the age of the internal nodes, where darker red colors represent older spread

events.

Serotype SAT2

As in SAT1, phylogeographic analyses for SAT2 indicated Uganda as the most likely origin of

the serotype, from which it spread to Botswana and The Gaza Strip later on time (Figures 7a

and 7b). From Botswana, the serotype moved to Ethiopia, Zimbabwe, and Zambia, continuing

spreading to surrounding countries also on the Eastern region of Africa (See Supplementary

Video S5 for detailed footage). Phylogenetic relationships identified two main sublineages, one

represented by Uganda and its further dispersion to Gaza Strip and the second group formed by

countries distributed in southeastern Africa. Spatiotemporal dynamics evidenced a slightly

higher proportion of long-distance movements (57%) over local dispersal events (Figure 7b).

Cattle and buffalo were also the main hosts found for SAT2, but in this case, buffalo was the

most representative host (65%), over 35% registered by cattle. However, cattle appeared in

most of the reported locations (except in South Africa), while buffalo was only described in

Uganda, Botswana and South Africa. As observed in SAT1, phylogenetic analysis showed a

higher influence of cattle as the host of the ancestral lineages of this serotype (Figures 7a and

7b).

FIGURE 7. Epidemiological statistics estimated for Foot-and-mouth disease (Serotype SAT2)

epidemic. (a) Maximum clade credibility phylogeny of the FMDV epidemic, colored according to

the countries where the disease has been registered. Dots at the end of the branches represent

the host species (buffalo, cattle, sheep, and swine). The probabilities of ancestral states

(inferred from the Bayesian MCMC discrete traits analysis) are shown in pie charts at each

node, where charts above branches represent locations and charts below branches most

probable hosts. (b) Reconstructed spatiotemporal diffusion of FMD serotype SAT2 spread,

where diameters of the colored circles are proportional to the square root of the number of MCC

branches, maintaining a particular location state at each time period. The color of the branches

represents the age of the internal nodes, where darker red colors represent older spread

events.

Serotype SAT 3

Following the previous SATs serotypes, SAT3 also presented its ancestral origin in Uganda,

from where it traveled to Zimbabwe and then spread to its neighboring countries (Figures 8a

and 8b, see Supplementary Video S6). Phylogenetic analyses identified two main sublineages,

one of them represented by Uganda and the second (and most diverse), by the countries that

are part of the southern spread (i.e., Botswana, South Africa, Zambia, and Zimbabwe).

Phylogeographic reconstruction reflected the importance of Zimbabwe for the spread of this

serotype, being this country the most common center of origin for the diffusion of the disease to

Zambia, Botswana and most recently to South Africa. Contrary to all the other serotypes,

spatiotemporal dynamics of serotype SAT3 showed that its spread has been dominated by local

events (75%). As in the other SATs serotypes, cattle and buffalo were the main hosts reported
for SAT3, appearing both in a similar proportion (cattle=59%, and buffalo=41%).
Phylogeographic analyses showed that, even when cattle was determined as the most likely
ancestral host, both species are almost equally responsible for the spread of the disease to
most of the countries (Figures 8a and 8b).

FIGURE 8. Epidemiological statistics estimated for Foot-and-mouth disease (Serotype SAT3)
epidemic. (a) Maximum clade credibility phylogeny of the FMDV epidemic, colored according to
the countries where the disease has been registered. Dots at the end of the branches represent
the host species (buffalo, cattle, sheep, and swine). The probabilities of ancestral states
(inferred from the Bayesian MCMC discrete traits analysis) are shown in pie charts at each
node, where charts above branches represent locations and charts below branches most
probable hosts. (b) Reconstructed spatiotemporal diffusion of FMD serotype SAT3 spread,
where diameters of the colored circles are proportional to the square root of the number of MCC
branches, maintaining a particular location state at each time period. The color of the branches
represents the age of the internal nodes, where darker red colors represent older spread
events.
359
360 **DISCUSSION**
361
Macroecological patterns in the spread of FMDV

Global patterns of FMDV spread showed considerable asymmetries in the spatiotemporal

dynamics exhibited due to differences in the local and global spread showed by each serotype,

these results are in concordance with the observed by Yoon et al., (2011). However, we had

discrepancies with previous studies regarding the phylogenetic relationship among the FMDV

serotypes, since some of them indicated that all FMDV serotypes were monophyletic (Cooke

and Westover, 2008). Lewis-Rogers et al. (2008) and Yoon et al. (2011) suggested that O, A,

Asia 1, C, and SAT 3 were monophyletic, while SAT 1 and SAT2 serotypes were polyphyletic.

Our results indicated only four monophyletic serotypes (O, A, Asia1, and C), whilst all SATs

serotypes appeared to have multiple ancestral origins which can be related to multiple points of

independent introduction of the virus in the affected countries.

In this study, we followed the historical spread of FMDV, serotypes, from the ones with

local distribution (Asia 1 and SATs) to the serotypes with widespread dispersion (O and A),

where the lineage diversification was accompanied by colonization of new areas, which in most

of the cases were considerable distant. Supporting previous studies (Sobrino et al., 2001; Fèvre

et al., 2006; Di Nardo et al., 2011; Jamal and Belsham, 2013; Knight-Jones and Rushton, 2013;

Brito et al., 2017) our results suggested that the success in the spread of this disease through

long distances has been mainly associated with movement of commercial hosts (i.e., cattle,

swine, and sheep) or/and international trade of animal products, while local movements could

be more related to wild hosts (i.e., African buffalos).

Our results also showed that, even though the general spatiotemporal dynamics that

characterize the spread of most serotypes is represented mostly by long distance movements

across countries that are not part of their international borders, local movements are also

important for some serotypes especially SATs. These results complement the observations

made by (Bouslikhane, 2015), who highlighted how nomadism and transhumance play a key

role in disease transmission, especially in African countries.

Long distance spread

The remarkable historical spread exhibited by serotype O was the biggest registered among all

FMDV serotypes. In half of a century, this serotype colonized almost all the continents.

Interestingly, this spread is characterized principally by dispersion across long geographically

distant regions (to regions not sharing international borders with the origin country), instead to

its neighboring countries. Our phylogeographic estimates were in agreement with the historical

records of the past outbreaks (Jamal and Belsham, 2013; Brito et al., 2017) where in the case of

serotype O, it was responsible for the most catastrophic economic impact of FMDV worldwide

(Knight-Jones and Rushton, 2013; Rushton and Knight-Jones, 2015). For example, the

outbreak in United Kingdom in 2001 (King et al., 1981; Thompson et al., 2002; Gloster et al.,

2006) Japan in 2010 (Muroga et al., 2011; Shimada et al., 2016), China 2010 (Zheng et al.,

2012) and South Korea 2010 (Jeong-Nam et al., 2014). One of the reasons for the success of

the evolutionary diversification of this serotype may be related to the diversity of hosts that it

affects, which is the highest among all serotypes.

Following the pandemic serotype O, the next large-scale potential of diffusion was found

in serotype A. This serotype has been reported in three continents, Asia, Africa and South

America, where it was reported as the causing agent of one of the biggest FMDV outbreaks

(Argentina, 2011), affecting a total of 2,126 herds (Perez et al., 2004b).

Local spread serotypes

Whereas our results showed that serotypes O and A have spread worldwide, serotypes Asia1

and SATs remained nonpandemic and confined in their endemic regions. The spread of the

serotype Asia1 was characterized mainly by local movements across the neighboring countries

surrounding India, China and Malaysia, where it is well known that free and unrestricted animal

movements across country borders play a key role in the spread of FMDV (Subramaniam et al.,

2013; Brito et al., 2017). The dispersion of the disease into Turkey in 2013 represents one of the

most recent and longer dispersion events for this serotype, which was directly related to an

Indian subgroup of the virus. Likewise, as reported by (Abdul-Hamid et al., 2011) the outbreak

occurred in Malaysia during 1999 seemed to be caused also by an independent subgroup from

the rest of the outbreaks observed in the region.

SAT serotypes (SAT1, SAT2, and SAT3) also exhibited a local spread, limited across

their endemic areas. This spread occurred mainly in Africa, where nomadic pastoralism across

international borders and animal trade in the sub-Saharan region is one of the most practiced

forms of livestock movements (Di Nardo. et al., 2011), which we suggest may be the main

transmission route for these serotypes. Previous studies have highlighted the importance of

African buffalo (*Syncerus caffer*) as one of the main hosts described for the SATs serotypes

(Vosloo et al., 1996; Thomson et al., 2003; Brito et al., 2017). Differences on phylogenetic

analyses showed that for SAT2, buffalo and cattle were both present as hosts of the ancestral

lineages, which could be a sign of an early crossed infection, contrary to SAT1 and SAT3,

where buffalo seemed to have appeared as a host later on time, mostly on the southern

countries of the continent. This could have conducted to a potential viral generalism between

these host species that would later disappear due to the tendency of the virus to evolve to

narrower niches, leading to the specificity of viral-host relationships currently reported (Elena et
al., 2009).

In general, considerable asymmetries in the spatiotemporal dynamics exhibited by the
different serotypes were observed. Serotypes such as Asia1 and SATs presented local spread
rates, mainly associated with cattle and sheep (with special importance of buffalo in the case of
SATs serotypes), while serotypes O and A showed long-distance spread, covering higher
extensions of territory between each outbreak. These serotypes presented the highest variety of
susceptible host, although we suggest that the main reason for their successful long-distance
spread relies mostly on the international movements of cattle and swine due to the intensive
commerce between countries.

Final remarks
Studies considering WGS should be preferred over specific regions, mainly to ensure the
detection of all informative mutations for phylogenetic analyses. FMD as a small-genome RNA
virus represents an excellent candidate for this approach, due to the well-known mutation rate
and the variability that RNA viruses possess (Gilchrist et al., 2015). However, for our study, the
use of WGS represented a limitation since partial sequences were available in some
underrepresented countries which were not included in our analysis. For that reason, here we
highlight the importance of including WGS in future studies. These projects would not only
increase the available information about the virus, but also have a direct impact on promoting
new and more specific measures for disease control (Schrag and Wiener, 1995; Duchatel et al.,
2018; Fountain-Jones et al., 2018), reducing drastically the economic impact caused by FMDV
and inefficient resource allocation. One example of this would be the consideration of the most
influencing host for the spread of FMDV in a given area, targeting these strategies to reduce the
main source of infection. This control would not only be beneficial for the targeted region, but
also for all the areas connected directly (i.e., through geographical limits) and indirectly (i.e.,
through commercial networks) with it. For that reason, we defend that understanding the
complex evolutionary history of FMDV is not only critical for endemic countries, but also for
regions in which FMDV has been eradicated and especially for countries in which vaccination
will be withdrawn in the coming years, such as in Brazil (Reuters, 2018). We also encourage
future efforts from regions and international organizations for the development of a better and
more comprehensive archive of virus circulation by including WGS and a complete metadata
repository to facilitate the understanding of the factors promoting the continual emergence and
rapid spatial dispersal of new viral lineages. This will be necessary to help future vaccine design

and better-informed strategies for viral control, providing insights into viral interactions and

disease control optimization based on scientific evidence.

CONCLUSION

In summary, in this study, we addressed the worldwide circulation of FMDV, by characterizing

its global macroevolutionary patterns at serotype scale. The macroevolutionary approach used

here relies on the principle that evolutionary processes are better understood when a broader

spatiotemporal vision is available, allowing us to understand the whole evolutionary history of a

given organism. Establishing the phylogeographic relationships among FMDV serotypes at the

global scale will shed light on their macroevolutionary patterns by reconstructing its historical

viral routes of spread, as well as the role of their major hosts. Besides, from this perspective, we

can conclude that as a response to the detection of an outbreak event, closing borders with

neighboring countries should not be the only measure applied, since some of the most virulent

FMDV lineages spread through the long-distance transport of animals or its natural products,

and therefore, control measures should take into account all the commercial network of the

affected country, especially from regions that have been historically considered as centers of

dispersal of the disease.

ACKNOWLEDGEMENTS

We acknowledge the Department of Population Health and Pathobiology- North Carolina State

University provided startup funds for G. Machado and M. Jara.

CONFLICT OF INTEREST

The authors declare that there are no conflict of interests.

REFERENCES

Abdul-Hamid, N.F., N.M. Hussein, J. Wadsworth, A.D. Radford, N.J. Knowles, and D.P. King,

2011: Phylogeography of foot-and-mouth disease virus types O and A in Malaysia and

surrounding countries. *Infect. Genet. Evol.* **11**, 320–328, DOI:

10.1016/j.meegid.2010.11.003.

Alkhamis, M.A., C. Gallardo, C. Jurado, A. Soler, M. Arias, and J.M. Sánchez-Vizcaíno, 2018:

Phylodynamics and evolutionary epidemiology of African swine fever p72-CVR genes in

Eurasia and Africa. (Ulrike Gertrud Munderloh, Ed.)*PLoS One* **13**, e0192565, DOI:

10.1371/journal.pone.0192565.

Auguste, A.J., J. Liria, N.L. Forrester, D. Giambalvo, M. Moncada, K.C. Long, D. Morón, N. de
Manzione, R.B. Tesh, E.S. Halsey, T.J. Kochel, R. Hernandez, J.C. Navarro, and S.C.
Weaver, 2015: Evolutionary and ecological characterization of mayaro virus strains isolated
during an outbreak, Venezuela, 2010. *Emerg. Infect. Dis.* **21**, 1742–1750, DOI:
10.3201/eid2110.141660.

Bachanek-Bankowska, K., A. Di Nardo, J. Wadsworth, V. Mioulet, G. Pezzoni, S. Grazioli, E.
Brocchi, S.C. Kafle, R. Hettiarachchi, P.L. Kumarawadu, I.M. Eldaghayes, A.S. Dayhum, D.
Meenowa, S. Sghaier, H. Madani, N. Abouchoaib, B.H. Hoang, P.P. Vu, K. Dukpa, R.B.
Gurung, S. Tenzin, U. Wernery, A. Panthumart, K.B. Seeyo, W. Linchongsubongkoch, A.
Relmy, L. Bakkali-Kassimi, A. Scherbakov, D.P. King, and N.J. Knowles, 2018:
Reconstructing the evolutionary history of pandemic foot-and-mouth disease viruses: the
impact of recombination within the emerging O/ME-SA/Ind-2001 lineage. *Sci. Rep.* **8**,
14693, DOI: 10.1038/s41598-018-32693-8.

Belsham, G., 1993: Distinctive features of foot-and-mouth disease virus, a member of the
picornavirus family; aspects of virus protein synthesis, protein processing and structure.
Prog. Biophys. Mol. Biol. **60**, 241–260.

Bielejec, F., G. Baele, B. Vrancken, M. Suchard, A. Rambaut, and P. Lemey, 2016: SpreaD3:
interactive visualization of spatiotemporal history and trait evolutionary processes. *Mol.
Biol. Evol.* **33**, 2167–2169.

Bouckaert, R., J. Heled, D. Kühnert, T. Vaughan, C.H. Wu, D. Xie, M.A. Suchard, A. Rambaut,
and A.J. Drummond, 2014: BEAST 2: A Software Platform for Bayesian Evolutionary
Analysis. (Andreas Prlic, Ed.)*PLoS Comput. Biol.* **10**, e1003537, DOI:
10.1371/journal.pcbi.1003537.

Bouslikhane, M., 2015: Cross border movements of animals and animal products and their
relevance to the epidemiology of animals disease in Africa. OIE Africa Regional
Commision.

Brito, B., S.J. Pauszek, M. Eschbaumer, C. Stenfeldt, H.C. De Carvalho Ferreira, L.T. Vu, N.T.
Phuong, B.H. Hoang, N.D. Tho, P. V. Dong, P.Q. Minh, N.T. Long, D.P. King, N.J.
Knowles, D.H. Dung, L.L. Rodriguez, and J. Arzt, 2017: Phylodynamics of foot-and-mouth
disease virus O/PanAsia in Vietnam 2010-2014. *Vet. Res.* **48**, 24, DOI: 10.1186/s13567-
017-0424-7.

Brito, B., S.J. Pauszek, E.J. Hartwig, G.R. Smoliga, L.T. Vu, P. V. Dong, C. Stenfeldt, L.L.
Rodriguez, D.P. King, N.J. Knowles, K. Bachanek-Bankowska, N.T. Long, D.H. Dung, and

533 J. Arzt, 2018: A traditional evolutionary history of foot-and-mouth disease viruses in

534  Southeast Asia challenged by analyses of non-structural protein coding sequences. *Sci.*

535  *Rep.* **8**, DOI: 10.1038/s41598-018-24870-6.

536 Brito, B., L. Rodriguez, J. Hammond, J. Pinto, and A. Perez, 2017 (April): Review of the Global

537  Distribution of Foot-and-Mouth Disease Virus from 2007 to 2014. *Transbound. Emerg. Dis.*

538  **64**, 316–332, DOI: 10.1111/tbed.12373.

539 Brown, F., 2003: The history of research in foot-and-mouth disease. *Virus Res.* **91**, 3–7.

540 Carrillo, C., E.R. Tulman, G. Delhon, Z. Lu, A. Carreno, A. Vagnozzi, G.F. Kutish, and D.L.

541  Rock, 2005: Comparative genomics of foot-and-mouth disease virus. *J. Virol.* **79**, 6487–

542  504, DOI: 10.1128/JVI.79.10.6487-6504.2005.

543 Cooke, J.N., and K.M. Westover, 2008: Serotype-specific differences in antigenic regions of

544  foot-and-mouth disease virus (FMDV): A comprehensive statistical analysis. *Infect. Genet.*

545  *Evol.* **8**, 855–863, DOI: 10.1016/j.meegid.2008.08.004.

546 Daszak, P., A.A. Cunningham, and A.D. Hyatt, 2001: Anthropogenic environmental change and

547  the emergence of infectious diseases in wildlife. *Acta Trop.* **78**, 103–116.

548 De Carvalho, L.M.F., L.B.L. Santos, N.R. Faria, and W. De Castro Silveira, 2013:

549  Phylogeography of foot-and-mouth disease virus serotype O in Ecuador. *Infect. Genet.*

550  *Evol.* **13**, 76–88, DOI: 10.1016/j.meegid.2012.08.016.

551 Dellicour, S., P. Lemey, R. Rose, N.R. Faria, O.G. Pybus, L.F.P. Vieira, H. Bourhy, and M.

552  Gilbert, 2017: Using Viral Gene Sequences to Compare and Explain the Heterogeneous

553  Spatial Dynamics of Virus Epidemics. *Mol. Biol. Evol.* **34**, 2563–2571.

554 Di Nardo., A., N. Knowles, and D. Paton, 2011: Combining livestock trade patterns with

555  phylogenetics to help understand the spread of foot and mouth disease in sub-Saharan

556  Africa, the Middle East and. *Rev. Sci. Tech.* **30**, 63.

557 Drummond, A.J., S.Y.W. Ho, M.J. Phillips, and A. Rambaut, 2006: Relaxed phylogenetics and

558  dating with confidence. (David Penny, Ed.)*PLoS Biol.* **4**, e88, DOI:

559  10.1371/journal.pbio.0040088.

560 Duchatel, F., M. Bronsvoort, and S. Lycett, 2018: Phylogeographic analysis and identification of

561  factors impacting the diffusion of Foot-and-Mouth disease virus in Africa. *BiorXiv*1–23.

562 Elena, S.F., P. Agudelo-Romero, and J. Lalic, 2009: The Evolution of Viruses in Multi-Host

563  Fitness Landscapes. *Open Virol. J.* **3**, 1–6, DOI: 10.2174/1874357900903010001.

564 Fares, M. a, and D.C. Tully, 2008: The tale of a modern animal plague: tracing the evolutionary

565  history and determining the time-scale for foot and mouth disease virus. *Virology* **382**, 250–

566  6, DOI: 10.1016/j.virol.2008.09.011.

567 Ferguson, N.M., C.A. Donnelly, and R.M. Anderson, 2001: Transmission intensity and impact of
568   control policies on the foot and mouth epidemic in Great Britain. *Nature* **413**, 542–548.

569 Fèvre, E., B. Bronsvoort, K. Hamilton, and S. Cleaveland, 2006: Animal movements and the
570   spread of infectious diseases. *Trends Microbiol.* **14**, 125–131, DOI:
571   10.1016/j.tim.2006.01.004.

572 Forni, D., R. Cagliani, M. Clerici, and M. Sironi, 2018: Origin and dispersal of Hepatitis e virus
573   article. *Emerg. Microbes Infect.* **7**, DOI: 10.1038/s41426-017-0009-6.

574 Fountain-Jones, N.M., W.D. Pearse, L.E. Escobar, A. Alba-Casals, S. Carver, T.J. Davies, S.
575   Kraberger, M. Papeş, K. Vandegrift, K. Worsley-Tonks, and M.E. Craft, 2018: Towards an
576   eco-phylogenetic framework for infectious disease ecology. *Biol. Rev.* **93**, 950–970, DOI:
577   10.1111/brv.12380.

578 Galvani, A.P., 2003: Epidemiology meets evolutionary ecology. *Trends Ecol. Evol.* **18**, 132–139,
579   DOI: 10.1016/S0169-5347(02)00050-2.

580 Ganji, V., S. Kontham, and M. Pottabathula, 2018: Understanding the Molecular Relationship
581   between Foot-and-Mouth Disease Virus Serotype O of Indian Vaccine Strain with Strains
582   across the World by. *Phylogenetic Anal. Int. J. Curr. Microbiol.* **7**, 99–105.

583 Gilchrist, C.A., S.D. Turner, M.F. Riley, W.A. Petri, and E.L. Hewlett, 2015: Whole-genome
584   sequencing in outbreak analysis. *Clin. Microbiol. Rev.* **28**, 541–63, DOI:
585   10.1128/CMR.00075-13.

586 Gloster, J., D.J. Paton, G.H. Hutchings, D.P. King, N.P. Ferris, D.T. Haydon, J.W. Wilesmith,
587   and E.M. Cottam, 2006: Molecular Epidemiology of the Foot-and-Mouth Disease Virus
588   Outbreak in the United Kingdom in 2001. *J. Virol.* **80**, 11274–11282, DOI:
589   10.1128/jvi.01236-06.

590 Grubman, M.J., and B. Baxt, 2004: Foot-and-mouth Disease. *Clin. Microbiol. Rev.* **17**, 465–493.

591 Hasegawa, M., H. Kishino, and T. Yano, 1985: Dating of the human-ape splitting by a molecular
592   clock of mitochondrial DNA. *J. Mol. Evol.* **22**, 160–174, DOI: 10.1007/BF02101694.

593 IUCN/SSC, I.S.S.G., 2018: Global Invasive Species Database version 2013-1 [Online] Available
594   at http://www.issg.org/database/welcome/ (accessed October 14, 2018).

595 Jacquot, M., K. Nomikou, M. Palmarini, P. Mertens, and R. Biek, 2017: Bluetongue virus spread
596   in Europe is a consequence of climatic, landscape and vertebrate host factors as revealed
597   by phylogeographic inference. *Proc. R. Soc. B Biol. Sci.* **284**, DOI:
598   10.1098/rspb.2017.0919.

599 Jamal, S.M., and G.J. Belsham, 2013: Foot-and-mouth disease: past, present and future. *Vet.*
600   *Res.* **44**, 116, DOI: 10.1186/1297-9716-44-116.

Jeong-Nam, P., L. Seo-Yong, C. Jia-qi, L. Yeo-Joo, K. Rae-Hyung, L. Kwang-Nyeong, K. Su-mi,
T. Dong-Seob, K. Byounghan, and P. Jong-Hyeon, 2014: Protection to homologous and
heterologous challenge in pigs immunized with vaccine against foot-and-mouth disease
type O caused an epidemic in East Asia during 2010/2011. *Vaccine* **32**, 1882–1889, DOI:
10.1016/j.vaccine.2014.01.067.

Kandeil, A., R. El-Shesheny, G. Kayali, Y. Moatasim, O. Bagato, M. Darwish, A. Gaffar, A.
Younes, T. Farag, M.A. Kutkat, and M.A. Ali, 2013: Characterization of the recent outbreak
of foot-and-mouth disease virus serotype SAT2 in Egypt. *Arch. Virol.* **158**, 619–627, DOI:
10.1007/s00705-012-1529-y.

Kilpatrick, A.M., 2011: Globalization, land use, and the invasion of West Nile virus. *Science (80-.*
). **334**, 323–327, DOI: 10.1126/science.1201010.

King, A.M.Q., B.O. Underwood, D. McCahon, J.W.I. Newman, and F. Brown, 1981: Biochemical
identification of viruses causing the 1981 outbreaks of foot and mouth disease in the UK.
Nature **293**, 479–480, DOI: 10.1038/293479a0.

Knight-Jones, T., and J. Rushton, 2013: The economic impacts of foot and mouth disease -
What are they, how big are they and where do they occur? *Prev. Vet. Med.* **112**, 162–173,
DOI: 10.1016/j.prevetmed.2013.07.013.

Kouba, V., 2003: Globalization of Communicable Animal Diseases - A Crisis of Veterinary
Medicine. *Acta Vet. Brno* **72**, 453–460, DOI: 10.2754/avb200372030453.

Kumar, S., M. Nei, J. Dudley, and K. Tamura, 2008: MEGA: A biologist-centric software for
evolutionary analysis of DNA and protein sequences. *Brief. Bioinform.* **9**, 299–306, DOI:
10.1093/bib/bbn017.

Lewis-Rogers, N., D.A. McClellan, and K.A. Crandall, 2008: The evolution of foot-and-mouth
disease virus: Impacts of recombination and selection. *Infect. Genet. Evol.* **8**, 786–798,
DOI: 10.1016/j.meegid.2008.07.009.

Lion, S., and S. Gandon, 2016: Spatial evolutionary epidemiology of spreading epidemics. *Proc.*
R. Soc. B Biol. Sci. **283**, DOI: 10.1098/rspb.2016.1170.

Magiorkinis, G., E. Magiorkinis, D. Paraskevis, S.Y.W. Ho, B. Shapiro, O.G. Pybus, J.P. Allain,
and A. Hatzakis, 2009: The global spread of hepatitis C virus 1a and 1b: A phylodynamic
and phylogeographic analysis. (Arthur Y. Kim, Ed.)*PLoS Med.* **6**, e1000198, DOI:
10.1371/journal.pmed.1000198.

Mideo, N., S. Alizon, and T. Day, 2008: Linking within- and between-host dynamics in the
evolutionary epidemiology of infectious diseases. *Trends Ecol. Evol.* **23**, 511–517, DOI:
10.1016/j.tree.2008.05.009.

Muroga, N., Y. Hayama, A. Kurogi, T. Tsuda, T. Tsutsui, and T. Yamamoto, 2011: The 2010
Foot-and-Mouth Disease Epidemic in Japan. *J. Vet. Med. Sci.* **74**, 399–404, DOI:
10.1292/jvms.11-0271.

Murray, G.G.R., F. Wang, E.M. Harrison, G.K. Paterson, A.E. Mather, S.R. Harris, M.A. Holmes,
A. Rambaut, and J.J. Welch, 2016: The effect of genetic structure on molecular dating and
tests for temporal signal. *Methods Ecol. Evol.* **7**, 80–89, DOI: 10.1111/2041-210X.12466.

Navascués, M., F. Depaulis, and B. Emerson, 2010: Combining contemporary and ancient DNA
in population genetic and phylogeographical studies. *Mol. Ecol. Resour.* **10**, 760–772, DOI:
10.1111/j.1755-0998.2010.02895.x.

OIE, T.M., 2009: Foot-And-Mouth Disease.

Orsel, K., A. Bouma, A. Dekker, J.A. Stegeman, and M.C.M. de Jong, 2009: Foot and mouth
disease virus transmission during the incubation period of the disease in piglets, lambs,
calves, and dairy cows. *Prev. Vet. Med.* **88**, 158–163, DOI:
10.1016/j.prevetmed.2008.09.001.

Perez, A.M., M.P. Ward, and T.E. Carpenter, 2004a: Control of a foot-and-mouth disease
epidemic in Argentina. *Prev. Vet. Med.* **65**, 217–226.

Perez, A.M., M.P. Ward, and T.E. Carpenter, 2004b: Epidemiological investigations of the 2001
foot-and-mouth disease outbreak in Argentina. *Vet. Rec.* **154**, 777–782.

Pigeault, R., J. Vézilier, S. Cornet, F. Zélé, A. Nicot, P. Perret, S. Gandon, and A. Rivero, 2015:
Avian malaria: A new lease of life for an old experimental model to study the evolutionary
ecology of Plasmodium. *Philos. Trans. R. Soc. B Biol. Sci.* **370**, DOI:
10.1098/rstb.2014.0300.

Priyadarshini, P., J.K. Mohapatra, D. Hemadri, S. Subramaniam, B. Pattnaik, L. Pandey, and A.
Sanyal, 2009: Analysis of the leader proteinase (Lpro) region of type A foot-and-mouth
disease virus with due emphasis on phylogeny and evolution of the emerging VP359-
deletion lineage from India. *Virus Res.* **141**, 34–46, DOI: 10.1016/j.virusres.2008.12.012.

Rambaut, A., 2017: FigTree-version 1.4. 3, a graphical viewer of phylogenetic trees. .

Rambaut, A., and A. Drummond, 2016: TreeAnnotator v.2.4. 3. . Part of the BEAST package.

Rambaut, A., T. Lam, L. Max Carvalho, and O. Pybus, 2016: Exploring the temporal structure of
heterochronous sequences using TempEst (formerly Path-O-Gen). *Virus Evol.* **2**, vew007,
DOI: 10.1093/ve/vew007.

Read, A.F., P. Aaby, R. Antia, D. Ebert, P.W. Ewald, S. Gupta, E.C. Holmes, A. Sasaki, D.C.
Shields, F. Taddei, and R.E. Moxon, 1999: What can evolutionary biology contribute to
understanding virulence?, pp. 205–216. In: Evolution in Health and Disease.

669 Reuters, 2018: Brazil expects to be declared free of foot-and-mouth with vaccination [Online]
670   Available at https://www.reuters.com/article/us-brazil-beef-disease/brazil-expects-to-be-
671   declared-free-of-foot-and-mouth-with-vaccination-idUSKBN1EU1WZ.

672 Rogalski, M.A., C.D. Gowler, C.L. Shaw, R.A. Hufbauer, and M.A. Duffy, 2017: Human drivers
673   of ecological and evolutionary dynamics in emerging and disappearing infectious disease
674   systems. *Philos. Trans. R. Soc. B Biol. Sci.* **372**, DOI: 10.1098/rstb.2016.0043.

675 Rushton, J., and T.J.D. Knight-Jones, 2015: The impact of foot and mouth disease. pp. 205–
676   209. In: *FAO/OIE, Glob. Conf. Foot Mouth Dis. Control.* Food and Agriculture Organization
677   of the United Nations and the World Organisation for Animal Health.

678 Schrag, S.J., and P. Wiener, 1995: Emerging infectious disease: what are the relative roles of
679   ecology and evolution. *Trends Ecol. Evol.* **10**, 319–324.

680 Schumann, K.R., N.J. Knowles, P.R. Davies, R.J. Midgley, J.-F. Valarcher, A.Q. Raoufi, T.S.
681   McKenna, W. Hurtle, J.P. Burans, B.M. Martin, L.L. Rodriguez, and T.R. Beckham, 2008:
682   Genetic characterization and molecular epidemiology of foot-and-mouth disease viruses
683   isolated from Afghanistan in 2003–2005. *Virus Genes* **36**, 401–413, DOI: 10.1007/s11262-
684   008-0206-4.

685 Shapiro, B., A. Rambaut, and A.J. Drummond, 2006: Choosing appropriate substitution models
686   for the phylogenetic analysis of protein-coding sequences. *academic.oup.com* **23**, 7–9,
687   DOI: 10.1093/molbev/msj021.

688 Shimada, N., K. Sakamoto, K. Yoshida, K. Fukai, K. Morioka, M. Yamada, T. Nishi, M.
689   Yamakawa, and T. Kanno, 2016: Genome variability of foot-and-mouth disease virus
690   during the short period of the 2010 epidemic in Japan. *Vet. Microbiol.* **199**, 62–67, DOI:
691   10.1016/j.vetmic.2016.12.025.

692 Smith, K.F., D.F. Sax, S.D. Gaines, V. Guernier, and J.F. Guégan, 2007: Globalization of
693   human infectious disease. *Ecology* **88**, 1903–1910, DOI: 10.1890/06-1052.1.

694 Sobrino, F., M. Sáiz, M.A. Jiménez-Clavero, J.I. Núñez, M.F. Rosas, E. Baranowski, and V. Ley,
695   2001: Foot-and-mouth disease virus: A long known virus, but a current threat. *Vet. Res.* **32**,
696   1–30, DOI: 10.1051/vetres:2001106.

697 Stearns, S.C., and D. Ebert, 2001: Evolution in Health and Disease: Work in Progress. *Q. Rev.*
698   *Biol.* **76**, 417–432, DOI: 10.1086/420539.

699 Stephens, P.R., S. Altizer, K.F. Smith, A. Alonso Aguirre, J.H. Brown, S.A. Budischak, J.E.
700   Byers, T.A. Dallas, T. Jonathan Davies, J.M. Drake, V.O. Ezenwa, M.J. Farrell, J.L.
701   Gittleman, B.A. Han, S. Huang, R.A. Hutchinson, P. Johnson, C.L. Nunn, D. Onstad, A.
702   Park, G.M. Vazquez-Prokopec, J.P. Schmidt, and R. Poulin, 2016: The macroecology of

infectious diseases: a new perspective on global-scale drivers of pathogen distributions
and impacts. (Hillary Young, Ed.)*Ecol. Lett.* **19**, 1159–1171, DOI: 10.1111/ele.12644.

Subramaniam, S., B. Pattnaik, A. Sanyal, J.K. Mohapatra, S.S. Pawar, G.K. Sharma, B. Das,
and B.B. Dash, 2013: Status of Foot-and-mouth Disease in India. *Transbound. Emerg. Dis.*
60, 197–203, DOI: 10.1111/j.1865-1682.2012.01332.x.

Thompson, D., P. Muriel, D. Russell, P. Osborne, A. Bromley, M. Rowland, and C. Brown, 2002:
Economic costs of the foot and mouth disease outbreak in the United Kingdom in 2001.
Rev. Sci. Tech. **21**, 675–687.

Thomson, G.R., W. Vosloo, and A.D. Bastos, 2003: Foot and mouth disease in wildlife. *Virus*
Res. **91**, 145–161, DOI: 10.1016/s0168-1702(02)00263-0.

Trevathan, W., 1999: Evolutionary medicine, pp. 1–4. In: The International Encyclopedia of
Biological Anthropology. Hoboken, NJ, USA: John Wiley & Sons, Inc.

USDA-APHIS, 2017: Planning and preparing for Foot-and-mouth disease: quick briefing [Online]
Available at
https://www.aphis.usda.gov/animal_health/emergency_management/downloads/fmd-
briefing.pdf.

Vosloo, W., A.D. Bastos, E. Kirkbride, J.J. Esterhuysen, D.J. van Rensburg, R.G. Bengis, D.W.
Keet, and G.R. Thomson, 1996: Persistent infection of African buffalo (Syncerus caffer)
with SAT-type foot-and-mouth disease viruses: rate of fixation of mutations, antigenic
change and interspecies transmission. *J. Gen. Virol.* **77**, 1457–1467, DOI: 10.1099/0022-
1317-77-7-1457.

Wang, T., Y. Sun, and H.-J. Qiu, 2018: African swine fever: an unprecedented disaster and
challenge to China. *Infect. Dis. poverty* **7**, 111, DOI: 10.1186/s40249-018-0495-3.

Weaver, S.C., and N.L. Forrester, 2015: Chikungunya: Evolutionary history and recent epidemic
spread. *Antiviral Res.* **120**, 32–39, DOI: 10.1016/j.antiviral.2015.04.016.

Yoon, S.H., W. Park, D.P. King, and H. Kim, 2011: Phylogenomics and molecular evolution of
foot-and-mouth disease virus. *Mol. Cells* **31**, 413–421, DOI: 10.1007/s10059-011-0249-6.

Zheng, H., J. He, J. Guo, Y. Jin, F. Yang, L. Lv, and X. Liu, 2012: Genetic characterization of a
new pandemic Southeast Asia topotype strain of serotype O foot-and-mouth disease virus
isolated in China during 2010. *Virus Genes* **44**, 80–88, DOI: 10.1007/s11262-011-0670-0.

SUPPLEMENTARY MATERIAL

TABLE S1. Sample information for all Foot and Mouth Disease virus complete genome

sequences used in this study.

TABLE S2. Root-to-tip regression analyses of phylogenetic temporal signal. Correlation and

determination coefficient (R2) were estimated with TempEst (Rambaut et al. 2016). P-values

were calculated using the approach of Murray et al. (2016) and were based on 1,000 random

permutations of the sequence sampling dates (Navascuès et al. 2010).

TABLE S3. Number of sequences and serotypes per country.

FIGURE S1. Spatial distribution of Foot-and-mouth disease virus showing the number of

serotypes per country.

Video S1. Reconstructed spatiotemporal diffusion of FMD serotype O spread, where diameters

of the colored circles are proportional to the square root of the number of MCC branches,

maintaining a particular location state at each time period. The color of the branches represents

the age of the internal nodes, where darker red colors represent older spread events, this

visualization match with the main time bar on top of the video.

Video S2. Reconstructed spatiotemporal diffusion of FMD serotype A spread, where diameters

of the colored circles are proportional to the square root of the number of MCC branches,

maintaining a particular location state at each time period. The color of the branches represents

the age of the internal nodes, where darker red colors represent older spread events, this

visualization match with the main time bar on top of the video.

Video S3. Reconstructed spatiotemporal diffusion of FMD serotype Asia1 spread, where

diameters of the colored circles are proportional to the square root of the number of MCC

branches, maintaining a particular location state at each time period. The color of the branches

represents the age of the internal nodes, where darker red colors represent older spread

events, this visualization match with the main time bar on top of the video.

Video S4. Reconstructed spatiotemporal diffusion of FMD serotype SAT1 spread, where

diameters of the colored circles are proportional to the square root of the number of MCC

branches, maintaining a particular location state at each time period. The color of the branches

represents the age of the internal nodes, where darker red colors represent older spread

events, this visualization match with the main time bar on top of the video.

Video S5. Reconstructed spatiotemporal diffusion of FMD serotype SAT2 spread, where

diameters of the colored circles are proportional to the square root of the number of MCC

branches, maintaining a particular location state at each time period. The color of the branches

represents the age of the internal nodes, where darker red colors represent older spread

events, this visualization match with the main time bar on top of the video.

Video S6. Reconstructed spatiotemporal diffusion of FMD serotype SAT3 spread, where

diameters of the colored circles are proportional to the square root of the number of MCC

branches, maintaining a particular location state at each time period. The color of the branches

represents the age of the internal nodes, where darker red colors represent older spread

events, this visualization match with the main time bar on top of the video.