Substantial genetic mixing among sexual and androgenetic lineages within the clam genus
Corbicula

Vastrade M. [1], Etoundi E.[1], Bournonville T.[2], Colinet M.[1], Debortoli N. [1,2], Hedtke S.M.[3], Nicolas E.[4],
Pigneur L.-M. [5], Virgo J.[1], Flot J.-F.[6,7], Marescaux J. [1,2] and Van Doninck K.[1,4]

[1] Laboratory of Evolutionary Genetics and Ecology; Research Unit in Environmental and Evolutionary
Biology; Institute of Life, Earth and Environment (ILEE); University of Namur, 5000 Namur,
Belgium

[2] e-biom SA, 5/7 Rue Godefroid, 5000 Namur, Belgium

[3] Department of Physiology, Anatomy and Microbiology, School of Life Sciences, La Trobe
University, Bundoora, VIC 3086, Australia

[4] Université libre de Bruxelles (ULB), Molecular Biology & Evolution, C.P. 160/16, Avenue F.D.
Roosevelt 50, 1050 Brussels, Belgium

[5] Université de Liège, Conservation Genetics Laboratory, Chemin de la vallée 4, 4000 Liège, Belgium

[6] Université libre de Bruxelles (ULB), Evolutionary Biology & Ecology, C.P. 160/12, Avenue F.D.
Roosevelt 50, 1050 Brussels, Belgium

[7] Interuniversity Institute of Bioinformatics in Brussels – (IB)[2], Brussels, Belgium

Corresponding authors: Martin Vastrade & Karine Van Doninck

Fax: +3281724362

E-mail: martin.vastrade@unamur.be, karine.van.doninck@ulb.be

Abstract

"Occasional" sexuality occurs when a species combines clonal reproduction and genetic mixing.
This strategy is predicted to combine the advantages of both asexuality and sexuality, but its actual
consequences on the genetic diversity and species longevity are poorly understood. Androgenesis, a
reproductive mode in which the offspring inherits its entire nuclear genome from the father, is often
reported as a strictly clonal reproductive mode. Androgenesis is the predominant reproductive mode
within the hermaphroditic, invasive lineages of the mollusk genus *Corbicula*. Their ability to
reproduce clonally through androgenesis has been determinant in their invasive success, having
colonized during the 20th century American and European freshwater systems, where they became
notorious invaders with a widespread, global distribution. However, in androgenetic *Corbicula* clams,
occasional genetic mixing between distinct lineages has also been observed when the sperm of one
lineage fertilizes the oocyte of another one. Because of these occasional introgressions, the genetic
relationships between *Corbicula* species remained unclear, and the biogeographic origins of the
invasive androgenetic lineages have been challenging to identify. To address these issues, we analyzed
the patterns of allele sharing for several nuclear and mitochondrial molecular markers among
Corbicula individuals collected across both the native and invasive range. Our results show the
occurrence of an allelic pool encompassing all *Corbicula* freshwater species worldwide, including
sexual and androgenetic ones, which highlights the substantial genetic mixing within this genus.
However, the differences in allele sharing patterns between invasive lineages, and the low diversity
within each lineage, suggest recent, distinct biogeographic origins of invasive *Corbicula* androgenetic
lineages. Finally, the polyploidy, high heterozygosity, and hybrid phenotypes and genotypes found in
our study probably originated from hybridization events following egg parasitism between distinct
Corbicula lineages. This extensive cross-lineage mixing found in *Corbicula* may generate nuclear
diversity in an otherwise asexually reproducing species.

Keywords: androgenesis, occasional sexuality, allele sharing, hybridization, invasion

Introduction

Sexual reproduction with meiotic recombination is the most widespread reproductive mode among eukaryotes (Dimijian 2005; Barton 2009; Hartfield & Keightley 2012). Sex, through genetic reshuffling, is predicted to promote new allele combinations and to reduce genetic associations of linked loci, increasing the efficiency of selection and the purging of deleterious mutations (Lenormand *et al.*, 2016; Bast *et al.* 2018). However, sex has many costs, including the two-fold cost of males (*i.e.,* the reduced fecundity when half of the individuals does not produce offspring), the cost of meiosis (*i.e.,* the reduced genetic contribution to the offspring), the cost of recombination (*i.e.,* the possible loss of favorable associations between alleles), and the cost of mate searching and mating (Hartfield & Keightley 2012; Lehtonen *et al.* 2012; Liegeois *et al.* 2020). The prevalence of sex in eukaryotes is therefore considered a major paradox in evolutionary biology.

Besides sexual reproduction *sensu stricto*, including meiosis and syngamy, a diversity of asexual reproductive mechanisms exists in which oocytes appear to be produced through a modified meiosis or mitotically (Lenormand *et al.*, 2016). In some organisms, asexual and sexual reproductions can alternate, referred to as occasional sexuality (or cyclical parthenogenesis in animals). In addition, there are several reproductive modes in which meiosis and fertilization still take place, but only the unreduced paternal or maternal genome is transmitted to the next generation. One such quasi-sexual reproductive mode is androgenesis, in which the offspring, after fertilization, inherits only the entire paternal nuclear genome (McKone & Halpern 2003, Schwander & Oldroyd, 2016). The maternal nuclear chromosomes are either discarded from the egg, or oocytes are non-nucleated prior to fertilization (reviewed in Schwander & Oldroyd, 2016). When androgenetic males or hermaphrodites use the eggs of other individuals to clone themselves, this reproductive mode is considered sexual parasitism or "egg parasitism" (Hedtke *et al.* 2008; Komaru *et al.* 2013, reviewed in Lehtonen *et al.* 2013), because one individual uses the female gamete of another individual to transmit his entire nuclear genome to the next generation, while the female host does not contribute with its nuclear genome. The processes that generate and maintain such a peculiar reproductive mode remain elusive. Moreover, androgenesis appears rare in nature or remains undetected, but is more prevalent in hermaphroditic species (McKone & Halpern 2003, Schwander & Oldroyd 2016). It has been observed as a main reproductive mode in only a few species, such as the conifer *Cupressus dupreziana* (Pichot *et al.* 2001), the ant *Wasmannia auropunctata* (Rey *et al.* 2013), and some stick-insect hybrid species (*e.g., Bacillus rossius-grandii,* Mantovani & Scali 1992, *Pijnackeria hispanica*, Milani *et al.* 2020). However, androgenesis is widespread in the clam genus *Corbicula,* with several species reproducing through obligate androgenesis (for a review on androgenetic species, see Schwander & Oldroyd, 2016).

The genus *Corbicula* is therefore one of the most suitable system to better understand the origin and maintenance of androgenesis and to study the genetic consequences of asexual reproduction while maintaining the sexual characteristics of meiosis and fertilization. Indeed, androgenesis can be considered as sexual from a functional and ecological perspective (eggs and sperm are required and fertilization does occur, albeit without karyogamy) but asexual from a genetic point of view (Pigneur *et al.* 2012). Moreover, within the clam genus *Corbicula,* besides hermaphroditic androgenetic species, dioecious sexual species are found, and it is the only metazoan clade known in which the hermaphroditic species reproduce through obligate androgenesis for which the underlying cytological mechanism is well described (Komaru *et al.* 2000; Ishibashi *et al.* 2002, 2003; Ishibashi & Komaru, 2006; Hotta & Komaru, 2018). Androgenetic *Corbicula* clams produce offspring after fertilization of a congeneric oocyte by a biflagellate, unreduced sperm. The oocyte continues meiosis, but the meiotic axis is parallel to the egg surface, while it is perpendicular to the cortex in the case of a canonical

meiosis (Komaru *et al.* 2000, Figure 1). As a result, having both asters associated to the egg surface in androgenetic *Corbicula*, all the maternal nuclear chromosomes in meiosis I are discarded from the egg as two polar bodies, instead of one polar body in canonical meiosis (Komaru *et al.* 2000; Hotta & Komaru, 2018; Figure 1). The zygote, therefore, inherits the mitochondria from the oocyte and the nuclear genome from the sperm (reviewed in Pigneur *et al.*, 2012). The unreduced, biflagellate sperm of androgenetic *Corbicula* is a presumed hallmark of their androgenesis, plausibly formed through an ameiotic process or modified meiosis in which ploidy reduction is absent (the details remain unknown). Besides the hermaphroditic androgenetic species, the sexual *Corbicula* species appear to be diploid, producing reduced, uniflagellate sperm (Glaubrecht *et al.,* 2006; Obata *et al.* 2006).

In *Corbicula,* androgenetically produced offspring are formed through self-fertilization or outcrossing (Kraemer *et al.*, 1986). When outcrossing occurs between distinct species, the nuclear genome of one species becomes associated with the mitochondria of another one, resulting in cytonuclear mismatches (Figure 1b) (Hedtke & Hillis 2011) and increased diversity of mitochondrial sequences within a species (Lee *et al.*, 2005; Hedtke *et al.*, 2008; Pigneur *et al.*, 2012). Moreover, in some cases, maternal nuclear chromosomes may be partially retained and both maternal and paternal nuclear chromosomes are inherited by the offspring, eventually resulting in an increased ploidy (see Komaru *et al.*, 1997; Park *et al.*, 2000; Qiu *et al.*, 2001). Such nuclear capture or hybridization events, if they occur among different *Corbicula* species, could in addition be a cause of genetic divergence between androgenetic species (Pigneur *et al.* 2014), generating nuclear diversity in otherwise asexually reproducing species (Figure 1c) (Hedtke *et al*., 2011). This process has been recorded in *Corbicula* (Komaru *et al.*, 2006) and was reported as "rare" (Hedtke *et al.* 2011), although its frequency has never been thoroughly quantified. In addition, because these introgression events appear to occur between distinct *Corbicula* species, hybrid genotypes might be the cause of observed intermediate phenotypes (Pfenninger *et al.* 2002; Pigneur *et al.* 2014) and polyphyletic gene trees that fail to resolve taxonomy in the genus *Corbicula*. The taxonomy of invasive *Corbicula* species is particularly contentious because it has been only based on morphology and mitochondrial COI barcodes with several species' names being given including *C. fluminea, C. fluminalis, C. largillierti, C. fluviatilis*, and *C. leana*, while lacking the nuclear genotype information (*e.g.,* Coughlan *et al.* 2019; Crespo *et al.* 2015; Gomes et al. 2020; López-Soriano *et al.* 2018; Minchin 2014; Patrick *et al.* 2017; Reyna *et al.* 2018; Voroshilova *et al.* 2021). Some of these species' names are likely to be synonymous and therefore using this species-based nomenclature can confuse lineage identification between studies.

Corbicula clams originated from Africa, Asia, Australia, and the Middle East (Araujo *et al.* 1993). In some of these locations, both sexual and androgenetic species co-occur but the reproductive mode of many native lineages remains unknown. The well-described sexual dioecious species have very restricted geographic distribution: *Corbicula sandai* is endemic to Lake Biwa in Japan, *Corbicula japonica* is restricted to East Asian brackish water, and a few sexual *Corbicula* species are from Indonesia (Glaubrecht *et al.* 2006; Yamada *et al.*, 2014, Bespalaya *et al.* 2020). These species were described as sexual based on sperm morphology (Konishi *et al.* 1998, Hedtke *et al.* 2011, Pigneur *et al.* 2014). The androgenetic *Corbicula* species have a more widespread distribution across Asia. While the sexual species *C. sandai* and *C. japonica* have been identified as distinct species based on reproductive, morphological and genetic parameters (Pigneur *et al.* 2014), the species delimitation of native androgenetic *Corbicula* is also uncertain. For example, Komaru *et al.* (2012) showed a sister relationship between native, androgenetic *C. leana* and sexual *C. sandai* in Lake Biwa (Japan)*,* but in a separate study *C. leana* clustered instead with androgenetic *C. fluminea* (Komaru *et al.* 2013). We

therefore refer in our previous studies and here to androgenetic *Corbicula* lineages or forms instead of
species (as in Pigneur *et al.* 2014).

All of the invasive *Corbicula* lineages found in Europe, North America, and South America
appear to be exclusively androgenetic (Lee *et al.*, 2005; Hedtke *et al.*, 2008; Pigneur *et al.*, 2014;
Gomes *et al.,* 2016). Based on the COI barcode and confirmed by nuclear microsatellite data (Pigneur
et al. 2014), four distinct androgenetic and invasive *Corbicula* lineages have been described: form A/R
being widespread and cosmopolitan in America and Europe (COI haplotype FW5, Table S1), form B
only known from America (COI haplotype FW1, Table S1), form C/S found in America and Europe
(COI haplotype FW17, Table S1), and form Rlc only described in Europe (COI haplotype FW4, Table
S1) (Siripattrawan *et al.* 2000; Renard *et al.* 2000, Pfenninger *et al.* 2002, Lee *et al.* 2005; Hedtke et
al. 2008; Marescaux *et al.* 2010, Pigneur *et al.* 2014). Peñarrubia *et al.* (2017) considered Rlc and B as
one form, because their COI haplotypes differ by only one mutation, but the nuclear data showed
significant differences between forms Rlc and B (Pigneur *et al.* 2014). More recently, a new invasive
form D was described in America based on mitochondrial and nuclear genetic data (Tiemann *et al.*
2017; Haponski and Foighil 2019). Within all these invasive forms virtually no genetic diversity was
found over their entire invasive range at the nuclear and mitochondrial markers studied (Hedtke *et al.*,
2008; Pigneur *et al.*, 2014; Gomes *et al.,* 2016; Haponski & Foighil, 2019). When distinct invasive
forms were found in sympatry, cytonuclear mismatches and hybrids (resulting from maternal genome
retention) have been detected (Lee *et al.,* 2005; Hedtke *et al.,* 2008; Pigneur *et al.,* 2014; Peñarrubia *et
al.,* 2017; Tiemann *et al.*, 2017; Bespalaya *et al.,* 2018; Haponski & Foighil, 2019; Sano *et al.* 2020).
For example, out of twelve sampled locations in Europe where form A/R occurs in sympatry with
another invasive form (C/S or Rlc), cytonuclear mismatches were detected at 8 locations and hybrids
at 2 locations (Pigneur *et al.* 2014; Etoundi, pers. obs.). However, a more thorough quantification
would be required.

The low diversity and the invasion by only a few *Corbicula* lineages in America and Europe are
probably the consequence of a recent introduction by only few hermaphroditic individuals (1920's in
America and 1980's in Europe; Mouthon, 1981, McMahon, 1892, Pigneur *et al.*, 2014; Gomes *et al.,*
2016; Bespalaya *et al.,* 2018) and their ability to reproduce through androgenesis and to self-fertilize
(Kraemer *et al.*, 1986). This indeed facilitates their rapid spread because only one hermaphroditic
individual is required to establish a new population. In addition, since the descendants are derived only
from the unreduced sperm, there is no genetic mixing (except if different lineages occur in sympatry),
no loss of heterozygosity with time (except if recombination takes place), and no inbreeding
depression in the invasive populations (Schwander & Oldroyd, 2016). In contrast, androgenetic
Corbicula from the native range have been characterized by high genetic diversity comparable to that
of sexual *Corbicula* lineages (Pigneur *et al.,* 2014). This high native diversity could be the
consequence of genetic captures by co-occurring androgenetic lineages or multiple transitions to
asexuality from sexually reproducing populations (*e.g.* in ostracods, Bode *et al.*, 2010; in *Timema* stick
insect, Schwander & Crespi, 2009; van der Kooi & Schwander, 2014), or both.

The non-monophyly of androgenetic *Corbicula* lineages could be explained either by a single
origin of androgenetic *Corbicula* lineages followed by independent nuclear capture events between
sympatric lineages, or by multiple, independent origins from diverse sexual lineages followed by
nuclear captures (Hedtke *et al.*, 2011; Pigneur *et al.* 2011a, 2012, 2014; Gomes *et al.* 2016; Peñarrubia
et al., 2017). Hedtke *et al.* (2011) proposed, based on phylogenetic reconstructions using nuclear
markers, a common origin for most invasive androgenetic *Corbicula* lineages from a *C. sandai*
ancestor but with a possible second origin of androgenesis for the invasive form C/S because it
grouped with *Corbicula moltkiana* from Indonesia across their phylogenies. However, in the COI

phylogenies of Pigneur *et al.*, form C/S clustered with *C. australis* (2011a) or with *C. fluminalis*
africana (2014). These phylogenetic discordances due to nuclear and mitochondrial captures among
Corbicula lineages have until now hampered the investigation of phylogenetic relationships between
androgenetic and sexual *Corbicula* (see Hedtke & Hillis 2011; Hedtke *et al.* 2011; Pigneur *et al.,*
2012, 2014).

In the present work, multiple nuclear markers combined with the COI mitochondrial barcode were
used to study the genetic relationships between *Corbicula* lineages from the native and invasive
distributions, including different reproductive modes. Because phylogenetic approaches are not able to
resolve lineage relatedness in *Corbicula*, we investigated how individuals and lineages cluster using an
allele sharing-based approach to species delimitation that can delineate species regardless of their
monophyly or lack thereof (haplowebs, Flot *et al.,* 2010). This is the first time this method is applied
to the genus *Corbicula*, providing novel insights into the origin of their invasive alleles. The
specimens used in this study were obtained from the worldwide distribution of *Corbicula,* focusing
our sampling efforts on regions where genetically diverse *Corbicula* species co-occur, including
sexual and/or endemic species (*C. sandai* from Lake Biwa, estuarine *C. japonica* in Japan, and *C.*
tobae from Indonesia). Populations where sexual and androgenetic *Corbicula* individuals are found
together (such as in Lake Biwa and South-East Asia) are a critical source of information since
sympatric populations that include sexuals are putative hotspots for the origin of new asexual lineages
(Simon *et al.*, 2003).

The main aims of this study are to identify i) the origin of alleles found in invasive *Corbicula*
lineages and ii) the genetic mixing that have occurred between distinct lineages, both from the native
and invasive range. Owing to the reconstruction method of allele sharing, we predict that one genetic
cluster of shared alleles, including a native population, would be observed if all invasive lineages had
the same biogeographic origin, while distinct clusters including different invasive lineages would be
detected in the case of multiple origins. Given the previously reported genetic relationship of *C.*
sandai with androgenetic lineages, the exact nature of shared alleles between this sexual species and
each invasive lineage should be identified. In addition, the occurrence of genetic captures between
distinct *Corbicula* lineages (*i.e.* cytonuclear mismatches and hybridization events) blurs the
delimitation of genetic clusters within *Corbicula* genus, unless it is rare. With the allele-sharing
method, we will identify the magnitude of genetic exchange. Moreover, if introgressions are frequent
but regionally restricted, a pool of shared alleles should be retrieved per geographic region. If this
genetic mixing occurs worldwide, we expect to detect one large genetic cluster of shared alleles
encompassing most sampled populations.

Material and methods

Specimen collection and reproductive mode characterization.

A total of 359 *Corbicula* individuals from 44 distinct localities were considered in this study,
collected across the geographic range of *Corbicula,* including both native and invasive areas (see
Table S2). Mollusks were collected using a scoop net and preserved in 96% ethanol.

For samples collected in *Corbicula*'s native range, the species identity was initially based on the
mitochondrial haplotype or on published species taxonomic identification in the same region where
the collection took place. Because taxonomic identification of native androgenetic *Corbicula* is often
uncertain, we refer to some taxa as *C. sp.* (putative species name) (Table S2-S3). The reproductive
mode was inferred from the sperm morphology when possible, from the literature available for the

sampled population (lineage assignment and/or reproductive morphology) or, if these characteristics
were indeterminate, classified as "undetermined".

Invasive *Corbicula* lineages encompass a wide geographic distribution both in Europe and
America, and can be discriminated by eye on the basis of morphological characteristics. We refer to
these invasive forms as previously described (A/R, B, Rlc, and C/S: Pigneur *et al.,* 2014; note form D
was not sampled in this study). Clams sampled from one invasive population ("Wa", Netherlands)
could not be identified as one of these forms because it displays an intermediate morphology. The
reproductive mode of invasive individuals was determined based on spermatozoa morphology (as
described in Pigneur *et al.,* 2014) under the assumption that androgenetic individuals produce
biflagellate sperm while sexual clams produce uniflagellate sperm (Komaru & Konishi, 1999). Owing
to the very little genetic diversity observed within invasive lineages in previous studies (Lee *et al.*,
2005; Hedtke *et al.*, 2008; Pigneur *et al.*, 2011a, 2014), we analyzed less than 10 individuals from
each invasive form for further genetic analyses.

Genetic data from previous studies were included only when the raw data (chromatograms) were
available, allowing a thorough check of the final sequences used in our genetic study.

Genetic analyses

DNA isolation, marker amplification, and sequencing.

We extracted DNA either from the adductor muscles or the foot using the DNeasy Blood and
Tissue Kit (QIAGEN GmbH, Hilden, Germany) following the manufacturer's protocol. We sequenced
four different markers: (1) a 651-bp fragment of the mitochondrial cytochrome c oxidase gene (COI);
(2) a 413 to 415-bp fragment of the nuclear 28S gene (28S); (3) a 551 to 660-bp fragment containing
the third intron of the nuclear α-amylase gene (*amy*), and (4) a 310 to 365-bp fragment of the
hypervariable putative intron of the α subunit of the adenosine triphosphate synthase (*ATPS*). PCR
reactions for the four markers were carried out in a mix comprising 1-4 µl DNA, 1X Taq buffer
(Promega Corp., Madison, Wisconsin, USA), 0.2 mM nucleotides, 0.3-0.5 µM of each primer, and 0.1
unit of *GoTaq G2* polymerase (Promega) in a total volume of 25 µl. The PCR conditions and primers
used for each marker are described in Supplemental Information 1. PCR products were purified and
sequenced on an automated sequencer at Genoscreen (Lille, France) or Beckman Coulter Genomics
(Bishop's Stortford, United Kingdom).

As polyploidy has been recorded in *Corbicula* (Park *et al.*, 2000; Qiu *et al.*, 2001) (Figure 1), the
occurrence of more than two alleles in each individual is possible. Therefore, direct sequencing data
were confirmed by cloning, using the NEB® PCR Cloning Kit with the Q5 High-Fidelity Taq
Polymerase and following the manufacturer's protocol. Between two and four individuals were picked
in each population, based on direct sequencing results, to confirm as many alleles as possible. Between
20 and 30 clones per individual were selected randomly and purified using the QIAprep Spin Miniprep
Kit (QIAGEN GmbH) in order to cover several times each represented allele. Sequences were
obtained from purified plasmids using the Genewiz (Leipzig, Germany) Sanger sequencing service. A
column was added to Table S2 for each marker and each individual to indicate the use of direct
sequencing, cloning, or both.

Sequence editing, alignment, and species delimitation methods.

All sequences were first assembled and cleaned using Sequencher 4.1.4 (Gene Codes Corp., Ann
Arbor, Michigan, USA). Alignments were made using MAFFT version 7 (Katoh *et al*. 2017) with the

iterative refinement method E-INS-I and indels treated as a 5[th] character state in downstream analyses.
Because mitochondrial and nuclear markers potentially have different inheritance patterns in
Corbicula genus, and because the allele sharing-based approach used in this study cannot be applied to
mitochondrial genes, two different clusterization methods were used: one based on the mitochondrial
marker using the K/θ method and one based on the nuclear markers using the haploweb and
conspecificity matrix approach.

COI sequences were manually edited using BioEdit (Hall 1999) and Sequencher, then tested for
cross-contamination using AutoConTAMPR (https://github.com/jnarayan81/autoConTAMPR). No
contamination was detected among these tested haplotypes. A median-joining haplotype network
(haplonet) was constructed using HaplowebMaker (Spöri *et al.* 2020) based on the COI sequence
alignment, including in total 251 COI *Corbicula* sequences. The K/θ method is based on a comparison
between intergroup and intragroup sequence diversity (Birky *et al.* 2010; Birky 2013; Birky &
Maughan, 2021) as implemented in the online tool KoT (Spöri & Flot, 2021). This method defines a
new species when the gap between two clades cannot be caused by genetic drift (based on a threshold)
and was shown efficient on asexual or clonal organisms (*e.g.:* Schön *et al.*, 2012) and using haploid
markers (Birky, 2013). The distinct *Corbicula* clusters (*i.e.*, putative species) defined by this KoT
method were color coded on the haplonet constructed for the COI gene and on the haplowebs
constructed for each nuclear gene.

For the three nuclear genes (28S, *amy* and *ATPS*), the allele validation based on both direct and
cloning sequencing data was done manually in a case-by-case process (Supplemental Information 2)
for the 359 individuals with the majority (more than 90%) being sequenced in both directions. Allele
sharing based on the three nuclear markers was investigated using the haploweb approach (Doyle,
1995; Flot *et al.*, 2010) (Figure S1). The online tool HaplowebMaker (Spöri *et al.*, 2020) was used
with default parameters to build for each nuclear marker a raw haploweb, which is a median-joining
haplotype network (or haplonet, as produced for the mitochondrial, haploid COI marker) on which
curves are added connecting nuclear haplotypes or alleles found co-occurring in heterozygous
individuals. A group of alleles linked together by heterozygotes creates an exclusive allele pool (AP)
(akin to species, Doyle, 1995). The haplowebs were edited manually for visualization purpose. Allele-
sharing patterns involving the four invasive forms A/R, C/S, B, and Rlc were manually numbered,
referring to non-exclusive invasive alleles (invasive alleles that are not specific to one given invasive
form) according to the Tables S2 and S3. The allele-sharing information on the three nuclear markers
provided by HaplowebMaker was used to build a "conspecificity matrix" (as in Debortoli *et al.*, 2016)
with the CoMa tool (Spöri *et al.*, 2020) with no calculation on heterospecific pairs (option 1: *i.e.*, no
value was given to the absence of sharing). In this matrix, the conspecificity score for each pair of
individuals is the number of markers for which these individuals belong to the same allelic pool
(ranging from 0 to 3 for the three nuclear markers taken together). After computing the matrix, rows
and columns were reordered to maximize the scores along the diagonal using the hierarchical
clustering method implemented in the R package ''heatmap3'' (Zhao *et al.* 2014). Fields For
Recombination (FFRs) appear as blocks along the diagonal with high conspecificity scores within
blocks, and low scores among them, except in case of introgressions. These introgressions between
FFRs were highlighted using a letter code. The final FFR definition was based on the dendrogram
(Ward clustering) built by "heatmap3", and each FFR was given a different color code. This color
code was used in each haploweb based on nuclear markers and on the haplonet constructed for the
COI marker to highlight the FFRs defined by the conspecificity matrix using the three nuclear
markers. The haplowebs provide details on the allele sharing patterns between FFRs for each marker
individually.

Alleles shared between *Corbicula* lineages were also summarized and visualized in a circular plot
using the command-line version of Circos (Krzywinski *et al.*, 2009). Fifteen distinct groups (namely
idiotypes) were manually defined in the Circos plot to highlight the allele-sharing patterns for all four
markers (COI, 28S, *amy*, and *ATPS*). More specifically, the four invasive forms (A/R, C/S, B, and Rlc)
were considered as four distinct idiotypes, with the invasive Wa population considered separately as it
could not be formally identified as one of these four invasive lineages. The brackish water sexual
lineage *C. japonica* was also considered a separate group. Other native idiotypes were defined based
on the sampled geographic localities (Lake Biwa in Japan including *C. sandai*, other sites in Japan,
China, Hawaii, Madagascar, Indonesia, Vietnam, and South Africa). Due to the low number of
individuals sampled, we combined the few individuals from Korea, Taiwan, Thailand, and the
Philippines together in a last group (or idiotype) named "South-East Asia". The occurrence of sexual
and androgenetic lineages was represented by mono- and biflagellate spermatozoa, respectively, for
each group when known. A color code was used to distinguish the invasive alleles, which fall into two
groups: the exclusive invasive alleles, shared between a native lineage and only one invasive form,
represented in red (form A/R), blue (form B), green (form Rlc), and orange (form C/S), and the non-
exclusive invasive alleles, each found in at least two different invasive forms and some native
populations, represented in purple. This same color code has been applied in the Table S2.

All alleles are summarized in Table S2, with unique names within and between the different
markers: for each marker, the name of the allele starts with the name of the marker followed by the
name of the individual it was retrieved from, with an additional letter (a, b, c, d) if this *Corbicula*
individual is heterozygous. If the same allele is found within several individuals, the name used is
arbitrarily based on one of those individuals. Alleles shared between distinct idiotypes, based on the
Circos analysis, are in bold type, and the same color code is used for invasive alleles. These alleles
shared between distinct invasive lineages were named IA (for Invasive Allele), are in purple and are
numbered (from 1 up to 3 according to the marker). These numbers are also used in the haplowebs
(Figure 2a, b, and c) to highlight these alleles.

All validated sequences of our study were compared to those available on the NCBI nucleotide
database (https://www.ncbi.nlm.nih.gov/) to avoid redundancy with previously submitted sequences
and new sequences were deposited (accession numbers listed in Table S1).

Genetic diversity.

Nucleotide diversity and haplotype diversity (π and H_d, respectively) were calculated on the entire
dataset (including all alleles even in case of polyploidy) for each marker as the percentage of variable
sites between 2 sequences (*i.e.,* uncorrected pairwise distance) using DNAsp (Librado & Rozas 2009).
Gaps, however, were not considered, underestimating the diversity among nuclear alleles.

Results

Alleles were checked independently using two methods: direct sequencing and allele inferences,
and sequencing of cloned amplicons. When the called alleles gave conflicting results between those
inferred from direct sequencing and cloned sequences, a manual verification was performed by
comparing chromatograms (see Supp. Info. 2). As a result, the majority of alleles obtained in the
present work were confirmed by several independent analyses and were identified within multiple
individual clams (see Supp. Info. 2). However, some markers were not successfully amplified in an
individual, or could not be confirmed with confidence, and were therefore excluded. Consequently,
many individuals lack information for one or several markers (see Table S2). For this reason, two
distinct datasets are considered in this study: one with all available sequences for all markers,

including a total of 359 individuals but with only 251, 266, 237, and 248 individuals for COI, 28S,
alpha-amylase, and ATPS markers respectively (Table S2); and a second dataset with only 141
individuals for which all four markers could be amplified and confidently validated (Table S3). These
two datasets are referred to as completed and filtered datasets, respectively.

High heterozygosity and haplotype diversity in Corbicula

All markers studied had a relatively high haplotype diversity (H_d) but a low nucleotide diversity
(π), indicating the presence of many low-frequency, genetically similar alleles in *Corbicula* (Table 1).
This was also true for the mitochondrial COI marker: 251 sequences revealed 44 distinct haplotypes
with 110 polymorphic sites. Moreover, for all three nuclear markers, most *Corbicula* individuals were
heterozygous: for 28S, 84% (224/266) were heterozygous, for *amy,* 85% (201/237), while for *ATPS*,
only 60% (148/248) were heterozygous (Table S2).

For the three nuclear markers studied, *amy* was the most diverse*,* with 74 alleles retrieved with 147
polymorphic sites in 237 individuals. For *ATPS*, 40 alleles were recovered with 118 polymorphic sites
in 248 individuals. 28S was the least diverse nuclear marker of this study, with only 43 polymorphic
sites giving 56 alleles in 266 sequenced individuals (complete dataset, Table S2).

Only four genetic clusters retrieved within the genus Corbicula *using nuclear markers with two*
distinct biogeographic origins of the invasive lineages

Haplowebs for the three nuclear markers (using heterozygosity to link alleles, see Fig. S1) and a
haplonet for the mitochondrial COI marker were produced for the filtered dataset (Fig. 2) as well as
for the complete dataset (Fig. S2). Each resulting graph is presented with two color codes based on the
distinct clusterization methods: (1) using the colors from the putative COI species suggested by KoT
and (2) using the colors from the consensus species delimitation obtained from the nuclear markers
using CoMa.

As the conspecificity matrix analysis gave consistent results between the filtered and complete
datasets, only the matrix built from the filtered dataset (n = 141) is presented (Fig. 3) for clarity. Based
on the CoMa clustering analysis using only the nuclear markers, a total of four genetic clusters (or
FFRs) were delimited in the genus *Corbicula* (including individuals from 17 distinct populations or
invasive lineages, see Table S3), suggesting a substantial allele sharing between putative species.

The first genetic cluster delineated by the analysis, FFR1, contained the androgenetic *Corbicula*
lineages from Vietnam, and was well supported by the dendrogram (Figure 3). In the haplowebs, this
cluster is clearly separated for all three nuclear markers (Fig. 2 and S2, dark pink clusters).

For all three nuclear markers, the brackish water, sexual species *C. japonica* clustered separately
from other clams in both the conspecificity matrix and haplowebs, with no sharing pattern with any
other cluster, and is defined as FFR2 based on the dendrogram (Fig.2, 3 and S2, grey clusters).

FFR3 contains *C. fluminalis africana* from South Africa and the invasive *C.* sp. form C/S. These
two lineages formed a unique block in the conspecificity matrix, but are independent lineages based on
the dendrogram (Figure 3). In the haplowebs, *C. fluminalis africana* and form C/S clustered together
for the genes *amy* and *ATPS*, but not for 28S, with the 28S alleles of these two lineages differing only
by 1 to 3 SNPs (Fig. 2a and S2a, orange cluster).

FFR4 was the largest FFR in all haploweb analyses and the conspecificity matrix. In the complete
dataset, it included the sexual lineage *C. sandai* from Lake Biwa as well as androgenetic individuals

across the entire geographic distribution of the genus *Corbicula*, notably China (28S, *ATPS*), Taiwan (*amy*), Korea (*ATPS*), Japan (28S, *amy*, *ATPS*) including Lake Biwa (*amy*, *ATPS*), Hawaii (28S, *ATPS*), the Philippines (*ATPS*), Thailand (*ATPS*), Madagascar (28S), and at least three out of the four studied invasive forms from Europe and America (forms A/R, B, and Rlc; Fig. 2 and S2, blue clusters). These species and lineages clustered together in the conspecificity matrix and therefore formed a fourth distinct group, FFR4, based on the dendrogram (Figure 3). In the filtered dataset, it included the three invasive forms and five populations from Japan (Fig. 2, Table S3).

Three of the four FFRs identified in the CoMa analysis were also supported by mitochondrial data. In both the filtered (Fig. 2d) and complete (Fig. S2d) datasets, FFR1, 2, and 3 corresponded to well-clustered sets of COI haplotypes, with a maximum of 4, 6, and 9 SNPs between two haplotypes, respectively, in the complete dataset. However, the FFR4 included highly variable COI haplotypes, with up to 43 SNPs (complete dataset) between two haplotypes from this FFR. The clusterization based on KoT analyses of the COI dataset was represented on both the corresponding mitochondrial haplonet and the nuclear haplowebs (Fig. 2 and S2). For FFR1 and FFR2, the delimitations made by the KoT analyses on the COI dataset (Table S4 and S5) are consistent with the ones from CoMa: all individuals from Vietnamese populations were in a distinct cluster (corresponding to FFR1), and all *C. japonica* individuals were another distinct cluster (corresponding to FFR2). FFR3 was however split into two genetic clusters (orange and brown on Fig. 2) based on COI, because the FW17 haplotype found in *C.* sp. form C/S differs by 7 to 9 SNPs from the sequences found in *C. fluminalis africana*. Finally, FFR4 found by the CoMa analysis is divided into four distinct lineages based on the KoT analysis of COI on the filtered dataset (Table S5) and … lineages based on the complete dataset, with all three invasive lineages (*C.* sp. Form A/R, B and Rlc) having distinct COI sequences and placed in distinct groups. The form A/R clustered with populations from Japan only (CFG, CFY, KMT, LB). In the analysis based on the filtered dataset (Table S5), the form Rlc clustered alone, the form B clustered with native populations from Japan only (EHM, CFG), and the Japanese Cl population clustered alone, while in the complete dataset (Table S4), forms B, Rlc and Japanese populations (EHM, CFG, Cl, LB) clustered together. Additional clusters delimited in Table S5 were the populations from Madagascar, China, Indonesia, with *C. sandai* clustering alone as an independent lineage.

In the end, we observed more genetic clusters within *Corbicula* defined by the COI marker than by the nuclear markers. The KoT analysis using the COI marker defined 11 and 8 putative species based on the complete (n=259, Table S4) and filtered (n=141, Table S5) datasets, respectively. Due to introgression events during androgenetic reproduction between different lineages, distinct COI haplotypes can be retrieved within a nuclear pool of shared alleles. The nuclear markers only delimited 4 genetic clusters in the present study, two of them including only geographically restricted lineages, the two others encompassing lineages from the native and invasive range that were previously described as distinct species. This suggests: i) at least two biogeographic origins of invasive lineages, and ii) a substantial genetic mixing between distinct *Corbicula* lineages.

Introgressions occurred between three out of the four FFRs of Corbicula

Alleles shared between FFRs (as defined by the conspecificity matrix) can be identified and visualized using the haploweb of each nuclear marker. The sexual *C. japonica,* FFR2, does not share alleles with other clusters (Figures 2, 3 and 4). However, alleles are shared between the three other FFRs.

First, shared alleles are observed between FFR1 and FFR3 for the *amy* marker (Fig. 2b, S2b and Fig. 4), because two Vietnamese individuals (FFR1: Vt03 and CR04) shared the amy-C1b allele with

the invasive form C/S and with all individuals from South Africa (FFR3) (Table S2, S3). In the conspecificity matrix, this sharing pattern of one marker was observed and is indicated as zone *a* (Fig. 3).

Second, several sharing patterns were observed between FFR3 and FFR4. The invasive forms A/R and C/S shared an allele for two distinct markers, 28S (Fig. 2a and S2a) and ATPS (Fig. 2b and S2b). In the conspecificity matrix, a sharing pattern for the two markers was therefore highlighted (Fig. 3, zone *b*). For the 28S marker, the shared allele between forms A/R and C/S is 28S-IA1 (Table S2, S3). As form A/R shared one or more of his 28S alleles (28S-IA1, 28S-IA2 and 28S-IA3) with all other populations from FFR4 (Table S3), a sharing pattern of one marker was observed between form C/S from FFR3 and all individuals from FFR4, with allele 28S-IA1 being the link (Fig. 3, zone *c*). For the ATPS marker, the shared allele between forms A/R and C/S is ATPS-IA1. Since form C/S shared its second ATPS allele (ATPS-C1a) with all sequenced individuals of *C. fluminalis africana* from FFR3, a sharing pattern of one marker was observed between form A/R from FFR4 and all *C. fluminalis africana* individuals from FFR3 (Fig. 3, zone *d*).

These sharing patterns between FFRs can also be visualized using a Circos plot (Fig. 4), where the color code of the four FFRs have been applied to the circle arcs representing the groups (idiotypes). Notably, only a very small fraction (2 out of 60) of the individuals from Vietnam (FFR1) were implied in the sharing with FFR3 (including form C/S and *C. fluminalis africana*), which represented 2 out of the 262 alleles retrieved from this location (total number, Fig. 4). In FFR3, all individuals for which the *amy* marker was successfully sequenced had this shared allele with FFR1. The possible introgressions between FFR3 and FFR4 can also be characterized based on the Circos plot, implying only non-exclusive invasive alleles (in purple, Fig. 4) which are never directly found in *C. fluminalis africana*.

Substantial allele sharing occurred between distinct Corbicula *lineages*

In addition to the previously described allele sharing with form C/S clustering in a distinct FFR, form A/R also shared alleles, for the three nuclear markers tested, with the two other invasive lineages, forms B and Rlc. In total, the four invasive forms A/R, B, C/S, and Rlc from Europe and America shared alleles with each other and with *Corbicula* individuals from 19 different locations in Asia and Africa (Table S2), including the sexual lineage *C. sandai* (Fig. 4, n = 359). Both exclusive and non-exclusive invasive alleles are shared.

In *Corbicula* form A/R, few variations between tested individuals were found within each marker at the nuclear and mitochondrial level, but 7 alleles (out of 12 for all markers) were shared with other invasive forms (non-exclusive invasive alleles, bold purple in Table S2 and Fig. 4). Three other alleles (exclusive invasive alleles) are shared only with native populations (bold red in Table S2 and Fig. 4) and two other alleles are private (red in Table S2). The invasive *Corbicula* form A/R therefore shared alleles with the three other invasive forms (C/S, B, and Rlc) and with freshwater lineages from all studied countries except Vietnam, Indonesia and South Africa (Fig. 4).

Corbicula form B also showed low variability between tested individuals and included only 11 distinct haplotypes (across all four markers). Among them, 3 are non-exclusive invasive alleles shared with form A/R or form Rlc (bold purple in Table S2 and Fig. 4), 7 alleles are shared with native populations only (bold blue in Table S2 and Fig. 4), and 1 is a private allele (blue in Table S2). Form B shared alleles with two invasive forms and all freshwater groups defined in this analysis, except populations from Waal (Netherlands), Madagascar, Vietnam, and South Africa.

A total of 8 haplotypes (for all 4 markers) were retrieved in *Corbicula* form Rlc with almost no
variability between tested individuals. Four non-exclusive invasive alleles (bold purple in Table S2
and Fig. 4, red for COI) and 4 exclusive invasive alleles (bold green in Table S2 and Fig. 4) were
found, but no private alleles. Invasive form Rlc shared alleles with forms A/R and form B but not form
C/S, as well as with all native freshwater groups defined in this analysis except the Vietnamese and
South African populations.

Within *Corbicula* form C/S, no variability was detected between tested individuals except for one
individual from Argentina and one from the Netherlands (Table S2). A total of 12 haplotypes (for all 4
markers) were recorded in form C/S with 2 non-exclusive invasive alleles (bold purple in Table S2 and
Fig. 4), 6 exclusive alleles shared with native populations (bold orange in Table S2 and Fig. 4), and 4
private sequences (orange in Table S2). This invasive form only shared alleles with form A/R and
Waal (Netherlands) populations, but not with any other invasive lineage. It also shared alleles with
individuals from Hawaii, China, and Japan, but never exclusive alleles, and it is the only invasive form
which does not share any haplotypes with several Japanese populations (EHM, KMT, CFG), including
Lake Biwa where sexual *C. sandai* is found. Form C/S is also the only invasive lineage sharing alleles
with Vietnamese populations and extensively with *C. fluminalis africana* (*i.e.,* these shared alleles are
carried by all individuals from both *C. fluminalis africana* and form C/S).

Finally, within the 2 *Corbicula* individuals collected from the Waal river in the Netherlands –
whose invasive form could not be identified by morphology – a total of 9 alleles (for all 3 nuclear
markers) were retrieved: 3 of them were non-exclusive invasive alleles (bold purple in Table S2 and
Fig. 4), and 6 were exclusive to form C/S (bold orange in Table S2 and Fig. 4). Although this
population shared alleles with forms A/R, Rlc, and with native populations, its genotype is most
closely related to form C/S: 8 out of its 9 shared alleles were identical to those from form C/S. For this
reason, it is represented in orange in Figure 4.

In general, among the seven non-exclusive invasive alleles retrieved for the three markers (3 in
28S, 3 in *amy*, 1 in *ATPS*), one (amy-IA3) was shared between form B and form Rlc while the six
others were found in form A/R shared with the three other forms (form C/S with 28S-IA1 and ATPS-
IA1; form B with amy-IA1 and amy-IA2; form Rlc with 28S-IA2 and 28S-IA3). Even the FW5 COI
haplotype, reported as unique to the form A/R, was found within Db25, whose morphology and
nuclear markers correspond to form Rlc. As a result, we found more non-exclusive haplotypes (70/89
if all individuals are considered) than exclusive ones within form A/R (Figure 4; Table S2). The
proportion of non-exclusive alleles were lower for the other invasive lineages: 21/61 in form B, 12/31
in form Rlc, and 19/87 in form C/S.

Corbicula fluminalis africana only shared exclusive invasive alleles (amy-C1b and ATPS-C1a)
with form C/S and the Waal (Netherlands) population. One of them (amy-C1b) was also identified in
two individuals (Vt03 and CR04) from Vietnam (Fig. 4, Table S2).

All native freshwater *Corbicula* lineages – except Vietnamese populations and *C. fluminalis*
africana, which were included in distinct FFRs – shared alleles with each other (Fig. 4). Interestingly,
this allele sharing between native *Corbicula* lineages also includes invasive alleles, notably 28S-IA3
which is responsible for the sharing observed between native populations from distinct countries such
as Madagascar, China, and Japan, including *C. sandai* (Table S2). In addition, the populations from
southeast Asia shared the amy-IA2 allele with Japan including Lake Biwa, and the ATPS-AA12c
allele with Hawaii (Table S2).

Discussion

Substantial allele sharing but distinct biogeographic origins of invasive alleles within Corbicula

Species delimitation in *Corbicula* has been fraught with conflict, in part because shell morphology alone is not a suitable parameter to define species due to its phenotypic plasticity (Glaubrecht *et al.* 2007). Moreover, species delimitation using genetic markers and phylogenetic analyses has similarly been challenged by conflicting signals between nuclear and mitochondrial markers (see Hedtke *et al.* 2008, 2011; Pigneur *et al.* 2011a, 2011b, 2014). *Corbicula* taxonomy and the specific status of most freshwater lineages are therefore still debated and there is no consensus on how to name a particular *Corbicula* species. Even the nomenclature for the invasive androgenetic lineages used here (form A/R, form B, form Rlc, and form C/S; Siripattrawan *et al.* 2000; Lee *et al.* 2005, Marescaux *et al.* 2010) could be challenged (*e.g.,* form Rlc/B, Peñarrubia *et al.*, 2017), and new forms are still described (form D, Tiemann *et al.* 2017; Haponski and Foighil 2019).

The haploweb approach used here delineated groups of shared alleles on a worldwide sampling dataset of *Corbicula* clams and found a very limited number of genetic clusters (or "fields for recombination", FFRs) within this genus, based on three nuclear markers tested (Fig. 2, 3, S2). We observed a single allelic pool encompassing most *Corbicula* freshwater lineages worldwide (Fig. 4). These results highlight the cross-species substantial genetic mixing and introgression events in the clam genus *Corbicula* that have hampered taxonomic species delimitation.

Phylogenetic trees have suggested separate geographic origins for the invasive lineages, with form A/R originating in Japan, forms B and Rlc from the Asian mainland (China, Korea, Vietnam), and form C/S from a *C. moltkiana*-like ancestor from Indonesia, although the latter also clustered with other *Corbicula* lineages depending on the marker used (Siripattrawan *et al.* 2000, Park & Kim 2003, Lee *et al.* 2005, Hedtke *et al.* 2011, Pigneur *et al.,* 2014). The allele sharing pattern observed in this study offers an alternative hypothesis.

On one hand, the invasive *C.* sp. forms A/R, B, and Rlc shared extensively their exclusive alleles (*i.e.*, alleles found in no other invasive lineage) with native populations, mainly in Japan and China but also in Korea, Thailand, the Philippines, and Madagascar (Fig. 4, Table S2). On the other hand, none of the exclusive alleles of form C/S were retrieved in these native samples. Form C/S shared exclusive alleles with nearly all studied individuals of *C. fluminalis africana* from South Africa, and with a few specimens from Vietnam (Fig. 4, Table S2), while no exclusive alleles of the other invasive forms were found in these populations. Moreover, the androgenetic individuals from the Vietnamese populations clustered separately, with the exception of two individuals that shared the amy-C4b allele with form C/S and *C. fluminalis africana* (Fig. 2, 4, S2). Given the extent of nuclear captures taking place between androgenetic *Corbicula* lineages, as highlighted in this study, it is impossible to disentangle the origin of the peculiar reproductive mode of androgenesis in *Corbicula*. Nevertheless, since the origins of the invasive lineages appear recent in comparison, we still observed a signal of distinct biogeographic origins. The allele sharing results obtained here suggest recent and at least three distinct biogeographic sources of androgenetic lineages within the genus *Corbicula*.

First, a southeastern Asian source for the Vietnamese androgenetic *Corbicula* lineages that clustered separately from the other studied individuals (FFR1, Fig. 2, 3, S2). Their sharing of an *α-amylase* allele with one individual of the Indonesian *C. tobiae* (which has unknown reproductive mode) is intriguing. Given the low number of individuals from Indonesia in the present study, the genetic relationships between Indonesian and Vietnamese clams are still unresolved. *Corbicula* populations from this geographic area need a more thorough study, especially since sexual Indonesian

Corbicula lineages have been reported (Glaubrecht *et al.* 2006). The sharing of one *amy* allele (amy-
C4b) between Vietnamese populations, form C/S, and *C. fluminalis africana* (Fig. 2, S2, Table S2, S3)
is probably the consequence of a nuclear capture event during an invasion process, although these
populations have not been reported to occur in sympatry.

Second, the invasive form C/S showed a clear sharing pattern with the South African *C. fluminalis*
africana individuals (whose reproductive mode could not be identified) (Fig. 3, 4). This would
confirm a distinct biogeographic invasion source for *Corbicula* form C/S from Africa, the current
native range of the species named *C. fluminalis* (Korniushin 2004). This putative African origin of
form C/S was already suggested by Pigneur *et al.* (2014) based on COI and 10 microsatellite markers.
An alternative hypothesis would be the invasion of South Africa by form C/S and the subsequent
establishment of an invasive population in this region. *C. fluminalis africana* would thus originate
from the European form C/S rather than the opposite. *Corbicula* genus already occurred in Africa
during Pleistocene (Meijer & Preece, 2000) and was still recorded in this continent approximately 50
years before its European invasion (Pilsbry & Bequaert, 1927). Historical data on *Corbicula*
distribution in Africa are thus not in favor with that hypothesis but do not exclude it. The form C/S
and/or *C. fluminalis africana* could also have originated from another unsampled population from the
native range ("ghost population"). More attention should be given to this African population in future
studies, with a more thorough sampling in the native range and identification of the reproductive mode
of these lineages.

Third, extensive allele sharing was observed between the Lake Biwa population in Japan,
including the sexual *C. sandai,* and most freshwater *Corbicula* individuals studied here, including
three out of the four invasive androgenetic lineages (forms A/R, B, and Rlc) (Fig. 3, 4). The alleles
found in these three invasive forms are found in numerous native locations, including several Asian
populations, Madagascar, and Hawaii, although the majority of them are found in Japan, including
Lake Biwa. Because these alleles are identical across a wide geographic distribution of *Corbicula*, the
shared history appears to be recent. Interestingly, within this cluster (FFR4), the population from Lake
Biwa is particularly genetically diverse, with numerous private alleles, but also encompassing most
alleles found in invasive lineages A/R, B, and Rlc (Table S2, S3). Our analysis therefore points to
Japan as a hotspot and distinct biogeographic source of most invasive androgenetic *Corbicula*
lineages, particularly Lake Biwa, where sexual *C. sandai* occurs (Houki *et al.*, 2011). Repeated origins
of asexual clones from a sexual relative have been observed in other taxa where sexual and asexual
individuals co-occur (Simon *et al.*, 2003; Bode *et al.*, 2010; Neiman *et al.*, 2010). Moreover, the
studies of Komaru *et al.* (2012, 2013) found that androgenetic hermaphroditic individuals of *C. leana*
shared 28S rDNA alleles with sexual *C. sandai*, both co-occurring in Lake Biwa. Nevertheless, we
cannot argue that *C. sandai* itself is the direct ancestor of the studied androgenetic lineages as a
number of morphological, cytological, and biological features (*e.g.* biflagellate *vs.* uniflagellate sperm,
dioecy *vs.* hermaphroditism, non-brooding *vs.* brooding) distinguish it from androgenetic clams
(Glaubrecht *et al.* 2006; Obata *et al.* 2006). More research in Lake Biwa populations, using genome-
wide markers, is required to i) confirm the recent biogeographic origin of invasive lineages suggested
by the present work and ii) understand the older origin of androgenetic reproductive mode in
Corbicula genus.

Genetic mixing between invasive Corbicula *lineages*

While the invasive form C/S found in Europe and South America may have a distinct
biogeographic origin in Africa, the sharing of alleles between form C/S and form A/R (Fig. 4, Table
S2) is probably the result of secondary nuclear captures when found in sympatry. Indeed, form C/S

shared two alleles (28S-IA1 and ATPS-IA1) with the invasive form A/R. This suggests potential post-
introductory nuclear genetic exchanges between forms A/R and C/S when found in sympatry (*e.g.* Lee
et al., 2005; Hedtke *et al.*, 2008; Pigneur *et al.*, 2011, 2014).

Individuals from invasive form B shared alleles with form A/R (amy-IA1 and 2) and form Rlc
(amy-IA3), the latter two also sharing two alleles with each other (28S-IA2 and 3). Allele sharing
between form A/R and B (Fig. 4, Table S2) probably represents introgression events when found in
sympatry, as observed by Hedtke *et al.* (2008). However, form B and form Rlc were almost never
reported to occur in sympatry: form B is collected in America and form Rlc in Europe. These forms
do, however, share a very similar COI haplotype, with only 1 SNP difference over 650 bp, suggesting
a plausible common origin of these forms that later diverged when invading America and Europe.
More recently, the FW1 COI haplotype from form B was found in Europe by Penarrubia *et al.* 2017.
These authors considered forms B and Rlc as a unique form ("RlcB"), hypothesizing that alleles
shared between these two forms could be due to a close ancestry. More research in the invasive range
of *Corbicula* genus, using more sampling sites and genome-wide markers and/or cytology, would
provide insights on the mechanism of hybridization between distinct lineages.

Furthermore, a sampling of invasive *Corbicula* lineages in the Iberian Peninsula revealed many
heterozygous individuals, with 28S alleles found in form A/R and either form Rlc or form C/S. In
some cases, these heterozygotes were associated with the COI haplotype of a third lineage (Peñarrubia
et al. 2017). Recently, in a man-made channel in Northern European Russia, all sampled *Corbicula*
individuals belonged to two distinct invasive lineages apparently fixed for a mismatch between the
mtDNA haplotype and morphotype. The first lineage belonged to morphotype Rlc associated with the
FW5 haplotype of form A/R, while the second lineage had morphotype A/R associated with FW17
COI haplotype of form C/S (Bespalaya *et al.* 2018). Such cytonuclear mismatches, but also hybrids,
have been retrieved in several locations in Europe where form A/R occurs in sympatry with another
invasive form (C/S or Rlc) (Pigneur *et al.* 2014; Etoundi, pers. obs.). All these results highlight that
nuclear genetic exchanges can occur between invasive *Corbicula* lineages when found in sympatry. In
the present study, the two *Corbicula* individuals collected from the Waal River in Europe showed an
intermediate phenotype between form A/R and form C/S (unpublished data) and contain alleles from
both forms, although a majority was from form C/S, suggesting introgressions occurred after these
lineages were introduced. A cytonuclear mismatch was also observed in Db25, carrying all nuclear
alleles of form Rlc associated with the COI haplotype of form A/R (Fig. 2d, S2d, Table S2, S3).

Previous studies on *Corbicula* that used phylogenetics to discriminate species observed
cytonuclear discordances that confounded species delimitation (*e.g.,* Hedtke *et al.* 2011). In contrast,
the approach used here identified distinct lineages that were consistent across mitochondrial and
nuclear markers. CoMa (nuclear) and KoT (mitochondrial) analyses both identified the clusters FF1
and FFR2. The two other FFRs, FFR3 and FFR4, were defined as clusters by the nuclear marker-based
analysis only (Figure 2a-c), but as independent lineages according to the COI-based analysis (Figure
2d) even though individuals within FFR3 are located in the same part of the mitochondrial haplotype
network, and individuals within FFR4 similarly share nodes with each other. Notably, the different
invasive lineages (forms A/R, B and Rlc) were delimited as distinct lineages based on the haploid
mitochondrial marker COI but were included within the same FFR because of massive sharing of
nuclear alleles between invasive forms and especially with native lineages. Similarly, form C/S and *C.*
fluminalis africana are considered distinct lineages based on COI but are included within the same
FFR. The debate that remains now is how to delineate species within a genus where frequent
introgression occurs between distinct lineages. We strongly recommend the continued use of both

mitochondrial and nuclear markers in *Corbicula* when studying genetic relationships and when
defining taxa.

Introgressions between invasive and native Corbicula *lineages*

The genetic diversity within the four studied androgenetic *Corbicula* lineages in the invaded range
in Europe and America is very low (reviewed in Pigneur *et al.*, 2012), while androgenetic *Corbicula*
lineages found in the native range, particularly Asia, are both genetically and morphologically diverse
(*e.g.* Glaubrecht *et al.*, 2003; Hedtke *et al.*, 2011; Pigneur *et al.*, 2014; Table S2), possibly due to
multiple origins from sexual species or multiple introgression events when occurring in sympatry.
Interestingly, the most widespread alleles (*e.g.:* 28S-IA3, 28S-AB11a, amy-Db22a, ATPS-AB11a)
shared between lineages and populations from FFR4 belong to the invasive forms A/R, B, and Rlc and
are also shared with *Corbicula* from Japan, China, Madagascar, and Hawaii. Two hypotheses could
explain this pattern: (i) these widespread alleles represent shared ancestral alleles present in the native
range before the invasion of other Asian countries, America, and Europe; or (ii) these alleles are
signatures of nuclear genetic exchanges in contact zones, or both.

A surprisingly higher rate of homozygosity was observed at the *ATPS* locus compared to the 28S
and *amy* markers, even in populations that are genetically diverse, such as Lake Biwa. Moreover, we
were not able to amplify sequences from the *C. sandai* population and a few others (Table S2),
suggesting that there could be mutations in the primer binding sites that prevented amplification (*i.e.,*
null alleles), which would result in an apparent homozygosity. However, at the other two loci (*amy*
and 28S), both invasive and native androgenetic *Corbicula* individuals are mostly heterozygotes (also
confirmed by Hedtke et al., 2011; Pigneur *et al.*, 2014; Peñarrubia *et al.* 2017). One explanation for
this observation might be the "Meselson effect", in which allelic diversity evolves over time in the
absence of recombination (Judson & Normark 1996; Birky 1996). However, this divergence would
probably impact all alleles in a lineage and would therefore degrade the signal of shared alleles among
lineages (Hedtke et al. 2011); we observe instead that identical sequences are shared across sexual and
androgenetic lineages. A more plausible explanation for this high heterozygosity is the occurrence of
nuclear captures between different lineages. Polyploidy is found across androgenetic *Corbicula*
populations (Okamoto & Arimoto 1986, Qiu *et al.* 2001, Ishibashi *et al.* 2003, Komaru *et al.* 2006;
Hedtke *et al.* 2008), and up to four different alleles per individual were observed in some populations
(Table S2). Egg parasitism by one androgenetic *Corbicula* lineage of a distinct lineage could be
accompanied not only by the capture of maternal mitochondrial DNA but also of nuclear DNA,
increasing the ploidy and the genetic diversity of hybrids (Figure 1) (Hedtke *et al.* 2011, Peñarrubia *et*
al. 2017, Bespalaya *et al*. 2018). We hypothesize that the independent androgenetic *Corbicula*
lineages confirmed here originated after hybridization from partial nuclear capture following egg
parasitism. These nuclear genetic capture events between lineages, which occurs because these asexual
organisms have retained sexual features such as fertilization (Figure 1), might facilitate the evolution
and maintenance of genetic diversity through enhanced heterozygosity, with alleles from different
genetic lineages being brought together within one individual. This hybridization can occur where
distinct androgenetic species co-occur, but also in populations where sexuals and asexuals are found in
sympatry (such as Lake Biwa in Japan).

The consequences of androgenetic egg parasitism for chromosome organization, and thus the
extent of allele capture, are still unclear in *Corbicula*. Nuclear captures between individual clams have
been observed to cause the retention of a complete haploid maternal genome, *i.e.* one copy of each
chromosome from the mother combined with the diploid or triploid paternal genome (Figure 1b;
Komaru et al. 2006). However, in the present study, results from three nuclear loci were not

completely consistent with each other, as they led to the description of different allelic pools (Figures 2a, b, and c). This might be explained by non-uniform nuclear capture, with only one or some chromosomes retained from the mother and others discarded during meiosis. Such non-uniform nuclear captures would result in aneuploidy (imbalanced chromosome numbers), but research on *Corbicula* karyotypes have only reported balanced chromosome numbers (two, three, or four copies for all chromosomes, see Okamoto & Arimoto 1986; Komaru *et al.*, 1997; Park *et al.*, 2000; Qiu *et al.*, 2001, Skuza *et al.* 2009). Another explanation for the incongruent nuclear allele pools between markers would be the occurrence of crossing-over and recombination during gamete (*i.e.,* sperm) production. Nuclear captures were detected between distinct lineages (namely hybridization), but probably also occurs within a lineage, in which case it would be harder to detect. The frequency of this process could thus be underestimated in *Corbicula*. Further research exploring chromosome diversity and gamete production is needed to disentangle the mechanism of nuclear captures in *Corbicula*. This could also shed light on other systems, as androgenesis is associated with peculiarities such as hybridization and/or polyploidy and in some systems the occurrence of closely related sexual species, not only within the genus *Corbicula* but also in very distinct organisms (*e.g.,* stick-insects *Clonopsis* and *Pijnackeria* genus, Milani *et al.* 2016, 2020).

Acknowledgments

We thank B. Hespeels, M. Terwagne, L. Bankers, K. McElroy, J. Sharbrough, M. Neiman, L. Keller and T. Schwander for advice on the analyses and proofreading, as well as N. Yasuda and theM. Yamaguchi for access to Japanese samples, particularly specimens from Lake Biwa. We are also thankful to all the colleagues who provided samples (C. Ayres, Y. Bagatini, L. Beran, D. Eekelers, J. Higuti, C. Ituarte, K. de Koch, P. Laforge, T. Lee, S. Milla, K. Martens, J. Mouthon, N. Perrin, M.-A. Pierrard, S. Schmidlin, R. Sousa, N. Spann, E. Tinti, M. Yamaguchi and the Illinois Natural History Survey). This work was supported by the Belgian National Fund for Scientific Research (Fonds National pour la Recherche Scientifique - FNRS) through a PhD thesis grant (FC91712) to E.E., by a PDR FNRS research grant (CARMA, 14596412) to K.V.D., by UNamur for the teaching assistant grant to M.V. and by an Horizon 2020 research and innovation program of the European Union under the European Research Council (ERC) grant agreement 725998 (RHEA) to KVD financing postdoctoral researcher E.N. and technician M.C.

Conflict of interest disclosure

The authors of this article declare that they have no conflict of interest with the content of this article.

References

Akaike, H. 1974. A new look at the statistical model identification. *IEEE Trans. Automat. Contr.* **19**:716-723.

Araujo, R., Moreno, D., Ramos, M.A. 1993. The Asiatic clam *Corbicula fluminea* (Müller, 1974) (Bivalvia: Corbiculidae) in Europe. *American Malacological Bulletin*. **10**(1):39-49.

Bah, T. 2011. Inkscape. Guide to a vector drawing program. 4th edition. Prentice Hall.

Bandelt HJ, Forster P. & Röhl A. 1999. Median-joining networks for inferring intraspecific phylogenies. *Molecular Biology and Evolution* **16**:37-48.

Barbuti, R., Mautner, S., Carnevale, G., Milazzo, P., Rama, A. & Sturmbauer, C. 2012. Population dynamics with a mixed type of sexual and asexual reproduction in a fluctuating environment. *BMC Evolutionary Biology* **12**:49.

Barton, N.H. 2009 Why sex and recombination? *Cold spring Harbor Symposia on Quantitative Biology* **74**: 187-195.

Bast, J., Parker, D.J., Dumas, Z., Jalvingh, K.M., Tran Van, P., Jaron, K.S., Figuet, E., Brandt, A., Galtier, N., Schwander, T. 2018. Consequences of asexuality in natural populations: Insights from stick insects. *Mol. Biol. Evol.* **35**, 1668–1677.

Birky, C.W. Jr., Adams, J., Gemmel, M., Perry, J. 2010. Using Population Genetic Theory and DNA Sequences for Species Detection and Identification in Asexual Organisms. *PLoS ONE* **5**(5): e10609. doi:10.1371/journal.pone.0010609.

Birky, C.W.Jr. 2013. Species Detection and Identification in Sexual Organisms Using Population Genetic Theory and DNA Sequences. *PLoS ONE* **8**(1): e52544. doi:10.1371/journal.pone.0052544.

Birky, C.W. Jr., Adams, Maughan, H. 2021. Evolutionary Genetic Species Detected in Prokaryotes by Applying the K/θ1 Ratio to DNA Sequences. bioRchiv. doi: https://doi.org/10.1101/2020.04.27.062828.

Bespalaya, Y.V., Bolotov, I.N., Aksenova, O.V., Kondakov, A.V., Gofarov, M.Y., Laenko, T.M., Sokolova, S.E., Shevchenko, A.R., Travina, O.V. 2018. Aliens are moving to the Arctic frontiers: an integrative approach reveals selective expansion of androgenic hybrid *Corbicula* lineages towards the North of Russia. *Biol. Invasions*. https://doi.org/10.1007/s10530-018-1698-z

Bespalaya, Y.V., Aksenova, O.V., Gofarov, M.Y., Kondakov, Kropotin, A.V., Kononov, O.D., Bolotov, I.N. 2020. Who inhabits the world's deepset crater lake? A taxomnomic review of *Corbicula* (Bivalvia: Cyrenidae) clams from Lake Toba, North Sumatra, Indonesia. *J. Zool. Syst. Res.* **00**:1-11.

Bode, S.N.S., Adolfsson, S., Lamatsch, D.K., Martins, M.J.F., Schmit, O., Vandekerkhove, J.,
Mezquita, F., Namiotko, T., Rossetti, G., Schön, I., Butlin, R.K., Martens, K. 2010.
Exceptional cryptic diversity and multiple origins of parthenogenesis in a freshwater ostracod.
Molecular Phylogenetics and Evolution **54**:542-552.

Britton, J.C. & Morton, B. 1986. Polymorphism in *Corbicula fluminea* (Bivalvia: Corbiculoidea) from
North *American Malacological Review* **19**:1–43.

Castagnone-Sereno, P. 2006. Genetic variability and adaptive evolution in parthenogenetic root-knot
nematodes. *Heredity* **96**:282–289.

Coughlan, N.E., Cuthbert, R.N., Potts, S., Cunningham, E.M., Crane, K., Caffrey, J.M., Lucy, F.E.,
Davis, E., Dick, J.T.A. 2019. Beds Are Burning: eradication and control of invasive Asian
clam, *Corbicula fluminea*, with rapid open-flame burn treatments. *Management of Biological*
Invasions **10**(3): 486–499, https://doi.org/10.3391/mbi.2019.10.3.06.

Crespo, D., Dolbeth, M., Leston, S., Sousa, R., Pardal, M.A. 2015. Distribution of *Corbicula fluminea*
(Müller, 1774) in the invaded range: a geographic approach with notes on species traits
variability. *Biol. Invasions* **17**:2087-2101. DOI 10.1007/s10530-015-0862-y

Darriba, D., Taboada, G.L., Doallo, R. & Posada, D. 2012. jModelTest 2: more models, new heuristics

and parallel computing. *Nature Methods* **9**, 772.

Debortoli, N., Li, X., Eyres, I., Fontaneto, D., Hespeels, B., Tang, C. Q., Flot, J.-F., & Van Doninck,
K. 2016. Genetic exchange among bdelloid rotifers is more likely due to horizontal gene
transfer than to meiotic sex. *Current Biology* **26**(6), 723-732.

Delmotte, F., Sabater-Muñoz, B., Prunier-Leterme, N., Latorre, A., Sunnucks, P., Rispe, C. & Simon,
J.-C. 2003. Phylogenetic evidence for hybrid origins of asexual lineages in an aphid species.
Evolution **57**:1291-1303.

Dimijian, G.G. 2005. Evolution of Sexuality: Biology and Behavior. *Baylor University Medical*
Center Proceedings **18**:3, 244-258. DOI: 10.1080/08998280.2005.11928075.

Doyle, J.J. 1995. The irrelevance of allele tree topologies for species delimitation, and a non-
topological alternative. *Systematic Botany* **20**:574-588.

Felsenstein, J. 1974. The evolutionary advantage of recombination. *Genetics* **1**:737-756.

Fisher, R.A. 1930. The genetical theory of natural selection. Clarendon, Oxdord.

Flot, J.F., Tillier, A., Samadi, S. & Tillier, S. 2006. Phase determination from direct sequencing of
length-variable DNA regions. *Molecular Ecology Resources* **6**:627-630.

Flot, J.F. 2007. Champuru 1.0: a computer software for unraveling mixtures of two DNA sequences of
unequal lengths. *Molecular Ecology Resources* **7**:974-977.

Flot, J.F. 2010. SeqPHASE: a web tool for interconverting PHASE input∕output files and FASTA
sequence alignments. *Molecular Ecology Resources* **10**:162–166.

Flot, J.F., Couloux, A. & Tillier, S. 2010. Haplowebs as a graphical tool for delimiting species: a
813   revival of Doyle's "field for recombination" approach and its application to the coral genus
814   *Pocillopora* in Clipperton. *BMC Evolutionary Biology* **10**:372.

815   Glaubrecht, M., von Rintelen, T. & Korniushin, A.V. 2003. Toward a systematic revision of brooding
816   freshwater Corbiculidae in Southeast Asia (Bilvavia, Veneroida): on shell morphology,
817   anatomy and molecular phylogenetics of endemic taxa from islands in Indonesia. *Malacologia*
818   **45**:1-10.

819   Glaubrecht, M., Fehér, Z., von Rintelen, T. 2006. Brooding in *Corbicula madagascariensis* (Bivalvia,
820   Corbiculidae) and the repeated evolution of viviparity in corbiculids. *Zoologica Scripta*
821   **35**:641-654.

822   Gomes, C., Sousa, R., Mendes, T., Borges, R., Vilares, P., Vasconcelos, V., Guilhermino, L., Antunes,
823   A. 2016 Low genetic diversity and high invasion success of *Corbicula fluminea* (Bivalvia,
824   Corbiculidae) (Müller, 1774) in Portugal. *PLoS ONE* **11**(7): e0158108.
825   doi:10.1371/journal.pone.0158108.

826   Gomes, C., Mendes, T., Borges, R., Guarneri, I., Marchi, I., Guilhermino, L., Vasconcelos, V.,
827   Riccardi, N., Antunes, A. 2020. The genetic diversity of two invasive sympatric bivalves
828   (*Corbicula fluminea* and *Dreissena polymorpha*) from Lakes Garda and Maggiore, Northern
829   Italy. *Journal of Great Lakes Research* **46**: 225-229.

830   Hall, T.A. 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis program
831   for Windows 95/98/NT. *Nucleic Acids Research* **41**:95-98.

832   Haponski, A.E. & Foighil, D.Ó. 2019. Phylogenomic analyses confirm a novel invasive North
833   American *Corbicula* (Bivalvia: Cyrenidae) lineage. *PeerJ* **7**:e7484 DOI 10.7717/peerj.7484.

834   Hartfield, M., Keightley, P.D. 2012. Current hypotheses for the evolution of sex and recombination.
835   *Integrative Zoology* **7**: 192–209. doi: 10.1111/j.1749-4877.2012.00284.x

836   Hedtke, S.M., Stanger-Hall, K., Baker, R.J. & Hillis, D.M. 2008. All-male asexuality: origin and
837   maintenance of androgenesis in the Asian clam *Corbicula*. *Evolution* **62**:1119-1136.

838   Hedtke, S.M. 2009. Origin and maintenance of androgenesis: male asexual reproduction in the clam
839   genus *Corbicula*. Thesis dissertation, University of Texas at Austin (129 pages).

840   Hedtke, S.M., Glaubrecht, M. & Hillis, D. 2011. Rare gene capture in predominantly androgenetic
841   species. *Proc. Natl. Acad. Sci. U.S.A.* **108**:9520-8524.

842   Hedtke, S.M. & Hillis, D. 2011. The potential role of androgenesis in cytoplasmic–nuclear
843   phylogenetic discordance. *Systematics Biology* **60**:87-109.

844   Hillis, D.M. & Patton, J.C. 1982. Morphological and electrophoretic evidence for two species of
845   *Corbicula* (Bivalvia: Corbiculidae) in North America. *American Midland Naturalist* **108**:74-
846   80.

847   Hotta M. & Komaru A. 2018. The process of first polar body formation in eggs of the androgenetic
848   clam *Corbicula fluminea*. *Journal of Shellfish Research* **37**: 131-137.

Houki, S., Yamada, M., Honda, T. & Komaru, A. 2011. Origin and possible role of males in
hermaphroditic androgenetic *Corbicula* clams. *Zoological Science* **28**:526-531.

Huelsenbeck, J.P. & Ronquist, F. 2001. MRBAYES: Bayesian inference of phylogenetic trees.

Bioinformatics **17**:754-755.

Hurvich, C.M. & Tsai, C.L. 1989. Regression and time series model selection in small samples.

Biometrika **76**:297-307.

Ishibashi, R., Komaru, A., Ookubo, K., Kiyomoto, M. 2002. The second meiosis occurs in
cytochalasin D-treated eggs of Corbicula leana even though it is not observed in control
androgenetic eggs because the maternal chromosomes and centrosomes are extruded at first
meiosis. *Dev. Biol.* **244:** 37–43.

Ishibashi, R., Ookubo, K., Aoki, M., Utaki, M., Komaru, A. & Kawamura, K. 2003. Androgenetic
reproduction in a freshwater diploid clam *Corbicula fluminea* (Bivalvia: Corbiculidae). *Zool.*
Sci. **20**:727-732.

Ishibashi, R. & Komaru, A. 2006. Abortive second meiosis detected in cytochalasin-treated eggs in
androgenetic diploid *Corbicula fluminea. Develop. Growth Differ.* **48**: 277-282.

Katoh, K., Rozewicki, J., Yamada, K.D. 2017. MAFFT online service: multiple sequence alignment,
interactive sequence choice and visualization. *Briefings in Bioinformatics*, bbx108,
https://doi.org/10.1093/bib/bbx108.

Komaru, A., Konishi, K., Nakayama I., Kobayashi, T., Sakai, H., & Kawamura, K. 1997.
Hermaphroditic freshwater clams in the genus *Corbicula* produce non-reductional
spermatozoa with somatic DNA content. *Biology Bulletin* **193**:320-323.

Komaru, A. & Konishi, K. 1999. Non-reductional spermatozoa in three shell color types of the
freshwater clam *Corbicula fluminea* in Taiwan. *Zoological Science* **16**:105-108.

Komaru, A., Ookubo, K. & Kiyomoto, M. 2000. All meiotic chromosomes and both centrosomes at
spindle pole in the zygotes discarded as two polar bodies in clam *Corbicula leana*: unusual
polar body formation observed by antitubulin immunofluorescence. *Developmental Genes and*
Evolution **210**:263-269.

Komaru, A., Kumamoto, A., Kato, T., Ishibashi, R., Obata, M. & Nemoto, T. 2006. A hypothesis of
ploidy elevation by formation of a female pronucleus in the androgenetic clam *Corbicula*
fluminea in the Tone River estuary, Japan. *Zoological Science* **23**: 529-532.

Komaru, A., Houki, S., Yamada, M., Miyake, T., Obata, M., Kawamura, K. 2012. 28S rDNA
haplotypes of males are distinct from those of androgenetic hermaphrodites in the clam
Corbicula leana. *Dev. Genes Evol*. **222**:181–187.

Komaru, A., Yamada, M., Houki, S. 2013. Relationship Between Two Androgenetic Clam Species,
Corbicula leana and *Corbicula fluminea*, Inferred from Mitochondrial Cytochrome b and
Nuclear 28S rRNA Markers. *Zoological Science*. **30**(5):360-365.

Kondrashov, A.S. 1988. Deleterious mutations and the evolution of sexual reproduction. *Nature*
336:435-440.

Korniushin, A.V. 2004. A revision of some Asian and African freshwater clams assigned to *Corbicula*
fluminalis (Müller, 1774) (Mollusca: Bivalvia: Corbiculidae), with a review of anatomical
characters and reproductive features based on museum collections. *Hydrobiologia* **529**:251-
270.

Kraemer, L.R., Swanson, C., Galloway, M. & Kraemer, R. 1986. Biological basis of behavior in
Corbicula fluminea. II. Functional morphology of reproduction and development and review
of evidence for self-fertilization. *American Malacolical Bulletin* Spec. Ed., **2**:193-202.

Krzywinski, M.I., Schein, J.E., Birol, I., Connors, J., Gascoyne R, Horsman, D., *et al.* 2009. Circos:
An information aesthetic for comparative genomics. *Genome Research* **19**:1639-1645.

Lee, T., Siripattrawan, S., Ituarte, C.F. & Ò Foighil, D. 2005. Invasion of the clonal clams: *Corbicula*
lineages in the New World. *American Malacological Bulletin.* **20**:113-122.

Lehtonen, J., Jennions, M.D., Kokko, H. 2012. The many costs of sex. *Trends in Ecology and*
Evolution **27**(3): 172-178.

Lehtonen, J., Schmidt, D.J., Heubel, K. & Kokko, H. 2013. Evolutionary and ecological implications
of sexual parasitism. *Trends in Ecology and Evolution* **28**: 297-306.

Lenormand, T., Engelstädter, J., Johnston, S.E., Wijnker, E., Haag, C.R. 2016. Evolutionary mysteries
in meiosis. *Phil. Trans. R. Soc. B* **371**: 20160001. http://dx.doi.org/10.1098/rstb.2016.0001.

Librado, P. & Rozas, J. 2009. DnaSP v5: A software for comprehensive analysis of DNA
polymorphism data. *Bioinformatics* **25**:1451-1452.

Liegeois, M., Sartori, M. & Schwander, T. 2020. Extremely Widespread Parthenogenesis and a Trade-
Off Between Alternative Forms of Reproduction in Mayflies (Ephemeroptera). *J. Hered.*
doi:10.1093/jhered/esaa027.

López-Soriano, J., Quiñonero-Salgado, S., Cappelletti, C., Faccenda, F., Ciutti, F. 2018. Unraveling
the complexity of *Corbicula* clams invasion in Lake Garda (Italy). *Advances in Oceanography*
and Limnology **9**(2): 97-104. DOI: 10.4081/aiol.2018.7857.

Mantovani, B., Passamonti, M. & Scali, V. 2001. The mitochondrial cytochrome oxidase II gene in
Bacillus stick insects: ancestry of hybrids, androgenesis, and phylogenetic relationships.
Molecular Phylogenetics and Evolution **19**:157–163.

Marescaux, J., Pigneur, L.M. & Van Doninck, K. 2010. New records of Asian clams *Corbicula* spp. in
French Rivers. *Aquatic Invasions* **5**:S35-S39.

Maynard Smith, J. 1978. The Evolution of Sex. *Cambridge University Press, Cambridge.*

McKone, M.J., Halpern, S.L. 2003. The Evolution of Androgenesis. *The American Naturalist* **161**(4):
641-656.

McMahon, R.F. 1982. The occurrence and spread of the introduced Asiatic freshwater clam,
Corbicula fluminea (Muller), in North America: 1924–1982. *Nautilus.* **96**:134–141.

Meijer, T. & R. C. Preece, 2000. A review of the occurrence of *Corbicula* in the Pleistocene of North–
West Europe. *Geologie en Mijnbouw (Netherlands Journal of Geo-sciences)* **79**(2/3): 241–
255.

Milani, L., Scali, V., Passamonti, M. 2016. Speciation through androgenesis in the stick insect genus
Clonopsis (Insecta Phasmatodea). *J. Zoolog. Syst. Evol. Res.* doi: 10.1111/jzs.12087.

Milani, L., Scali, V., Punzi, E., Luchetti, A., Ghiselli, F. 2020. The puzzling taxonomic rank of
Pijnackeria hispanica, a chimerical hybrid androgen (Insecta, Phasmida). *Organisms Diversity*
& Evolution. **20**:285–297.

Minchin, D. 2014. The distribution of the Asian clam *Corbicula fluminea* and its potential to spread in
Ireland. *Management of Biological Invasions* **5**(2): 165–177.

Mouthon, J. 1981. Sur la présence en France et au Portugal de *Corbicula* (Bivalvia, Corbiculidae)
originaire d'Asie. *Basteria*. **45**:109–116.

Muller, H.J. 1932. Some genetic aspects of sex. *American Naturalist* **66**:118-138.

Muller, H.J. 1964. The relation of recombination to mutational advance. *Mutation Research.* **1**:2–9.

Neiman, M., Hehman, G., Miller, J.T., Logsdon J.M., Jr. & Taylor, D.R. 2010. Accelerated mutation
accumulation in asexual lineages of a freshwater snail. *Molecular Biology and*
*Evol*ution**27**:954–963.

Neiman, M., Schwander, T. 2011. Using parthenogenetic lineages to identify advantages of sex.
*Evolutionary Biol*ogy **38**: 115-123.

Obata, M., Nishimori, K., Komaru, A. 2006. Change of Centrosome Attachment Site Causes
Androgenesis in the Freshwater Clam *Corbicula fluminea*: Comparison with *C. sandai. Venus*
65(3): 247-257.

Okamoto, A. & Arimoto, B. 1986. Chromosomes of *Corbicula japonica, C. sandai* and *C.*
(Corbiculina) leana (Bivalvia: Corbiculidae). *Venus Jpn J. Malacol.* **45**:194–202.

Otto, S.P., 2008. Sexual reproduction and the evolution of sex. *Nature Education* **1**:182.

Paczesniak, D., Jokela, J., Larkin, K. & Neiman, M. 2013. Discordance between nuclear and
mitochondrial genomes in sexual and asexual lineages of the freshwater snail *Potamopyrgus*
antipodarum. Molecular Ecology **22**: 4695-4710.

Park, G., Yong, T., Im, K., Chung, E. 2000. Karyotypes of three species of *Corbicula* (Bivalvia:
Veneroida) In Korea. *Journal of Shellfish Research* **19**(2): 979-982.

Park, J.K. & Kim, W. 2003. Two *Corbicula* (Corbiculidae: Bivalvia) mitochondrial lineages are
widely distributed in Asian freshwater environment. *Molecular Phylogenetics and Evolution*
29: 529-539.

Patrick, C.H., Waters, M.N., Golladay, S.W. 2017. The distribution and ecological role of *Corbicula*
fluminea (Müller, 1774) in a large and shallow reservoir. *BioInvasions Records* **6**(1): 39–48.

Peñarrubia, L., Araguas, R.M., Vidal, O., Pla, C., Viñas, J., Sanz, N. 2017. Genetic characterization of
the Asian clam species complex (*Corbicula*) invasion in the Iberian Peninsula. *Hydrobiologia*.
DOI 10.1007/s10750-016-2888-2.

Pfenninger, M., Reinhardt, F., Streit, B. 2002. Evidence for cryptic hybridization between different
evolutionary lineages of the invasive clam genus *Corbicula* (Veneroida, Bivalvia). *Journal of
Evolutionary Biology* **15**: 818-829.

Pichot, C., El Maâtaoui, M., Raddi, S., Raddi, P. 2001. Surrogate mother for endangered *Cupressus.*
Nature **412**(5): 39.

Pigneur, L.-M., Marescaux, J., Roland, K., Etoundi, E., Descy J.-P. & Van Doninck, K. 2011a.
Phylogeny and androgenesis in the invasive *Corbicula* clams (Bivalvia, Corbiculidae) in
Western-Europe. *BMC Evolutionary Biology.* **11**:147.

Pigneur, L.-M., Risterucci, A.-M., Dauchot, N., Li, X., Van Doninck, K. 2011b. Development of novel
microsatellite markers to identify the different invasive lineages in the *Corbicula* complex and
to assess androgenesis. *Molecular Ecology Resources*, **11**: 573–577.

Pigneur, L.-M., Hedtke, S.M., Etoundi, E. & Van Doninck, K. 2012. Androgenesis: a review through
the study of the selfish shellfish *Corbicula* spp. *Heredity* **108**:581-591.

Pigneur, L.-M., Etoundi, E., Aldridge, D.C., Marescaux, J., Yasuda, N. & Van Doninck, K. 2014.
Genetic uniformity and long-distance clonal dispersal in the invasive androgenetic *Corbicula*
clams. *Molecular Ecology* **23**:5102-16.

Pilsbury, H.A., & Bequaert, J. (1927) The aquatic molluscs of the Belgian Congo, with a geographical
and ecological account of Congo malacology. *Bull. Amer. Mus. Nat. Hist.* **53:** 69-602.

Qiu, A., Shi, A., Komaru, A. 2001. Yellow and Brown Shell Color Morphs of *Corbicula fluminea*
(Bivalva: Corbiculidae) from Sichuan Province, China, are triploids and tetraploids. *Journal of
Shellfish Research* **20**(1): 323-328.

R Development Core Team. 2008. R: A language and environment for statistical computing. R
Foundation for Statistical Computing, Vienna, Austria.

Renard, E., Bachmann, V., Cariou, M.L., Moreteau, J.C. 2000. Morphological and molecular
differentiation of invasive freshwater species of the genus *Corbicula* (Bivalvia, Corbiculidea)
suggest the presence of three taxa in French rivers. *Molecular Ecology.* **9**:2009–2016.

Rey, O., Facon, B., Foucaud, J., Loiseau, A., Estoup, A. 2013. Androgenesis is a maternal trait in the
invasive ant *Wasmannia auropunctata. Proc R Soc B* **280**: 20131181.
http://dx.doi.org/10.1098/rspb.2013.1181.

Reyna, P., Nori, J., Ballesteros, M.L., Hued, A.C., Tatián, M. 2018. Targeting clams: insights into the
invasive potential and current and future distribution of Asian clams. *Environmental
Conservation* 1-9. doi:10.1017/S0376892918000139.

Rintelen, T. v. & Glaubrecht, M. 2006. Rapid evolution of sessility in an endemic species flock of the
freshwater bivalve *Corbicula* from ancient lakes on Sulawesi, Indonesia. *Biology Letters.*
2:73–77.

Ronquist, F., Huelsenbeck, J. & Teslenko, M. 2001. Draft MrBayes version 3.2 Manual: tutorials and
model summaries.

Sagata, N. 1996. Meiotic metaphase arrest in animal oocytes: its mechanisms and biological
significance. *Trends in Cell Biology* **6**: 22-28.

Sano, N., Houki, S., Kodan, A., Kawamura, K., Yamada, M., Komaru, A. 2020. Genetic confirmation
of "egg parasitism" in androgenetic freshwater *Corbicula* clams by paternity testing using
microsatellite DNA markers. *Plankton Benthos Res.* **15**(1): 58-62.

Simon, J.C., Delmotte, F., Rispe, C. & Crease, T. 2003. Phylogenetic relationships between
parthenogens and their sexual relatives: the possible routes to parthenogenesis in animals.
Biological Journal of the Linnean Society **1**: 151-163.

Schön, I., Pinto, R.L., Halse, S., Smith, A.J., Martens, K., Birky, C.W.Jr. 2012. Cryptic Species in
Putative Ancient Asexual Darwinulids (Crustacea, Ostracoda). *PLoS ONE* **7**(7): e39844.
doi:10.1371/journal.pone.0039844.

Schwander, T., & Crespi B.J. 2009. Multiple direct transitions from sexual reproduction to apomictic
parthenogenesis in *Timema* stick insects. *Evolution* **63**:84-103.

Schwander, T., Henry, L. & Crespi B.J. 2011. Molecular evidence for ancient asexuality in *Timema*
stick insects. *Current Biology* **21**: 1129-1134.

Schwander, T., Oldroyd, B.P. 2016. Androgenesis: where males hijack eggs to clone themselves. *Phil.*
Trans. R. Soc. B 371:20150534. http://dx.doi.org/10.1098/rstb.2015.0534.

Sequencher® version 4.1.4 DNA sequence analysis software, Gene Codes Corporation, Ann Arbor,
MI USA http://www.genecodes.com

Siripattrawan, S., Park, J.-K. & Ò Foighil, D. 2000. Two lineages of the introduced Asian freshwater
clam *Corbicula* occur in North America. *Journal of Molluscan Studies* **66**: 423–429.

Skuza, L., Labecka, A.M., Domagala, J. 2009. Cytogenetic and Morphological Characterization of
Corbicula fluminalis (O. F. Müller, 1774) (Bivalvia: Veneroida: Corbiculidae): Taxonomic
Status Assessment of a Freshwater Clam. *Folia biologica (Kraków)* **57**: 3-4
doi:10.3409/fb57_3-4.177-185

Spöri, Y., Flot, J.F. 2020. HaplowebMaker and CoMa: Two web tools to delimit species using
haplowebs and conspecificity matrices. *Methods Ecol. Evol.* **00**:1-5.

Spöri, Y., Stoch, F., Dellicour, S., Birky, C.W.Jr., Flot, J.F. 2021. KoT: an automatic implementation
of the K/theta method for species delimitation. Bioarchiv. Doi:
https://doi.org/10.1101/2021.08.17.454531.

Stephens, M., Smith, N. & Donnelly, P. 2001. A new statistical method for haplotype reconstruction
from population data. *American* Journal of *Human Genetics* **68**: 978-989.

Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M. & Kumar, S. 2011. MEGA5: Molecular
Evolutionary Genetics Analysis using Maximum Likelihood, Evolutionary Distance, and
Maximum Parsimony Methods. *Molecular Biology and Evolution* **28**: 2731-2739.

Tiemann, J.S., Haponski, A.E., Douglass, D.A., Lee, T., Cummings, K.S., Davis, M.A., Foighil, D.Ó.
2017. First record of a putative novel invasive Corbicula lineage discovered in the Illinois
River, Illinois, USA. *BioInvasions Records*. **6**(2):159-166.

van der Kooi, C.J. & Schwander, T. 2014. On the fate of sexual traits under sexuality. *Biol. Rev.*
89:805-819.

Voroshilova, I.S., Pryanichnikova, E.G., Prokin, A.A., Sabitova, R.Z., Karabanow, D.P., Pavlov, D.D.,
Kurina, E.M. 2021. Morphological and Genetic Traits of the First Invasive Population of the
Asiatic Clam *Corbicula fluminea* (O.F. Müller, 1774) Naturalized in the Volga River Basin.
Russian Journal of Biological Invasions. **12**(1): 36-43.

Yamada, M., Ishibashi, R., Toyoda, K., Kawamura, K. & Komaru, A. 2014. Phylogeography of the
brackish water clam *Corbicula japonica* around the Japanese archipelago inferred from
mitochondrial COII gene sequences. *Zoological Science* **31**: 168-179.

Zhao, S., Guo, Y., Sheng, Q., & Shyr, Y. 2014 Advanced Heat Map and Clustering Analysis Using
Heatmap3. *BioMed Research International*. Article ID 986048, 6p.
https://doi.org/10.1155/2014/986048.

Data Accessibility

- Sample information: Table S1

- DNA sequences: Genbank accessions (Table S2)

Author contribution

E.E., M.V. and K.V.D. designed the questions and research. E.E., E.N., J.V., M.C., M.V. and T.B. performed the research. E.E., J.M., N.D., M.V. and K.V.D. analyzed the data. J.F.F. contributed to new analytical tool development and edited drafts. L.-M.P. and S.M.H. provided extra sequences and samples and edited drafts. M.V. made the figures and tables. E.E., J.M., L.-M.P., M.V. and K.V.D. wrote the paper. K.V.D. supervised the work and provided funding.

 Figure captions

 Figure 1: Possible outcomes of nuclear DNA through meiosis in *Corbicula* genus. Although this is

 variable in metazoans (Sagata 1996), fertilization occurs during meiosis I in Corbicula (Obata *et al*.

 2006, Komaru *et al.* 2006). In all cases, the mitochondrial DNA is inherited maternally. **A.** In sexual

 species, a two-step meiosis leads to the production of a reduced haploid egg. Its fertilization by

 uniflagellate reduced sperm gives diploid progeny from both maternal (light blue) and paternal (dark

 blue) nuclear DNA (Obata *et al.* 2006). **B.** In androgenetic lineages, all the maternal nuclear DNA is

 usually extruded within polar bodies during meiosis I (meiotic axis parallel to the adjacent cortex, see

 Komaru *et al.* 1998), and the sperm is biflagellate and unreduced. The progeny is thus diploid from

 paternal nuclear DNA only. If this occurs between two distinct lineages (here represented in blue and

 red), this leads to a *mitochondrial mismatch*. **C.** In androgenetic lineages, the maternal DNA exclusion

 can also be incomplete, producing a triploid progeny with both maternal and paternal nuclear DNA

 (Komaru *et al.* 2006). If occurring between distinct lineages (blue and red), this leads to *hybridization*.

 For clarity, only two chromosome sets are represented (different size and edge thickness) and the

 gamete sizes are not to scale.

 Figure 2: Haplotype webs (haplowebs, see Fig. S1) based on nuclear sequence alignments and

 haplotype network (haplonet) based on mitochondrial sequence data, all including only the 141

 individuals for which all four markers are available. The circle size is proportional to the frequency of

 the represented alleles/haplotypes. The number of mutation steps inferred by the median-joining

 algorithm is displayed on the lines connecting the haplotypes. For each marker, [top] alleles on the

 graph are colored by their nuclear field for recombination (FFR) delimitation (Figure 3) and [bottom]

 alleles on the identical graph are colored by delimitation based on KoT analysis of the mitochondrial

 data (Table S5). For haplowebs, the curves connecting each pair of alleles co-occurring in

 heterozygous individuals are drawn (width proportional to the number of individuals in which the two

 alleles co-occur), and non-exclusive invasive alleles are numbered from 1 to maximum 3. The name

 and origin of each *Corbicula* lineage or allele follows the nomenclature of Table S3. **a)** 28S haploweb,

 b) alpha-amylase haploweb, **c)** ATPS haploweb, **d)** COI haplonet.

Figure 3: Conspecificity matrix based on alleles shared between individuals across the three nuclear markers (28S, alpha-amylase, ATPS). In this matrix, the axes correspond to the individuals (n = 141) and the red signal corresponds to the number of times individuals were found within the same allelic pool (score from 0 to 3 for all tested markers). Based on the dendrogram (Ward clustering), four FFRs were defined. All individuals are represented using the same color scheme as used for haplowebs and the haplonet (see Figure 2). Signals of allele sharing between these FFRs, corresponding to introgression events, are framed with dotted lines and labelled with letters from *a* to *d*. Zone *a* corresponds to alleles shared between FFR1 and 3, while zones *b*, *c*, and *d* all correspond to sharing between FFR3 and 4. Individual names and geographic origins are detailed in Table S3. Dendrogram and sharing zones are only represented once for clarity.

Figure 4: Circos plot representing allele sharing among Corbicula (n = 359). Groups (namely idiotypes) are represented by circle arcs with a size determined by the total number of alleles found in each group. Their colors refer to the FFR delimitation from the haplowebs (Fig. 2 and S2) and conspecificity matrix (Fig. 3) analyses. Linking lines between these groups represent shared alleles, with the thickness proportional to the number of occurrences of this allele in each population. The line color refers to alleles found in invasive lineages (Table S2), with exclusive alleles from *C.* sp. form A/R, B, C/S, and Rlc in red, blue, orange, and green, respectively, non-exclusive alleles in purple, and alleles restricted to the native range in grey. When it is known, the occurrence of sexual and androgenetic lineages is represented by mono- and biflagellate spermatozoa, respectively, for each group. For Hawaii, China, Madagascar, and South Africa idiotypes, the reproductive mode is unknown.

Table 1: Genetic diversity indices: haplotype diversity (H_d) and nucleotide diversity (π) for all *Corbicula* samples based on COI, 28S, alpha-amylase, and ATPS.

Supporting information

Supplemental information 1: PCR conditions for all markers used in this study.

Supplemental information 2: details on phasing resolution method on triploid individuals.

Figure S1: Schematic description of the haploweb method.

Figure S2: Haplotype webs (haplowebs, see Fig. S1) based on nuclear sequence alignments and
haplotype network (haplonet) based on mitochondrial sequence data. The circle size is proportional to
the frequency of the represented alleles/haplotypes. The number of mutation steps inferred by the
median-joining algorithm is displayed on the lines connecting the haplotypes. For each marker, [top]
alleles on the graph are colored by their nuclear field for recombination (FFR) delimitation (Figure 3)
and [bottom] alleles on the identical graph are colored by delimitation based on KoT analysis of the
mitochondrial data (Table S5). For haplowebs, the curves connecting each pair of alleles co-occurring
in heterozygous individuals are drawn (width proportional to the number of individuals in which the
two alleles co-occur), and non-exclusive invasive alleles are numbered from 1 to maximum 3. The
name and origin of each *Corbicula* lineage or allele follows the nomenclature of Table S3. **a)** 28S
haploweb (n = 266), **b)** alpha-amylase haploweb (n = 237), **c)** ATPS haploweb (n = 248), **d)** COI
haplonet (n = 251).

Table S1: GenBank accession numbers, haplotype designation, taxon, reproductive mode, and
localities of the *Corbicula* spp. sequences used in the present study.

Table S2: List of the 359 individuals collected for the present study and genetic results obtained. The
color code corresponds to the four invasive lineages: exclusive alleles retrieved from *C.* sp. form A/R
are in red, these from *C.* sp. form B are in blue, these from *C.* sp. form C/S are in orange and these
from *C.* sp. form Rlc are in green. Non-exclusive invasive alleles are in purple. Shared allele from one
idiotype to another are in bold.

Table S3: List of the 141 individuals for which all markers are available and genetic results obtained,
using the same representation as Table S2.

Table S4: KoT results on COI marker for the complete dataset (n=359). A group number is assigned
to each individual as in KoT output (n=251). This clusterisation was inferred for individual with no
COI available but belonging to the same populations. In case of ambiguity, the COI was not inferred,

these individuals are labelled in black. The 12 corresponding colors were applied on haplowebs

(Figure S2).

Table S5: KoT results on COI marker for the filtered dataset (n=141). A group number is assigned to

each individual as in KoT output. The 8 corresponding colors were applied on haplowebs (Figure 2).

A. Sexual reproduction   B. Classical androgenesis   C. Androgenesis with nuclear capture

Meiosis I

Meiosis II

Progeny

□ Female from lineage 1   ■ Male from lineage 1   ■ Male from lineage 2   ⚭ Duplicated chromosome   ‖ Unduplicated chromosome   🦠 Mitochondrion

FFR4: *C.* sp. form A/R, form B, form Rlc (America, Europe), *C.* sp. (Japan)

FFR3: *C. fluminalis africana* (South Africa)

FFR3: *C.* sp. form C/S (America, Europe)

IA1

IA2

IA3

FFR2: *C. japonica* (Japan)

FFR1: *C.* sp. (Vietnam)

C. sp. CFY, Cl (Japan)

C. sp. form A/R (America, Europe)
C. sp. CFG, CFY, KMT (Japan)

C. fluminalis africana (South Africa)

C. sp. form C/S (America, Europe)

C. sp. form Rlc (America, Europe)

C. sp. form B (America, Europe)
C. sp. CFG, EHM (Japan)

C. japonica (Japan)

C. sp. (Vietnam)

Figure 2a: 28S haploweb (n = 141).

Figure 2b: alpha-amylase haploweb (n=141).

Figure 2c: ATPS haploweb (n=141).

Figure 2d: COI haplonet (n=141).

Figure 3: Conspecificity matrix.

Figure 4: Circos plot.

Table 1

Genetic diversity indices: haplotype diversity (H_d) and nucleotide diversity (π) for all *Corbicula* samples based on COI, 28S, alpha-amylase, and ATPS.

	COI	28S	*amy*	*atps*
Haplotype diversity (H_d)	0.906 ± 0.011	0.947 ± 0.003	0.933 ± 0.005	0.928 ± 0.003
Nucleotide diversity (π)	0.0296 ± 0.0021	0.0145 ± 0.0003	0.0405 ± 0.0032	0.0774 ± 0.0060

Figure S1. Schematic description of the haploweb method (Flot *et al.* 2010).

Let us imagine a hypothetical sampling of 6 diploid individuals, among which five are heterozygous and one is homozygous for a marker of interest, with each allele/haplotype sequence represented by a specific color below.

A first step is to draw a median-joining network of the alleles (*i.e.,* a distance-based network representing sequence similarities), in which each haplotype is represented by a circle of diameter proportional to the number of individuals harboring this haplotype in the dataset.

As a second step, connections are added between each pair of alleles found co-occurring in heterozygous individuals.

Based on allele sharing, the last step is to delimit fields for recombination (FFRs *sensu* Doyle 1995), *i.e.* groups of individuals belonging to the same allelic pool.

Three pools of alleles   Three fields for recombination

FFR3: *C. fluminalis africana* (South Africa)

FFR3: *C. sp.* form C/S, *C. sp.* (America, Europe)

1A1

1A2

1A3

FFR4: *C. sp.* form A/R, form B, form Rlc (America, Europe), *C. sandai* (Japan), *C. spp.* (China, Japan, Hawaii, Madagascar)

FFR2: *C. japonica* (Japan)

FFR1: *C. sp.* (Vietnam)

C. sp. (Madagascar)

C. sp. form B, form Rlc (America, Europe) *C. sp.* CFG, Cl, EHM, LB (Japan)

C. fluminalis africana (South Africa)

C. sandai, C. sp. LB (Japan)

C. sp. form A/R (America, Europe) *C. sp.* CFG, CFY, KMT (Japan)

C. sp. (China)

C. sp. form C/S (America, Europe)

C. japonica (Japan)

C. sp. (Vietnam)

Individuals in black were not available for COI analysis This includes populations from Hawaii and Waal

Figure S2a: 28S haploweb (n = 266).

FFR1: *C.* sp. (Vietnam), *C. tobae* (Indonesia)

FFR4: *C.* sp. form A/R, form B, form Rlc (America, Europe), *C. sandai* (Japan), *C.* sp. (Japan, Taiwan)

IA2

IA1

IA3

FFR3: *C. fluminalis africana* (South Africa) C. sp. form C/S, C. sp. (America, Europe)

FFR2: *C. japonica* (Japan)

C. sp. (Vietnam)

C. sandai, *C.* sp. LB (Japan)

C. sp. form A/R (America, Europe) *C.* sp. CFG, CFY, KMT (Japan) *C.* sp. (Taiwan)

C. sp. form B, form Rlc (America, Europe) *C.* sp. CFG, Cl, EHM, LB (Japan)

C. fluminalis africana (South Africa)

C. sp. form C/S (America, Europe)

C. japonica (Japan)

Individuals in black were not available for COI analysis **This includes populations from Indonesia and Waal**

Figure S2b: alpha-amylase haploweb (n=237).

FFR4: *C.* sp. form A/R, form B, form Rlc (America, Europe),
C. sp. (China, Hawaii, Japan, Korea, Philippines, Thailand)

FFR2: *C. japonica* (Japan)

FFR3: *C. fluminalis africana* (South Africa)
C. sp. form C/S, *C.* sp. (America, Europe)

IA1

FFR1: *C.* sp. (Vietnam)

C. sp. form A/R (America, Europe)
C. sp. CFG, CFY, KMT, LB (Japan)
C. sp. (Taiwan)

C. sp. LB (Japan)

C. sp. form B, form Rlc (America, Europe)
C. sp. CFG, CFY, Cl (Japan)

C. sp. (China)

C. sp. form C/S (America, Europe)

C. fluminalis africana
(South Africa)

C. japonica (Japan)

C. sp. (Vietnam)

Individuals in black were not available for COI analysis
This includes populations from Hawaii, Korea,
Thailand, Philippines and Waal

Figure S2c: ATPS haploweb (n=248).

FFR4: *C.* sp. form A/R, form B, form Rlc (America, Europe), *C. sandai* (Japan), *C.* sp. (China, Japan, Madagascar, Taiwan)

FFR2: *C. japonica* (Japan)

fricana (South Africa)
America, Europe)

FFR1: *C.* sp. (Vietnam)

Individuals in black were not available for nuclear marker analysis
This includes populations from Indonesia

C. sandai, C. sp. LB (Japan)

C. matannensis
(Indonesia)

C. loehensis
(Indonesia)

C. sp. (Madagascar)

C. sp. form B, form Rlc (America, Europe)
C. sp. CFG, CFY, Cl, EHM, LB (Japan)

FW1

FW4

C. fluminalis africana
(South Africa)

FW17

C. sp. form C/S
(America, Europe)

C. sp. (Vietnam)

FW5

C. sp. form A/R (America, Europe)
C. sp. CFG, CFY, KMT, LB (Japan)

C. sp. (China)

C. japonica (Japan)

Figure S2d: COI haplonet (n = 251).