Title: Intergenerational effects of early adversity on survival in wild baboons

Matthew N. Zipple[1], Elizabeth A. Archie[2,3], Jenny Tung[1,3,4,5], Jeanne Altmann[3,6], Susan C.

Alberts[1,3,4]

[1] Department of Biology, Duke University

[2] Department of Biological Sciences, University of Notre Dame

[3] Institute of Primate Research, National Museums of Kenya

[4] Department of Evolutionary Anthropology, Duke University

[5] Duke Population Research Institute, Duke University

[6] Department of Ecology and Evolutionary Biology, Princeton University

Corresponding Author: Susan Alberts, alberts@duke.edu

Abstract: In humans and nonhuman animals, early life adversity can affect an individual's health, survival, and fertility for many years after the adverse experience. However, whether early life adversity also imposes intergenerational effects on the exposed individual's offspring is not well understood. Here, we fill this gap by leveraging prospective, longitudinal data on a wild, long-lived primate. We find that juveniles whose mothers experienced early life adversity exhibit high mortality before age 4, and this effect is independent of the juvenile's own experience of early adversity. Furthermore, our results point towards a strong role for classic parental effects in driving these effects: mothers that experienced early life adversity displayed reduced viability in adulthood, which in turn led to reductions in offspring survival. Importantly, these mothers' juvenile offspring often preceded them in death by 1 to 2 years, indicating that, for high adversity mothers, the quality of maternal care declines near the end of life. While we cannot exclude direct effects of a parent's environment on offspring quality (e.g., transgenerational epigenetic changes), our results are most consistent with a classic parental effect, in which the environment experienced by a parent affects its future phenotype and therefore its offspring's phenotype. Together, our findings demonstrate that adversity experienced by individuals in one generation can have strong effects on the survival of offspring in the next generation, even if those offspring did not themselves experience early adversity.

An individual's health, survival, and fertility can be profoundly shaped by its early life environment (1). For example, in humans, low early life socioeconomic status predicts increased risk of coronary heart disease (2–4), stroke (2, 5, 6), type II diabetes (7), poor perceived health (8), and all-cause mortality (9, 10) in adulthood. Similarly, numerous studies of wild mammals (11–14) and birds (15–17) find that adult fecundity is reduced in animals that experienced

adverse early life environments, and a few have also found an effect of early life adversity on

adult survival (13–15, 18).

If the effects of early adversity extend to the descendants of exposed individuals, the

epidemiological and evolutionary impact of these effects would be further amplified. However,

evidence from humans for intergenerational effects that result directly from the early life

experience of the parent is mixed, as studies have produced somewhat contradictory results (19–

22). For example, a study of the Överkalix population in Sweden identified strong, contrasting

effects of grandparents' exposure to early-life food scarcity on grand-offspring survival,

depending on small differences in the age at which the grandparent was exposed to scarcity (22).

Similarly, two studies of the same population exposed *in utero* to the Dutch hunger winter (a

well-studied famine that resulted from a German blockade of the Netherlands during the winter

of 1944-1945) found contradictory, sex-specific intergenerational effects, in one case suggesting

an intergenerational effect that depended only upon the mother's early experience (20), and in

the other case an effect that depended only upon the father's early experience (19). Furthermore,

any possible effects of parental or grandparental adversity on future generations are assumed to

be transgenerational, operating as a result of inherited epigenetic changes (19–22). Yet no

genetic validation of this assumption has been carried out, and it remains possible that a simpler

intergenerational pathway explains such results. Specifically, early adversity experienced by a

parent may act as a classic parental effect by changing the parent's phenotype, which in turn

influences the offspring's phenotype (23–26).

The best evidence for intergenerational effects of early adversity comes from several

laboratory studies of short-lived animals, which find strong relationships between a female's

early life environment and the body size of her offspring [(27–36), reviewed in (37), but see (38)

58 for a rare example in wild house wrens]. These findings provide important proof-of-principle that

59 intergenerational effects of early adversity can occur. However, these studies do not address

60 whether intergenerational effects of early adversity occur in natural populations of long-lived

61 animals. And while a few studies of captive animals have demonstrated a relationship between a

62 female's early environment and her offspring's survival or reproduction (39–41), the ecological

63 validity of these findings has yet to be verified by studying intergenerational fitness effects in a

64 population of wild and/or long-lived animals. By working in a natural population, we are able to

65 guarantee that animals are exposed only to natural levels of early adversity, and are also subject

66 to any social factors which might mitigate or aggravate the influence of those early adverse

67 events.

68   Addressing whether the effects of early adversity in one generation affect reproduction or

69 survival in the next is challenging because of the difficulties of linking high-quality data on early

70 adversity in one generation to health and survival outcomes in the next. Here, we overcome these

71 challenges by taking advantage of a prospective longitudinal dataset from a natural primate

72 population: the baboons of the Amboseli ecosystem in southern Kenya (42). This dataset

73 includes 45 years of individual-based data on early adversity, and real-time observations of later-

74 life survival outcomes for hundreds of subjects with known maternities and grand maternities.

75 Moreover, unlike many human populations, we do not observe inter-generational transmission of

76 adverse conditions; that is, offspring of females who experienced early life adversity are not

77 more likely to experience early life adversity themselves—allowing us to avoid this common

78 confound in human societies.

79   To test for intergenerational effects of early adversity, we focused on early adversity

80 experienced by female baboons who later became mothers, and whose offspring were also in our

data set. We asked whether the early adversity experienced by these females ("maternal early

adversity") predicted the survival of their juvenile offspring in the next generation, after

controlling for the early adversity directly experienced by the offspring themselves.

We considered five types of early adverse conditions (Table 1), based on previous work

in our study population that demonstrated effects of these conditions on a female baboon's own

adult survival (18). These included: (i) maternal death during development (0-4 years of age),

which indicates the loss of an important source of social support, physical protection, and

nutrition (43, 44), (ii) being born to a low-ranking mother, which influences growth rates and age

at maturation (45–47) (iii) being born into a large social group (and thus experiencing high

density conditions and high levels of within-group competition) (11, 45, 48) (iv) being born

during a drought, which reduces fertility in adulthood (11, 49), and (v) experiencing the birth of

a close-in-age younger sibling, which may reduce maternal investment received during

development (50). Importantly—and in contrast to human studies (51)—sources of early

adversity are not strongly correlated in our population (Table S1).

Results

We built a mixed effects Cox proportional hazards model of offspring survival during the

juvenile period that included early adversity measures present in the mother's and the offspring's

early life as binary fixed effects. We defined the juvenile period based on survival until age 4,

near the age of menarche for females and earliest dispersal for males in this population (52). We

included data on maternal early adversity for all five adverse early life conditions, and we

included data on offspring early adversity for four of the five conditions. We excluded the birth

of a close-in-age younger sibling for the offspring generation because the survival of the focal

offspring strongly affects the length of the subsequent birth interval (i.e., offspring that die

shortly after birth also have the closest-in-age younger siblings). We included maternal and

grandmaternal ID as random effects. In total, we used data collected from 1973-2017 to analyze

the survival of 687 offspring (46.5% males) born to 169 females (mean 4.1 offspring per female,

range 1-12) for whom we had data on all five adverse conditions in the mother's early life, and

all four adverse conditions in the offspring's early life.

Each adverse condition was scored as present or absent for each subject, and each one

affected a minority of our study subjects (range 6%-34%). Mothers and offspring had similar

chances of experiencing adverse conditions, except for social density: offspring were more likely

than mothers to be born into large social groups because of population growth over the 5-decade

study period (Table 1). Unlike typical patterns of early adversity in human populations (51),

sources of early life adversity in our population were not strongly correlated: with the exception

of maternal rank in the mother's and offspring's generations (p<0.0001, r=0.40), no adverse

condition explained more than 4% of the variance in any other condition, either within or

between generations (Table S1).

Maternal Early Life Adversity and Offspring Survival: Our full multivariate Cox

proportional hazards model for offspring survival (Table S2) included all 9 early adverse

conditions (five for mothers and four for offspring). We found strong negative effects of two

characteristics of the *mother's* early life environment on their offspring's survival during the first

4 years of life: maternal loss (hazard ratio = 1.48, p=0.006) and presence of a close-in-age

younger sibling (HR = 1.39, p=0.03). Following backwards model selection, these two

characteristics remained the only significant maternal early life predictors of offspring survival

(Table 2, Figure 1, along with two conditions in the offspring's early life environment: see

below). Specifically, offspring whose mothers experienced early maternal loss experienced a

48% higher probability of dying throughout the first four years of life than unaffected offspring,

and offspring whose mothers had a close-in-age sibling experienced a 39% higher probability of

dying than unaffected offspring. This effect is striking especially considering that a median of

7.0 and 8.0 years separated the offspring's own birth from the mother's experience of maternal

loss or birth of a close-in-age sibling, respectively. Notably, previous work in our population

found that these two sources of adversity—maternal loss and the presence of a close-in-age

younger sibling during early life—are also sources of mortality risk once females reach

adulthood, and in fact are the two strongest predictors of adult survival among six different early-

life conditions considered (18). Hence, early-life conditions that are especially adverse for

females when they reach adulthood also negatively affect the survival of their offspring.

Both the full and reduced models of offspring survival also included two conditions in the

offspring's early life environment as significant predictors of juvenile survival. Specifically,

maternal loss experienced by the offspring and low maternal rank during the offspring's juvenile

period had strong negative effects on offspring survival (Table 2, maternal death: HR = 1.95,

$p=5 \times 10^{-7}$, low maternal rank: HR=1.43, p=0.025). Thus, maternal loss in the offspring's

generation had a stronger effect on offspring survival (nearly doubling offspring mortality risk)

than maternal loss in the mother's generation. In contrast, the effect of having a low-ranking

mother, which was associated with a 43% increase in offspring mortality risk, was comparable in

its effect size to the two significant predictors from the maternal generation (maternal loss and

close-in-age sibling for the mother, 48% and 39% increase in offspring mortality, respectively).

Thus, two adverse conditions in a mother's early life had as large or larger of an impact on her

offspring's survival than all but one adverse condition experienced by the offspring directly.

Maternal Viability and Offspring Survival: The strong effect of the mother's death on the

survival of offspring to age four years (Table 2) could arise if offspring die *after* their mothers

die: even after weaning (approximately 1.5 years of age), juvenile baboons rely on their mothers

for social support and social learning (53). Alternatively, these offspring may die *before* their

mothers die if those mothers are themselves in poor condition. To distinguish these alternatives,

we modeled offspring survival to age 2 years (halfway through the juvenile period) as a function

of maternal death during years 2-4 after an offspring's birth (i.e., the two years that *followed* the

offspring survival period modeled in the response variable). In this analysis we considered only

the subset of offspring in our dataset whose mothers survived the entirety of the first two years of

the offspring's life, and for whom we were able to evaluate the four significant predictors of

offspring survival identified above and in Table 2 (N=671). Our results showed that offspring

were less likely to survive during the first two years of life if they were born to mothers who died

2-4 years after their birth. In other words, these offspring were more likely to die even when their

mother was still alive (hazard ratio=1.50 [1.01-2.23], p=0.045).

To test whether this link between offspring survival and maternal viability was driven by

maternal early adversity, we next partitioned our analysis of offspring survival to age 2 based on

whether the mother experienced either maternal loss or a close-in-age younger sibling (i.e., either

or both of the two maternal early life conditions that significantly predicted their offspring's

survival; Table 2). We found that, among offspring whose mothers experienced either or both of

these two conditions (N=247), maternal death in years 2-4 after the offspring's birth significantly

predicted reduced offspring survival to age 2 years (Figure 2a, hazards ratio=1.78, 95% CI =

[1.05-3.01], p=0.03). Maternal death in the same period did not, however, predict reduced

offspring survival when mothers had not experienced maternal loss or a close-in-age younger

sibling (N=424; Figure 2b, hazard ratio=1.21, 95% CI = [0.7-2.2], p=0.53). This finding is

consistent with the hypothesis that maternal early life adversity results in low maternal viability

in adulthood, which in turn results in both earlier death for adult females and a reduction in their

ability to successfully raise offspring towards the end of their lives.

Maternal Early Life Adversity and Quantity of Maternal Care: We hypothesized that

developmental constraints imposed on females by early life adversity could lead to reduced

survival in their offspring as a result of two non-mutually exclusive mechanisms. First, mothers

that experienced early life adversity might provide lower levels of maternal care to their

offspring than other mothers, through differences in either maternal behavior or physiology (e.g.

reduced nutrient content in milk). Second, females who experienced early adversity might also

exhibit reduced egg and/or amniotic environmental quality (an established mechanism for

transmission of maternal effects (37, 54)). We were able to partially test the first hypothesis by

drawing on longitudinal behavioral data for this population. Specifically, we built linear mixed

effects models to test whether maternal early adversity affected the proportion of time during 10-

minute focal follows that a mother spent either carrying or suckling her dependent infants. Fixed

effects in the model included maternal viability (a binary variable indicating whether the mother

survived for four years after offspring birth), the two early adverse circumstances experienced by

the mother that affected offspring survival (the mother's maternal loss and close-in-age sibling),

offspring age (both as a linear and as a quadratic term), maternal rank, maternal age, the number

of adult females in the social group, and season. We also included maternal ID, offspring ID,

group ID, and observer ID as random effects.

Neither maternal viability nor either of the two maternal early adversity effects was a

significant predictor of the proportion of time that mothers spent carrying or suckling their

infants in either full multivariate models or models resulting from backwards selection (p>0.1 in

all cases, Tables S3-S6). However, the proportion of time that mothers spend carrying and

suckling offspring are relatively coarse metrics of maternal care, and it could be that more fine-

grained measures of maternal care (which we do not regularly collect) could reveal a relationship

between maternal early adversity and future maternal care.

Discussion

We have demonstrated that adverse environmental conditions during the early life of a

female baboon, which are already known to negatively affect both her survival (18) and her

reproduction (11) in adulthood, also reduce the survival of her offspring. Importantly, this effect

is independent of the environment experienced by those offspring themselves (Figure 1). The

reduction in offspring survival is likely linked to reductions in maternal viability: mothers that

experienced early life adversity are significantly less able to successfully raise offspring born

near the ends of their lives, while the same is not true for mothers that did not experience early

life adversity (Figure 2). Together, these findings support the hypothesis that early life adversity

produces constraints during development that lead not only to reduced adult survival and lifetime

reproductive success (18) but also to a reduced ability to successfully raise those offspring that

are produced.

The results reported here help to fill a key gap in the literature concerning the

intergenerational effects of early life adversity on survival. Results from human studies have

yielded inconsistent results on this topic thus far: different studies on the same populations have

reported contradictory sex-specific effects on health (19, 20) or have found that small differences

in the age at which probands' parents or grandparents were exposed to adversity can lead to a

reversal in the direction of these effects (21, 22). Among studies in non-human animals, several

studies in fish (55, 56), reptiles (57), birds (58, 59), and ungulates (60–64) have found that

parental body condition at the time of offspring birth influences offspring survival, but none have

linked parents' early adverse experiences to offspring survival. Additionally, while previous

studies have identified effects of parental early adversity on offspring traits in a limited number

of systems (37, 39, 40), ours is the first to link parental early adversity to offspring fitness

outcomes in a wild, long-lived animal.

Our findings help to explain the persistence of health deficits across generations (65–67),

by revealing that in long-lived primates, the early life experiences of mothers have important

implications for offspring health and survival. Recent studies in humans have demonstrated that

conditions experienced by mothers during pregnancy (e.g., low SES, psychosocial stress, mood

dysregulation, prenatal smoking) can affect HPA axis regulation (68, 69) and birthweight (65,

66) in her offspring. These and other maternal characteristics present during pregnancy are

influenced not only by mothers' experiences in adulthood, but also by the long-term effects of

environmental conditions experienced in mothers' early lives (54, 66). Our findings therefore

motivate future work to test for comparable intergenerational fitness effects of early adversity in

humans and other non-human animals.

Our findings are consistent with the hypothesis that early adversity results in

intergenerational effects of developmental constraints (11, 70–72) and are not consistent with an

intergenerational predictive adaptive response hypothesis (71, 73, 74). Rather than being

buffered against the effects of maternal loss, those offspring that experienced maternal loss and

whose mothers had also experienced maternal loss were more likely, not less likely, to die, as

compared to offspring that experienced maternal loss but whose mothers did not. Thus, offspring

experience constraints not only as a result of their own early environment, but also as a result of

their mother's developmental history, including events that occurred years before the offspring's

own conception. Our results are consistent with the hypothesis that a female's condition at the

time of her offspring's conception and/or birth reflects her previous experiences, and that her

condition thereby influences the development and survival of her offspring (54, 75, 76).

246   Finally, our study provides insight into how the effects of early adversity may be

transmitted from parent to offspring. Intergenerational transmission of adversity has often been

viewed as a potential transgenerational effect – i.e., environmental exposures in one generation

that directly alter the biology of animals born at least one generation later, perhaps via the

inheritance of environmentally-induced epigenetic changes (77). While we cannot exclude this

possibility, our results suggest that a simpler mechanism may be operating. The intergenerational

effects of early adversity on offspring survival that we present here are consistent with a classic

parental effect (19-22) in which early life adversity affects the phenotypic quality of the mother

during adulthood, and her resulting deficits directly affect her offspring's development. This

mode of transmission – in which intergenerational transmission of parental early adversity

operates via differences in parental phenotype, rather than via transgenerational effects that

determine offspring phenotypes – may be more widespread than is typically thought (78). This

insight may have important implications for understanding the persistence of human health

deficits across generations and inform the approaches taken to intervene in this transmission

(79).

Methods

Study system: The Amboseli Baboon Research Project is a long-term longitudinal study of

wild baboons living in and around Amboseli National Park, Kenya. A detailed description of the

study system can be found elsewhere (42). Researchers have continuously collected behavioral,

environmental, and demographic data from the population since 1971. All subjects are visually

recognized, and near-daily censuses allow us to precisely document the timing of demographic

events, including the birth and death of study individuals. Critical to this study, we have

continuously collected near-daily measures of group size, daily rainfall levels, and monthly

calculations of social dominance rank (80), and we can accurately assign the dates of birth of all

mothers and offspring born into study groups as well as the dates of all juvenile and adult female

deaths.

Study Subjects: In our analyses of offspring survival, we included individuals who met two

criteria: (i) we were able to evaluate each of the five sources of maternal early life adversity and

four sources of offspring early life adversity outlined below; and (ii) they lived in social groups

that fed exclusively on wild foods rather than having their diet supplemented with human-

sourced refuse. Although transmission of paternal early adversity may also occur in our

population, we did not consider it here because we knew paternal identities for only a subset of

our study subjects and had early life data on only a limited number of fathers. Our analysis

ultimately relied on data spanning more than four decades, from 1973 to 2017.

Measuring Early Life Adversity: Previous work in the Amboseli population defined six

binary indicators of early life adversity and considered a single index of cumulative adversity

based on the sum of these indicators (18). This cumulative adversity index is a strong predictor

of adult lifespan: females that experienced high levels of early life adversity (i.e., a greater

number of adverse early life conditions) but still survived to adulthood lived dramatically shorter

lives compared to females that did not experience adversity (18). In addition to the five sources

of early adversity discussed above, this previous analysis also considered early social

connectedness (social integration versus social isolation) as a sixth source of adversity (18).

Social connectedness data are missing for some mothers who were born relatively early in the

long-term study. To maximize our sample size, we therefore did not include measures of social

connectedness in this analysis. Our operational definitions for each source of adversity mirrored

those used by Tung *et al* (18) for the remaining five conditions, except that here we employed

measures of proportional rather than ordinal dominance rank (i.e., rank measured as a proportion

of females that the focal individual dominates, rather than her ordinal rank number). We also

built an index of cumulative maternal adversity, but because that model did not fit the data better

than our reduced multivariate model (in contrast to the results for adult female survival (18)) we

report the multivariate model in the main text. The alternative model based on cumulative

maternal adversity is presented in Table S7.

Statistical Analysis: We built a mixed effects Cox proportional hazards model of offspring

survival during the first four years of life using the R package coxme (81, 82). The response

variable in our model was the age at which offspring death occurred (if at all) during the first 4

years of life. We considered offspring survival to age 4 as the key survival period of interest

because it roughly corresponds to the end of the juvenile period for baboons (52). Offspring that

survived beyond age 4 were treated as censored individuals who survived until at least age 4. In

our models of offspring survival as a function of maternal viability (Figure 2), we altered the first

model to predict survival during the first two years of life as a function of maternal survival

during years 2-4 after offspring birth. To test for effects of maternal adversity on the quantity of

maternal care we built linear mixed effects models using the R package lme4 (83) (see Table S8

for model syntax).

References:

1.   Uller T, Nakagawa S, English S (2013) Weak evidence for anticipatory parental effects in
plants and animals. *J Evol Biol* 26(10):2161–2170.
2.   Naess Ø, Claussen B, Davey Smith G (2004) Relative impact of childhood and adulthood
socioeconomic conditions on cause specific mortality in men. *J Epidemiol Community*
Health 58(7):597–8.
3.   Beebe-Dimmer J, et al. (2004) Childhood and Adult Socioeconomic Conditions and 31-
Year Mortality Risk in Women. *Am J Epidemiol* 159(5):481–490.
4.   Kittleson MM, et al. (2006) Association of Childhood Socioeconomic Status With
Subsequent Coronary Heart Disease in Physicians. *Arch Intern Med* 166(21):2356.
5.   Davey Smith G, Hart C, Blane D, Hole D (1998) Adverse socioeconomic conditions in
childhood and cause specific adult mortality: prospective observational study. *BMJ*
316(7145):1631–5.
6.   Frankel S, Smith GD, Gunnell D (1999) Childhood Socioeconomic Position and Adult
Cardiovascular Mortality: The Boyd Orr Cohort. *Am J Epidemiol* 150(10):1081–1084.
7.   Lidfeldt J, Li TY, Hu FB, Manson JE, Kawachi I (2007) A Prospective Study of
Childhood and Adult Socioeconomic Status and Incidence of Type 2 Diabetes in Women.
Am J Epidemiol 165(8):882–889.
8.   Van De Mheen H, Stronks K, Looman C, Mackenbach J (1998) Does childhood
sodoeconomic status influence adult health through behavioural factors? *C Int*
Eptdemlological Assoc Gt Britain Int J ojEpidemiology 27:431–137.
9.   Kuh D, Hardy R, Langenberg C, Richards M, Wadsworth MEJ (2002) Mortality in adults
aged 26-54 years related to socioeconomic conditions in childhood and adulthood: post
war birth cohort study. *BMJ* 325(7372):1076–80.
10.   Galobardes B, Lynch JW, Davey Smith G (2004) Childhood Socioeconomic
Circumstances and Cause-specific Mortality in Adulthood: Systematic Review and
Interpretation. *Epidemiol Rev* 26(1):7–21.
11.   Lea AJ, Altmann J, Alberts SC, Tung J (2015) Developmental Constraints in a Wild
Primate. *Source Am Nat* 185178111(6):809–821.
12.   Douhard M, et al. (2014) Fitness consequences of environmental conditions at different
life stages in a long-lived vertebrate. *Proceedings Biol Sci* 281(1785):20140276.
13.   Nussey DH, Kruuk LEB, Morris A, Clutton-Brock TH (2007) Environmental conditions
in early life influence ageing rates in a wild population of red deer. *Curr Biol*
17(23):R1000–R1001.
14.   Pigeon G, Pelletier F (2018) Direct and indirect effects of early-life environment on
lifetime fitness of bighorn ewes. *Proceedings Biol Sci* 285(1870):20171935.
15.   Herfindal I, van de Pol M, Nielsen JT, Saether B-E, Møller AP (2015) Climatic conditions
cause complex patterns of covariation between demographic traits in a long-lived raptor. *J*
Anim Ecol 84(3):702–711.

16. Balbontín J, Møller AP (2015) Environmental conditions during early life accelerate the rate of senescence in a short-lived passerine bird. *Ecology* 96(4):948–959.

17. Millon A, Petty SJ, Little B, Lambin X (2011) Natal conditions alter age-specific reproduction but not survival or senescence in a long-lived bird of prey. *J Anim Ecol* 80(5):968–975.

18. Tung J, Archie EA, Altmann J, Alberts SC (2016) Cumulative early life adversity predicts longevity in wild baboons. *Nat Commun* 7:11181.

19. Veenendaal M, et al. (2013) Transgenerational effects of prenatal exposure to the 1944-45 Dutch famine. *BJOG An Int J Obstet Gynaecol* 120(5):548–554.

20. Painter R, et al. (2008) Transgenerational effects of prenatal exposure to the Dutch famine on neonatal adiposity and health in later life. *BJOG An Int J Obstet Gynaecol* 115(10):1243–1249.

21. Kaati G, Bygren LO, Pembrey M, Sjöström M (2007) Transgenerational response to nutrition, early life circumstances and longevity. *Eur J Hum Genet* 15(7):784–790.

22. Pembrey ME, et al. (2006) Sex-specific, male-line transgenerational responses in humans. *Eur J Hum Genet* 14(2):159–166.

23. Mousseau T, Fox C eds. (1998) *Maternal effects as adaptations* (Oxford University Press).

24. Mousseau TA, Fox CW (1998) The adaptive significance of maternal effects. *Trends Ecol Evol* 13(10):403–407.

25. Russell AF, Lummaa V (2009) Maternal effects in cooperative breeders: from hymenopterans to humans. *Philos Trans R Soc B Biol Sci* 364(1520):1143–1167.

26. Badyaev A V, Uller T (2009) Parental effects in ecology and evolution: mechanisms, processes and implications. *Philos Trans R Soc Lond B Biol Sci* 364(1520):1169–77.

27. William Huck U, Labov JB, Lisk RD (1986) Food Restricting Young Hamsters (Mesocricetus Auratus) Affects Sex Ratio and Growth of Subsequent Offspring1. *Biol Reprod* 35(3):592–598.

28. Alonso-Alvarez C, Bertrand S, Sorci G (2007) Sex-specific transgenerational effects of early developmental conditions in a passerine. *Biol J Linn Soc* 91(3):469–474.

29. Helle H, Koskela E, Mappes T (2012) Life in varying environments: experimental evidence for delayed effects of juvenile environment on adult life history. *J Anim Ecol* 81(3):573–582.

30. Goerlich VC, Nätt D, Elfwing M, Macdonald B, Jensen P (2012) Transgenerational effects of early experience on behavioral, hormonal and gene expression responses to acute stress in the precocial chicken. *Horm Behav* 61:711–718.

31. Taborsky B (2006) Mothers determine offspring size in response to own juvenile growth conditions. *Biol Lett* 2(2):225–8.

32. Beckerman AP, Benton TG, Lapsley CT, Koesters N (2003) Talkin' 'bout my generation: environmental variability and cohort effects. *Am Nat* 162(6):754–67.

33. Saastamoinen M, Hirai N, van Nouhuys S (2013) Direct and trans-generational responses to food deprivation during development in the Glanville fritillary butterfly. *Oecologia* 171(1):93–104.

34. Vijendravarma RK, Narasimha S, Kawecki TJ (2010) Effects of parental larval diet on egg size and offspring traits in Drosophila. *Biol Lett* 6(2):238–41.

35. Fischer K, Eenhoorn E, Bot ANM, Brakefield PM, Zwaan BJ (2003) Cooler butterflies lay larger eggs: developmental plasticity versus acclimation. *Proceedings Biol Sci*

270(1528):2051–6.

36. Jobson MA, et al. (2015) Transgenerational Effects of Early Life Starvation on Growth,
Reproduction, and Stress Resistance in Caenorhabditis elegans. *Genetics* 201(1):201–12.

37. Burton T, Metcalfe NB (2014) Can environmental conditions experienced in early life
influence future generations? *Proceedings Biol Sci* 281(1785):20140311.

38. Bowers EK, Thompson CF, Sakaluk SK (2017) Maternal Natal Environment and
Breeding Territory Predict the Condition and Sex Ratio of Offspring. *Evol Biol* 44(1):11–
20.

39. William Huck U, Labov JB, Lisk RD (1987) Food-Restricting First Generation Juvenile
Female Hamsters (Mesocricetus Auratus) Affects Sex Ratio and Growth of Third
Generation Offspring1. *Biol Reprod* 37(3):612–617.

40. Naguib M, Nemitz A, Gil D (2006) Maternal developmental stress reduces reproductive
success of female offspring in zebra finches. *Proceedings Biol Sci* 273(1596):1901–5.

41. Marcil-ferland D, Festa-bianchet M, Martin AM, Pelletier F (2013) Despite Catch-Up ,
Prolonged Growth Has Detrimental Fitness Consequences in a Long-Lived Vertebrate.
Am Nat 182(6):775–782.

42. Alberts SC, Altmann J (2012) The Amboseli baboon research project: 40 years of
continuity and change. *Long-Term Field Studies of Primates*, pp 261–287.

43. Altmann J (1980) *Baboon Mothers and Infants* (Harvard University Press, Cambridge,
Massachussets).

44. Lea AJ, Learn NH, Theus MJ, Altmann J, Alberts SC (2014) Complex sources of variance
in female dominance rank in a nepotistic society. *Anim Behav* 94:87–99.

45. Charpentier MJE, Tung J, Altmann J, Alberts SC (2008) Age at maturity in wild baboons:
genetic, environmental and demographic influences. *Mol Ecol* 17:2026–2040.

46. Altmann J, Alberts SC (2003) Intraspecific Variability in Fertility and Offspring Survival
in a Nonhuman Primate: Behavioral Control of Ecological and Social Sources. *Offspring:*
The Biodemography of Fertility and Family Behavior, eds Wachter K.W., Bulatao R.A
(National Academy Press, Washington, D.C.). Available at:
https://amboselibaboons.nd.edu/assets/81618/162.pdf [Accessed February 27, 2018].

47. Altmann J, Hausfater G, Altmann SA (1988) Determinants of Reproductive Success in
Savannah Baboons, Papio cynocephalus. *Reproductive Success*, ed Clutton-Brock TH
(University of Chicago Press, Chicago). Available at:
https://amboselibaboons.nd.edu/assets/219836/determinants_of_reproductive_success_in_
savannah_baboons_paio_cynocephalus.pdf [Accessed February 27, 2018].

48. Altmann J, Alberts SC (2003) Variability in reproductive success viewed from a life-
history perspective in baboons. *Am J Hum Biol* 15(3):401–409.

49. Beehner JC, Onderdonk DA, Alberts SC, Altmann J (2006) The ecology of conception
and pregnancy failure in wild baboons. *Behav Ecol* 17:741–750.

50. Altmann J, Altmann SA, Hausfater G, Gould JL (2015) Primate Infant ' s Effects on
Mother ' s Future Reproduction Linked references are available on JSTOR for this article :
Primate Infant ' s Effects on Mother ' s Future Reproduction. 201(4360):1028–1030.

51. Felitti VJ, et al. (1998) Relationship of Childhood Abuse and Household Dysfunction to
Many of the Leading Causes of Death in Adults: The Adverse Childhood Experiences
(ACE) Study. *Am J Prev Med* 14(4):245–258.

52. Charpentier MJE, Tung J, Altmann J, Alberts SC (2008) Age at maturity in wild baboons:
Genetic, environmental and demographic influences. *Mol Ecol* 17(8):2026–2040.

53. van Noordwijk MA (2012) From maternal investment to lifetime maternal care. *The*
Evolution of Primate Societies, eds Mitani JC, Call J, Kappeler PM, Palombit RA, Silk JB
(University of Chicago Press, Chicago, Illinois).
54. Kuzawa CW (2005) Fetal origins of developmental plasticity: Are fetal cues reliable
predictors of future nutritional environments? *Am J Hum Biol* 17(1):5–21.
55. Donelson JM, McCormick MI, Munday PL (2008) Parental condition affects early life-
history of a coral reef fish. *J Exp Mar Bio Ecol* 360(2):109–116.
56. Venturelli PA, et al. (2010) Maternal influences on population dynamics: evidence from
an exploited freshwater fish. *Ecology* 91(7):2003–2012.
57. Warner DA, Lovern MB (2014) The Maternal Environment Affects Offspring Viability
via an Indirect Effect of Yolk Investment on Offspring Size. *Physiol Biochem Zool*
87(2):276–287.
58. Blomqvist D, Johansson OC, Götmark F (1997) Parental quality and egg size affect chick
survival in a precocial bird, the lapwing Vanellus vanellus. *Oecologia* 110(1):18–24.
59. RIDLEY AR (2007) Factors affecting offspring survival and development in a
cooperative bird: social, maternal and environmental effects. *J Anim Ecol* 76(4):750–760.
60. Cameron RD, Smith WT, Fancy SG, Gerhart A N D Robert G White KL (1987) Calving
success of female caribou in relation to body weight. Available at:
http://www.nrcresearchpress.com/doi/pdf/10.1139/z93-069 [Accessed February 26, 2018].
61. Théoret-Gosselin R, Hamel S, Côté SD (2015) The role of maternal behavior and
offspring development in the survival of mountain goat kids. *Oecologia* 178(1):175–186.
62. Keech MA, et al. (2000) Life-History Consequences of Maternal Condition in Alaskan
Moose. *J Wildl Manage* 64(2):450.
63. Clutton-Brock TH, Major M, Albon SD, Guinness FE (1987) Early Development and
Population Dynamics in Red Deer. I. Density-Dependent Effects on. *Source J Anim Ecol J*
Anim Ecol 56(1):53–67.
64. Clutton-Brock TH, Albon SD, Guinness FE (1984) Maternal dominance, breeding success
and birth sex ratios in red deer. *Nature* 308(5957):358–360.
65. Aizer A, Currie J (2014) The intergenerational transmission of inequality: maternal
disadvantage and health at birth. *Science* 344(6186):856–61.
66. Kane JB, Harris KM, Siega-Riz AM (2018) Intergenerational pathways linking maternal
early life adversity to offspring birthweight. *Soc Sci Med* 207:89–96.
67. Cnattingius S, Villamor E, Lagerros YT, Wikström A-K, Granath F (2012) High birth
weight and obesity—a vicious circle across generations. *Int J Obes* 36(10):1320–1324.
68. Thayer ZM, Kuzawa CW (2014) Early origins of health disparities: Material deprivation
predicts maternal evening cortisol in pregnancy and offspring cortisol reactivity in the first
few weeks of life. *Am J Hum Biol* 26(6):723–730.
69. Entringer S, Kumsta R, Hellhammer DH, Wadhwa PD, Wüst S (2009) Prenatal exposure
to maternal psychosocial stress and HPA axis regulation in young adults. *Horm Behav*
55(2):292–298.
70. Grafen A (1988) On the uses of data on lifetime reproductive success. *Reproductive*
Success, ed Clutton-brock TH (University of Chicago Press, Chicago, IL).
71. Monaghan P (2007) Early growth conditions, phenotypic development and environmental
change. *Philos Trans R Soc B Biol Sci* 363(1497):1635–1645.
72. Lea AJ, Tung J, Archie EA, Alberts SC (2017) Developmental plasticity: bridging
research in evolution and human health. *Evol Med Public Heal* 2017(1):162–175.

73.   Gluckman PD, Hanson MA, Spencer HG (2005) Predictive adaptive responses and human
evolution. *Trends Ecol Evol* 20(10):527–533.
74.   Herman JJ, Spencer HG, Donohue K, Sultan SE (2014) How stable "should" epigenetic
modifications be? Insights from adaptive plasticity and bet hedging. *Evolution (N Y)*
68(3):632–643.
75.   Kuzawa CW (2017) Which environments matter in studies of early life developmental
plasticity? *Evol Med Public Heal* 2017(1):188–190.
76.   Lea AJ, Tung J, Archie EA, Alberts SC, Kenya of (2018) Developmental plasticity
research in evolution and human health Response to commentaries. *Evolution (N Y)*:201–
205.
77.   Heard E, Martienssen RA (2014) Transgenerational Epigenetic Inheritance: Myths and
Mechanisms. *Cell* 157(1):95–109.
78.   Drake A, Walker B (2004) The intergenerational effects of fetal programming: non-
genomic mechanisms for the inheritance of low birth weight and cardiovascular risk. *J
Endocrinol* 180. Available at:
https://joe.bioscientifica.com/abstract/journals/joe/180/1/1.xml [Accessed March 11,
2019].
79.   Godfrey K, Gluckman P, Hanson M (2010) Developmental origins of metabolic disease:
life course and intergenerational perspectives. *Trends Endocrinol Metab* 21. Available at:
https://ac.els-cdn.com/S1043276009002197/1-s2.0-S1043276009002197-
main.pdf?_tid=e9b743c6-ff83-42f3-af31-
5952fbc576c9&acdnat=1552315902_02b505e3a3df2b73ea473967441f4523 [Accessed
March 11, 2019].
80.   Hausfater G (1975) Dominance and reproduction in Baboons (Papio cynocephalus).
Contrib Primatol 7:1–150.
81.   Therneau TM (2012) coxme: Mixed Effects Cox Models, R Package Version 2.3-3.
82.   R_Core_Team (2013) R: A language and environment for statistical computing.
83.   Bates D, Maechler M, Bolker B, Walker S (2014) lme4: Linear mixed-effects models
using Eigen and S4.
84.   Silk JB, Alberts SC, Altmann J (2003) Social Bonds of Female Baboons Enhance Infant
Survival. *Source Sci New Ser* 302(5648):1231–1234.
519
520

1 **Figure 1. Offspring survival was influenced by characteristics of their mothers' early-life**
2 **environments.** Offspring survived relatively less well during the juvenile period if (A) their
3 mother lost her own mother during her early life and/or (B) their mother experienced a close-in-
4 age younger sibling.

Figure 2. Effects of maternal adversity on offspring survival are explained by reduced maternal viability. (A) Among those offspring whose mothers experienced significant early life adversity (maternal loss and/or a competing younger sibling), maternal death in years 2-4 was associated with poor offspring survival in years 0-2 after birth, while the mother was still alive. (B) In contrast, among those offspring whose mothers did not experience early life adversity, maternal death in years 2-4 was not associated with reduced offspring survival during the first 2 years of life.

Table 1. Early adverse conditions and the frequencies with which they occur in maternal and offspring generations of our dataset.

Adverse Condition[a]	Criterion	Frequency	
		Maternal Generation	*Offspring Generation*
Drought	During the first year of life, the focal individual experienced less than 200 mm of rainfall (i.e., drought conditions (49)).	0.09	0.15
High Social Density	The individual was born into a group with a high social density (>35 adults), indicating high levels of within-group competition.	0.06	0.32
Maternal Loss	The mother of the focal individual died within four years of the individual's birth.	0.21	0.25
Low Maternal Rank[b]	The focal individual was born to a mother with a low social rank (mother's rank fell in the bottom quartile of the group's dominance hierarchy).	0.17	0.23
Close-In-Age Younger Sibling	The focal individual had a younger sibling born to its mother within 18 months of the focal's birth.	0.20	--

[a]These mirror criteria used in a previous analysis in our population (18), with the exception of rank, which is evaluated here as a proportional measure.

[b]Proportional rank is the proportion of other adult females in a group that an individual's mother outranks. An animal therefore has a low maternal rank if her mother's rank is <0.25. The reduced frequency with which low maternal rank appears in the maternal generation is a likely a result of offspring of low-ranking mothers surviving less well (84), and therefore not surviving to appear as mothers in our dataset.

1   **Table 2.** Reduced model of the effects of maternal and offspring early adversity on offspring
2   survival during early life ($R^2=0.07$).

Generation	Parameter[a]	Coefficient	Hazard Ratio (95% CI)	p value	Interpretation
Maternal	Maternal Loss	0.37	1.44 (1.10-1.90)	**0.009**	Offspring survive less well if their mother experienced maternal loss during her early life.
	Close-in-age Younger Sibling	0.35	1.42 (1.06-1.90)	**0.018**	Offspring survive less well if their mother had a close in age younger sibling during her early life.
Offspring	Maternal Loss	0.68	1.98 (1.53-2.56)	**3×10^{-7}**	Offspring survive less well if they experienced maternal loss within four years of their birth.
	Low Maternal Rank	0.42	1.54 (1.17-2.01)	**0.002**	Offspring survive less well if they were born to a low-ranking mother.

3   [a]An alternative model that considered cumulative maternal adversity was not a better or worse fit
4   than the reduced multivariate maternal adversity model (see Table S7. For both the model
5   presented here and that in Table S7, $R^2=0.07$, log likelihood = -1598).

7