1   **Resting networks and personality predict attack speed in social spiders**

2   Edmund R. Hunt[1*], Brian Mi[2], Rediet Geremew[2], Camila Fernandez[2], Brandyn M. Wong[2],

3   Jonathan N. Pruitt[3], and Noa Pinter-Wollman[1]

[1] Department of Ecology and Evolutionary Biology, University of California, Los Angeles, CA
90095
[2] BioCircuits Institute, University of California, San Diego, La Jolla CA 92093
[3] Department of Ecology, Evolution, and Marine Biology, University of California, Santa
Barbara, CA 93106
*Corresponding author: Edmund R. Hunt, edmund.hunt@bristol.ac.uk, +44 (0)117 331 4760

ORCIDS: ERH, 0000-0002-9647-124X; JNP, 0000-0003-0326-6755; NP-W, 0000-0002-0448-
8037

Keywords: Collective behaviour, Foraging, Keystone individual, Boldness, Social network

analysis, *Stegodyphus dumicola*

Significance statement

Certain animals in a group, such as leaders, may have a more important role than other

group members in determining their collective behavior. Often these individuals are defined

by their behavioral attributes, for example, being bolder than others. We show that in social

spiders both the behavioral traits of the influential individual, and its interactions with other

group members, shape its role in affecting how quickly the group collectively attacks prey.

Abstract

Groups of social predators capture large prey items collectively, and their social interaction

patterns may impact how quickly they can respond to time-sensitive predation

opportunities. We investigated whether various organizational levels of resting interactions

(individual, sub-group, group), observed at different intervals leading up to a collective prey

attack, impacted the predation speed of colonies of the social spider *Stegodyphus dumicola*.

We found that in adult spiders overall group connectivity (average degree) increased group

attack speed. However, this effect was detected only immediately before the predation

event; connectivity two and four days before prey capture had little impact on the collective

dynamics. Significantly, lower social proximity of the group's boldest individual to other

group members (closeness centrality) immediately prior and two days before prey capture

was associated with faster attack speeds. These results suggest that for adult spiders, the

long-lasting effects of the boldest individual on the group's attack dynamics are mediated by

its role in the social network, and not only by its boldness. This suggests that behavioural

traits and social network relationships should be considered together when defining

keystone individuals in some contexts. By contrast, for subadult spiders, while the group

maximum boldness was negatively correlated with latency to attack, no significant resting

network predictors of latency to attack were found. Thus, separate behavioural mechanisms

might play distinctive roles in determining collective outcomes at different developmental

stages, timescales, and levels of social organization.

Introduction

Group living can benefit group members through access to mates, protection from

predators, and increased foraging opportunities (Krause and Ruxton, 2002). A variety of

animals engage in cooperative hunting to capture prey that is larger than what they could

capture alone. Examples of cooperative hunting can be seen in chimpanzees (Boesch, 2002),

lions (Stander, 1992), wild dogs (Creel and Creel, 1995), hawks (Bednarz, 1988), killer whales

(Pitman and Durban, 2012) and invertebrates such as ants (Witte et al., 2010), and social

spiders (Whitehouse and Lubin, 1999). During collective prey capture, individuals often

coordinate their behaviour through social interactions, to maximize their capture success

(Boesch, 2002; Pitman and Durban, 2012; Stander, 1992). A shorter latency to attack can

reduce the time and effort needed to capture prey, and increases the probability of success,

thus conferring important fitness benefits to all group members (Pasquet and Krafft, 1992).

At the same time, social groups often rely on particular individuals to expedite group

dynamics; such individuals have been termed 'keystone' individuals (Modlmeier et al.,

2014b).

It is still unknown how keystone individuals influence the performance of a group.

Generally speaking, keystone individuals can either perform the work itself or catalyse the

work of other group members (Robson and Traniello, 1999), for example through social

interactions. A recent model predicts that when the behaviour of group members is

persistent, social interactions play a larger role than behaviour in shaping collective

outcomes. This prediction emerges because who interacts with whom can change more

rapidly than behaviour, and behaviour does not necessarily determine who interacts with

whom. In contrast, if behaviour is plastic, social interactions may play a smaller role in

determining collective outcomes because behaviour is malleable (Pinter-Wollman et al.,

2016). Younger animals are often more behaviourally plastic than adults (Scott, 1962). Thus,

by comparing the social behaviour of both subadult, or juvenile, and adult organisms, we

can test when social interactions and when behavioural traits have a larger impact on

collective outcomes (Pinter-Wollman et al., 2016).

Social network analysis provides tools to quantify interaction patterns and has been

instrumental in understanding the dynamics and outcomes of interactions within groups

(Krause et al., 2015; Kurvers et al., 2014; Pinter-Wollman et al., 2014; Wey et al., 2008) and

in predicting group success (Royle et al., 2012). Different network measures can describe

interactions occurring at various organizational levels, such as an individual's direct

interactions with its neighbours, links within a subgroup, or interactions at the whole group

level (Krause et al., 2007; Lusseau and Newman, 2004; Wey et al., 2008; Wittemyer et al.,

2005). The social structure of animal groups often changes over time, and interaction

patterns occurring in one period can impact the group later (Blonder et al., 2012; Krause et

al., 2015; Pinter-Wollman et al., 2014). For example, social connectivity early in life can

predict male mating success several years later in long-tailed manakins (McDonald, 2007),

social connections to relatives can persist for longer than relationships with non-kin in

spotted hyenas (Holekamp et al., 2012), and social stability of subgroups can be maintained

over years in sparrows (Shizuka et al., 2014). Thus, temporal dynamics of interactions may

occur at different rates at different organization levels (Blonder et al., 2012).

Animals within a society often differ from one another in their behaviour, and these

differences can be consistent over time, a phenomenon that has been referred to as

'personality' (Bell et al., 2009; Jandt et al., 2014; Sih et al., 2004). Behavioural variation

among individuals in a social group can have a considerable impact on group functions

(Pinter-Wollman, 2012). Only a small amount of variation among individuals may be

necessary to have large impacts on the entire group. In the most extreme situations just one

'keystone' individual, such as a leader or a tutor, may have a disproportionate impact on the

group (Conradt and Roper, 2003; Modlmeier et al., 2014b). A keystone can be either a

particular individual or a role that different individuals assume at different times

(Modlmeier et al., 2014b). Keystone individuals can have an increased interaction rate with

other group members (Lloyd-Smith et al., 2005) and their behavioural tendencies can

influence interaction patterns (Firth et al., 2015; Krause et al., 2010; Pike et al., 2008; Pinter-

Wollman et al., 2011; Sih et al., 2009), thus in turn impacting collective actions (Bansal et al.,

2007; Brown and Irving, 2014).

Individual differences that are consistent over short time frames may change over

longer periods (Stamps and Groothuis, 2010). Such changes can alter the relationship

between a group's personality composition and collective outcomes over time. For example,

in social insects the task that each individual performs can change with age (Seeley, 1982;

Tripet and Nonacs, 2004) potentially altering the distribution of task performance in the

colony. Changes to an animal's personality may arise from changes in physiological

processes such as growth (Biro and Stamps, 2008), metamorphosis (Hedrick and Kortet,

2012; Wilson and Krause, 2012), or sexual maturation (Stamps and Groothuis, 2010).

Personality may also develop due to changes in the external physical or social environment

over time, i.e. experiential factors (Stamps and Groothuis, 2010). For example, spiders

placed in a group of unfamiliar individuals show initial high individual variation in boldness

that becomes more consistent over time, as group members become familiar with one

another (Laskowski et al., 2016; Laskowski and Pruitt, 2014; Modlmeier et al., 2014c).

In the social spider *Stegodyphus dumicola* (Araneae, Eresidae) individuals vary in

their boldness, and the boldest individual in the colony (referred to as the keystone)

determines the foraging intensity and mass gain of the entire group (Keiser and Pruitt, 2014;

Pruitt and Keiser, 2014). It is not known how the influence of keystone individuals is

imparted in this species, only that keystones have long lasting effects, and that the duration

of their impact is proportional to the tenure of the keystone in the group (Pruitt and Pinter-

Wollman, 2015). *S. dumicola* spiders live in colonies of up to several hundred individuals

that exhibit cooperative behaviours such as prey capture and allo-maternal care (Bilde et al.,

2007; Junghanns et al., 2017). Enhanced foraging success is thought to be a primary driver

of sociality in social spiders (Whitehouse and Lubin, 2005), to enable subduing of large prey

items (Guevara et al., 2011; Harwood and Avilés, 2013). More frequent co-feeding

interactions of the same prey item has been observed in sibling groups compared with non-

sibling groups in group-foraging subsocial spiders, suggesting that social network structure

may play a role in the evolution of social behaviour in spiders (Ruch et al., 2015). Colonies

composed of bolder spiders attack more rapidly (Keiser et al., 2014; Keiser and Pruitt, 2014;

Pruitt and Keiser, 2014) and with more individuals (Grinsted et al., 2013; Keiser and Pruitt,

2014; Laskowski and Pruitt, 2014; Pruitt and Keiser, 2014) than colonies with shy individuals.

Here we use social network analysis to determine the temporal scale and the social

organization level at which interactions between group members have the most impact on

collective prey capture dynamics. We evaluate if the keystone individual influences the

group through its role in the group's social network. We also consider whether these social

interactions have long-lasting effects on prey capture success, or if the impact of

interactions is immediate and ephemeral. Furthermore, we examine if developmental stage

(subadults vs. adults) affects the relationships between personality, social interactions, and

collective prey attack. Comparing subadults, with more plastic, emergent personalities, to

adults, with more established personalities, allows us to test the predictions of the model of

keystone influence detailed above (Pinter-Wollman et al., 2016).

In sum, we consider whether the speed at which *S. dumicola* colonies of either

subadult or adult spiders collectively attack prey depends on: *(i)* interactions occurring at

different levels of social organization; *(ii)* temporal changes in social structure; and *(iii)* the

behavioural and social attributes of the keystone individual.

Methods

Animal collection and maintenance

Colonies of *S. dumicola* were collected from roadside *Acacia* trees in the Northern Cape of

South Africa in November 2015 (subadults) and March 2016 (adults), transported to the

laboratory, and fed crickets *ad lib*. The size of collected colonies ranged between 70-300

individuals and contained only females - males are short-lived and rare (12%) in natural

colonies (Henschel et al., 1995). We created 15 groups of 26-30 female spiders, from 4

source colonies of subadults, and 24 groups of 10 adult female spiders each, from 3 other

source colonies. Group sizes were larger for subadults because of the small size of those

individuals, and because it potentially requires more small individuals to execute a

successful attack on large prey (see supplementary Figure S2 for differences in sizes

between adults and subadults). Groups were housed in large round containers (18cm

diameter, 8cm depth for subadults and 11cm diameter, 10cm depth for adults) with vertical

wire meshes (two 9x6cm sheets positioned 10cm apart for subadults and a 5x5cm sheet for

adults) to allow the spiders to build both a retreat and a capture web. Trials were conducted

from January until August, 2016.

Experimental procedure

To determine the effect of interaction patterns at different time scales on prey attack we

observed groups over time. Each group was observed for 6.5 weeks. Boldness and prey

capture were measured once a week and resting interactions, as detailed below, were

observed three times a week with 2-3 days separating each observation. The first resting

network was obtained immediately before measuring boldness on Day 4, numbered as four

days before measuring prey attack. The second resting network was observed on Day 2, two

days after measuring boldness and two days before measuring prey attack. The third resting

network was observed immediately before testing prey attack speed, on Day 0. This spacing

of measures allowed ample time for the spiders to recover from disturbances due to

removing them from their web to determine boldness. Each group was fed with a single 4-

week-old cricket once a week, which provides *ad lib* food, after the prey assay (described

below), hence all colonies had an equal opportunity to consume prey, and were at a similar

state of hunger. We obtained 7 boldness measures for each individual spider, 6 collective

prey capture response measures for each group, and 18 resting networks for each group (19

including a final boldness/network observation not used here). We compared the predictive

power of the resting networks observed four days, two days, and immediately before each

prey capture trial, for explaining the speed of prey attack. This allows us to differentiate

between short term (immediately before prey attack), medium (two days), and long (four

days) term influences of spider interaction networks.

Boldness

To determine individuals' boldness, each spider was tested once a week using an

established assay that measured the recovery of a spider from exposure to air puffs, which

mimic the approach of an avian predator (Riechert and Hedrick, 1993). Spiders react to the

air puffs by huddling and remaining still. The faster the spiders resume movement after this

simulated threat, i.e., move one body length away from where they were huddled, the

bolder they are considered. Boldness is a repeatable behaviour in this species when spiders

are kept isolated over days, with a repeatability of 0.63 (Keiser et al., 2014). However, in a

social context, boldness is much more plastic and changes as a function of the boldness of

the individuals with whom one interacts (Hunt et al., 2018). To test boldness, spiders were

placed individually in a plastic container (15x15cm) and after 30s of acclimation, two puffs

of air were administered to the anterior prosoma using an infant nose-cleaning bulb.

Boldness was measured as the latency to resume movement and move one body length.

Because bolder individuals resume movement faster, we subtracted the time to resume

movement from the maximum duration of the procedure (600s) to create a metric that

increases with boldness. We designated as 'shy' individuals those with a latency to resume

movement of 400-600s, while 'bold' individuals were those with a latency to resume

movement of 0-200s. The abdomen of each spider was marked uniquely with acrylic paint

to track their behaviour over time (Figure 1). We examined how many individuals occupied

the role of keystone (boldest) in each experimental group over the 7 weeks of the

experiment – this ranged from 1 to 7 individuals (same individual throughout, or a different

individual each week). A priori, turnover is expected to be higher for the subadults because

they have larger group sizes and therefore more individuals that might replace the keystone

individual. We compare the keystone turnover for the two developmental stages using a

Wilcoxon rank sum (Mann-Whitney) test. We used linear interpolation to obtain an

estimate of boldness on Days 0 and 2 and identify a putative boldest individual on those

days, which were in between the weekly boldness measurement taken on Day 4.

Group composition

To examine the effect of group composition on collective behaviour we assigned spiders to

one of three group compositions: all shy, all bold, and 'keystone' (all shy individuals plus one

bold individual). For subadults, we established five groups of each composition and for

adults we established five groups of all bold, nine of all shy, and ten keystone groups.

Individuals that were not assigned to experimental groups, including those with a boldness

score of 200-400, were returned to their source colony. After the first week of our study,

the average boldness of all group compositions converged, with the 'all bold' groups

reducing their average boldness substantially (Figure S1 A, B). We used a one-way ANOVA to

compare mean group boldness of the three different group compositions, for the latter five

weeks of our experiment, excluding week one. Because we did not detect a significant

difference in average boldness between the three compositions for the latter five weeks

(subadults: $F_{2,71}=0.739$, p=0.48; adults: $F_{2,117}=2.076$, p=0.13), we excluded week one and

pooled the remaining data across compositions in all analyses (see also distribution analysis

in Table S1). Because boldness was found to be more plastic in a social context compared

with isolation, and because the artificially manipulated boldness distributions were quickly

returned to their natural skewed state by the spiders' collective boldness dynamics, we did

not find evidence that our treatments had any long-term effects. For further information on

changes in boldness over time in a social context see Hunt et al., 2018.

Social interactions

The physical contacts among spiders were manually recorded three times a week:

immediately (1-2 hours) before the prey-capture assay, two days prior to the prey capture

assay, and four days before to the prey capture assay. Resting interactions were defined as

a physical contact between any body parts of two spiders (Figure 1). Interactions were

observed during the day, when spiders are resting and inactive. Care was taken to note each

spider in the colony so that all interactions are recorded. These interactions were used to

construct social networks and calculate network variables that indicate individual, sub-

group, and group level dynamics. The network variables measuring individual level

behaviours were keystone degree, keystone closeness, and maximum boldness in the group

(i.e., boldness of the keystone); for the sub-group level, modularity; and for the group level,

average degree and degree distribution skewness, as detailed in Table 1.

To calculate the 2 individual-level network measures (Table 1), we had to first

identify which individual was the boldest, i.e. occupying the (keystone role each week. In

this system, the individual with the highest boldness assumes the keystone role, regardless

of its identity (Pinter-Wollman et al., 2017b). Thus, the role of keystone is not necessarily

maintained by a specific individual. In a social setting, individuals change their boldness over

time (Hunt et al., 2018) and so boldness ranks among individuals may also shift. When more

than one spider exhibited the same maximum value, we took the average network value of

those individuals (this happened 2 out of 74 times for the subadult spiders and 1 out of 120

times for the adults). When all spiders had zero boldness, we identified the keystone

individual as the boldest spider in the previous week (this did not occur for the subadults,

and 13 out of 120 times for the adults); where this was not possible, we calculated network

values as an average across all individuals in the group that week (7 out of 120 times for the

adults).

Prey response

To determine the speed at which groups attack prey, we examined the spiders' latency to

respond to vibrations on their capture web (Grinsted et al., 2013). We used a custom-made

vibratory device assembled from an Arduino Uno board, a vibratory motor, and a metal

wire, directed at a 1x1cm piece of paper placed in the capture web (Pinter-Wollman et al.,

2017b). The stimulus was always placed on the capture web at the same distance (4cm)

from the nest retreat, where most spiders were gathered, to control for any effects the

distance of the stimulus might have on the response of the spiders. The Arduino board was

programmed to vibrate the piece of paper in pulses that varied randomly between 0.5-1.5

sec in both the duration of the vibration and the pauses between vibrations, to simulate the

irregular vibrations that a prey makes when captured in the web (Hedrick and Riechert,

1989). The paper was vibrated until a spider touched it, to avoid habituation to our stimulus,

or until 10 minutes elapsed, in which case the trial was stopped (Pinter-Wollman et al.,

2017b). As the first individual left the retreat, others followed, creating a collective

response. The first individual to leave the retreat was not necessarily the one closest to the

simulated prey (personal observations). When no attack took place, we set the latency to

attack to ten minutes. We noted the identity of the first individual(s) to touch the stimulus,

as well as the identity of all the individuals that left the nest during the attack as

participants, so that we could assess whether the keystone (boldest) individual participated

in prey attack. Both adult and subadult groups responded to the simulated prey in a similar

manner (Figure S3).

Data analysis

To examine the relationship between social network structure, boldness, and prey attack we

used censored mixed regression models. We considered six variables as predictors of

latency to attack, as detailed in Table 1. These were included as fixed effects interacting

with the effect 'Day' which accounted for the number of days before the attack assay (4, 2,

or 0). This approach allowed us to determine the timescale on which social interactions act

on prey capture. We constructed separate models for adult and subadult spider behaviour,

and each model included 5 weeks of data (weeks 2-6, excluding week 1). For the adult

spiders these included N=360 resting network observations (24 colonies x 5 weeks x 3

observations per week) and for the subadults N=224 (15 colonies x 5 x 3, except N=74 for

Day 0 because of 1 missing network observation). Because latency to attack was right-

censored at 600 s, with 76% of subadult trials and 65% of adult trials resulting in an attack,

we used censored regression (Tobit) models with the R package 'censReg' (Henningsen,

2017). The response variable, latency to attack, was log-transformed to adhere to the model

assumption that the error term is normally distributed. To account for variation among

groups and source colonies we included group identity as a random effect and source

colony identity as a fixed effect. We further included a time-varying residual component as a

random effect to account for changes over the 5 weeks. Because we did not have an *a priori*

prediction regarding which network variable would best explain collective prey attack, we

identified a suitable model for the adults and subadults by first estimating all 63 possible

models that linearly combined one or more of the 6 fixed effects listed in Table 1. We

calculated the Akaike weight of the 63 models (Burnham et al., 2011). These model weights

were then used to estimate the relative importance of the six predictor variables under

consideration (Table 1). The importance of each predictor is determined by summing the

Akaike weights of each model in which it appears (Symonds and Moussalli, 2011). If a

certain predictor appears in many of the top models, its summed Akaike weight will tend

toward 1. On the other hand, if it only appears in the weaker models, its weight will tend

toward 0 (Symonds and Moussalli, 2011). This procedure results in a ranking of predictor

variables in terms of performance, and one can interpret the predictor weight as a

probability that it is a component of the best model (Burnham and Anderson, 2002;

Symonds and Moussalli, 2011). We selected the top-ranked predictor variable at the

individual level, and the top-ranked predictor at the group or subgroup level, to obtain a

parsimonious model for the adult and subadult behaviour with two main predictors in each

model (Tables 2, 3, 4). We find this predictor-ranking approach to be preferable to full-

model averaging because it allows us to identify a particular model with properly estimated

predictor standard errors and without excessive model complexity (full-model averaging

results are available in Table S7, S8). We checked for multicollinearity between predictors in

the final models by calculating their corrected generalized variance inflation factors (Fox and

Monette, 1992), using the R package 'car' (Fox and Weisberg, 2011). There was low

collinearity in both the adult model (Table S3) and subadult model (Table S5). To assess

whether the keystone spider's role in the interactions network (closeness and degree) was

correlated with its boldness we calculated the Pearson's correlation between these

measures.

Results

The adult spider model included average degree (predictor weight 46%), as the group or

subgroup-level fixed effect, and closeness of the keystone individual (predictor weight 81%),

as the individual-level fixed effect (Tables 2, 3). The subadult spider model included average

degree (predictor weight 47%), as the group or subgroup-level fixed effect, and maximum

boldness (predictor weight 97%), as the individual-level fixed effect (Tables 2, 4).

Network measures impacted collective prey attack in groups of adult spiders. Adult

spider groups that were overall more connected (high average degree) attacked the prey

stimulus more quickly. This relationship was found only for resting networks observed

immediately before testing collective prey attack, but not for networks measured earlier

(Table 3). At the level of the individual, when the groups' boldest (keystone) individual was

more closely connected to all other individuals (closeness centrality), prey attack was

slower. This relationship between prey attack and the closeness centrality of the boldest

individual was retained for resting networks obtained two days prior to prey attack (Table

3).

No relationship between the network measure average degree and attack speed was

found for the groups of subadult spiders (Tables 4). We also ran an alternative model that

included degree of keystone as a second individual-level, network effect (predictor weight

52%) but this was also not significant (Table S6). However, the boldness of the boldest

individual in the group, i.e., the keystone's boldness, was a significant predictor of latency to

attack; groups with a bolder keystone attacked more quickly (Table 4).

In the adult spiders, the boldness of the boldest (keystone) individual was not

associated with its degree in the before-prey resting network (Pearson's correlation: r =

0.039, t = 0.428, df = 118, p = 0.670), or its closeness (Pearson's correlation: r = -0.007, t = -

0.074, df = 118, p-value = 0.941). This was also the case for the subadults' boldest spiders

and their degree (Pearson's correlation: r = -0.212, t = -1.837, df = 72, p-value = 0.070) and

closeness (Pearson's correlation: r = -0.069, t = -0.5909, df = 72, p = 0.556). There was high

turnover in the role of keystone (boldest) individual in both in the adult (4.38 $\pm$ 0.68, mean

$\pm$ standard deviation) and subadult (5.67 $\pm$ 0.87) groups (number of different individuals in

the role out of a maximum of 7). There was a significant difference between the adult and

subadult distributions of keystone turnover (Wilcoxon rank sum test: W=230, p=0.0003).

Keystone individuals participated in 7-29% of prey attacks. In the adult groups on the

first week, 24 trials resulted in 21 prey attacks. 17 of these attacking groups had a keystone

individual (the remainder of the groups were set up as all-shy individuals), and 5 (29.4%) of

the attacks by these groups included the keystone individual. In contrast, on weeks 2-6, 73

of 120 trials resulted in prey attacks. In 70 of these groups there was an individual that had

the highest boldness (i.e., was a keystone), and in only 5 (7.1%) of these groups, the

keystone individual participated in the prey attack. This decrease in keystone participation

in prey attack after week 1 is consistent with previous findings (Figure S2 of Pruitt and

Pinter-Wollman, 2015). In the subadult groups, on the first week, 15 trials resulted in 12

prey attacks and only in one (8.3%) of these attacks a keystone individual participated. In

weeks 2-6, there were 75 trials, 56 resulting in prey attack, 4 (7.1%) of which had keystone

participation.

Discussion

The structure of resting interaction networks measured immediately before prey

attack predicted the attack speed of adult social spiders. Higher overall connectivity

(average degree), a group-level measure, led to faster attack, though only in the resting

interactions measured pre-stimulus. Furthermore, the more connected to others (greater

closeness centrality) was the boldest individual (keystone), the slower the collective prey

attack speeds, when considering both the networks observed immediately and two days

prior to the prey attack. Therefore, the social connectivity of the keystone to the rest of its

group, but not its boldness, was found to be significant in groups of adults. The opposite

finding was observed for the subadult spiders: groups with a bolder keystone individual

attacked the prey stimulus more quickly, but we did not detect any significant associations

between subadult resting network measures and attack speeds. These findings support our

hypothesis that subadult spiders will rely on behavioural traits likely because they have

greater behavioural plasticity than adults, owing to their relative lack of prior exposure to

social contact, over their lifetime, not just within the experiment. Our results further

support the hypothesis that adult groups rely on social interactions more than on

behavioural changes to modify collective dynamics. We also found that the individual-level

predictor had a noticeably higher predictor weight compared to the group-level predictor in

both the adults and subadults: 81% vs 46% in the adults, and 97% vs 47% in the subadults

(Table 3). This seems to confirm the relevance of the 'keystone individual' concept in this

species.

Influential 'keystone' individuals in social groups have a large effect on their social

environment relative to their abundance (Modlmeier et al., 2014b). While this influence can

be mediated through the behavioural traits of the keystone individual, it may also depend

on its social interactions. Here we found that the behavioural traits of the keystone are not

the only feature that impacts group success. When the boldest individual in an adult spider

group had a lower closeness, i.e., was less connected with other resting spiders, the colony

attacked prey more quickly. Thus, we find that the impact of the keystone's defining trait –

high boldness – was mediated by its interactions with the rest of the social group. Other

studies define keystone individuals according to their centrality in an interaction network

(Lusseau and Newman, 2004; Modlmeier et al., 2014b; Vital and Martins, 2013) and deem

central individuals as critical for the stability of their society. Our findings suggest that *both*

the behavioural traits and role in the social network should be considered when defining

keystone individuals in some contexts. The boldness of the keystone individuals was not

associated with network closeness or degree in either adults or subadults, which suggests

that centrality in an interaction network is not merely a direct consequence of individual

behavioural characteristics but may result from a different process.

Our results suggest that the effect of the keystone individual on group dynamics that

have been previously observed (Keiser and Pruitt, 2014; Pruitt and Keiser, 2014; Pruitt and

Pinter-Wollman, 2015), could be made directly through its own behaviour, or through

influencing the formation and dissolution of social interactions. Here, there was low direct

participation of keystone individuals in prey attacks (7-29%), which is similar to keystone

participation seen in other studies of this system (Pruitt and Pinter-Wollman, 2015). Direct

keystone participation in prey attack is most common during the first three days after *S.*

dumicola colony establishment (Pruitt and Pinter-Wollman, 2015) or after a new boldest

individual is introduced (Pinter-Wollman et al., 2017b). Keystones rarely participate in prey

capture directly in established colonies (Pruitt and Keiser, 2014). Seismic recruitment signals

have been indicated in *S. sarasinorum* (Bradoo, 1980) and other spider genera such as

Theridion saxatile (Norgaard, 1956). Likewise, adult *S. dumicola* are observed to catalyse

foraging participation by juvenile spiders without becoming directly involved (Modlmeier et

al., 2015). Thus, faster attack speeds may result from keystone individuals signalling

information to the rest of the group about the presence of prey through, for example,

vibrations on the web. Such signalling does not necessarily require physical proximity, only

connection through the capture web, thus it is consistent with the peripheral position of the

keystone individual in the proximity networks. In addition, the observed fast attacks in

groups with high connectivity that we observed could point to another mechanism

underlying the effects of interactions on behaviour. Recent work shows that in this study

system boldness is impacted by proximity interactions (Hunt et al., 2018). Thus, it is possible

that individuals in groups with high connectivity have more opportunities to modify each

other's behaviour and shape overall group boldness composition, which impacts prey

attack.

The closeness of the keystone was a significant predictor of attack speed up to two

days before the stimulus. This effect duration may be a result of persistence in occupying

certain social roles in the interaction network or, as previous research indicates, because

keystone spiders have legacy effects on their social environment (Pruitt and Pinter-

Wollman, 2015). A keystone individual's long-term influence on its group can result from

influencing the behaviour of others or their interactions. For example, in primates the

removal of certain individuals who engage in policing behaviour affects the group's social

network structure, even in situations, such as play, in which the removed individuals did not

participate (Flack et al., 2006). *S. dumicola* colonies containing bolder keystones attack with

more spiders, even after the keystone has been removed, suggesting that keystone

individuals shape the overall structure of the social network, and/or change the group

members' individual characteristics (Pruitt and Pinter-Wollman, 2015). For example, bold

spiders may actively contribute to a process of social niche construction, whereby repeated

social interactions reinforce and enhance inter-individual differences in behaviour such as

boldness and prey attack speed (Laskowski and Pruitt, 2014).

The impact of overall group connectivity (average degree) on the adult spiders'

response to prey suggests that most group members have a role to play in coordinating

collective predation, even without needing to participate directly in prey attack. For

instance, the presence of fellow colony members may catalyse increased foraging

aggressiveness in individuals that are already predisposed to participate in prey capture

(e.g., bold individuals (Wright et al., 2015, 2016)). Such catalytic effects have been observed

in the context of nightly web repair in *S. dumicola* (Keiser et al., 2016). Furthermore, group

level interactions impacted collective prey attack only on a short-time scale. The significant

effect of group connectivity (average degree) immediately before, but not two and four

days prior to prey capture on its speed, suggests that group level measures are important

for immediate dynamics, such as the coordination of motion during prey capture (Krafft and

Pasquet, 1991). Such immediate effects are consistent with observations that small and

confined artificial nest scaffolds facilitate rapid prey capture (Modlmeier et al., 2014a).

Tighter proximity among group members may increase the frequency of social interactions

among all group members (Modlmeier et al., 2014a). Future work on the impact of nest

architecture on social interactions could uncover a potential mechanism underlying social

interaction patterns (Pinter-Wollman et al., 2017a).

If individuals are behaviourally flexible, the group may be less reliant on social

interactions to coordinate activities. Subadult spider groups attacked the prey stimulus

more quickly if they had a bolder keystone individual, however, no association was found

between their resting networks and attack speeds. These findings are consistent with

previous work showing that groups with bolder keystones attack prey faster and with more

attackers (Keiser and Pruitt, 2014). The lower importance of social network structure in

subadults compared with adult spider groups could be related to greater behavioural

plasticity in the subadults. In a new social environment *Stegodyphus* spiders show higher

within-individual variation in boldness before becoming more behaviourally consistent with

time (Laskowski et al., 2016; Laskowski and Pruitt, 2014; Modlmeier et al., 2014c). Earlier

developmental stages show more behavioural flexibility in a variety of animal systems

(Stamps and Groothuis, 2010). Here we found high turnover in keystone identity in both

adults and subadults, with higher turnover in the subadults. Because adult spiders are

perhaps more constrained in adjusting their behavioural traits, they may be more reliant on

social interactions to achieve a beneficial group-level behavioural phenotype, because who

interacts with whom can be changed more easily. Thus, effective group-level hunting

behaviour, which is a key function of sociality in social spiders (Whitehouse and Lubin,

2005), could depend on a balance between the plasticity of individual-level behavioural

traits and the ease of modifying social interactions (Pinter-Wollman et al., 2016), which may

vary according to developmental stage.

Finally, group size may have played a role in the differences we observed between

adult and subadult groups. In our experiments, adult groups had 10 individuals and

subadults 26-30 and so development stage and group size are not independent of each

other in this study. Individuals in smaller groups have been found to participate more in

collective prey capture, probably as a consequence of necessity rather than a change in

behaviour (Wright et al., 2015). Thus, it is possible that, if not all individuals are required for

prey capture in large groups, interactions among all group members are less important than

in small groups. Furthermore, larger groups are more likely to contain skilled or experienced

individuals, assuming they have greater diversity in ability or experience; this is referred to

as the 'pool of competence' hypothesis (Morand-Ferron and Quinn, 2011). However,

subadults are all likely equally inexperienced and unskilled at hunting and so variation in

experience in a large group may not be a suitable explanation for the differences we

observed between the adult and subadult groups. Finally, the containers occupied by

subadults were somewhat larger than those of the adults (to accommodate the larger

number of individuals). However, subadult group members tended to all cluster in one small

area and not utilize more space than the adult spiders in their smaller containers.

Therefore, we expect group size to have a smaller impact on the differences between

subadult and adult groups than the difference in developmental stage.

Conclusion

Our study shows that social interactions are important in determining the speed of

collective predation in social spiders and that the impact of interactions may differ

according to developmental stage, time scale, and level of social organization.

Developmental stages may vary in the importance of social interactions versus shifts in

individuals' behavioural tendencies (Pinter-Wollman et al., 2016). Furthermore, different

types of interactions on different social organizational levels may vary in the duration of

their effects. For example, we found that an individual-based network measure had longer

lasting effects than a group-level measure. Finally, our results indicate that the social role of

keystone individuals within a group can have a longer-term effect on collective hunting

success. This finding emphasizes the importance of observing both the social structure of

individuals in a group and noting inter-individual differences in personality characteristics,

for understanding the success of some animal societies. Although keystone individuals can

occupy a wide variety of roles across different taxa (Modlmeier et al., 2014b), our work

suggests that a combination of network and behavioural trait characteristics could provide a

more precise definition of keystones in many contexts in future work.

Acknowledgements

We thank the South Africa Department of Tourism, Environment, and Conservation for

providing permits for animal collection (FAUNA 1072/2013 and 1691/2015) and Colin

Wright and James Lichtenstein for collecting spiders in the field. We further thank

Arne Henningsen for guidance on the 'censReg' R package.

Data availability

The datasets generated and analysed during the study will be available in an online data

repository upon acceptance of the paper.

Funding

This work was supported by the National Science Foundation IOS grants 1456010 to NPW

and 1455895 to JNP, and National Institutes of Health grant GM115509 to NPW and JNP.

Author contributions

ERH analyzed the data and drafted the manuscript, NPW and JNP designed the study, BM,

RG, CF, BW, and NPW collected the data, and all authors contributed to the final version of

the manuscript.

References

Bansal S, Grenfell BT, Meyers LA, 2007. When individual behaviour matters: homogeneous
533   and network models in epidemiology. Journal of The Royal Society Interface 4:879-
534   891. doi: 10.1098/rsif.2007.1100.
535   Bednarz JC, 1988. Cooperative hunting in Harris' hawks (Parabuteo unicinctus). Science
536   239:1525.
537   Bell AM, Hankison SJ, Laskowski KL, 2009. The repeatability of behaviour: a meta-analysis.
538   Anim Behav 77:771-783. doi: DOI 10.1016/j.anbehav.2008.12.022.
539   Bilde T, Coates KS, Birkhofer K, Bird T, Maklakov AA, Lubin Y, Aviles L, 2007. Survival benefits
540   select for group living in a social spider despite reproductive costs. Journal of
541   Evolutionary Biology 20:2412-2426. doi: 10.1111/j.1420-9101.2007.01407.x.
542   Biro PA, Stamps JA, 2008. Are animal personality traits linked to life-history productivity?
543   Trends in Ecology & Evolution 23:361-368. doi: 10.1016/j.tree.2008.04.003.
544   Blonder B, Wey TW, Dornhaus A, James R, Sih A, 2012. Temporal dynamics and network
545   analysis. Methods in Ecology and Evolution 3:958-972. doi: 10.1111/j.2041-
546   210X.2012.00236.x.

Boesch C, 2002. Cooperative hunting roles among taï chimpanzees. Human Nature 13:27-46.
doi: 10.1007/s12110-002-1013-6.
Bradoo B, 1980. Feeding behaviour and recruitment display in the social spider *Stegodyphus*
sarasinorum Karsch (Araneae, Eresidae). Tijdschrift voor entomologie 123.
Brandes U, Delling D, Gaertler M, Gorke R, Hoefer M, Nikoloski Z, Wagner D, 2008. On
Modularity Clustering. IEEE Transactions on Knowledge and Data Engineering
20:172-188. doi: 10.1109/TKDE.2007.190689.
Brown C, Irving E, 2014. Individual personality traits influence group exploration in a feral
guppy population. Behav Ecol 25:95-101. doi: DOI 10.1093/beheco/art090.
Burnham KP, Anderson DR, 2002. Model selection and multimodel inference, 2nd edn.
Springer, New York.
Burnham KP, Anderson DR, Huyvaert KP, 2011. AIC model selection and multimodel
inference in behavioral ecology: some background, observations, and comparisons.
Behavioral Ecology and Sociobiology 65:23-35. doi: 10.1007/s00265-010-1029-6.
Conradt L, Roper TJ, 2003. Group decision-making in animals. Nature 421:155-158. doi:
10.1038/Nature01294.
Creel S, Creel NM, 1995. Communal hunting and pack size in African wild dogs, Lycaon
pictus. Animal Behaviour 50:1325-1339. doi: 10.1016/0003-3472(95)80048-4.
Firth JA, Voelkl B, Farine DR, Sheldon BC, 2015. Experimental Evidence that Social
Relationships Determine Individual Foraging Behavior. Curr Biol 25:3138-3143. doi:
10.1016/j.cub.2015.09.075.
Flack JC, Girvan M, de Waal FBM, Krakauer DC, 2006. Policing stabilizes construction of
social niches in primates. Nature 439:426-429. doi: 10.1038/nature04326.
Fox J, Monette G, 1992. Generalized Collinearity Diagnostics. Journal of the American
Statistical Association 87:178-183. doi: 10.1080/01621459.1992.10475190.
Fox J, Weisberg S, 2011. An R Companion to Applied Regression, Second Edition. Thousand
Oaks CA: Sage.
Grinsted L, Pruitt JN, Settepani V, Bilde T, 2013. Individual personalities shape task
differentiation in a social spider. Proceedings of the Royal Society B: Biological
Sciences 280. doi: 10.1098/rspb.2013.1407.
Guevara J, Gonzaga MO, Vasconcellos-Neto J, Avilés L, 2011. Sociality and resource use:
insights from a community of social spiders in Brazil. Behavioral Ecology 22:630-638.
doi: 10.1093/beheco/arr022.
Harwood G, Avilés L, 2013. Differences in group size and the extent of individual
participation in group hunting may contribute to differential prey-size use among
social spiders. Biology Letters 9. doi: 10.1098/rsbl.2013.0621.
Hedrick AV, Kortet R, 2012. Sex differences in the repeatability of boldness over
metamorphosis. Behavioral Ecology and Sociobiology 66:407-412. doi:
10.1007/s00265-011-1286-z.
Hedrick AV, Riechert SE, 1989. Genetically-based variation between two spider populations
in foraging behavior. Oecologia 80:533-539. doi: 10.1007/bf00380078.
Henningsen A, 2017. censReg: Censored Regression (Tobit) Models.
Henschel JR, Lubin YD, Schneider J, 1995. Sexual competition in an inbreeding social spider,
Stegodyphus dumicola (Araneae: Eresidae). Insectes Sociaux 42:419-426. doi:
10.1007/BF01242170.

Holekamp KE, Smith JE, Strelioff CC, Van Horn RC, Watts HE, 2012. Society, demography and
genetic structure in the spotted hyena. Molecular Ecology 21:613-632. doi:
10.1111/j.1365-294X.2011.05240.x.
Hunt ER, Mi B, Fernandez C, Wong BM, Pruitt JN, Pinter-Wollman N, 2018. Social
interactions shape individual and collective personality in social spiders. P R Soc B
285. doi: 10.1098/rspb.2018.1366.
Jandt JM, Bengston S, Pinter-Wollman N, Pruitt JN, Raine NE, Dornhaus A, Sih A, 2014.
Behavioural syndromes and social insects: personality at multiple levels. Biological
Reviews 89:48-67. doi: Doi 10.1111/Brv.12042.
Junghanns A, Holm C, Schou MF, Sørensen AB, Uhl G, Bilde T, 2017. Extreme allomaternal
care and unequal task participation by unmated females in a cooperatively breeding
spider. Animal Behaviour 132:101-107. doi: 10.1016/j.anbehav.2017.08.006.
Keiser CN, Jones DK, Modlmeier AP, Pruitt JN, 2014. Exploring the effects of individual traits
and within-colony variation on task differentiation and collective behavior in a desert
social spider. Behavioral Ecology and Sociobiology 68:839-850. doi: DOI
10.1007/s00265-014-1696-9.
Keiser CN, Pruitt JN, 2014. Personality composition is more important than group size in
determining collective foraging behaviour in the wild. P R Soc B 281:20141424. doi:
10.1098/rspb.2014.1424.
Keiser CN, Wright CM, Pruitt JN, 2016. Increased bacterial load can reduce or negate the
effects of keystone individuals on group collective behaviour. Animal Behaviour
114:211-218. doi: 10.1016/j.anbehav.2016.02.010.
Krafft B, Pasquet A, 1991. Synchronized and rhythmical activity during the prey capture in
the social spider *Anelosimus eximius* (Araneae, Theridiidae). Insectes Sociaux 38:83-
90. doi: 10.1007/bf01242716.
Krause J, Croft DP, James R, 2007. Social network theory in the behavioural sciences:
potential applications. Behavioral Ecology and Sociobiology 62:15-27.
Krause J, James R, Croft DP, 2010. Personality in the context of social networks. Philos T R
Soc B 365:4099-4106. doi: 10.1098/rstb.2010.0216.
Krause J, James R, Franks DW, Croft DP, 2015. Animal social networks: Oxford University
Press, USA.
Krause J, Ruxton GD, 2002. Living in groups: Oxford University Press.
Kurvers RHJM, Krause J, Croft DP, Wilson ADM, Wolf M, 2014. The evolutionary and
ecological consequences of animal social networks: emerging issues. Trends in
Ecology & Evolution 29:326-335. doi: 10.1016/j.tree.2014.04.002.
Laskowski KL, Montiglio P-O, Pruitt JN, 2016. Individual and Group Performance Suffers from
Social Niche Disruption. The American Naturalist 187:776-785. doi: 10.1086/686220.
Laskowski KL, Pruitt JN, 2014. Evidence of social niche construction: persistent and repeated
social interactions generate stronger personalities in a social spider. P Roy Soc B-Biol
Sci 281. doi: 10.1098/rspb.2013.3166.
Lloyd-Smith JO, Schreiber SJ, Kopp PE, Getz WM, 2005. Superspreading and the effect of
individual variation on disease emergence. Nature 438:355-359. doi: Doi
10.1038/Nature04153.
Lusseau D, Newman MEJ, 2004. Identifying the role that animals play in their social
networks. Proceedings Biological sciences / The Royal Society 271 Suppl:S477-481.
doi: 10.1098/rsbl.2004.0225.

McDonald DB, 2007. Predicting fate from early connectivity in a social network. Proceedings
of the National Academy of Sciences 104:10910-10914. doi:
10.1073/pnas.0701159104.
Modlmeier AP, Forrester NJ, Pruitt JN, 2014a. Habitat structure helps guide the emergence
of colony-level personality in social spiders. Behavioral Ecology and Sociobiology
68:1965-1972. doi: 10.1007/s00265-014-1802-z.
Modlmeier AP, Keiser CN, Watters JV, Sih A, Pruitt JN, 2014b. The keystone individual
concept: an ecological and evolutionary overview. Anim Behav 89:53-62.
Modlmeier AP, Laskowski KL, Brittingham HA, Coleman A, Knutson KA, Kuo C, McGuirk M,
Zhao K, Keiser CN, Pruitt JN, 2015. Adult presence augments juvenile collective
foraging in social spiders. Animal Behaviour 109:9-14. doi:
10.1016/j.anbehav.2015.07.033.
Modlmeier AP, Laskowski KL, DeMarco AE, Coleman A, Zhao K, Brittingham HA, McDermott
DR, Pruitt JN, 2014c. Persistent social interactions beget more pronounced
personalities in a desert-dwelling social spider. Biology Letters 10. doi:
10.1098/rsbl.2014.0419.
Morand-Ferron J, Quinn JL, 2011. Larger groups of passerines are more efficient problem
solvers in the wild. Proceedings of the National Academy of Sciences 108:15898-
15903. doi: 10.1073/pnas.1111560108.
Norgaard E, 1956. Environment and Behaviour of Theridion Saxatile. Oikos 7:159-192. doi:
10.2307/3564917.
O'Brien RM, 2007. A Caution Regarding Rules of Thumb for Variance Inflation Factors.
Quality & Quantity 41:673-690. doi: 10.1007/s11135-006-9018-6.
Pasquet A, Krafft B, 1992. Cooperation and Prey Capture Efficiency in a Social Spider,
Anelosimus eximius (Araneae, Theridiidae). Ethology 90:121-133. doi:
10.1111/j.1439-0310.1992.tb00826.x.
Pike TW, Samanta M, Lindstrom J, Royle NJ, 2008. Behavioural phenotype affects social
interactions in an animal network. P R Soc B 275:2515-2520. doi: DOI
10.1098/rspb.2008.0744.
Pinter-Wollman N, 2012. Personality in social insects: How does worker personality
determine colony personality? Curr Zool 58:580-588. doi: 10.1093/czoolo/58.4.580.
Pinter-Wollman N, Fiore SM, Theraulaz G, 2017a. The impact of architecture on collective
behaviour. Nature Ecology & Evolution:0111. doi: DOI: 10.1038/s41559-017-0111.
Pinter-Wollman N, Hobson EA, Smith JE, Edelman AJ, Shizuka D, de Silva S, Waters JS, Prager
SD, Sasaki T, Wittemyer G, Fewell J, McDonald DB, 2014. The dynamics of animal
social networks: analytical, conceptual, and theoretical advances. Behavioural
Ecology 25:242–255.
Pinter-Wollman N, Hubler J, Holley J-A, Franks NR, Dornhaus A, 2012. How is activity
distributed among and within tasks in Temnothorax ants? Behavioral Ecology and
Sociobiology 66:1407-1420. doi: 10.1007/s00265-012-1396-2.
Pinter-Wollman N, Keiser CN, Wollman R, Pruitt JN, 2016. The Effect of Keystone Individuals
on Collective Outcomes Can Be Mediated through Interactions or Behavioral
Persistence. American Naturalist 188:240-252. doi: 10.1086/687235.
Pinter-Wollman N, Mi B, Pruitt JN, 2017b. Replacing bold individuals has a smaller impact on
group performance than replacing shy individuals. Behavioral Ecology 28:883-889.
doi: 10.1093/beheco/arx054.

Pinter-Wollman N, Wollman R, Guetz A, Holmes S, Gordon DM, 2011. The effect of
individual variation on the structure and function of interaction networks in
harvester ants. Journal of the Royal Society Interface 8:1562-1573. doi: DOI
10.1098/rsif.2011.0059.
Pitman RL, Durban JW, 2012. Cooperative hunting behavior, prey selectivity and prey
handling by pack ice killer whales (*Orcinus orca*), type B, in Antarctic Peninsula
waters. Marine Mammal Science 28:16-36. doi: 10.1111/j.1748-7692.2010.00453.x.
Pruitt JN, Keiser CN, 2014. The personality types of key catalytic individuals shape colonies'
collective behaviour and success. Anim Behav 93:87–95.
Pruitt JN, Pinter-Wollman N, 2015. The legacy effects of keystone individuals on collective
behaviour scale to how long they remain within a group. P R Soc B 282:89-96. doi:
10.1098/rspb.2015.1766.
Riechert SE, Hedrick AV, 1993. A test for correlations among fitness-linked behavioural traits
in the spider *Agelenopsis aperta* (Araneae, Agelenidae). Animal Behaviour 46:669-
675. doi: 10.1006/anbe.1993.1243.
Robson SK, Traniello JFA, 1999. Key individuals and the organization of labor in ants. In:
Detrain C, Deneubourg JL, Pasteels JM, editors. Information processing in social
insects Basel Boston, Berlin: Birkhauser. p. 239-260.
Royle NJ, Pike TW, Heeb P, Richner H, Kolliker M, 2012. Offspring social network structure
predicts fitness in families. P Roy Soc B-Biol Sci 279:4914-4922. doi:
10.1098/rspb.2012.1701.
Scott JP, 1962. Critical Periods in Behavioral Development. Science 138:949-958.
Seeley TD, 1982. Adaptive significance of the age polyethism schedule in honeybee colonies.
Behavioral Ecology and Sociobiology 11:287-293. doi: 10.1007/bf00299306.
Shizuka D, Chaine AS, Anderson J, Johnson O, Laursen IM, Lyon BE, 2014. Across-year social
stability shapes network structure in wintering migrant sparrows. Ecology Letters
17:998-1007. doi: 10.1111/ele.12304.
Sih A, Bell A, Johnson JC, 2004. Behavioral syndromes: an ecological and evolutionary
overview. Trends Ecol Evol 19:372-378. doi: DOI 10.1016/j.tree.2004.04.009.
Sih A, Hanser SF, McHugh KA, 2009. Social network theory: new insights and issues for
behavioral ecologists. Behavioral Ecology and Sociobiology 63:975-988. doi: DOI
10.1007/s00265-009-0725-6.
Stamps J, Groothuis TGG, 2010. The development of animal personality: relevance, concepts
and perspectives. Biological Reviews 85:301-325. doi: 10.1111/j.1469-
185X.2009.00103.x.
Stander PE, 1992. Cooperative hunting in lions: the role of the individual. Behavioral Ecology
and Sociobiology 29:445-454. doi: 10.1007/bf00170175.
Symonds MRE, Moussalli A, 2011. A brief guide to model selection, multimodel inference
and model averaging in behavioural ecology using Akaike's information criterion.
Behavioral Ecology and Sociobiology 65:13-21. doi: 10.1007/s00265-010-1037-6.
Tripet F, Nonacs P, 2004. Foraging for Work and Age-Based Polyethism: The Roles of Age
and Previous Experience on Task Choice in Ants. Ethology 110:863-877. doi:
10.1111/j.1439-0310.2004.01023.x.
Vital C, Martins EP, 2013. Socially-Central Zebrafish Influence Group Behavior More than
Those on the Social Periphery. PLOS ONE 8:e55503. doi:
10.1371/journal.pone.0055503.

Wey T, Blumstein DT, Shen W, Jordan F, 2008. Social network analysis of animal behaviour:
a promising tool for the study of sociality. Anim Behav 75:333-344. doi: DOI
10.1016/j.anbehav.2007.06.020.
Whitehouse MEA, Lubin Y, 1999. Competitive foraging in the social spider Stegodyphus
dumicola. Animal Behaviour 58:677-688. doi: 10.1006/anbe.1999.1168.
Whitehouse MEA, Lubin Y, 2005. The functions of societies and the evolution of group living:
spider societies as a test case. Biological Reviews 80:347-361. doi:
10.1017/S1464793104006694.
Wilson ADM, Krause J, 2012. Personality and metamorphosis: is behavioral variation
consistent across ontogenetic niche shifts? Behavioral Ecology 23:1316-1323. doi:
10.1093/beheco/ars123.
Witte V, Schliessmann D, Hashim R, 2010. Attack or call for help? Rapid individual decisions
in a group-hunting ant. Behavioral Ecology 21:1040-1047. doi:
10.1093/beheco/arq100.
Wittemyer G, Douglas-Hamilton I, Getz WM, 2005. The socioecology of elephants: analysis
of the processes creating multitiered social structures. Animal Behaviour 69:1357-
1371. doi: 10.1016/j.anbehav.2004.08.018.
Wright CM, Keiser CN, Pruitt JN, 2015. Personality and morphology shape task participation,
collective foraging and escape behaviour in the social spider *Stegodyphus dumicola*.
Animal Behaviour 105:47-54. doi: 10.1016/j.anbehav.2015.04.001.
Wright CM, Keiser CN, Pruitt JN, 2016. Colony personality composition alters colony-level
plasticity and magnitude of defensive behaviour in a social spider. Animal Behaviour
115:175-183. doi: 10.1016/j.anbehav.2016.03.002.
753
754

 Figures and Tables

 Fig. 1. Close-up photograph of resting spiders in the first week of the experiment. The abdomen of

 each spider is marked uniquely with acrylic paint to track their behaviour over time. The resting

 network here corresponds to one connected chain (left) and one unconnected node (right).

762
763

Latency to attack = 567s   Latency to attack = 165s   Latency to attack = 46s
Average degree = 2.6   Average degree = 3.4   Average degree = 2.6
Keystone closeness = 0.045   Keystone closeness = 0.034   Keystone closeness = 0.023

764
765
766   **Fig. 2.** Interaction networks of three sample adult spider groups, immediately before prey attack

767   speed was examined. Each node represents an individual spider and colour represents boldness –

768   redder indicates higher boldness, with the boldest individual marked as a star. Lower keystone

769   closeness and higher average degree are associated with faster latency to attack. Latencies to attack,

770   average degree, and keystone closeness are noted below each network.

771
772

773 Table 1. Network measures examined

Level of analysis	Network variable	Description
Individual	Keystone degree	Number of individuals interacting with the boldest individual
	Keystone closeness	The sum of reciprocal shortest paths from the boldest individual to all other nodes. Measures how well integrated an individual is in the overall network.
	Maximum boldness	The boldness of the boldest (keystone) individual in the group for that week (not a network measure).
Sub-group	Modularity	The extent to which the network is divided into modules, measured by the ratio between links within a module and links outward to other modules. Modules were defined based on the Optimal Community Structure algorithm (Brandes et al., 2008).
Group	Average degree	The average number of individuals that each individual interacts with. Quantifies overall network connectivity.
	Skewness of degree distribution	The skewness of the degree distribution. Larger absolute values indicate that degree is less evenly distributed among individuals in the group.

775
776
777

778 Table 2. The Akaike weights for the attack speed predictors, as calculated across all possible models
779 (see Tables S3 and S4). The top two predictors are selected for the final models, one individual and
780 one group or subgroup level.

Level of analysis	Network variable	Predictor weights	
		Adults	Subadults
Individual	Keystone degree	0.14	0.52
	Keystone closeness	**0.81**	0.15
	Maximum boldness	0.29	**0.97**
Subgroup	Modularity	0.36	0.42
Group	Average degree	**0.46**	**0.47**
	Skewness of degree distribution	0.06	0.18

789 Table 3. Statistics of the selected model for attack speed in adult spiders. *denotes significance at p
790 < 0.05. See Table S2 for information on random effects and t values.

Level of Analysis	Coefficient	Days before	Estimate log(seconds)	Standard error	p
Individual	Closeness of keystone	4	10.715	6.682	0.109
		2	12.766	6.269	**0.042 ***
		0	18.842	7.205	**0.0089 ***
Group	Average degree	4	-0.172	0.129	0.182
		2	-0.239	0.150	0.111
		0	-0.367	0.175	**0.036 ***

798 Table 4. Statistics of the selected model for attack speed in subadult spiders. *denotes significance
799 at p < 0.05. See Table S4 for information on random effects and t values.

Level of Analysis	Coefficient	Days before	Estimate log(seconds)	Standard error	p
Individual	Maximum boldness	4	-0.0050	0.0015	**< 0.001 ***
		2	-0.0049	0.0016	**0.002 ***
		0	-0.0053	0.0016	**< 0.001 ***
Group	Average degree	4	0.005	0.0247	0.836
		2	-0.041	0.0314	0.192
		0	-0.025	0.0245	0.317