Characterization of novel *Chrysosporium morrisgordonii* sp. nov., from bat white-nose syndrome (WNS) affected mines, northeastern United States

Tao Zhang[1, 2], Ping Ren[1, 4], XiaoJiang Li[1], Sudha Chaturvedi[1, 3*], and Vishnu Chaturvedi[1, 3*]

[1]Mycology Laboratory, Wadsworth Center, New York State Department of Health, Albany, New York, USA

[2] Institute of Medicinal Biotechnology, Chinese Academy of Medical Sciences & Peking Union Medical College, Beijing 100050, PR China

[3] Department of Biomedical Sciences, School of Public Health, University at Albany, Albany, New York, USA

[4]Department of Pathology, University of Texas Medical Branch, Galveston, Texas, USA

*Corresponding authors:

Sudha Chaturvedi, sudha.chaturvedi@health.ny.gov;

Vishnu Chaturvedi, vishnu.chaturvedi@health.ny.gov.

Key words: *Chrysosporium morrisgordonii*, White-nose syndrome, mine soils, taxonomy

Abstract

A new fungal species, *Chrysosporium morrisgordonii sp.* nov., is described from bat's white-nose syndrome afflicted Graphite mine in Warren county, New York. This taxon is characterized by its ability to grow at low temperature spectra from 6°C to 25°C and by the presence of tuberculate and thick-walled arthroconidia and aleurioconidia, spherical or broadly ellipsoidal, unicellular, intercalary, 3.5-4.6 μm ×3.5-4.6 μm, sessile or borne on side branches or at the ends of narrow stalks. The phylogenetic tree inferred from the analysis of the sequences of the large subunit D1/D2 and ITS1-5.8S-ITS2 regions confirm the separation of this new species from other taxa of the genus *Chrysosporium*. *Chrysosporium morrisgordonii* is distinguished by morphology and physiology characteristics from the closest phylogenetic species or taxa. It is described here as a new species in *Chrysosporium* genus.

Introduction

During a survey of mycobiome from bat's White-nose syndrome affected caves and mines in northeastern region of United States [1], we isolated one interesting psychorotolerant hyphomycete which belong to the genus *Chrysosporium*. This isolate was recovered from Graphite mine in Warren County, New York, USA.

The genus *Chrysosporium*, belonging to the subphylum Pezizomycotina, comprises a large collection of cosmopolitan saprotrophs, which are commonly recovered from soil, marine sediments, decaying wood, skin or feet of reptiles [2-5]. The genus *Chrysosporium* was first introduced by Corda in 1833. Currently, a total of seventy species have been reported[3]. *Chrysosporium* species are often keratinolytic, although occasionally cause skin infections and onychomycosis in humans and animals [2, 6-10]. The resemblance of *Chrysosporium* to dermatophytes, and their pathogenic potential, displaying similarities to *Emmonsia*, *Botryotrichum*, *Trichosporiella*, *Myceliophthora*, *Sepedonium*, *Ovadendron*, *Zymonema*, *Sporotrichum* and associated teleomorhps, is of immense diagnostic importance [11-15]. The genus is readily identified by mostly white to yellowish colonies, tree-like conidiophores structure, and terminal or intercalary conidia [16]. The conidia are usually subglobose, pyriform, or clavate and are released rhexolytically [2, 16].

In this study, based on the mentioned isolate, we propose a new species of *Chrysosporium*. A comparative study employing phenotypic, microscopic, physiologic, molecular of this taxa and other members of *Gymnoascaceae* provide further support for this proposal.

Materials and methods

Sample origin and isolation of fungus

Site description and sample collection were described previously [1]. Environmental samples including sediment or swab sample were performed according to the method described previously [1]. Culture and purification process were followed the studies performed previously [1].

Fungal strains and media preparation

A total of five strains from different reference culture collections were included in the study. Among them, three including *Neogymnomyces demonbreunii* (Ajello et Cheng) CBS 122.67 (ATCC®18028™), *Renispora flavissima* (Sigler *et al*.,) CBS 708.79 (ATCC®38504™), and *Amauroascus volatilis-patellus* (Eidam) Schroeter CBS 249.72 (ATCC®18898™) were provided by the Centraalbureau voor Schimmelcultures (CBS) Utrecht, Netherlands. *Chrysosporium speluncarum* CCF 3760 and *Neogymnomyces virgineus* JN038186 were generous gifts from Prof. Alena Novakova (Institute of Microbiology, Czech Republic) and Prof. Sabrina Sarrocco (Università di Pisa, Italy), respectively. The strains were cultured on potato dextrose agar (PDA),

sabouraud dextrose agar (SDA; peptone 10 g, dextrose 40 g, agar 20 g L^{-1}), and malt extract agar (MEA; malt extract 20 g, peptone 1.0 g, dextrose 20 g, agar 20 g L^{-1}), respectively. The basal medium (1 L volume) constituents included $(NH_4)_2SO_4$ 3.0 g, K_2HPO_4 1.0 g, KH_2PO_4 1.0 g, $MgSO_4.7H_2O$ 0.5 g, 10 mL trace elements solution [17]. All strains were generally incubated at 20°C or 30°C for 14 days.

Morphological identification

The growth rate of the isolate was determined at 6°C, 10°C, 15°C, 20°C, 25°C, 30°C on SDA and MEA. The petri dishes were inoculated in the center and the colony diameters (in millimeters) were measured daily. The strains were identified using the traditional methods based on morphological characteristics [2]. The morphological structures were measured in warmed lactophenol mounts. Photomicrographs were obtained with an Olympus Provis AX70 microscope. Scanning electron microscopy techniques were described previously by Figueras and Guarro [18]. The pure culture is deposited in the Mycology Laboratory Culture Collection Repository, Wadsworth Center, New York State Department of Health, Albany, New York, USA.

DNA extraction and PCR amplification

Genomic DNA from fungal mycelium grown on SDA plates were extracted according to the manufacturer's instructions (Epicentre, Madison, WI, USA, Cat. No. MC85200). Complete ITS1-5.8S-ITS2 (ITS) and partial 28S nrDNA

were amplified using universal primers. ITS1 and ITS4 [19] were used to amplify ITS, and NL1 and NL4 [20] were used for partial 28S (large subunit, LSU) amplification. Each PCR reaction mixture (30 μL) contained 3.0 μL 10×PCR buffer, 0.9 μL sense primer (10 μM), 0.9 μL antisense primer (10 μM), 1.0 μL dNTP (10 mM each), 2.0 μL DNA template, 0.3 μL Q5$^®$ High-Fidelity DNA Polymerase (2.0 U/ μL) (New England Biolabs, Inc.) and 21.9 μL sterile water. The PCR reactions were performed in an C1000 Touch^{TM} Thermal Cycler (Bio-Rad) with cycling conditions of an initial activation at 98°C for 30 sec, followed by 28 cycles of denaturation at 98°C for 10 sec, annealing at 55°C (ITS) and 58°C (28S) for 30 s, and elongation at 72°C for 25 sec, with a final extension step at 72°C for 2 min. PCR products were purified with the Qiaquick$^®$ PCR Purification Kit (Qiagen) and sequenced with the ABI BigDye terminator chemistry in an ABI 3730 DNA sequencer (Applied Genomic Technologies Core, Wadsworth Center). The same primers used in PCR were employed as sequencing primers. The PCR amplicons were assembled, edited using Sequencher 4.6 software software (Gene Codes Corp., Ann Arbor, MI, USA) and BLAST searched against two databases, *i.e.* GenBank (www.ncbi.nlm.nih.gov/) and CBS-KNAW (www.cbs.knaw.nl/).

Phylogenetic analysis

All sequences were aligned with Clustal X version 1.8 [21] and BioEdit v7.0.9 [22], and the multiple alignments were adjusted manually following visual inspection and the areas of sequence ambiguity were eliminated when

necessary. Databases of ITS and 28S were combined for sequence analyses. Phylogenetic analyses using the maximum parsimony analysis (MP) was performed with the MEGA 5.1 computer program [23]. The robustness of branches was assessed by bootstrap analysis with 1,000 replicates. The Dictionary of the Fungi and UniProt (http://www.uniprot.org/taxonomy/) served as the source of taxonomic references for fungal species [24].

Physiological study

Growth on dermatophyte test medium agar (DTM) and color changes from yellow (acidic) to red (basic) were recorded. Lipase activity was tested by growing on Tween 80 opacity test medium (TOTM) according to Slifkin [25]. Production of urease was determined on Christensen's urea agar after incubation for 7 d [26]. Amylase, chitinase, proteinase, cellulase, and β-glucosidase activity was verified by growing strains on induction media including basal medium with 0.5% starch, 0.5% chitin, 0.5% casein, 0.5% CM-cellulose, and 0.5% D-(+)-cellobiose, respectively. Cycloheximide tolerance was evaluated by growing strains on SDA supplemented with 0.2% and 0.5% of cycloheximide (Sigma, USA). Tolerance of NaCl was evaluated by growth of strains on SDA, amended with 3% and 5% NaCl. Hemolysis was evaluated by culturing strains on blood agar. All detections were performed at 20°C for 14 d. Temperature sensitivity tests are incubated at 6°C and 30°C, respectively.

Nucleotide Sequence Accession Numbers

All nucleotide sequences of the case isolate were deposited in GenBank under accession numbers: ITS sequence (accession number: KM273791); for 28S nrRNA gene partial sequence (accession number: KM273792).

Results

Morphology

Colonies were moderately fast growing, attaining a diameter of 38-42 mm on MEA, and 32-36 mm on PDA medium after 20-day. Optimal growth was observed at 20°C. At 25°C colonies reached 10-14 mm on MEA, and 9-12 mm on PDA in 20 days. Although no growth was observed at 30°C, initial growth was detected at 6°C after 14 days of incubation with colonies attaining diameters of 17-22 mm after four weeks. Colonies on MEA at 20 °C were white to yellowish white, powdery and dense, with a thin margin, slightly cottony at center, with reverse pale yellow to brown (Figure 1A, B). On PDA medium, colonies were yellow to sand colored, powdery, suede-like or slightly granular, raised in the center, with a poorly defined margin with reverse brown to grey in age (Figure 1C, D).

Microscopically, the isolate was distinguishable. Conidia were tuberculate and thick-walled arthroconidia (intercalary conidia) and aleurioconidia, globose to subglobose, unicellular, 3.5-4.6 μm ×3.5-4.6 μm, intercalary, sessile or borne on side branches or at the ends of narrow stalks (Figure 2A, 2B). The stalks

occur at right angles to the hypha and are straight-sided. Arthroconidia were sometimes formed by swelling and enlargement within hyphae. Hyphae septate, hyaline, smooth-walled, 1-3 µm wide, straight or branched (Figure 2C, 2D, 2E). The presence of aleurioconidia and arthroconidia were observed in all the cultures examined (Figure 2F, 2G).

Phylogeny

The ITS1-5.8S-ITS2 region sequenced from the fungal isolate included 557 base pairs. A search on GenBank database using BLAST revealed only insignificant hits from GenBank, the closest match to an identified *Chrysosporium* strain was *Chrysosporium speluncarum* CCF 3760 with 94% identity (accession no. AM949568), followed by *Gymnoascus demonbreunii* CBS 122.67 (accession no. AJ315842, 84% coverage/84% identity), *Amauroascus volatilis-patellus* CBS 249.72 (accession no. AJ390378, 73% coverage/85% identity), and *Renispora flavissima* CBS 708.79 (accession no. AF299348, 61% coverage/86% identity). None of which were considered significant for a conspecific isolate (>97% identity). The sequence of the strain *C. morrisgordonii* differed from the sequence of *C. speluncarum* CCF 3760 at 37 nucleotide positions. The ITS phylogenetic tree inferred from the analysis showed this isolate was grouped in highly supported clade with *C. speluncarum* CCF 3760 as included in the study, representing a clear genetic separation from other *Chrysosporium* spp. (Figure 3).

The D1/D2 region (partial sequence) also did not return a significant BLAST hit. The closest sequence in GenBank was *Chrysosporium speluncarum* CCF 3760 (accession no. AM949568, 94% coverage/96% identity) sequence using BLAST search, followed by *Neogymnomyces virgineus* JN038186 (accession no. JN038186), *Amauroascus volatilis-patellus* CBS 249.72 (accession no. AB075324), *Gymnoascus demonbreunii* CBS 122.67 (accession no. AY176716) all displaying 95% similarity. The phylogenetic tree inferred from the analysis of the D1/D2 sequences showed a similar topology, confirming that the isolate formed a well-supported monophyletic group with *Chrysosporium speluncarum* CCF 3760 as the closest neighbor (Figure 4).

Physiology

The esterase test was strain dependent, however the majority of the isolates were positive. All the fungi tested grew on DTM changing the color of the medium from yellow to red (data not shown), while the strain *Chrysosporium morrisgordonii* displays weak activity. The results of the other physiological tests are summarized in Table 1. The urease and glucosidase test were positive for all the isolates. All the strains in the study produced hemolysis on blood agar, showed lipolytic, proteolytic, amylatic, cellulytic, chitinolytic activity. On SDA medium with 3% and 10% NaCl, growth was scarce for *A. volatilis-patellus* and *R. flavissima*, while the ohter four isolates showed good growth (Table 1). Cycloheximide tolerance and different temperature

tolerance are listed in the Table 1, *Chrysosporium morrisgordonii* and *C. speluncarum* CCF 3760 are psychrotolerant strains, which could not grow above 30°C.

Phylogenetic analysis showed that *C. morrisgordonii* is closely related to *C. speluncarum* CCF 3760, *G. demonbreunii* CBS 122.67, *A. volatilis-patellus* CBS 249.72, *R. flavissima* CBS 708.79, and *N. virgineus* JN038186. However, *C. morrisgordonii*can is distinguished from *A. volatilis-patellus*, *R. flavissima*, and *N. virgineus* by its ability to grow at 6°C and exhibited no growth at 30°C. In addition, the *C. speluncarum* could produce much more soluble pigments in media (Figure 5).

Taxonomy

The combined morphological, molecular and physiological features of the strain isolated from the Graphite mine soil samples does not correspond to any of the species described to date within the genus *Chrysosporium* or closely related genera. Thus, we propose here as new species.

***Chrysospor*ium *morrisgordonii* Zhang, Ren, Li, Chaturvedi, Chaturvedi, sp. nov. MycoBank MB 830215**

Etymology: The epithet is derived from late Dr. Morris A. Gordon, eminent mycologist and ex-director of Mycology Laboratory, New York State Department of Health, where the isolate *Chrysosporium morrisgordonii* was characterized.

Holotypus: USA, New York, Warren County, soil of Graphite mine, January 2011, ex Morrisgordonii, Isotypus 6762-S5.

Colonies on MEA at 20°C attaining a diameter of 35-42 mm after 7 d, whitish to yellowish white, powdery and dense, with rich aerial mycelium, elevated at the center and radially folded, compact; reverse pale yellow to brown. *Hyphae* hyaline, septate, smooth-walled, straight or branched, 1-3 µm in diameter. *Conidia* (aleurioconidia) unicellular, tuberculate and thick-walled, globose to subglobose, sessile or borne on side branches or at the ends of narrow stalks, 3.5-4.6 µm ×3.5-4.6 µm; arthroconidia (intercalary conidia), formed by swelling and enlargement within hyphae, tuberculate and thick-walled, globose to subglobose, 3.5-4.6 µm ×3.5-4.6 µm.

Colonies on PDA showed features similar to those on MEA, but was more floccose and sand colored, with more sporulation. The optimum growth temperature was 20°C, and the minimum was 6°C. The fungus grew at 25°C did not grow at 30°C.

Discussion

Considering the morphological, physiological and phylogenetic species concepts, we present evidence for a novel species within *Chrysosporium*, *Chrysosporium morrisgordonii sp.* nov. *C. morrisgordonii* is distinguished by its ability to grow at low temperature range from 6°C-25°C and by the combined presence of arthroconidia and globose or subglobose aleurioconidia,

sessile or borne at the ends of narrow stalks. In the nrLSU and nrITS phylogentic trees, *Chrysosporium morrisgordonii*, *Chrysosporium speluncarum*, *Renispora flavissima*, together with two species, *Neogymnomyces virgineus* and *Gymnoascus demonbreunii* are belong to a well-defined group also known from analyses of previous studies [14, 27-30]. *C*. *speluncarum* and *R*. *flavissima* were recovered from cave samples, have affinity to bat guano, and exhibit a similar phenotype to *C. morrisgordonii* with yellowish colonies and similar tuberculate conidia [14]. The strain *C. morrisgordonii* differs from *C*. *speluncarum* by the presence of the much less pigment, powdery growth, and two different conidia formation (Figure 4). In contrast, *R*. *flavissima* differentiates from *C. morrisgordonii* for conidial size (6-8×5-8 μm) and better growth at 30°C [31, 32]. *N*. *virgineus* and *G*. *demonbreunii* have a broader ecological distribution and a *Chrysosporium* state with smooth conidia [33, 34].

There are few reports of the ability of *Chrysosporium* to adapt to low temperature, in which *C. magnasporum* and *C. oceanitesii* grow in Antarctic soil [35], *C. evolceanui* in alpine soil [36], and *C. longisporum* from snake [3], but can be easily differentiated from the new species by their morphology.

Acknowledgements

Special thanks are grateful to Profs. Alena Novakova (Institute of Microbiology, Czech Republic) and Sabrina Sarrocco (Università di Pisa, Italy) for providing the strains *C. speluncarum* CCF 3760 and *N. virgineus* JN038186. We are thankful to the Wadsworth Center Media & Tissue Culture Core for the preparation of various media and reagents and Wadsworth Center Advanced Genomics Core for sequencing of DNA samples. This study was supported financially partly with funds from the National Science Foundation (16-0039-01), as well as National Natural Science Foundation of China (No.31872617), and the Fundamental Research Funds for the Central Universities (No.3332018097).

Figure legends

Fig. 1 *Chrysosporium morrisgordonii* Zhang, Ren, Li, Chaturvedi, Chaturvedi, sp. nov. A, B) Colony on MEA agar (surface and reverse); C, D) Colony and PDA agar (surface and reverse).

Fig. 2 *Chrysosporium morrisgordonii* Zhang, Ren, Li, Chaturvedi, Chaturvedi, sp. nov. A-D) Fertile hyphae bearing sessile conidia; E) Two singly intercalary conidia and a terminal chain of arthroconidia (white arrows show the septa); F) sessile (some 2-celled) conidia. — Scale bars = 10 µm (A–F, differential interference contrast) or 1 µm (G).

Fig. 3 Neighbor-joining tree based on Kimura 2-p corrected nucleotide distances among ITS1-5.8S-ITS2 rRNA gene sequences of the species related with *Chrysosporium morrisgordonii*. Branch lengths are proportional to distance. Bootstrap replication frequencies over 60% (1,000 replications) are indicated at the nodes. The sequences were retrieved from CBS (www.cbs.knaw.nl/) or GenBank (www.ncbi.nlm.nih.gov). *Aspergillus niveoglaucus* CBS 101750 and *Aspergillus alliaceus* NRRL 20602 were used as outgroup.

Fig. 4 Neighbor-joining tree based on Kimura 2-p corrected nucleotide distances among the D1 and D2 domains of the 28S rRNA gene sequences of the species related with *Chrysosporium morrisgordonii*. Branch lengths are proportional to distance. Bootstrap replication frequencies over 60% (1,000 replications) are indicated at the nodes. The sequences were retrieved from CBS (www.cbs.knaw.nl/) or GenBank (www.ncbi.nlm.nih.gov). *Aspergillus acristata* NRRL 2394 and *Aspergillus fumigatus* CBS 286.95 were used as outgroup.

Fig. 5 Radial growth determination at temperatures ranging from 6^{o}C to 30^{o}C on Sabouraud dextrose agar (SAB). Type strains of *C. speluncarum* CCF 3760, *G. demonbreunii* CBS 122.67, *A. volatilis-patellus* CBS 249.72, *R. flavissima* CBS 708.79, and *N. virgineus* DCDSL7716.

Reference

1. Zhang T, Victor TR, Rajkumar SS, Li X, Okoniewski JC, Hicks AC, Davis AD, Broussard K, LaDeau SL, Chaturvedi S *et al*: **Mycobiome of the bat white nose syndrome affected caves and mines reveals diversity of fungi and local adaptation by the fungal pathogen Pseudogymnoascus (Geomyces) destructans**. *PLoS One* 2014, **9**(9):e108714.

2. de Hoog GS GJ, Gene J, Figueras MJ. (ed.): **Atlas of Clinical Fungi.**, 2nd ed. edn: Utrecht, The Netherlands: Centraalbureau voor Schimmelcultures. ; 2000.

3. Stchigel AM, Sutton DA, Cano-Lira JF, Cabanes FJ, Abarca L, Tintelnot K, Wickes BL, Garcia D, Guarro J: **Phylogeny of chrysosporia infecting reptiles: proposal of the new family Nannizziopsiaceae and five new species**. *Persoonia* 2013, **31**:86-100.

4. Pilar Vidal, María de los Angeles Vinuesa, José María Sánchez-Puelles, Guarro J: **Phylogeny of the anamorphic genus Chrysosporium and related taxa based on rDNA internal transcribed spacer sequences**. In: *Biology of Dermatophytes and other Keratinophilic Fungi.* edn. Edited by Kushwaha RKS GJE: Revista Iberoamericana de Micología, Bilbao, 2000 2000.

5. Abarca ML, Castella G, Martorell J, Cabanes FJ: **Chrysosporium guarroi sp. nov. a new emerging pathogen of pet green iguanas (Iguana iguana)**. *Med Mycol* 2009, **48**(2):365-372.

6. Gan GG, Kamarulzaman A, Goh KY, Ng KP, Na SL, Soo-Hoo TS: **Non-sporulating Chrysosporium: an opportunistic fungal infection in a neutropenic patient**. *Med J Malaysia* 2002, **57**(1):118-122.

7. Anstead GM, Sutton DA, Graybill JR: **Adiaspiromycosis causing respiratory failure and a review of human infections due to Emmonsia and Chrysosporium spp**. *J Clin Microbiol* 2012, **50**(4):1346-1354.

8. Shekhar H, Gogia V, Vashisht N, Gupta S, Kakkar A, Venkatesh P: **Endogenous endophthalmitis by chrysosporium: an opportunistic pathogen**. *Ocul Immunol Inflamm* 2014, **22**(2):158-160.

9. Abarca ML, Martorell J, Castella G, Ramis A, Cabanes FJ: **Cutaneous hyalohyphomycosis caused by a Chrysosporium species related to Nannizziopsis vriesii in two green iguanas (Iguana iguana)**. *Med Mycol* 2008, **46**(4):349-354.

10. Cabanes FJ, Sutton DA, Guarro J: **Chrysosporium-related fungi and reptiles: a fatal attraction**. *PLoS Pathog* 2014, **10**(10):e1004367.

11. Rajeev S, Sutton DA, Wickes BL, Miller DL, Giri D, Van Meter M, Thompson EH, Rinaldi MG, Romanelli AM, Cano JF *et al*: **Isolation and characterization of a new fungal species, Chrysosporium ophiodiicola, from a mycotic granuloma of a black rat snake (Elaphe obsoleta obsoleta)**. *J Clin Microbiol* 2009, **47**(4):1264-1268.

12. Allender MC, Dreslik M, Wylie S, Phillips C, Wylie DB, Maddox C, Delaney MA, Kinsel MJ: **Chrysosporium sp. infection in eastern massasauga rattlesnakes**. *Emerg Infect Dis* 2011, **17**(12):2383-2384.

13. Sigler L, Hambleton S, Pare JA: **Molecular characterization of reptile pathogens currently known as members of the chrysosporium anamorph of Nannizziopsis vriesii complex and relationship with some human-associated isolates**. *J Clin Microbiol* 2013, **51**(10):3338-3357.

14. Nováková A, Kolařík M: **Chrysosporium speluncarum, a new species resembling Ajellomyces capsulatus, obtained from bat guano in caves of temperate Europe**. *Mycological Progress* 2010, **9**(2):253-260.

15. Kamath PM, Prasad V, Shenoy VS, Mukundan A, Shenoy S: **Chrysosporium: an uncommon fungus in chronic rhinosinusitis**. *J Clin Diagn Res* 2015, **9**(3):MD01-02.

16. Guy St Germain , Summerbell R: **Identifying Filamenytous Fungi: A Clinical Laboratory Handbook**: Star Pub Co; 1 edition; 1996.

17. Zhang T, Zhuo Y, Jia X, Liu J, Gao H, Song F, Liu M, Zhang L: **Cloning and characterization of the gene cluster required for beauvericin biosynthesis in Fusarium proliferatum**. *Sci China Life Sci* 2013, **56**(7):628-637.

18. Figueras MJ, Guarro JA: **scanning electron microscopic study of ascoma development in Chaetomium malaysiense.** *Mycologia* 1988, **80**:298-306.

19. White TJ, T. D. Bruns, S. Lee, and J. W. Taylor: **Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics.** In: *PCR protocols: a guide to methods and applications.* edn. Edited by M. A. Innis DHG, J. J. Sninsky, and T. J. White (ed.),. New York, N.Y.: Academic Press Inc.,; 1990: 315–324.

20. O'Donnell K. In: *The Fungal holomorph: mitotic, meiotic and pleomorphic speciation in fungal systematics : proceedings of an international symposium.* edn. Edited by Taylor. DRRJW. Wallingford (UK): CAB International; 1993: 225-233.

21. Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG: **The CLUSTAL_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools**. *Nucleic Acids Res* 1997, **25**(24):4876-4882.

22. Hall TA: **BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT.** *Nucleic Acids Symposium Series* 1999, **41**(41):95-98.

23. Tamura K PD, Peterson N, Stecher G, Nei M, and Kumar S.: **MEGA5: Molecular Evolutionary Genetics Analysis using Maximum Likelihood, Evolutionary Distance, and Maximum Parsimony Methods.** *Molecular Biology and Evolution* 2011, **28**(10):2731-2739.

24. Kirk PM CP, Minter DW, Stalpers JA: **Ainsworth & Bisby's Dictionary of the Fungi.**, 10th Edition. edn: Wallingford: CAB International. UK; 2008.

25. Slifkin M: **Tween 80 opacity test responses of various Candida species**. *J Clin Microbiol* 2000, **38**(12):4626-4628.

26. Zhang T, Ren P, Chaturvedi V, Chaturvedi S: **Development of an Agrobacterium-mediated transformation system for the cold-adapted fungi Pseudogymnoascus destructans and P. pannorum**. *Fungal Genet Biol* 2015, **81**:73-81.

27. Doveri F, Pecchia S, Vergara M, Sarrocco S, Vannacci G: **A comparative study of Neogymnomyces virgineus , a new keratinolytic species from dung, and its relationships with the Onygenales**. *Fungal Diversity* 2012, **52**(1):13-34

28. Untereiner WA, Scott JA, Naveau FA, Currah RS, Bachewich J: **Phylogeny of Ajellomyces, Polytolypa and Spiromastix (Onygenaceae) inferred from rDNA sequence and non-molecular data**. *Studies in Mycology* 2002, **45**(47):25-35.

29. Untereiner WA, Scott JA, Naveau FA, Sigler L, Bachewich J, Angus A: **The Ajellomycetaceae, a new family of vertebrate-associated Onygenales**. *Mycologia* 2004, **96**(4):812-821

30. Vidal P, Vinuesa MA, Sanchez-Puelles JA, Guarro J: **Phylogeny of the anamorphic genus Chrysosporium and related taxa based on rDNA internal transcribed spacer sequences. In: Kushwaha RKS, Guarro J (eds),**

Biology of dermatophytes and other keratinophilic fungi: Revista Iberoamericana de Micología, Bilbao, Spain.; 2000.

31.   Garrison RG, Mirikitani FK: **Fine structural studies of the Chrysosporium state of Renispora flavissima**. *Mycopathologia* 1983, **81**(1):3-7.

32.   Sigler L, Gaur PK, Lichtwardt RW, Carmichael JW: **Renispora flavissima, a new gymnoascaceous fungus with tuberculate Chrysosporium conidia** *Mycotaxon* 1979, **10**(1):133-141

33.   Orr GF: **Neogymnomyces, a New Genus of Gymnoascaceae**. *Canadian Journal of Botany* 1970, **48**(6):1061-+.

34.   Smith CD, Furcolow ML: **Fifteen isolations of Gymnoascus demonbreunii from canine and soil sources**. *Sabouraudia* 1969, **7**(2):142.

35.   Stchigel A, Cano JF, Guarro J, Cormack WPM, Archubi DI: **Antarctic Chrysosporia: Chrysosporium magnasporum Stchigel, Cano, Mac Cormack & Guarro, sp nov. and Chrysosporium oceanitesii Stchigel, Cano, Archubi & Guarro, sp nov** *Persoonia - Molecular Phylogeny and Evolution of Fungi* 2013(31):268-269.

36.   Otcenasek M, Hubalek Z, Dvorak J, Kunert J: **Chrysosporium indicum-like fungi and Chrysosporium evolceanui from the soil of Cuba**. *Mykosen* 1968, **11**(1):19-24.

Table 1 Key physiological features of fungi included in this study

Fungi / Physiology	*Chrysosporium morrisgordonii*	*C. speluncarum* CCF 3760	*G. demonbreunii* CBS 427.70	*N. virgineus* DCDSL7716	*A. volatilis-patellus* CBS 249.72	*R. flavissima* CBS 708.79
casein agar	++	-	++	++	+++	+
BHI agar	+++	+++	+++	+++	++	+++
lipase	++	+	++	++	+++	+
ureae	+++	+++	++	+++	+++	+
chitinase	++	++	+	+++	+++	+
cellulase	++	+++	+++	++	++	+
β-Glucosidase	++	+	++	++	+	+
DTM agar	R/B, +	R/B, +++	R/B, ++	R/B, ++	R/B, +++	R/B, ++
SDA+3% NaCl	+++	+++	+++	+++	+	+
SDA+10% NaCl	+	+	+	+	-	-
SDA+0.2% Cyc	+++	++	++	+++	++	++
SDA+0.5% Cyc	+++	++	+	++	+	++
SDA at 6°C	++	++	-	-	-	-
SDA at 30°C	-	-	+++	+++	+++	++

Note: *C*, *Chrysosporium*; *N*, *Neogymnomyces*; *G*, *Gymnoascus*; *A*, *Amauroascus*; *R*, *Renispora*. BHI agar, Brain Heart Infusion agar; Cyc, cycloheximide; SDA, sabouraud dextrose agar; +++, positve or strong activity; +, scarse positive, moderate activity; -, absence or no activity. Y/A, yellow (acidic); R/B, red (basic).

Figure 1

MEA

PDA

Figure 2

Figure 3

Figure 4

Figure 5