Human influences on antipredator behaviour in Darwin's finches

Kiyoko M. Gotanda[a]*

[a] Department of Zoology, University of Cambridge, Cambridge, United Kingdom CB2 3EJ

Telephone: +44 (0) 1223-767130, Fax: +44(0) 1223 336676, Email: kg419@cam.ac.uk

* Corresponding author

ORCID ID: https://orcid.org/0000-0002-3666-0700

Abstract

1) Humans exert dramatic influences upon the environment, creating novel selective pressures to which organisms must adapt. On the Galapagos, humans have established a permanent presence and have altered selective pressures through influences such as invasive predators and urbanization, affecting iconic species such as Darwin's finches.

2) Here, I ask two key questions: (i) does antipredator behaviour (e.g. FID) change depending on whether invasive predators are historically absent, present, or eradicated? and (ii) to what degree does urbanization affect antipredator behaviour? This study is one of the first to quantify antipredator behaviour in endemic species *after* the eradication of invasive predators. This will help to understand the consequences of invasive predator eradication and inform conservation measures.

3) I quantified flight initiation distance (FID), an antipredator behaviour, in Darwin's finches, across multiple islands in the Galapagos that varied in the presence, absence, or successful eradication of invasive predators. On islands with human populations, I quantified FID in urban and non-urban populations of finches.

4) FID was higher on islands with invasive predators compared to islands with no predators. On islands from which invasive predators were eradicated ~11 years previously, FID was also higher than on islands with no invasive predators. Within islands that had both urban and non-urban populations of finches, FID was lower in urban finch populations, but only above a threshold human population size. FID in larger urban areas on islands with invasive predators was similar to or *lower* than FID on islands with no history of invasive predators.

5) Overall, these results suggest that invasive predators can have a lasting effect on antipredator behaviour, even after eradication. Furthermore, the effect of urbanization can strongly oppose the

effect of invasive predators, reducing antipredator behaviour to levels lower than found on

pristine islands with no human influences. These results improve our understanding of human

influences on antipredator behaviour which can help inform future conservation and

management efforts on islands.

Keywords: Galapagos, invasive predators, flight initiation distance, urbanization, post-

eradication

Introduction

"All of [the terrestrial birds] are often approached sufficiently near to be killed with a

switch, and sometimes, as I myself tried, with a cap or a hat." – Charles Darwin in "The Voyage

of the Beagle"

Human influences such as invasive species and urbanization can strongly affect the process of

local adaptation (Alberti et al., 2017; Johnson & Munshi-South, 2017; Low, 2002; Vitousek,

Mooney, Lubchenco, & Melillo, 1997). Such effects are amplified on islands such as the

Galapagos Islands, where small population sizes and strong isolation increase the vulnerability of

local flora and fauna to human influences, often resulting in loss of island biodiversity through

extinctions (Blackburn, Cassey, Duncan, Evans, & Gaston, 2004; Fritts & Rodda, 1998;

MacArthur & Wilson, 1967; Sax & Gaines, 2008). Among the endemic species on the Galapagos

Islands are Darwin's finches, an iconic example of an adaptive radiation in which a single

founding species has evolved into several species, each with different adaptions (e.g. beak shapes

and body sizes) to exploit different ecological niches (Grant & Grant, 2008, 2014). Humans

began establishing settlements on the Galapagos in the early 19[th] century (Fitter, Fitter, &

Hosking, 2016), and since then, human influences such as invasive predators and urbanization

have affected several islands on the Galapagos. Many organisms initially respond to such human

influences through behavioural adaptations. Here, I consider how two human influences –

invasive predators and urbanization – might alter antipredator behaviour in Darwin's finches on

the Galapagos Islands.

Invasive predators have strong ecological and evolutionary effects (Clavero, Brotons,

Pons, & Sol, 2009; Lever, 1994; Low, 2002; Vitousek et al., 1997), and this impact is known to

be correlated with local extinction events (Clavero & García-Berthou, 2005; McKinney &

Lockwood, 1999). On islands, the lack of predators and correlated relaxed selection can result in

reduced antipredator behaviour (Beauchamp, 2004; Blumstein, Daniel, & Springett, 2004;

Blumstein, 2002; Jolly, Webb, & Phillips, 2018). This evolutionary naïveté of isolated animals

that have evolved without major predators can contribute to the extirpation of island species

(Blumstein et al., 2004; Blumstein, 2002; Blumstein & Daniel, 2005; Cooper, Pyron, & Garland,

2014). In particular, feral and domestic house cats (*Felis silvestris catus*) are of concern for

island biodiversity because cats target small animals such as birds and reptiles (Blancher, 2013;

Loss, Will, & Marra, 2013; Woinarski et al., 2018; Woods, Mcdonald, & Har Ris, 2003), and

invasive house cats now exist on four islands of the Galapagos (Phillips, Wiedenfeld, & Snell,

2012), presenting a critical threat for Galapagos biodiversity (Stone, Snell, & Snell, 1994;

Wiedenfeld & Jiménez-Uzcátegui, 2008). Past research on the effects of invasive predators in the

Galapagos has focused on behavioural adaptations in reptiles (e.g. Berger, Wikelski, Romero,

Kalko, & Roedl, 2007; Stone et al., 1994), and thus, little is known about the effect of novel

mammalian predators on endemic land birds. Given the resulting selective pressures, natural

selection should favour an increase in antipredator behaviour after the introduction of an invasive

predator to reduce mortality.

Effective conservation management, especially on islands, often involve eradication of

invasive predators to protect the local and endemic species (Jones et al., 2016; Nogales et al.,

2013). Post-eradication research typically follows local and endemic species population recovery

(Côté & Sutherland, 1997; Hughes, Martin, & Reynolds, 2008; Lavers, Wilcox, & Donlan,

2010), monitors the re-introduction of extirpated species to previously abandoned breeding

grounds (Campbell et al., 2011), or focus on major ecological effects such as changes in food

web dynamics (Hughes et al., 2008; Zavaleta, Hobbs, & Mooney, 2001). All this research

contributes to the growing need to understand post-eradication effects (Côté & Sutherland, 1997;

Lavers et al., 2010; Phillips, 2010), yet surprisingly little research has focused on post-

eradication behavioural adaptations, nor how quickly such behavioural adaptations might occur.

Post-eradication behavioural adaptations could have population-level consequences on fitness.

For example, increased antipredator behaviour can have associated costs due to the reallocation

of energy and time away from other important behaviours such as foraging, reproduction, and

rearing of young (Cooper & Frederick, 2007; Cooper & Blumstein, 2015; Ydenberg & Dill,

1986). Thus, if increased antipredator behaviour is maintained after eradication, this might result

in a decrease in fitness for local and endemic species. Understanding how local and endemic

species will behaviourally adapt post-eradication could help improve conservation efforts. On the

Galapagos, some islands have invasive house cats, some have remained free of invasive

predators, and some islands have successfully eradicated invasive predators. This allows for

among-island comparisons of antipredator behaviour in relation to the current and historical

invasive-predator regime.

Urbanization has rapidly increased in the past century, with more than half the world's

population occupying urban settlements, severely altering patterns of selection and adaptation

(Alberti et al., 2017; Hendry, Gotanda, & Svensson, 2017; Johnson & Munshi-South, 2017). In

general, animals such as birds show decreased antipredator behaviour in urban areas compared to

rural areas, likely due to habituation to humans (Díaz et al., 2013; Møller, 2009; Møller &

Tryjanowski, 2014; Samia et al., 2017). However, such a reduction in antipredator behaviour in

urban areas could make organisms more vulnerable to different threats, such as invasive

predators. Quantifying the degree to which urbanization can reduce antipredator behaviours can

inform our understanding of the impacts of urbanization. Can urbanization reduce antipredator

behaviour to levels *before* the introduction of predators?

The Galapagos Islands represent an excellent opportunity to study the effects on invasive

predators and urbanization for two key reasons that few, if any, other systems offer. First, few

archipelagos in the world have islands that vary not only in the presence or absence of invasive

predators, but also have islands that have successfully eradicated invasive predators. Second,

islands differ not only in the presence or absence of urban centres, but also in the size of the

urban centres, representing a novel opportunity to compare antipredator behaviour along a

gradient of urbanization as well as among islands that differ in the presence or absence of urban

centres. The isolation of the Galapagos Islands removes potentially confounding factors such as

high gene flow or continued influxes of introduced predators, allowing me to ask two key

questions regarding human influences and antipredator behaviour. First, I ask how will

antipredator behaviour change depending on whether invasive predators are present, absent, or

perhaps more importantly, eradicated from an island? Very little research has been done on

behavioural adaptations in endemic species post invasive predator eradication. Second, I ask how

much can urbanization reduce antipredator behaviour – can it be reduced to levels found on

islands with no history of invasive predators? Together, these two questions can inform how

human influences are affecting antipredator behaviour on isolated islands.

Materials and methods

Site descriptions

The Galapagos Islands are a volcanic archipelago located ~1,000 km off the coast of Ecuador.

Local or endemic predators such as owls or snakes are found on all islands (Supplemental Table

1; Swash & Still 2005). Snakes are thought to prey on the nestlings of ground finches and are

thus an unlikely predator. However, short-eared owls, found on all islands surveyed in this study,

are known predators of adult land birds such as Darwin's finches (Groot, 1982). Galapagos

Hawks, also predators of ground finches (Vries, 1976, 2015), are found on four (Santa Fe,

Española, Isabela, and Santa Cruz) of the eight islands surveyed in this study (Swash & Still,

2005; Vries, 1989). Unfortunately, little data are available about the current densities of local and

endemic predators on these islands; however, the ecology of the local and endemic predators

(e.g. owls) is well documented and can thus be assumed to be predators of finches. The invasive-

predator regime (presence of house cats) on the islands (Supplemental Table 1) were classified

as: present (Floreana, Isabela, San Cristobal, Santa Cruz), pristine (Santa Fe, Española), or

successful eradication (Baltra, North Seymour). House cats and rodents were successfully

eradicated from Baltra in 2003 (R. B. Phillips et al., 2005; R.Brand Phillips et al., 2012), and rats

from North Seymour in 2008 (Carrión, Sevilla, & Tapia, 2008; Harper & Carrión, 2011); I will

refer to these as "eradicated" below for brevity. The two pristine islands and two eradicated

islands have no permanent human populations. On the four islands with human populations

(Floreana, Isabela, San Cristobal, Santa Cruz), site urbanization categories were classified as:

urban (in town) or non-urban (remote areas several kilometres away from town and not visited

by tourists). No islands with permanent human populations and no presence of invasive predators

nor pristine islands with permanent human populations exist in the Galapagos archipelago, and

thus, I am restricted to the among island comparisons outlined above.

Data collection

Flight initiation distance (FID), the distance at which a prey flees an approaching

predator, is a metric used to quantify antipredator behaviour (Cooper & Frederick, 2007; Cooper

& Blumstein, 2015; Ydenberg & Dill, 1986). An individual's decision to flee is influenced by the

perceived costs and benefits of remaining or taking flight, which means FID can be an indicator

of how an organism assesses risk, and thus, antipredator behaviour. Data were collected from

2015 to 2018, generally between February and April (some data on San Cristobal were collected

in November 2017), on eight islands of the Galapagos archipelago: four islands with the

presence of invasive predators, two islands with no history of invasive predators, and two island

that have successfully eradicated invasive predators (Supplemental Figure 1). Due to finch

species distributions varying across islands as well as Darwin's finch's species being closely

related, data from the only species that occurs on all islands, the small ground finch, *Geospiza*

fuliginosa, are presented.

FID measurements were performed with a human stimulus following methods from

Blumstein (2006). A focal finch was located by walking and searching the landscape at a slow

walking pace, and the finch's initial behaviour was noted. To minimize the possibility of

pseudoreplication, in a given day, each trial ensured the focal finch was of a different sex or age

class (for males) than finches that had previously been approached. However, it is possible the

same bird might have been approached on different days or years because the finches were not

individually banded. Birds were located in areas that had relatively open habitat to ensure a

straight approach by the human and a clear sightline from the human to the finch. The human

would then approach the focal finch at a standardized speed (~0.85 m/s). Human stimuli always

wore neutral-coloured clothing and looked at the focal individual while approaching.

Flight was considered to have been initiated if the finch extended its wings and flew; the

distance flown could be short (<0.5 m) or substantial (out of sight). Finches that hopped away

instead of taking flight were omitted from the study, though this was an extremely rare

occurrence (two or three occurrences across all sampling periods). A marker was placed where

the finch originally was and where the stimulus was when the finch took flight, and the distance

between these markers was the FID. Because of the complexity of the landscape, the distance at

which the stimulus started from could not be standardized (Blumstein, 2003, 2006; Samia,

Nomura, & Blumstein, 2013), and so I noted the distance from where the human started to the

flight-initiation marker (starting distance). Alert distance, the distance at which an individual is

aware of the approaching stimulus (Cooper, Samia, & Blumstein, 2015), could not be quantified

because the focal individual was often foraging on the ground and would repeatedly raise its

head, and so normal foraging behaviour was indistinguishable from an alert reaction to an

approaching human.

Each data point collected included the island, invasive-predator regime, urbanization

category (on islands with permanent human populations), sex, time of day, and group size

(defined as at least one other finch within a 1m radius of the focal finch and if so, how many

other finches were within 1m of each other). A minimum of six finches were sampled at each site

(Supplementary Table 1). Finch sex was identified by plumage, and for ambiguous cases sex was

denoted as unknown. Time of day was noted because birds are most active at dawn and at dusk,

so baseline activity levels and behaviours can vary throughout the day. Group size was noted

because it could increase FID because larger groups mean more observers and thus, detection of

a potential threat will occur when the threat is still a longer distance away (Ydenberg & Dill,

1986). Conversely, group size could decrease FID through the dilution effect where the

probability a predator will target a specific individual decreases as group size increases (Cooper

& Blumstein, 2015).

Statistical analysis

All analyses were done in R (version 3.4.3). To meet assumptions of normal distributions, FID

and starting distance (the distance between the focal finch and the stimulus starting position)

were log-transformed. Then, FID and starting distance were centered by subtracting the mean

from each value and then dividing by the standard deviation. Because starting distance could not

be standardized and is known to affect FID, it was included in the analyses, and group size and

time of day were included as covariates. Lastly, sex was included as a fixed factor and island was

included as a random factor as appropriate. Further analysis details are below, but in general,

linear mixed models were performed using lmer() from the lme4 package (Bates, Maechler,

Bolker, & Walker, 2015) and Anova() from the car package (Fox & Weisberg, 2011). Random-

factor significance was determined with ranova(), and post-hoc pairwise comparisons for fixed

factors in the linear mixed models used difflsmeans(), both from the lmerTest package

(Kuznetsova, Brockhoff, & Christensen, 2017). R^2 values for mixed models were calculated with

the r2beta() from the r2glmm package (version 0.1.2) and η^2 values for linear models were

calculated with etaSquared() from the lsr package (version 0.5).

Does antipredator vary depending on whether invasive predators are historically absent,

present, or have been eradicated?

On islands that had both urban and non-urban populations, only data from non-urban sites

were used for this analysis. For this analysis, the number of finches sampled on each island

ranged from 9 finches to 29 finches. A linear mixed model was first performed with invasive-

predator regime, sex, and starting distance (and their interactions) as fixed factors, island a as

random factor, and time of day and group size as covariates. None of the two- or three-way

interactions were significant ($0.634 < P < 0.924$) and the best-fit model had the interactions

removed. Post-hoc comparisons focused on pairwise comparisons between invasive-predator

regimes.

To determine whether FID varied among islands within a given invasive-predator regime,

I ran a linear model with island, sex, and starting distance (and their interactions) as fixed factors

and time of day and group size as covariates. The only significant interaction was between island

and starting distance. The best-fit model had the non-significant interactions removed. Post-hoc

comparisons were focused on pairwise comparisons between islands within a given invasive-

predator regime. A Tukey's HSD (honestly significant difference) test was used for post-hoc

comparisons for this analysis.

How much can urbanization affect antipredator behaviour?

Only data collected from the four islands with permanent populations were used for this analysis.

For this analysis, the number of finches sampled in each population ranged from 6 finches to 29

finches. A linear mixed model was performed with site urbanization, sex, and starting distance

(and their interactions) as fixed factors, island as a random factor, and time of day and group size

as covariates. None of the two- or three-way interactions were significant ($0.244 < P < 0.560$)

and model with no interactions was a better fit and is reported here. To look at within island

differences in FID among urban and non-urban finches, for each island, a linear model was

performed with the same factors as above without island as a random factor.

Results

Does antipredator vary depending on whether invasive predators are historically absent,

present, or have been eradicated?

On islands with invasive predators and islands where predators have been eradicated, FID

in finches was significantly higher than on pristine islands (Table 1, Figure 1). Post-hoc

comparisons showed that finches on pristine islands (Santa Fe and Española) had lower FID

when compared to finches on islands with invasive predators (Figure 1; $p = 0.009$) and

eradicated islands (Figure 1; $p = 0.005$), and finches on eradicated islands did not differ in FID

when compared to finches on islands with invasive predators (Figure 1, $p = 0.353$). When

considering the effect of island independently of invasive-predator regime, island had a

significant effect on FID (Figure 1; Table 2). A significant interaction between island and

starting distance was found (Supplemental Figure 2; Table 2). Post-hoc comparisons of FID

within an invasive-predator regime showed no significant difference in FID between finches on

the two pristine islands ($p = 1.000$), among the four islands with invasive predators ($0.458 < p <$

1.000), or between the two eradicated islands ($p = 0.994$).

How much can urbanization affect antipredator behaviour?

Urban finches had significantly lower FID as compared to non-urban finches (Table 3, Figs. 1

and 2). A post-hoc linear model analysing FID in finches on Floreana showed no significant

difference between urban and non-urban populations ($F = 0.158$, $p = 0.514$) whereas FID in

finches was significantly lower in urban areas as compared to non-urban areas on Isabela ($F =$

9.005, $p = 0.011$), San Cristobal ($F = 28.000$, $p < 0.001$), and Santa Cruz ($F = 7.555$, $p = 0.008$).

On San Cristobal and Santa Cruz, finches in urban populations had lower FID than found on

pristine islands (Figure 1). Group size was positively correlated with FID (Table 3, Supplemental

Figure 3) for both urban and non-urban finches.

Discussion

Increased antipredator behaviour is maintained after eradication of invasive predators

Finches exhibited increased antipredator behaviour on islands with invasive predators (Table 1; Figure 1). More interestingly, this increased antipredator behaviour was also observed on islands where invasive predators had been eradicated (in 2003 and 2008, 13 and 8 years prior to data collection; the mean generation time of finches is one year, by comparison). Since all islands have naturally occurring local or endemic predators of finches (Supplemental Table 1), this observation was most likely due to the presence of invasive predators, even after eradication. This is one of the first studies to show that increased antipredator behaviour has been maintained on islands that have had invasive predators removed. Several possible reasons exist for these observations, especially the apparent maintenance of elevated FID on eradicated islands. First, it is possible increased antipredator behaviour has evolved on islands that have and used to have invasive predators. However, without knowledge of heritability, and thus actual evolution, this cannot be confirmed, but would be an area for future research. Second, perhaps the expected costs of increased antipredator behaviour are not high enough to cause a reversion to pre-predator levels. Increased FID can have associated costs (Cooper & Frederick, 2007; Cooper & Blumstein, 2015; Ydenberg & Dill, 1986), suggesting that if this behavioural adaptation were costly then finches on eradicated islands would have FID comparable to finches on pristine islands. It could also be that not enough time has elapsed for reversion in antipredator behaviour. Third, the multi-predator hypothesis (Blumstein, 2006) suggests that increased antipredator behaviour can be maintained by the presence of remaining local and endemic predators. However, the lower antipredator behaviour on islands with no history of invasive predators

suggest that it is the invasive predators that have caused the increase in antipredator behaviour,

and a different mechanism is maintaining the increased antipredator behaviour. Fourth, cultural

transmission of increased FID (Aplin et al., 2015; Aplin, 2018) with learned behaviour

transmitted from generation to generation could maintain the increased FID. Lastly, the increase

in FID could be due to something other than predation (e.g. parasitism, life history, or habitat;

Cooper & Blumstein, 2015), which could still be present on eradicated islands, or that for

unknown historical reasons, antipredator behaviour on the eradicated islands have historically

been high. The last reason is possible, but it would be a quite a coincidence if the eradicated

islands that have this elevated FID for some reason unrelated to predation just happen to be

exactly the islands where predation was introduced and then eradicated.

Regardless of the mechanism, the fact that antipredator behaviour levels did not revert

post-eradication (when comparing FID on eradicated islands to FID on pristine islands; Figure 1)

has potential consequences for evaluating the efficacy of eradication efforts. Recent studies of

local animal populations post-eradication have focused on demographic parameters such as

population recovery (Côté & Sutherland, 1997; Hughes et al., 2008; Lavers et al., 2010) or on

ecological parameters such as food-web dynamics (Hughes et al., 2008; Zavaleta et al., 2001).

However, such phenomena will be influenced by behavioural shifts. For example, increased

antipredator behaviour correlates with decreased time and energy for behaviours such as

foraging, courting, defending territories, or caring for offspring (Cooper & Frederick, 2007;

Cooper & Blumstein, 2015; Ydenberg & Dill, 1986), which could affect population recovery

and/or food-web dynamics. Thus, understanding how the eradication of invasive predators will

affect the behaviour of local or endemic animals should be central to future conservation efforts

(Côté & Sutherland, 1997; Lavers et al., 2010; R. A. Phillips, 2010).

Urbanization can decrease antipredator behaviour to levels lower than before the introduction

of predators

Finches in urban areas had lower FID than finches in non-urban areas, supporting previous

findings (Díaz et al., 2013; Møller, 2009; Møller & Tryjanowski, 2014; Samia et al., 2017).

However, two interesting points are found in this general trend. First, the *degree* of urbanization

appears to determine just how much lower FID is for urban finches when compared to non-urban

finches. Finches in the town of Puerto Velasco Ibarra on Floreana had the highest FID compared

to finches in other towns, and Puerto Velasco Ibarra is also the smallest town, with a permanent

population of only 111 (Supplemental Table 1). The significantly lower FID of finches in larger

towns as compared to non-urban finches suggests an urbanization "threshold", such that the

degree of urbanization needs to be high enough to exert sufficient selective pressure on finches

to drive behavioural adaptation; in short, perhaps Puerto Velasco Ibarra is simply too small to be

an "urban" site for the purposes of finch behavioural adaptation. The next largest town, Puerto

Villamil on Isabela, with a population of 2,164 (Supplemental Table 1), had significantly lower

FID than Puerto Velasco Ibarra on Floreana. Puerto Villamil is still a relatively small town

(Supplemental Table 1), showing that the threshold amount of urbanization sufficient to produce

differences in antipredator behaviour is not very high.

The second interesting point is that on some islands, urbanization can result in FID that is

lower than FID on islands that have never been exposed to predators (Figure 1), even though

urban areas invariably contain invasive predators such as cats and rats. In other words, in some

towns, such as Puerto Baquizo Moreno on San Cristobal and Puerto Ayora on Santa Cruz, FID

has been reduced to levels below what was observed on islands with no history of invasive

predators. Such reductions in FID in urban finches is likely due initially to habituation

(Blumstein, Fernandez-Juricic, Zollner, & Garity, 2005; Cavalli, Baladrón, Isacch, Biondi, & Bó,

2018; Møller, 2008; Samia, Nakagawa, Nomura, Rangel, & Blumstein, 2016) which can lead to

lower FID being an evolved adaptation (van Dongen, Robinson, Weston, Mulder, & Guay, 2015)

This suggests pressures from urbanization on antipredator behaviour can be so strong that it

results in FID lower than the baseline FID quantified on pristine islands, counteracting any

increase in FID due to the presence of invasive predators. For both this and the previous point,

more research needs to be conducted on more systems with town of varying size given the

relatively small number of islands and populations in this study. Altogether, this suggests that the

effects of urbanization on organisms can be quite strong, with likely evolutionary and ecological

consequences (Alberti, 2015; Alberti et al., 2017; Johnson & Munshi-South, 2017).

Group size and starting distance

FID significantly increased with increasing finch group size for urban and non-urban

populations on islands with permanent human populations (Supplemental Figure 3). This

supports the "many-eyes hypothesis" that detection of predators occurs earlier in large groups,

when the predator is further away, due to the larger number of individuals watching (Ydenberg

& Dill, 1986). However, an effect of group size on FID in relation to invasive predation regime

was not found, indicating that on islands with no history of or having successfully eradicated

invasive predators, group size does not affect FID in individual finches. Perhaps the vigilance

achieved with "many-eyes" is necessary on islands with predators, but not in the absence

(historically or eradicated) of invasive predators.

Interestingly, starting distance was not significantly positively correlated with FID as

expected based on previous research (Blumstein, 2003; Samia et al., 2013). In fact, on one

island, San Cristobal, the relationship was negative with longer starting distances resulting in

shorter FIDs (Supplementary Figure 2), and this relationship could be explored with further

research. However, the effect of starting distance on FID is highly variable among species and a

some species do not have a positive correlation between starting distance and FID (Blumstein,

2003). Thus. it is possible that Darwin's finches are one of the very few species where starting

distance has a very low impact on FID and that is why a positive correlation between starting

distance and FID was not detected in this study.

Conclusions

The Galapagos Islands represent an opportune system to study the effects of human influences

due to the among island differences in invasive-predator regime as well as the differences in the

amount of urbanization. Such systems do not readily exist elsewhere. Here, I showed how

antipredator behaviour increased in response to invasive predators and was maintained even after

eradication of those invasive predators, and this can have possible demographic and ecological

effects. This is one of the first studies to look at post-eradication behaviours in local and endemic

species, which can have implications for future conservation efforts. I also found that

urbanization can reduce antipredator behaviour to levels at or below what was found on pristine

islands, attesting to the strength of the effect urbanization can have on behavioural traits.

Understanding the effects of different human influences will help us predict how organisms

might respond to their rapidly changing environments.

Acknowledgements

I wish to thank Marc-Olivier Beausoleil, Lara Bernasconi, Carlos Camacho, Sofia Carvajal,

Angela Hansen, Célina Leuba, and Ashley Saulsberry for their assistance in the field. Permits

were obtained with the assistance of Sofia Carvajal, Jaime Chaves, Jeff Podos, the Charles

Darwin Research Centre, the Galápagos Science Centre, the Galápagos National Park, and the

Universidad de San Francisco (Permits Nos PC-25-15, PC-42-16, PC-03-18). I thank Sarah

Knutie for numerous helpful suggestions on ideas and methods, Ben Haller, Andrew Hendry, and

Claire Spottiswoode for extensive comments and friendly reviews of the manuscript, and Tom

Houslay for guidance with statistics. I thank Dan Blumstein and an anonymous reviewer for

comments that greatly improved the manuscript. My research has been generously supported by

the Natural Sciences and Engineering Research Council of Canada (Banting Postdoctoral

Fellowship), Le Fonds Québécois de la Recherche sur la Nature et les Technologies

(postdoctoral research fellowship), Clare Hall (Whitehead Fund), Christ's College (Galápagos

Islands Visiting Scholarship Scheme), Newnham College (Phyllis and Eileen Gibbs Travelling

Research Fellowship), the British Ornithologists' Union, and McGill University (Department of

Biology Vineberg Fellowship). I have no conflicts to declare. Lastly, many thanks to Andrew

Hendry, Jeff Podos, and Joost Raeymaekers for introducing me to the magical place that

happened to be visited, some 185 years ago, by someone named Charles Darwin.

Author's contributions

KMG conceived and designed this research project, collected and analysed the data, and wrote

the manuscript.

Data Accessibility

Data will be archived on Dryad and on the author's academic website.

References

Alberti, M. (2015). Eco-evolutionary dynamics in an urbanizing planet. *Trends in Ecology & Evolution*, *30*(2), 114–126. doi:10.1016/j.tree.2014.11.007

Alberti, M., Correa, C., Marzluff, J. M., Hendry, A. P., Palkovacs, E. P., Gotanda, K. M., … Zhou, Y. (2017). Global urban signatures of phenotypic change in animal and plant populations. *Proceedings of the National Academy of Sciences*, *114*(34), 8651–8956. doi:10.1073/pnas.1606034114

Aplin, L M, Farine, D. R., Morand-Ferron, J., Cockburn, A., Thornton, A., & Sheldon, B. C. (2015). Experimentally induced innovations lead to persistent culture via conformity in wild birds. *Nature*, *518*(7540), 538–541. doi:10.1038/nature13998

Aplin, Lucy M. (2018). Culture and cultural evolution in birds: a review of the evidence. *Animal Behaviour*. doi:10.1016/j.anbehav.2018.05.001

Bates, D. M., Maechler, M., Bolker, B. M., & Walker, S. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, *67*(1), 1–48.

Beauchamp, G. (2004). Reduced flocking by birds on islands with relaxed predation. *Proceedings of the Royal Society B: Biological Sciences*, *271*(1543), 1039–1042. doi:10.1098/rspb.2004.2703

Berger, S., Wikelski, M., Romero, L. M., Kalko, E. K., & Roedl, T. (2007). Behavioral and physiological adjustments to new predators in an endemic island species, the Galápagos marine iguana. *Hormones and Behavior*, *52*(5), 653–663. doi:10.1016/j.yhbeh.2007.08.004

Blackburn, T. M., Cassey, P., Duncan, R. P., Evans, K. L., & Gaston, K. J. (2004). Avian extinction and mammalian introductions on oceanic islands. *Science*, *305*(5692), 1955–1958. doi:10.2307/3837875

Blancher, P. (2013). Estimated number of birds killed by house cats (*Felis catus*) in Canada. *Avian Conservation and Ecology*, *8*(2), 3. doi:10.5751/ACE-00557-080203

Blumstein, D T. (2003). Flight-initiation distance in birds is dependent on intruder starting distance. *Journal of Wildlife Management*, *67*, 852–857.

Blumstein, D T. (2006). Developing an evolutionary ecology of fear: how life history and natural history traits affect disturbance tolerance in birds. *Animal Behaviour*, *71*, 389–399.

Blumstein, D T, Daniel, J. C., & Springett, B. P. (2004). A test of the multi-predator hypothesis: rapid loss of antipredator behavior after 130 years of isolation. *Ethology*, *110*(11), 919–934. doi:10.1111/j.1439-0310.2004.01033.x

Blumstein, D T, Fernandez-Juricic, E., Zollner, P. A., & Garity, S. C. (2005). Inter-specific variation in avian responses to human disturbance. *Journal of Applied Ecology*, *42*(5), 943–953. doi:10.1111/j.1365-2664.2005.01071.x

Blumstein, Daniel T. (2002). Moving to suburbia: ontogenetic and evolutionary consequences of life on predator-free islands. *Journal of Biogeography*, *29*(5–6), 685–692. doi:10.1046/j.1365-2699.2002.00717.x

Blumstein, Daniel T. (2006). The multipredator hypothesis and the evolutionary persistence of antipredator behavior. *Ethology*, *112*(3), 209–217. doi:10.1111/j.1439-0310.2006.01209.x

Blumstein, Daniel T, & Daniel, J. C. (2005). The loss of anti-predator behaviour following isolation on islands. *Proceedings of the Royal Society B: Biological Sciences*, *272*(1573), 1663–1668. doi:10.1098/rspb.2005.3147

Campbell, K. J., Harper, G., Algar, D., Hanson, C. C., Keitt, B. S., & Robinson, S. (2011). Review of feral cat eradication on islands. In C. R. Veitch, M. N. Clout, & D. R. Towns (Eds.), *Island Invasives: Eradication and Management*. Sitzerland and New Zealand: IUCN

and CBB.

Carrión, V., Sevilla, C., & Tapia, W. (2008). Management of introduced animals in Galapagos.
Galapagos Research, *65*, 46–48. Retrieved from
http://aquaticcommons.org/21365/1/GR_65_p46-48.pdf

Cavalli, M., Baladrón, A. V., Isacch, J. P., Biondi, L. M., & Bó, M. S. (2018). The role of
habituation in the adjustment to urban life: An experimental approach with burrowing owls.
Behavioural Processes, *157*(September), 250–255. doi:10.1016/j.beproc.2018.10.011

Clavero, M., Brotons, L., Pons, P., & Sol, D. (2009). Prominent role of invasive species in avian
biodiversity loss. *Biological Conservation*, *142*(10), 2043–2049.
doi:http://dx.doi.org/10.1016/j.biocon.2009.03.034

Clavero, M., & García-Berthou, E. (2005). Invasive species are a leading cause of animal
extinctions. *Trends in Ecology & Evolution*, *20*(3), 110.
doi:http://dx.doi.org/10.1016/j.tree.2005.01.003

Cooper, W E, & Frederick, W. G. (2007). Optimal flight initiation distance. *Journal of
Theoretical Biology*, *244*(1), 59–67.

Cooper, William E., & Blumstein, D. T. (Eds.). (2015). *Escaping From Predators: An
Integrative View of Escape Decisions*. Cambridge, UK: Cambridge University Press.

Cooper, William E, Pyron, R. A., & Garland, T. (2014). Island tameness: living on islands
reduces flight initiation distance. *Proceedings of the Royal Society B: Biological Sciences*,
281(1777), 20133019. doi:10.1098/rspb.2013.3019

Cooper, William E, Samia, D. S. M., & Blumstein, D. T. (2015). FEAR, Spontaneity, and
Artifact in Economic Escape Theory: A Review and Prospectus. In *Advances in the Study of
Behavior* (Vol. 47, pp. 147–197).

Côté, I. M., & Sutherland, W. J. (1997). The effectiveness of removing predators to protect bird populations. *Conservation Biology*, *11*(2), 395–405.

Díaz, M., Møller, A. P., Flensted-Jensen, E., Grim, T., Ibáñez-Álamo, J. D., Jokimäki, J., … Tryjanowski, P. (2013). The geography of fear: a latitudinal gradient in anti-predator escape distances of birds across Europe. *PLoS ONE*, *8*(5), e64634. doi:10.1371/journal.pone.0064634

Fitter, J., Fitter, D., & Hosking, D. (2016). *Wildlife of the Galápagos* (2nd ed.). Princeton, New Jersey: Princeton University Press.

Fox, J., & Weisberg, S. (2011). *An {R} Companion to Applied Regression* (Second). Thousand Oaks, CA: Sage. Retrieved from http://socserv.socsci.mcmaster.ca/jfox/Books/Companion

Fritts, T. H., & Rodda, G. H. (1998). The role of introduced species in the degredation of island ecosystems: a case history of Guam. *Annual Review of Ecology and Systematics*, *29*(1), 113–140. doi:doi:10.1146/annurev.ecolsys.29.1.113

Grant, P. R., & Grant, R. B. (2008). *How and Why Species Multiply: The Radiation of Darwin's Finches*. Princeton, New Jersey, USA: Princeton University Press.

Grant, P. R., & Grant, R. B. (2014). *40 Years of Evolution: Darwin's Finches on Daphne Major Island*. Princeton, New Jersey, USA: Princeton University Press.

Groot, R. S. D. E. (1982). Origin, status and ecology of the owls in Galapagos. *Ardea*, *71*, 167–182.

Harper, G. A., & Carrión, V. (2011). Introduced rodents in the Galapagos: colonisation, removal and the future. In C. R. Veitch, M. N. Clout, & D. R. Towns (Eds.), *Island Invasives: Eradication and Management* (pp. 63–66). Gland, Switerzland: IUCN.

Hendry, A. P., Gotanda, K. M., & Svensson, E. I. (2017). Human influences on evolution, and

the ecological and societal consequences. *Philosophical Transactions of the Royal Society B*, *372*(1712), 20160028. doi:10.1098/rstb.2016.0028

Hughes, B. J., Martin, G. R., & Reynolds, S. J. (2008). Cats and seabirds: effects of feral domestic cat *Felis silvestris catus* eradication on the population of Sooty Terns *Onychoprion fuscata* on Ascension Island, South Atlantic. *Ibis*, *150*, 122–131. doi:10.1111/j.1474-919X.2008.00838.x

Johnson, M. T. J., & Munshi-South, J. (2017). Evolution of life in urban environments. *Science*, *358*(607), eaam8237. doi:10.1126/science.aam8327

Jolly, C. J., Webb, J. K., & Phillips, B. L. (2018). The perils of paradise: an endangered species conserved on an island loses antipredator behaviours within 13 generations. *Biology Letters*, *14*, 20180222. doi:10.1098/RSBL.2018.0222

Jones, H. P., Holmes, N. D., Butchart, S. H. M., Tershy, B. R., Kappes, P. J., Corkery, I., … Croll, D. A. (2016). Invasive mammal eradication on islands results in substantial conservation gains. *Proceedings of the National Academy of Sciences*, *113*(15), 4033–4038. doi:10.1073/pnas.1521179113

Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). lmerTest package: tests in linear mixed effects models. *Journal of Statistical Software*, *82*(13), 1–26. doi:10.18637/jss.v082.i13

Lavers, J. L., Wilcox, C., & Donlan, C. J. (2010). Bird demographic responses to predator removal programs. *Biological Invasions*, *12*(11), 3839–3859. doi:10.1002/ajmg.a.36454

Lever, C. (1994). *Naturalized Animals: The Ecology of Successfully Introduced Species*. London: Poyser Natural History.

Loss, S. R., Will, T., & Marra, P. P. (2013). The impact of free-ranging domestic cats on wildlife

of the United States. *Nature Communications*, *4*, 2961.

doi:http://dx.doi.org/10.1038/ncomms3961

Low, T. (2002). *Feral Future: The Untold Story of Australia's Exotic Invaders*. Chicago:

University of Chicago Press.

MacArthur, R. H., & Wilson, E. O. (1967). *The Theory of Island Biogeography*. Princeton, N. J.:

Princeton University Press.

McKinney, M. L., & Lockwood, J. L. (1999). Biotic homogenization: a few winners replacing

many losers in the next mass extinction. *Trends in Ecology & Evolution*, *14*(11), 450–453.

doi:http://dx.doi.org/10.1016/S0169-5347(99)01679-1

Møller, A. P. (2008). Flight distance of urban birds, predation, and selection for urban life.

Behavioral Ecology and Sociobiology, *63*, 63–75. doi:10.1007/s00265-008-0636-y

Møller, A. P. (2009). Successful city dwellers: a comparative study of the ecological

characteristics of urban birds in the Western Palearctic. *Oecologia*, *159*, 849–858.

doi:10.1007/s00442-008-1259-8

Møller, A. P., & Tryjanowski, P. (2014). Direction of approach by predators and flight initiation

distance of urban and rural populations of birds. *Behavioral Ecology*, *25*(4), 960–966.

doi:10.1093/beheco/aru073

Nogales, M., Vidal, E., Medina, F. M., Bonnaud, E., Tershy, B. R., Campbell, K. J., & Zavaleta,

E. S. (2013). Feral cats and biodiversity conservation: the urgent prioritization of island

management. *BioScience*, *63*(10), 804–810. doi:10.1525/bio.2013.63.10.7

Phillips, R. A. (2010). Eradications of invasive mammals from islands: Why, where, how and

what next? *Emu*, *110*(4). doi:10.1071/MUv110n4_ED

Phillips, R. B., Cooke, B. D., Campbell, K., Carrion, V., Marquez, C., & Snell, H. L. (2005).

Eradicating feral cats to protect Galapagos Land Iguanas: methods and strategies. *Pacific Conservation Biology*, *11*(4), 257–267. doi:10.1071/PC050257

Phillips, R.Brand, Wiedenfeld, D. A., & Snell, H. L. (2012). Current status of alien vertebrates in the Galápagos Islands: invasion history, distribution, and potential impacts. *Biological Invasions*, *14*(2), 461–480. doi:10.1007/s10530-011-0090-z

Samia, D. S. M., Nomura, F., & Blumstein, D. T. (2013). Do animals generally flush early and avoid the rush? A meta-analysis. *Biology Letters*, *9*(2), 20130016–20130016. doi:10.1098/rsbl.2013.0016

Samia, Diogo S. M., Blumstein, D. T., Diaz, M., Grim, T., Ibáñez-Álamo, J. D., Jokima□ki, J., … Moller, A. P. (2017). Rural-urban differences in escape behavior of European birds across a latitudinal gradient. *Frontiers in Ecology and Evolution*, *5*(June), 1–13. doi:10.3389/fevo.2017.00066

Samia, Diogo S. M., Nakagawa, S., Nomura, F., Rangel, T. F., & Blumstein, D. T. (2016). Increased tolerance to humans among disturbed wildlife. *Nature Communications*, *6*(1), 1–8. doi:10.1007/s13398-014-0173-7.2

Sax, D. F., & Gaines, S. D. (2008). Species invasions and extinction: the future of native biodiversity on islands. *Proceedings of the National Academy of Sciences*, *105*(Supplement 1), 11490–11497. doi:10.1073/pnas.0802290105

Stone, P. A., Snell, H. L., & Snell, H. M. (1994). Behavioral diversity as biological diversity: introduced cats and lava lizard wariness. *Conservation Biology*, *8*(2), 569–573. doi:10.2307/2386482

Swash, A., & Still, R. (2005). *Birds, Mammals, and Reptiles of the Galápagos Islands - An Identification Guide* (2nd ed.). New Haven: Yale University Press.

van Dongen, W. F. D., Robinson, R. W., Weston, M. A., Mulder, R. A., & Guay, P.-J. (2015). Variation at the DRD4 locus is associated with wariness and local site selection in urban black swans. *BMC Evolutionary Biology*, *15*(1), 253. doi:10.1186/s12862-015-0533-8

Vitousek, P. M., Mooney, H. A., Lubchenco, J., & Melillo, J. M. (1997). Human domination of Earth's ecosystems. *Science*, *277*(5325), 494–499. doi:10.1126/science.277.5325.494

Vries, T. De. (1976). Prey selection and hunting methods of the Galapagos Hawk, *Buteo galapagoensis*. *Le Gerfaut*, *66*, 3–43.

Vries, T. De. (1989). Conservation, Status and Ecological Importance of the Galapagos Hawk Buteo galapagoensis. In B.-U. Meyburg & R. D. Chancellor (Eds.), *Raptors in the Modern World* (pp. 327–330). Berlin, London, & Paris: WWGBP.

Vries, T. De. (2015). *The Galapagos Hawk/El gavilán de Galápagos* (Monografía). Castellón: Tundra Ediciones.

Wiedenfeld, D. A., & Jiménez-Uzcátegui. (2008). Critical problems for bird conservation in the Galápagos Islands. *Cotinga*, *29*, 22–27.

Woinarski, J. C. Z., Murphy, B. P., Palmer, R., Legge, S. M., Dickman, C. R., Doherty, T. S. E., … Stokeld, D. I. (2018). How many reptiles are killed by cats in Australia? *Wildlife Research*, *45*(3), 247–266.

Woods, M., Mcdonald, R. A., & Har Ris, S. (2003). Predation of wildlife by domestic cats *Felis catus* in Great Britain. *Mammal Rev . Mammal Review Blackwell Science*, *33*(33), 174–188. doi:10.1046/j.1365-2907.2003.00017.x

Ydenberg, R. C., & Dill, L. M. (1986). The economics of fleeing from predators. *Advances in the Study of Behavior*, *16*, 229–249.

Zavaleta, E. S., Hobbs, R. J., & Mooney, H. A. (2001). Viewing invasive species removal in a

whole-ecosystem context. *Trends in Ecology & Evolution*, *16*(8), 454–459.

doi:10.1016/S0169-5347(01)02194-2

Table 1. Results looking at the effects of invasive predator regime (present, pristine, or eradicated) on flight initiation distance (FID). A general linear mixed model was performed with FID as the dependent variable, invasive-predator regime, sex, and starting distance as fixed factors, time of day and group size as covariates, and island as a random factor (denoted by *). Non-significant interactions were removed from the analysis. R^2 for the full model was 0.142. Data used for this analysis were from eight islands that varied in invasive-predator regime. For data collected on islands with permanent human populations, only data from non-urban sites used here. Bold indicates significant P values.

	χ^2	d.f.	R^2	P
Predation	16.770	2	0.093	**<0.001**
Sex	3.505	2	0.008	0.173
Time of Day	1.215	1	0.007	0.270
Starting Distance	0.004	1	0.000	0.949
Group Size	0.207	1	0.010	0.207
Island*	0.182	1	NA	0.669

Table 2. Results looking at the effect of island on flight initiation distance (FID). A linear mixed model was performed with FID as the dependent variable, island, sex, and starting distance as fixed factors and time of day and group size as covariates. Non-significant interactions were removed from the analysis. Data used for this analysis were from eight islands that varied in invasive-predator regime. On islands with human populations, only data from non-urban sites are included. Bold indicates significant P values.

	F	d.f.	η^2	P
Island	4.087	7	0.140	**<0.001**
Sex	1.832	2	0.018	0.164
Time of Day	2.204	1	0.011	0.140
Starting Distance	0.330	1	0.000	0.567
Group Size	3.862	1	0.019	0.051
Island * Starting Distance	3.522	7	0.120	**0.002**

Table 3. Results looking at the effect of urbanization on flight initiation distance (FID). A general linear mixed model was performed with FID as the dependent variable, site urbanization category, sex, and starting distance as fixed factors, time of day and group size as covariates, and island as a random factor (denoted by *). Non-significant interactions were removed from the analysis. R^2 for full model was 0.258. Data used in this analysis were from urban and non-urban sites on the four islands with permanent human populations. Bold indicates significant P values.

	χ^2	d.f.	R^2	P
Site Category	43.307	1	0.215	**<0.001**
Sex	0.041	1	0.033	0.839
Time of Day	2.371	1	0.015	0.124
Starting Distance	2.000	1	0.069	0.157
Group Size	5.420	1	0.033	**0.020**
Island[*]	2.284	1	NA	0.131

Figure Legends

Figure 1. Mean FID and standard error on the eight islands data were collected from. Symbol shape denotes the invasive-predator regime (pristine, eradicated, or present). Island and site categorization are listed on the top, and numbers indicate sample sizes. Symbol colour indicates the site urbanization category on the four islands with permanent human populations and invasive predators (Floreana, Isabela, San Cristobal, and Santa Cruz). Grey colour indicates non-urban finches, and black indicates urban finches. Open symbols indicate islands with no permanent human populations. The two far left islands, Española and Santa Fe are pristine islands with no history of invasive predators. The next two islands, Baltra and North Seymour are where invasive predators have been successfully eradicated in 2003 and 2008 respectively, and show maintained increased antipredator behaviour. Islands with predators are ordered by population size with Floreana having the smallest urban population and Santa Cruz the largest urban population. In non-urban sites that have invasive predators, antipredator behaviour is increased, and in urban sites, antipredator is significantly decreased on all islands except Floreana. On San Cristobal and Santa Cruz, the antipredator behaviour is lower than on islands untouched by invasive predators and humans.

Figure 2. Mean FID and standard error contrasting urban finches with non-urban finches from the four islands that have permanent human populations. Colour indicates island. Above the 1:1 line indicates FID in non-urban areas is higher than FID in urban areas. Floreana, the island with the smallest human population size, has the least difference in FID between urban and non-urban finches.

Figure 1.

Figure 2.