Identification of novel neuroprotective loci modulating ischemic stroke volume in a cross between wild-derived inbred mouse strains

Han Kyu Lee*, Samuel J. Widmayer†, Min-Nung Huang‡, David L. Aylor†, Douglas A. Marchuk*

* Department of Molecular Genetics and Microbiology, Duke University Medical Center, Durham, North Carolina, USA

† Department of Biological Sciences, North Carolina State University, Raleigh, North Carolina, USA

‡ Division of Cardiology, Department of Medicine, Duke University Medical Center, Durham, North Carolina, USA

Running title: Novel neuroprotective loci for stroke

Key words: Ischemic stroke, wild-derived mouse strains, Neuroprotection, quantitative

trait mapping

Correspondence should be addressed to:

Douglas A. Marchuk, PhD

213 Research Drive

Duke University Medical Center, BOX 3175

Durham, NC 27710

Phone / Fax: +1-919-684-1945 / +1-919-684-2790

E-mail: douglas.marchuk@duke.edu

Abstract

To identify genes involved in cerebral infarction we have employed a forward genetic

approach in inbred mouse strains, using quantitative trait locus (QTL) mapping for

cerebral infarct volume after middle cerebral artery occlusion. We had previously

observed that infarct volume is inversely correlated with cerebral collateral vessel density

in the strains. In this study, we expanded the pool of allelic variation among classical

inbred mouse strains by utilizing the eight founder strains of the Collaborative Cross and

found a wild-derived strain, WSB/EiJ, that breaks this general rule that collateral vessel

density inversely correlates with infarct volume. WSB/EiJ and another wild-derived strain,

CAST/EiJ, show the highest collateral vessel densities of any inbred strain, but infarct

volume of WSB/EiJ mice is 8.7-fold larger than that of CAST/EiJ mice. QTL mapping

between these strains identified four new neuroprotective loci modulating cerebral infarct

volume while not affecting collateral vessel phenotypes. To identify causative variants in

genes we surveyed non-synonymous coding SNPs between CAST/EiJ and WSB/EiJ and

found 96 genes harboring coding SNPs predicted to be damaging and mapping within

one of the four intervals. In addition, we performed RNA sequencing for brain tissue of

CAST/EiJ and WSB/EiJ mice and identified 79 candidate genes mapping in one of the

four intervals showing strain-specific differences in expression. The identification of the

genes underlying these neuroprotective loci will provide new understanding of genetic

risk factors of ischemic stroke which may provide novel targets for future therapeutic

intervention of human ischemic stroke.

1 Introduction

Stroke, the sudden death of the brain tissue occurring when the blood flow to the brain is

lost by blockage or rupture, is a leading cause of death in the United States (the fourth)

and the world (the second) (Roger *et al.* 2012; Johnson *et al.* 2016). The majority (over

80%) of strokes are ischemic in origin and results in irreversible death of brain tissue

(infarction). Studies of genetic risk factors for stroke *susceptibility* in the human have

employed both family-based linkage (Gretarsdottir *et al.* 2003; Helgadottir *et al.* 2004)

and population-based genome-wide association studies (GWAS) (Matarin *et al.* 2007;

Matarin *et al.* 2008a; Matarin *et al.* 2008b). Unfortunately, these studies have not yet

uncovered druggable targets to treat stroke and modify clinical outcomes. These studies

were designed to identify stroke *susceptibility* risk factors, and not factors that modulate

infarct size or neurological outcomes once a stroke has occurred. More relevant to this

goal would be a study of anatomic or neurological outcomes in ischemic stroke patients.

A GWAS has recently been published on neurological or behavioral outcomes of 6,165

ischemic stroke patients (Soderholm *et al.* 2019) but only a single variant passed the

threshold of genome-wide significance. Furthermore, despite an explosion of the human

GWAS for disease phenotypes, no published GWAS for infarct volume in ischemic stroke

has been published. The paucity of human genetic variants shown to modulate stroke

outcomes is perhaps not surprising. Genetic studies in the human related to infarct

volume or neurological outcomes are intrinsically problematic due to uncontrollable

variation in the extent and location of the occluded vessel, and especially, variation in the

critical time between first recognized symptoms of stroke and medical intervention.

Instead, current understanding of the mechanisms underlying infarct damage is primarily

based on experimental animal models.

Animal models of focal cerebral ischemia have been established to investigate the

pathophysiologic events occurring after ischemic stroke. Most stroke models induce

cerebral ischemia within the middle cerebral artery (MCA) territory as most relevant for

thrombo-embolic stroke. MCA occlusion models vary both in the extent of occlusion

(permanent *vs.* transient occlusion) and the site of occlusion (proximal *vs.* distal portion

of the vessel). Several reports have determined that permanent occlusion of distal MCA

method produces more restricted and reproducible damage to the cerebral hemisphere

(Majid *et al.* 2000; Lambertsen *et al.* 2002; Carmichael 2005). Furthermore, previous

studies have also demonstrated that different inbred mouse strains show robust

differences in stroke outcomes, providing evidence that the innate response to permanent

focal cerebral ischemia is under strong genetic control (Barone *et al.* 1993; Majid *et al.*

2000; Lambertsen *et al.* 2002; Sugimori *et al.* 2004).

Therefore, we have taken a forward genetic approach (quantitative trait locus - QTL -

mapping) to identify novel genes/pathways (genetic components) involved in modulating

infarct volume in ischemic stroke using the well-established model of permanent

occlusion of distal MCA (pMCAO). Previously, we uncovered several genetic loci that are

involved in regulating ischemic stroke and identified several genes modulating infarct

volume within the genetic loci (Keum and Marchuk 2009; Chu *et al.* 2013; Keum *et al.*

2013; Lee *et al.* 2016; Lee *et al.* 2018).

One of these loci located on distal chromosome 7 (cerebral infarct volume QTL 1 (*Civq1*))

is the strongest and most significant locus modulating infarct volume, found in multiple

pairwise crosses of inbred mouse strains (Keum and Marchuk 2009; Keum *et al.* 2013).

Intriguingly, *Civq1* overlaps with a locus on chromosome 7 that modulates cerebral

collateral vessel number (collateral artery number QTL 1 (*Canq1*)) (Keum and Marchuk

2009; Zhang *et al.* 2010; Keum *et al.* 2013; Lee *et al.* 2016). We and others have noted

a strong inverse correlation between the volume of the infarct after pMCAO and the

number of collateral vessel connections in the cerebral vasculature (Keum and Marchuk

2009; Zhang *et al.* 2010). These vessels connect portions of the same or different arteries,

and upon occlusion of an artery, provide a circulatory shunt, enabling reperfusion of the

ischemic cerebral territory. *Canq1* was recently shown to be due to variation in the

Rabep2 (Lucitti *et al.* 2016), encoding a protein that modulates endosomal recycling of

VEGFR2, a receptor for vascular endothelial growth factor (Kofler *et al.* 2018). This locus

and gene exert an overwhelming effect on the size of the infarct after vessel occlusion

across a wide spectrum of inbred mouse strains.

Given the strong influence that the collateral circulation plays in the modulation infarct

volume across most inbred mouse strains, we sought strains that break the inverse

correlation between collateral vessel density and infarct volume. Previously, we found

one such strain, C3H, which breaks this inverse correlation. Genome-wide QTL mapping

for infarct volume after pMCAO in F2 progeny of a cross between B6 and C3H identified

a neuroprotective locus, *Civq4*, on chromosome 8 (Chu *et al.* 2013). *Civq4* was identified

in a cross between two classical inbred mouse strains that overall exhibit relatively low

sequence variation compared to that found within the entire species. These and other

commonly used inbred mouse strains contain 94% of their genetic background from *Mus*

musculus domesticus, 5% from *Mus musculus musculus*, and less than 1% from *Mus*

musculus castaneus (Yang *et al.* 2011). Thus, genetic mapping using these inbred mouse

strains may systemically miss potential candidate loci that might cause phenotypic

variation within the entire species. Therefore, in this study, we utilize the eight founder

mouse strains of the collaborative cross (CC) mouse to expand the scope of genetic

variation that we survey in the mouse genome. We use this information to select strains

for QTL mapping that will uncover novel loci that modulate infarct size via a collateral-

independent mechanism.

Materials and methods

Animals. All inbred mouse strains were obtained from the Jackson Laboratory (Bar

Harbor, ME), and then bred locally to obtain mice used in all experiments. Mice (both

male and female animals) were age matched (P21 for collateral vessel perfusion and 12

± 1 week for pMCAO) for all experiments. All animal procedures were conducted under

protocols approved by the Duke University IACUC in accordance with NIH guidelines.

Collateral vessel density measurement. As collateral vessel traits are determined by 3

weeks of age and remain constant for many months (Clayton *et al.* 2008), the collateral

vessel phenotype was measured at P21 as previously described (Lee *et al.* 2016; Lee *et*

al. 2018). Mice were anesthetized with ketamine (100 mg/kg) and xylazine (5 mg/kg), and

the ascending thoracic aorta was cannulated. The animals were perfused with freshly

made buffer (1 mg/ml adenosine, 40 g/ml papaverine, and 25 mg/ml heparin in PBS) to

remove the blood. The pial circulation was then exposed after removal of the dorsal

calvarium and adherent dura mater. The cardiac left ventricle was cannulated and a

polyurethane solution with a viscosity sufficient to minimize capillary transit (1:1 resin to

2-butanone, PU4ii, VasQtec) was slowly infused; cerebral circulation was visualized

under a stereomicroscope during infusion. The brain surface was topically rinsed with

10% PBS-buffered formalin and the dye solidified for 20 min. After post-fixation with 10%

PBS-buffered formalin, pial circulation was imaged. All collaterals interconnecting the

anterior- and middle cerebral artery trees of both hemispheres were counted.

Permanent MCAO. Focal cerebral ischemia was induced by direct permanent occlusion

of the distal MCA as previously described (Lee *et al.* 2016; Lee *et al.* 2018). Briefly, adult

mice were anesthetized with ketamine (100 mg/kg) and xylazine (5 mg/kg), and then 0.5%

bupivacaine (5 mg/ml) was also administrated by injection at the incision site. The right

MCA was exposed by a 0.5 cm vertical skin incision midway between the right eye and

ear under a dissecting microscope. After the temporalis muscle was split, a 2-mm burr

hole was made with a high-speed micro drill at the junction of the zygomatic arch and the

squamous bone through the outer surface of the semi-translucent skull. The MCA was

clearly visible at the level of the inferior cerebral vein. The inner layer of the skull was

removed with fine forceps, and the dura was opened with a 32-gauge needle. While

visualizing under an operating microscope, the right MCA was electrocauterized. The

cauterized MCA segment was then transected with microscissors to verify permanent

occlusion. The surgical site was closed with 6-0 sterile nylon sutures. The temperature of

each mouse was maintained at 37°C with a heating pad during the surgery and then was

placed in an animal recovery chamber until the animal was fully recovered from the

anesthetic. Mice were then returned to their cages and allowed free access to food and

water in an air-ventilated room with the ambient temperature set to 25°C.

Infarct volume measurement. Cerebral infarct volumes were measured 24 h after

surgery because the size of the cortical infarct is largest and stable at 24 h after distal

permanent MCA occlusion (Lambertsen *et al.* 2005). Twenty-four hours after pMCAO

surgery, the animals were euthanized, and the brains were carefully removed. The brains

were placed in a brain matrix and sliced into 1 mm coronal sections after being chilled at

-80°C for 4 min to slightly harden the tissue. Each brain slice was placed in 1 well of a 24-

well plate and incubated for 20 min in a solution of 2% 2,3,5-triphenyltetrazolium chloride

(TTC) in PBS at 37°C in the dark. The sections were then washed once with PBS and

fixed with 10% PBS-buffered formalin at 4°C. Then, 24 h after fixation, the caudal face of

each section was scanned using a flatbed color scanner. The scanned images were used

to determine infarct volume (Wexler *et al.* 2002). Image-Pro software (Media Cybernetics)

was used to calculate the infarcted area of each slice by subtracting the infarcted area of

the hemisphere from the non-infarcted area of the hemisphere to minimize error

introduced by edema. The total infarct volume was calculated by summing the individual

slices from each animal.

Flow cytometry. Adult (8 – 12 week) mouse brain cortex of 3 CAST and 6 WSB mice

were digested with collagenase A (1.5 mg/ml) and DNase I (0.4 mg/ml) in HBSS

containing 5% FBS and 10 mM HEPES at 37°C for 45 min. Pelleted post-digested brain

preparations were suspended in 30% Percoll® and spun at room temperature to separate

cells from myelin. Isolated cells were stained with fluorophore-conjugated antibodies for

30 min at room temperature in PBS containing 3% FBS, 5 µg/ml of anti-CD16/32, 5%

normal rat serum, 5% normal mouse serum and 10 mM EDTA. The antibodies were

obtained from BioLegend (San Diego, CA) (anti-CD3-AF488 (145-2C11), anti-CD49b-PE

(HMα2), anti-F4/80-PE-Cy7 (BM8), anti-B220-AF647 (RA3-6B2), anti-CD64-BV421

(X54-5/7.1), anti-CD103-BV510 (2E7), anti-CD45-BV605 (30-F11), anti-IA/IE-BV650

(M5/114.15.2), and anti-CD11c-BV785 (N418) or BD Biosciences (San Jose, CA) (Anti-

CD31-PerCP-Cy5.5 (MEC13.3), anti-SiglecF-PE-CF594 (E50-2440), anti-Ly6G-AF700

(1A8), anti-CD11b-APC-Cy7 (M1/70), and anti-CD24-BV711 (M1/69), or Invitrogen

(Carlsbad, CA) (Anti-Ly6C-PerCP-Cy5.5 (HK1.4). Dead cells were positively stained with

LIVE/DEAD Fixable Yellow Dead Cell Stains (Molecular Probes, Waltham, MA). Flow

cytometric analysis was performed using a LSRII flow cytometer (BD Biosciences, San

Jose, CA) and results analyzed using FlowJo software (BD, Franklin Lakes, NJ).

SNP genotyping. Genomic DNA was isolated from tails of F2 intercross between

CAST/EiJ (CAST) and WSB/EiJ (WSB) mice using DNeasy Tissue kit (Qiagen, Hilden,

Germany). Genome-wide SNP genotyping was performed with a Mouse Universal

Genotyping Array (MUGA, 8K SNPs). Array hybridization including sample preparation

was performed by Neogen/GeneSeek, Inc (Lincoln, NE).

Linkage (QTL) analysis. Genome-wide scans were performed using R/qtl software.

Genotypes from MUGA were prepared for QTL mapping as follows. A total of 2460

informative markers for CAST and WSB across the mouse genome were used for genetic

mapping. The significance thresholds for LOD scores were determined by 1000

permutations using all informative markers. Each QTL was indicated as a significant when

its LOD score exceeded 95% ($p < 0.05$) of the permutation distribution and QTL intervals

exceeded 99% ($p < 0.01$) of the permutation distribution were considered as candidate

intervals. The 95% confidence interval (CI) of each peak was determined by 1.5-LOD

support interval. The physical (Mb) map positions based on the genomic sequence from

the GRCm38/mm10 were calculated using Mouse Map Converter tool of the Jackson

Laboratory (http://cgd.jax.org/mousemapconverter/).

Haplotype analysis. For the intervals of the identified 4 neuroprotective loci (*Civq8* on

chromosome 1, *Civq9* on chromosome 6, *Civq10* on chromosome 13, and *Civq11* on

chromosome 17), SNP data were obtained from the Mouse Phenome Database

(http://phenome.jax.org/). An amino acid change affected detrimental to a protein was

examined by 3 independents *in silico* prediction algorithms, Polymorphism Phenotyping

v2 (PolyPhen-2, http://genetics.bwh.harvard.edu/pph2/index.shtml), Sorting Intolerant

From Tolerant (SIFT, http://sift.jcvi.org), and Protein Variation Effect Analyzer (PROVEAN, http://provean.jcvi.org).

RNA sequencing analysis. Paired-end, 150 bp sequencing reads were generated from adult (8 – 12 week) brain tissue mRNA of 6 CAST (3 males and 3 females) and 6 WSB (3 males and 3 females) mice on the Illumina HiSeq 2500 platform. Adapters for all paired-end sequencing reads were trimmed using Cutadapt (Martin 2011). Differential gene expression analysis is occasionally confounded by differences in alignment rate of reads obtained from strains that are highly diverged from the reference genome. Furthermore, there are millions of variant sites between the CAST and WSB strains, which would make these effects substantially pronounced. To alleviate these effects, we performed our alignment to pseudo-reference genomes (http://csbio.unc.edu/CCstatus/index.py?run= Pseudo), produced by altering the mouse reference genome (GRCm38.p4) using variant calls for the CAST and WSB strains. Reads from each strain were aligned to their respective pseudo-reference genome using TopHat (v. 2.1.1) (Trapnell *et al.* 2009) under default settings and re-mapped to reference coordinates using the Lapels pipeline (Holt *et al.* 2013; Huang *et al.* 2014).

Differential gene expression. Gene expression counts for each sample were obtained using HTseq (Anders *et al.* 2015) under default settings and subsequently imported into DESeq2 (Love *et al.* 2014), where gene counts for each mouse were normalized for differences in sequencing effort and dispersions were calculated for each sample. Differential expression was determined using a two-sided Wald test comparing two negative binomial distributions. *p*-values were adjusted using a false-discovery rate of 5% by Benjamini-Hochberg correction (Benjamin and Hochberg 1995).

Statistics analysis. Results were represented as the mean ± SEM. Significant

differences between datasets were determined using *p*-values *<.05* were considered

significant.

Data availability. Strain used in this study are available through the Jackson Laboratory

catalog. Supplemental data contain two figures (Figure S1 and S2) and three tables

(Table S1 – S3). Figure S1 shows flow cytometry data used to determine the expression

level of CD45$^+$ hematopoietic cells from brain cortex of CAST and WSB mice. Figure S2

shows both the number of collateral vessel connections and infarct volume after pMCAO

for all individual F2 animals. Table S1 contains genotype and phenotype information of

251 F2 (CAST x WSB) animals used for QTL mapping analysis. Table S2 contains

detailed information of all coding SNPs including *in silico* prediction of three independent

algorithms. Table S3 contains detailed information of RNA sequencing analysis within

identified neuroprotective loci. RNA sequencing data are available at GEO with the

accession number: GSE129379. Supplemental material available at Figshare.

Results

Analysis of collateral vessel density and infarct volume after permanent MCAO for

all eight founder strains of Collaborative Cross mouse strain.

Using the eight founder mouse strains of the CC, we first examined the number of

collateral vessel connections between the anterior cerebral artery (ACA) and middle

cerebral artery (MCA) (Figure 1A and B). Classical inbred mouse strains B6, NOD/ShiLtJ

(NOD), NZO/HILtJ (NZO), and 129S1/SvlmJ (129S1), and a wild-derived strain, PWK/PhJ

(PWK) show similar level of vessel connections (B6 (20.4), NOD (21.0), NZO (21.4),

129S1 (22.4), and PWK (20.4)), whereas A/J (10.2) showed a lower level. Interestingly,

two wild-derived strains, CAST and WSB, showed the highest number of collateral vessel

connections (CAST (28.8) and WSB (27.3)) that we have ever observed in any inbred

strain.

To determine whether collateral vessel density is inversely correlated with infarct volume

in these new strains, we next examined ischemic infarct volume in the permanent

occlusion model. We performed pMCAO and measured infarct volume for the eight CC

founder mouse strains (Figure 1C and D). Infarct volume in these mouse strains displayed

various levels after ischemic stroke induction; A/J (33.4 mm^3), B6 (8.6 mm^3), PWK (8.7

mm^3), NOD (12.3 mm^3), NZO (6.8 mm^3), 129S1 (9.0 mm^3), WSB (22.6 mm^3), and CAST

(2.6 mm^3). As expected, most CC founder mouse strains showed an inverse correlation

between collateral vessel connections and infarct volume. However, one strain, WSB,

breaks this inverse correlation. Although two wild-derived mouse strains, CAST and WSB,

exhibit the highest number of vessel connections we have seen in any inbred strain

(Figure 1B), WSB consistently displayed a dramatically larger infarct volume (8.7-fold)

after pMCAO compared to CAST (22.6 mm^3 vs. 2.6 mm^3) (Figure 1D). This suggests that

at least in part, WSB modulates infarct volume after cerebral ischemia through a

collateral-independent mechanism.

It is conceivable that the strain-dependent infarct volume differences are caused by

different composition of cells involved in host defense mechanisms in the brains of the

two strains (eg, CD45$^+$ hematopoietic cells). Thus, we performed flow cytometric analysis

of CD45$^+$ hematopoietic cells in the cerebral cortex of the two strains. We found no

difference in the fraction of microglia, lymphocytes (B and T cells), or CD11b$^+$ myeloid

cells (macrophages) in the brain tissue of the two strains (Figure S1 A and B).

Four novel loci contribute infarct volume differences between CAST and WSB.

In order to identify the genetic loci modulating infarct volume between these two strains,

we generated F1 and F2 progeny between CAST and WSB and examined both collateral

vessel density and infarct volume. Consistent with the number of collateral vessel

connections between CAST and WSB, collateral vessel connections in the F1 and F2

generations exhibited a very tight distribution; WSB (27.3), CAST (28.8), F1 (27.7), and

F2 (27.6) (Figure 2A). Moreover, the number of collateral vessel connections in individual

F2 animals falls within a rather tight range (from 25 to 31 connections between the ACA

and MCA), similar in number and range to the two parental strains (Figure S2A). Thus,

genetic variation within the two strains does not appear to modulate the collateral vessel

phenotype. Next, using F1 and F2 progeny, we measured infarct volume after pMCAO.

In contrast to the vascular phenotype, infarct volume in the F2 generation was widely

distributed, ranging from 1.1 to 38.9 mm^3 (Figure 2B and Figure S2B) and covering nearly

the entire phenotypic spectrum observed across all inbred strains (Keum and Marchuk

2009; Keum *et al.* 2013).

To discover genetic loci that modulate infarct volume we performed genome-wide QTL

mapping analysis. A total of 251 F2 mice (Figure 2B and Figure S2B) were used for

genome-wide SNP genotyping using a Mouse Universal Genotyping Array (MUGA, 8K

SNPs). For genome-wide QTL mapping analysis, 2460 informative SNP markers were

selected across the mouse genome (Table S1), and we identified four new QTL peaks

that display highly significant linkage to the infarct volume trait. These were located on

chromosomes 1, 6, 13, and 17 (Figure 3). These new loci were designated as *Civq8*

through *Civq11* (Cerebral infarct volume QTL) based on our previous work that identified

seven distinct loci for this trait (Keum and Marchuk 2009; Chu *et al.* 2013). The novel loci

are *Civq8* (chromosome 1, LOD 5.09 (JAX00250952)), *Civq9* (chromosome 6, LOD 7.68

(UNC_rs51474193)), *Civq10* (chromosome 13, LOD 6.15 (UNC130410312)), and *Civq11*

(chromosome 17, LOD 5.45 (UNC170802188)) (Table 1).

New loci uncover potential candidate genes.

For the precise definition of the candidate interval for each locus, we extended each

interval an additional 1.5 Mb flanking each arm of 1.5-LOD support interval. One hundred

seventy-three informative SNP markers were used for the definition of *Civq8* mapping to

chromosome 1. The resulting candidate interval for *Civq8* is from 40.66 Mb to 68.27 Mb

(Figure 4A and Table 1). Interestingly, *Civq8* harbors a transgressive allele (or alleles)

that modulates the phenotype in the opposite direction of the parental strains. As shown

in Figure 4B, the animals with the CAST allele are more sensitive to infarction, while those

with the WSB allele are more resistant to infarction. For *Civq9,* mapping to chromosome

6, one hundred forty-four informative SNP markers were used and its candidate interval

is from 22.41 Mb to 27.88 Mb (Figure 5A and Table 1). At *Civq9*, the CAST allele exhibits

a protective effect on infarct volume in agreement with the phenotype of the parental

CAST strain (Figure 5B). For *Civq10*, mapping to chromosome 13, one hundred twenty

informative SNP markers were used and its candidate interval is from 106.02 Mb to

120.42 Mb (Figure 6A and Table 1). *Civq10* also harbors a transgressive allele (or alleles)

since the WSB allele at *Civq10* is protective, in contrast to the overall phenotype of the

parental strain (Figure 6B). For *Civq11*, mapping to chromosome 17, seventy-eight

informative SNP markers were used and its candidate interval is from 56.20 Mb to 63.71

Mb (Figure 7A and Table 1). The CAST allele in *Civq11* exhibits a protective effect in

agreement with the parental strain (Figure 7B).

To identify possible candidate gene(s) modulating infarct volume in these loci, we first

surveyed candidate genes within the candidate interval of each QTLs and then sought

the presence of non-synonymous coding SNPs (hereafter, coding SNP) in these genes.

We identified a total of 330 coding SNPs in 90 coding genes in *Civq8*, 77 coding SNPs in

20 coding genes in *Civq9*, 109 coding SNPs in 34 coding genes in *Civq10*, and 157 coding

genes in 47 coding genes in *Civq11*. Within these intervals, the vast majority of these

genes harbor multiple coding SNPs (Table S2). To determine whether any of these non-

synonymous amino acid substitutions might affect protein function, all the coding SNPs

were subjected to three independent *in silico* algorithms that predict their functional

consequences; SIFT, PolyPhen-2, and PROVEAN. In *Civq8*, 46 genes have a coding

SNP (or SNPs) predicted to be damaging by at least one of the prediction algorithms.

However, only 11 genes harbor coding SNPs predicted to be damaging by all three

programs and another 15 harbor damaging SNPs if the threshold is relaxed that only two

of the three algorithms predict damaging consequences. In *Civq9*, 7 genes have coding

SNPs predicted to be damaging by at least one algorithm, with only one predicted to be

damaging by all three programs and another 3 genes by only two algorithms. In *Civq10*,

a total 16 genes have coding SNPs predicted to be damaging with only 4 predicted to be

damaging by all three programs and another 1 gene by only two algorithms. In *Civq11*, a

total 27 genes have coding SNPs predicted to be damaging with only 3 predicted to be

damaging by all three programs and another 7 genes by only two algorithms (Table 2 and

Table S2). Although genes containing coding SNPs predicted to be damaging by all 3

independents *in silico* algorithms are the best candidate genes, even genes containing

coding SNPs predicted to be damaging by only one *in silico* algorithm are still potential

candidates. The details of all coding SNPs and functional predictions are listed in Table

S2.

Strain-specific differential gene expression discovers candidate genes in the four

new loci.

To identify other candidate genes underlying these loci, we sought evidence of strain-

specific differential transcript levels between two wild-derived mouse strains, CAST and

WSB. Strain-specific differences in transcript levels could potentially be caused by

regulatory sequence variation acting in *cis* by any number of potential molecular

mechanisms. We employed RNA sequencing data analysis using adult brain tissues

between CAST and WSB. As shown in Figure 8, a total 9,563 genes were differentially

expressed between strains but only 220 of these genes are located within one of the four

new QLTs, with 135 in *Civq8*, 10 in *Civq9*, 47 in *Civq10* and 28 in *Civq11*. Assuming that

the most likely candidate genes would show at least a 2-fold difference in strain-specific

expression reduces the overall list to only 79 candidate genes; 47 genes in *Civq8*, 3 genes

in *Civq9*, 19 genes in *Civq10*, and 10 genes in *Civq11* (Table 3A and B). All 220 genes

are displayed in Table S3 which includes information on direction of difference, that is,

which inbred strain of the pair exhibits the higher expression. Genes harboring either

coding SNPs predicted to be damaging or those showing differential expression between

CAST and WSB are potential candidate genes that modulate infarct volume after

ischemic stroke via a collateral vessel-independent mechanism.

Discussion

The long-term objective of this work is to identify novel targets for stroke therapy, focusing

on those factors that modulate the size of the cerebral infarct in a collateral vessel-

independent manner. In support of this goal, a recent study of a meta-analysis of multiple

clinical stroke cohorts showed that collateral vessel anatomy did not correlate with

neurological outcomes (de Havenon *et al.* 2019). Furthermore, the collateral vasculature

is developmentally established and consequently, may be difficult to modify

therapeutically.

Here, to increase the genetic diversity of our mapping approach, we utilized eight founder

mouse strains of the CC recombinant inbred mapping panel. These strains include five

classical inbred strains (A/J, B6, NOD, NZO, and 129S1) and three wild-derived strains

from Mus musculus subspecies (PWK, WSB, and CAST). Specifically, these wild-derived

strains, PWK (*musculus*), WSB (*domesticus*), and CAST (*castaneous*), are more

genetically diverse than the classical inbred strains (Churchill *et al.* 2004; Threadgill 2004;

Aylor *et al.* 2011; Collaborative Cross Consortium 2012). Using all eight founder strains

of the CC mouse, we surveyed both the number of collateral vessel connections as well

as infarct volume after pMCAO. The number of collateral vessel connection of two wild-

derived strains, CAST and WSB, were the highest we have observed in any inbred mouse

strains. Consistent with the role of collateral vessels in reperfusion of the ischemic territory

following pMCAO, CAST mice exhibit the lowest infarct volume that we have ever

measured. However, although CAST and WSB have the largest number of collateral

vessel connections, unlike CAST, upon pMCAO, WSB exhibits a large infarct volume.

Therefore, this wild-derived WSB strain also breaks the inverse correlation between

collateral vessel connections and infarct volume. Supportive evidence for a non-collateral

mechanism is the observation that infarct volume in the F2 animals shows a broad range

of values while the collateral vessel connections are nearly invariant among F2 animals.

Based on the ability to reperfuse the ischemic tissue due to more than sufficient collateral

vessel connections, but displaying a much larger infarct volume than would be predicted,

WSB likely contains loci that upon ischemic insult, contribute to neuronal death. One

compelling hypothesis is that most strains contain natural neuroprotective loci that are

defective in WSB. In this F2 cross, we identified the most significant four novel

neuroprotective loci (*Civq8* on chromosome 1, *Civq9* on chromosome 6, *Civq10* on

chromosome 13, and *Civq11* on chromosome 17). Importantly, these loci identified in an

F2 cross between CAST and WSB did not overlap with other loci that we previously

identified with other F2 crosses. More importantly, the *Civq1* / *Canq1* locus, the strongest

locus controlling collateral vessel number, is absent in this cross. Interestingly but not

surprisingly, we found transgressive alleles among the loci, alleles that operate in the

opposite direction of the parental strain phenotypes. Transgressive segregation is not

uncommon in intraspecific crosses (Palijan *et al.* 2003; Roper *et al.* 2003), and one of our

other, previously-mapped neuroprotective loci, *Civq4*, also harbors a transgressive allele

(Chu *et al.* 2013).

From the QTL mapping analysis for infarct volume, we identified four candidate intervals

regulating infarct volume through a collateral-independent mechanism. In order to identify

candidate genes within the 1.5 LOD-support intervals, we searched for coding SNP

differences between CAST and WSB focusing on genes where the functional

consequence of the variants were predicted using three different *in silico* prediction

algorithms (Ng and Henikoff 2003; Adzhubei *et al.* 2010; Choi *et al.* 2012). Coding SNPs

predicted as "damaging" by all 3 algorithms were considered the highest priority

candidates, with full realization that any of the other genes harboring coding SNPs cannot

be ignored.

As the genes underlying these four loci might harbor *cis*-regulatory sequence variation

modulating differences in mRNA levels, we determined strain-specific differential

transcript levels between CAST and WSB using RNA sequencing analysis. Genes

exhibiting a statistically significant, 2-fold or greater difference in transcript levels are

considered our highest priority candidate genes.

Through these two independent approaches, we have narrowed our list of candidate

genes for all four novel loci to ninety-six genes harboring potentially damaging coding

SNP variation and seventy-nine genes exhibiting greater than two-fold, strain-specific

differential gene expression between CAST and WSB. To identify the causative genes,

further functional studies are required. However, with additional bioinformatics analyses

and a search of the relevant literature, two genes stand out.

Civq8 harbors a protective WSB allele. *Casp8* (caspase 8), mapping within the interval,

harbors multiple coding SNPs for which the WSB allele of one SNP in particular,

(rs226995171), is predicted to be damaged by all three algorithms used in our analysis.

Moreover, among 37 inbred mouse strains (Sanger4: Sanger SNP and indel data, 89+

million locations, 37 inbred strains of mice (2017)), only the CAST and SPRET/EiJ strains

harbor this rare allele. *Casp8* initiates an extrinsic pathway during apoptotic cell death

(Kischkel *et al.* 1995; Thornberry and Lazebnik, 1998) and it is possible that a poorly

functioning or non-functional *Casp8* would protect brain tissue against ischemia-induced

apoptotic cell death.

The other gene is *Map3k1*, a member of the mitogen-activated protein kinase kinase

kinase (MAP3K) superfamily controlling the MAPKK-MAPK signaling cascade. Although

MAP3K1 plays an important role in multiple aspects of cell physiology, MAP3K1 also

regulates apoptosis in response to multiple cellular stresses (Xia *et al.* 1995; Cardone *et*

al. 1997; Deak *et al.* 1998; Widmann *et al.* 1998). Similar to *Civq8*, *Civq10* is also a

transgressive locus where the WSB allele is protective. One of the coding SNPs

(rs257092960) in *Map3k1* is predicted to be damaged by all three algorithms and WSB is

the only strain harboring this allele among 37 mouse inbred strain (Sanger4). As with

Casp8, it is possible that a non-functional *Map3k1* protects ischemia-induced apoptotic

cell death.

In summary, we have identified a wild-derived mouse strain, WSB, which breaks the

inverse correlation between collateral vessel connections and infarct volume after

pMCAO. By selecting two wild-derived mouse strains, CAST and WSB, that contain

similarly high numbers of collateral vessel connections but which display large differences

in infarct volume, we identified a strain pair that might uncover novel loci involved in

neuroprotection. In a cross between these strains, we discovered four novel

neuroprotective genetic loci. Using RNA sequencing of brain tissue and *in silico* analyses

of coding SNPs, we further prioritized the genes mapping within these intervals. The

identification of such neuroprotective genes may provide novel targets for future

therapeutic intervention for human ischemic stroke.

Acknowledgments

The authors thank Dr. Sena Bae and Mr. Daniel A. Snellings for helpful discussion

concerning data sorting and analysis. This work was supported by a grant from NIH grant

5R01HL097281 and the Foundation Leducq Transatlantic Network of Excellence in

Neurovascular Disease (17 CVD 03).

References

Adzhubei, I.A., S. Schmidt, L. Peshkin, V. E. Ramensky, A. Gerasimova, *et al.*, 2010 A

method and server for predicting damaging missense mutations. Nat Methods.

7(4):248-9.

Anders, S., P. T. Pyl, and W. Huber, 2015 HTSeq--a Python framework to work with high-

throughput sequencing data. Bioinformatics. 31(2):166-9.

Aylor, D. L., W. Valdar, W. Foulds-Mathes, R. J. Buus, R. A. Verdugo, *et al.*, 2011 Genetic

analysis of complex traits in the emerging Collaborative Cross. Genome Res.

21(8):1213–22.

Barone, F.C., D. J. Knudsen, A. H. Nelson, G. Z. Feuerstein, and R. N. Willette, 1993

Mouse strain differences in susceptibility to cerebral ischemia are related to cerebral

vascular anatomy. J Cereb Blood Flow Metab. 13(4):683-92.

Benjamini, Y., and Y. Hochberg, 1995 Controlling the false discovery rate: a practical and

powerful approach to multiple testing. J R Statist Soc B. 57(1): 289-300.

Cardone, M. H., G. S. Salvesen, C. Widmann, G. Johnson, and S. M. Frisch, 1997 The

regulation of anoikis: MEKK-1 activation requires cleavage by caspases. Cell.

90(2):315-23

Carmichael, S.T., 2005 Rodent models of focal stroke: size, mechanism, and purpose.

NeuroRx. 2(3), 396-409.

Choi, Y., G. E. Sims, S. Murphy, J. R. Miller, and A. P. Chan, 2012 Predicting the

Functional Effect of Amino Acid Substitutions and Indels. PLoS One. 7(10):e46688.

Chu, P.L., S. Keum, and D. A. Marchuk, 2013 A novel genetic locus modulates infarct
volume independently of the extent of collateral circulation. Physiol Genomics.
45(17):751-63.

Churchill, G. A., D. C. Airey, H. Allayee, J. M. Angel, A. D. Attie *et al.*, 2004 The
Collaborative Cross, a community resource for the genetic analysis of complex traits.
Nat. Genet. 36(11):1133-7.

Clayton, J. A., D. Chalothorn, and J. E. Faber, 2008 Vascular Endothelial Growth Factor-
A Specifies Formation of Native Collaterals and Regulates Collateral Growth in
Ischemia. Circ Res. 103(9):1027-36.

Collaborative Cross Consortium, 2012 The genome architecture of the Collaborative
Cross mouse genetic reference population. Genetics. 190(2):389-401.

De Havenon, A., M. Mlynash, M. A. Kim-Tenser, M. G. Lansberg, T. Leslie-Maazwi, et
al., 2019 Results From DEFUSE 3: Good Collaterals Are Associated With Reduced
Ischemic Core Growth but Not Neurologic Outcome. Stroke. 50(3):632-638.

Deak, J. C., J. V. Cross, M. Lewis, Y. Qian, L. A. Parrott, *et al.*, 1998 Fas-induced
proteolytic activation and intracellular redistribution of the stress-signaling kinase
MEKK1. Proc Natl Acad Sci U S A. 95(10):5595-600.

Gretarsdottir, S., G. Thorleifsson, S. T. Reynisdottir, A. Manolescu, S. Jonsdottir, *et al.*,
2003 The gene encoding phosphodiesterase 4D confers risk of ischemic stroke. Nat
Genet. 35(2):131-138.

Helgadottir, A., A. Manolescu, G. Thorleifsson, S. Gretarsdottir, H. Jonsdottir, *et al.*, 2004 The gene encoding 5-lipoxygenase activating protein confers risk of myocardial infarction and stroke. Nat Genet. 36(3):233-9.

Holt, J., S. Huang, L. McMillan, and W. Wang, 2013 Read annotation pipeline for high-throughput sequencing data, pp. 605-612 in International Conference on Bioinformatics, Computational Biology and Biomedical Informatics, edited by ACM, New York.

Huang, S., J. Holt, C. Y. Kao, L. McMillan, and W. Wang, 2014 A novel multi-alignment pipeline for high-throughput sequencing data. Database (Oxford). bau057.

Johnson, W., O. Onuma, M. Owolabi, and S. Sachdev, 2016 Stroke: a global response is needed. Bull. World Health Organ. 94(9):634-634A.

Keum, S., and D. A. Marchuk, 2009 A locus mapping to mouse chromosome 7 determines infarct volume in a mouse model of ischemic stroke. Circ Cardiovasc Genet. 2(6):591-8.

Keum, S., H. K. Lee, P. L. Chu, M. J. Kan, M. N. Huang, *et al.*, 2013 Natural genetic variation of integrin alpha L (Itgal) modulates ischemic brain injury in stroke. PLoS Genet. 9(10):e1003807.

Kischkel, F. C., S. Hellbardt, I. Behrmann, M. Germer, M. Pawlita, *et al.*, 1995 Cytotoxicity-dependent APO-1 (Fas/CD95)-associated proteins forma death-inducing signaling complex (DISC) with the receptor. EMBO J. 14(22):5579-88.

Kofler, N., F. Corti, F. Rivera-Molina, Y. Deng, D. Toomre, *et al.*, 2018 The Rab-effector

protein RABEP2 regulates endosomal trafficking to mediate vascular endothelial

growth factor receptor-2 (VEGFR2)-dependent signaling. J Biol Chem. 293(13):4805-

4817.

Lambertsen, K. L., R. Gregersen, and B. Finsen, 2002 Microglial-macrophage synthesis

of tumor necrosis factor after focal cerebral ischemia in mice is strain dependent. J

Cereb Blood Flow Metab. 22(7), 785-97.

Lambertsen, K.L., M. Meldgaard, R. Ladeby, and B. Finsen, 2005 A quantitative study of

microglial-macrophage synthesis of tumor necrosis factor during acute and late focal

cerebral ischemia in mice. J Cereb Blood Flow Metab. 25(1):119-35.

Lee, H.K., S. Keum, H. Sheng, D. S. Warner, D. C. Lo, *et al.*, 2016 Natural allelic variation

of the IL-21 receptor modulates ischemic stroke infarct volume. J Clin Invest.

126(8):2827-38.

Lee, H.K., S. Koh, D. C. Lo, and D. A. Marchuk, 2018 Neuronal IL-4Rα modulates

neuronal apoptosis and cell viability during the acute phases of cerebral ischemia.

FEBS J. 285(15):2785-2798.

Love, M. I., W. Huber, and S. Anders, 2014 Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. Genome Biol. 15(12):550.

Lucitti, J.L., R. Sealock, B. K. Buckley, H. Zhang, L. Xiao, *et al.*, 2016 Variants of Rab

GTPase-Effector Binding Protein-2 Cause Variation in the Collateral Circulation and

Severity of Stroke. Stroke. 47(12):3022-3031.

Majid, A., Y. Y. He, J. M. Gidday, S. S. Kaplan, E. R. Gonzales, *et al.*, 2000 Differences

in vulnerability to permanent focal cerebral ischemia among 3 common mouse strains.

Stroke. 31(11):2707-14.

Martin, M., 2011 Cutadapt removes adapter sequences from high-throughput sequencing

reads. EMBnet.journal 17:10-12.

Matarin, M., W. M. Brown, S. Scholz, J. Simon-Sanchez, H. C. Fung, *et al.*, 2007 A

genome-wide genotyping study in patients with ischaemic stroke: initial analysis and

data release. Lancet Neurol. 6(5):414-20.

Matarin, M., W. M. Brown, A. Singleton, J. A. Hardy, and J. F. Meschia, 2008a Whole

genome analyses suggest ischemic stroke and heart disease share an association

with polymorphisms on chromosome 9p21. Stroke. 39(5):1586-9.

Matarin, M., J. Simon-Sanchez, H. C. Fung, S. Scholz, J. R. Gibbs, *et al.*, 2008b Structural

genomic variation in ischemic stroke. Neurogenetics. 9(2):101-8.

Ng, P. C., and S. Henikoff, 2003 SIFT: predicting amino acid changes that affect protein

function. Nucleic Acids Res. 31(13):3812–4.

Palijan, A., J. Dutil, and A. Y. Deng, 2003 Quantitative trait loci with opposing blood

pressure effects demonstrating epistasis on Dahl rat chromosome 3. Physiol

Genomics. 15(1):1-8.

Roger, V.L., A. S. Go, D. M. Lloyd-Jones, E. J. Benjamin, J. D. Berry, *et al.*, 2012 Heart

disease and stroke statistics--2012 update: a report from the American Heart

Association. Circulation. 125(1):e2-e220.

Roper, R. J., R. D. McAllister, J. E. Biggins, S. D. Michael, S. H. Min, *et al.*, 2003 Aod1
   controlling day 3 thymectomy-induced autoimmune ovarian dysgenesis in mice
   encompasses two linked quantitative trait loci with opposing allelic effects on disease
   susceptibility. J Immunol. 170(12):5886-91.

Soderholm, M., A. Pedersen, E. Lorentzen, T. M. Stanne, S. Beven, *et al.*, 2019 Genome-
   wide association meta-analysis of functional outcome after ischemic stroke.
   Neurology. 92(12):e1271-83.

Sugimori, H., H. Yao, H. Ooboshi, S. Ibayashi, and M. Iida, 2004 Krypton laser-induced
   photothrombotic distal middle cerebral artery occlusion without craniectomy in mice.
   Brain Res Brain Res Protoc. 13(3):189-96.

Thornberry N. A., and Y. Lazebnik, 1998 Caspase: enemies within. Science.
   281(5381)1312-6.

Threadgill, D. W., 2006 Meeting report for the 4th annual Complex Trait Consortium
   meeting: from QTLs to systems genetics. Mamm. Genome. 17(1): 2–4.

Trapnell, C., L. Pachter, and S. L. Salzberg, 2009 TopHat: discovering splice junctions
   with RNA-Seq. Bioinformatics. 25(9):1105-11.

Wexler, E. J., E. E. Peters, A. Gonzales, M. L. Gonzales, A. M. Slee, and J. S. Kerr, 2002
   An objective procedure for ischemic area evaluation of the stroke intraluminal thread
   model in the mouse and rat. J Neurosci Methods. 113(1):51-8.

Widmann, C., P. Gerwins, N. L. Johnson, M. B. Jarpe, and G. L. Johnson, 1998 MEK

kinase 1, a substrate for DEVD-directed caspases, is involved in genotoxin-induced

apoptosis. Mol Cell Biol. 18(4):2416-29.

Xia, Z., M. Dickens, J. Raingeaud, R. J. Davis, and M. E. Greenberg, 1995 Opposing

effects of ERK and JNK-p38 MAP kinases on apoptosis. Science. 270(5240):1326-

31.

Yang, H., J. R. Wang, J. P. Didion, R. J. Buus, T. A. Bell, *et al.*, 2011. Subspecific origin

and haplotype diversity in the laboratory mouse. Nat Genet. 43(7): 648–55.

Zhang, H., P. Prabhakar, R. Sealock, and J. E. Faber, 2010 Wide genetic variation in the

native pial collateral circulation is a major determinant of variation in severity of stroke. J

Cereb Blood Flow Metab. 30(5):923-34.

Figure 1

Collateral vessel density and infarct volume after pMCAO in 8 founder strains of

the Collaborative Cross mouse. (A) Representative images of the brains for the 8

founder strains of the CC recombinant inbred mapping panel; A/J, B6, PWK, NOD, NZO,

129S1, WSB, and CAST. For each strain, the area outlined in a red box in the leftmost

image is 3-fold magnified in the image to the right, where the red arrowheads indicate

collateral vessel connections between the ACA and the MCA. Scale bar: 1mm. **(B)** The

graph indicates the average number of collateral vessel connections between ACA and

MCA in the brain. The total number of animals for; A/J, B6, PWK, NOD, NZO, 129S1,

WSB, and CAST were 21, 37, 24, 9, 14, 31, 13, and 32 mice, respectively. Data represent

the mean ± SEM. **(C)** Serial brain sections (1 mm) for each founder strain of the CC

mouse 24 h after pMCAO. The infarct appears as white tissue after 2% TTC staining.

Scale bar: 5 mm. **(D)** The graph shows the infarct volume for each founder strain of the

CC mouse. The total number of animals for; A/J, B6, PWK, NOD, NZO, 129S1, WSB,

and CAST were 19, 32, 27, 9, 10, 32, 18, and 32 animals, respectively. Data represent

the mean ± SEM.

Figure 2

F2 intercross animals between CAST and WSB exhibit a tight distribution of

collateral vessel connections but a wide distribution for pMCAO-induced infarct

volume. (A) The dot plot graph shows the number of collateral vessel connections

between the ACA and MCA for parental strains, WSB and CAST, and their F1 and F2

intercross progeny. The box plot indicates the degree of dispersion and skewness. The

number of animals for WSB, CAST, F1 and F2 was 13, 32, 18, 24 animals, respectively.

Data represent the mean ± SEM. **(B)** Dot and box plot graphs show the distribution of

pMCAO-induced infarct volume for WSB, CAST, and their F1 and F2 intercross progeny.

The number of animals for the infarct volume measurements was 18, 32, 14, and 251

animals, respectively. Data represent the mean ± SEM.

Figure 3

Identification of four novel *Civq* loci modulating infarct volume in the F2 intercross

between CAST and WSB. The graph represents the analysis of a genome-wide QTL

mapping scan for infarct volume measured 24 h after pMCAO using 251 F2 progeny

(CAST and WSB). Chromosomes 1 through X are indicated numerically on the x axis.

The y axis shows a logarithm of the odds (LOD) score and the significant levels ($p < 0.05$

and 0.01) were determined by 1000 permutation tests. Four regions of the genome

mapping to chromosomes 1, 6, 13, and 17 display highly significant QTL mapping to the

infarct volume trait with LOD scores of 5.09, 7.68, 6.15, and 5.45, respectively.

Figure 4

The *Civq8* locus mapping to chromosome 1. (A) The graph shows the QTL mapping

across chromosome 1 using 173 informative SNP markers, highlighting the *Civq8* locus.

The LOD score at the peak is 5.09 (JAX00250952), and the 1.5-LOD support interval is

from 40.66 to 68.27 Mb, indicated by the red bar on the graph. **(B)** Genotype-phenotype

correlation of the F2 cohort at JAX00250952. The alleles are transgressive with the CAST

allele conferring increased susceptibility to infarction and the WSB allele conferring

protection. Data represent the mean ± SEM. ∗∗∗ *P < .001 vs.* CAST/WSB and WSB/WSB,

1-way ANOVA followed by Scheffe's test.

Figure 5

The *Civq9* locus mapping to chromosome 6. (A) The graph shows the QTL mapping

across chromosome 6 using 144 informative SNP markers, highlighting the *Civq9* locus.

The LOD score at the peak is 7.68 (UNC_rs51474193), and the 1.5-LOD support interval

is from 22.41 to 27.88 Mb, indicated by the red bar on the graph. **(B)** Genotype-phenotype

correlation of the F2 cohort at UNC_rs51474193. The CAST allele at *Civq9* is protective

for infarction. Data represent the mean ± SEM. ∗∗∗ *P < .001 vs.* CAST/CAST and

CAST/WSB, 1-way ANOVA followed by Scheffe's test.

Figure 6

The *Civq10* locus mapping to chromosome 13. (A) The graph shows the QTL mapping

across chromosome 13 using 120 informative SNP markers, highlighting the *Civq10*

locus. The LOD score at the peak is 6.15 (UNC130410312), and the 1.5-LOD support

interval is from 106.02 to 120.42 Mb, indicated by the red bar on the graph. **(B)** Genotype-

phenotype correlation of the F2 cohort at UNC130410312. The alleles are transgressive

with the CAST allele conferring increased susceptibility to infarction and the WSB allele

conferring protection. Data represent the mean ± SEM. ∗∗∗ *P < .001 vs.* CAST/WSB and

WSB/WSB and ∗∗ *P < .01 vs.* WSB/WSB, 1-way ANOVA followed by Scheffe's test.

Figure 7

The *Civq11* locus mapping to chromosome 17. (A) The graph shows the QTL mapping

across chromosome 17 using 78 informative SNP markers, highlighting the *Civq11* locus.

The LOD score at the peak is 5.45 (UNC170802188), and the 1.5-LOD support interval

is from 56.20 to 63.71 Mb, indicated by the red bar on the graph. **(B)** Genotype-phenotype

correlation of the F2 cohort at UNC170802188. The CAST allele at *Civq11* is protective

for infarction. Data represent the mean ± SEM. ∗∗∗ *P < .001 vs.* CAST/CAST and

CAST/WSB, 1-way ANOVA followed by Scheffe's test.

Figure 8

RNA sequencing data identifies genes showing strain-specific differences in

cerebral cortex gene expression. (A) The volcano plot shows differential gene

expression between CAST and WSB from brain cortex determined by RNA sequencing

analysis. Each dot represents a different gene with the $\log_2$ fold change plotted against

$\log_{10}$ *p*-value. The total number of significantly differential expressed genes across the

mouse genome is 9,563 genes but only 220 genes map within one of the four QTL

intervals. Differentially-expressed genes in the red box in the leftmost plot are separated

for each interval on the right; green dots indicate 135 genes mapping within *Civq8*, blue

dots indicate 10 genes mapping within *Civq9*, brown dots indicate 47 genes mapping

within *Civq10*, and orange dots indicate 28 genes mapping within *Civq11*.

1 **Table 1**

2 **Characteristics of four novel QTLs for ischemia-induced infarct volume**

QTL	Location	LOD Score	1.5 LOD Interval (Mb) (GRCm38/mm10)	Marker at Peak Peak Position (Mb)	Additive Effect*	Dominance Effect**	Protective Allele
Civq8	Chr1	5.09	1: 40.66 - 68.27 (27.61 Mb)	JAX00250952 53.22 Mb	6.25	-0.77	WSB
Civq9	Chr6	7.68	6: 22.41 - 27.88 (5.47 Mb)	UNC_rs51474193 26.49 Mb	-4.2	-2.74	CAST
Civq10	Chr13	6.15	13: 106.02 - 120.42 (14.40 Mb)	UNC130410312 118.60 Mb	4.17	-0.35	WSB
Civq11	Chr17	5.45	17: 56.20 - 72.82 (16.62 Mb)	UNC170802188 61.66 Mb	-3.93	-1.7	CAST

* Direction of additive effect corresponds to additive effect of CAST allele

** Direction of dominance effect corresponds to deviation from additive relationship between CAST/CAST and WSB/WSB genotypes

Table 2

Candidate genes harboring coding SNPs that are predicted to be damaging by three different *in silico* algorithms.

QTL	Predicted damaging by all 3 *in silico* algorithms			Predicted damaging by only 2 *in silico* algorithms			Predicted damaging by only 1 *in silico* algorithm		
Civq8	Aox2	Aox3	Aox4	Aox1	Bmpr2	Ccdc150	Boll	Carf	Ccdc168
	Casp8	Crygb	Dnah7a	Ercc5	Gm973	Hibch	Clfar	Col5a2	Cyp20a1
	Dytn	Il1r1	Mpp4	Il18rap	Il1r2	Il1rl1	D630023F18Rik	Dnah7b	Il18r1
	Nabp1	Nbeal1		Pard3b	Pms1	Sgol2a	Il1rl2	Map4k4	Mett21c
				Timem237	Trak2	Wdr75	Mettl21e	Mfsd9	Plcl1
							Rftn2	Slc39a10	Spag16
							Spats2l	Zdbf2	
Civq9	Iqub			Hyal4	Pax4	Ptprz1	Cadps2	Cped1	Pot1a
Civq10	Gzma	Itga1	Itga2	Mrps30			4833420G17Rik	Cdc20b	Dhx29
	Map3k1						Emb	Gapt	Hcn1
							Il31ra	Mcidas	Nnt
							Pde4d	Tcstv3	
Civq11	Adgre1	Arrdc5	Ticam1	Cntnap5c	Dennd1c	Pdzph1	1700061G19Rik	4930583I09Rik	Acer1
				Plin4	Pspn	Shd	Adgre4	Chaf1a	Clpp
				Zfp119b			Ebi3	Fer	Fsd1
							Mpnd	Plin5	Safb2
							Stap2	Tnfsf9	Vmn2r120
							Zfp119a	Zfp959	

Detailed coding SNP information for each gene is available in Table S2.

1 **Table 3**

2 **Differential gene expression between CAST and WSB determined by RNA**

3 **sequencing analysis.**

QTL	Symbol	Description	Gene type	log₂ FC (fold change)
	Gm37373	Predicted gene, 37373	TEC*	-4.36
	Gm15759	Predicted gene 15759	antisense_RNA	-3.07
	Gm28802	Predicted gene 28802	unprocessed_pseudogene	-2.75
	Als2cr12	Amyotrophic lateral sclerosis 2 chromosome region 12	protein_coding	-2.55
	4930558J18Rik	RIKEN cDNA 4930558J18 gene	lincRNA	-2.31
	Gm37915	Predicted gene, 37915	antisense_RNA	-2.28
	Dnah7a	Dynein, axonemal, heavy chain 7A	protein_coding	-2.24
	Aox2	Aldehyde oxidase 2	protein_coding	-2.23
	Gm26813	Predicted gene, 26813	lincRNA	-1.80
	Gm8241	Predicted gene 8241	processed_pseudogene	-1.72
	Boll	Boule homolog, RNA binding protein	protein_coding	-1.69
Civq8	Gm8419	Predicted gene 8419	processed_pseudogene	-1.65
	Cps1	Carbamoyl-phosphate synthetase 1	protein_coding	-1.64
	C2cd6b	C2 calcium dependent domain containing 6B	lincRNA	-1.60
	E330011M16Rik	RIKEN cDNA E330011M16 gene	TEC*	-1.59
	Gm27607	Predicted gene, 27607	snoRNA	-1.55
	Osgepl1	O-sialoglycoprotein endopeptidase-like 1	protein_coding	-1.52
	Stat4	Signal transducer and activator of transcription 4	protein_coding	-1.46
	Mstn	Myostatin	protein_coding	-1.40
	Gm35801	Predicted gene, 35801	antisense_RNA	-1.28
	Tmem182	Transmembrane protein 182	protein_coding	-1.24
	Gm11600	Predicted gene 11600	processed_pseudogene	-1.24
	Aox3	Aldehyde oxidase 3	protein_coding	-1.05
Civq9	Hyal5	Hyaluronoglucosaminidase 5	protein_coding	-6.24
	Iqub	IQ motif and ubiquitin domain containing	protein_coding	-1.10
	Gm47776	Predicted gene, 47776	processed_pseudogene	-3.67
	Gm21188	Predicted gene, 21188	protein_coding	-3.66
	Mcidas	Multiciliate differentiation and DNA synthesis associated cell cycle protein	protein_coding	-2.10
Civq10	Itga2	Integrin alpha 2	protein_coding	-1.97
	Gm6270	Predicted gene 6270	lincRNA	-1.84
	Gm15290	Predicted gene 15290	lincRNA	-1.69
	Gm15488	Predicted gene 15488	unitary_pseudogene	-1.69
	Gm15285	Predicted gene 15285	processed_pseudogene	-1.25
	Vmn2r120	Vomeronasal 2, receptor 120	protein_coding	-5.23
Civq11	Nudt12os	Nucleoside diphosphate linked moiety X-type motif 12, opposite strand	antisense_RNA	-2.98
	Cd70	CD70 antigen	protein_coding	-2.68
	Catsperd	Cation channel sperm associated auxiliary subunit delta	protein_coding	-1.59

4 * To be experimentally confirmed

6 **(A)** The table lists all genes showing at least 2-fold differential expression where the CAST

7 allele is higher than WSB allele.

QTL	Symbol	Description	Gene type	log$_2$ FC (fold change)
	Rps27a-ps1	Ribosomal protein S27A, pseudogene 1	processed_pseudogene	3.82
	1700066B17Rik	RIKEN cDNA 1700066B17 gene	lincRNA	3.64
	Gm553	Predicted gene 553	lincRNA	3.07
	Gm29018	Predicted gene 29018	antisense_RNA	2.77
	Gm28177	Predicted gene 28177	processed_transcript	2.49
	BC055402	CDNA sequence BC055402	lincRNA	2.19
	Il1rl2	Interleukin 1 receptor-like 2	protein_coding	1.94
	D930019O06Rik	RIKEN cDNA D930019O06	antisense_RNA	1.91
	Sgo2a	Shugoshin 2A	protein_coding	1.78
	Rfx8	Regulatory factor X 8	protein_coding	1.63
	9530026F06Rik	RIKEN cDNA 9530026F06 gene	lincRNA	1.60
Civq8	4933402D24Rik	RIKEN cDNA 4933402D24 gene	protein_coding	1.43
	Dnah7c	Dynein, axonemal, heavy chain 7C	protein_coding	1.39
	Gm15832	Predicted gene 15832	processed_transcript	1.39
	Hibch	3-hydroxyisobutyryl-Coenzyme A hydrolase	protein_coding	1.30
	Myl1	Myosin, light polypeptide 1	protein_coding	1.24
	1700019D03Rik	RIKEN cDNA 1700019D03 gene	protein_coding	1.22
	Gm28826	Predicted gene 28826	lincRNA	1.21
	Il1r2	Interleukin 1 receptor, type II	protein_coding	1.07
	Gm28449	Predicted gene 28449	lincRNA	1.07
	Gm28982	Predicted gene 28982	lincRNA	1.05
	2810408I11Rik	RIKEN cDNA 2810408I11 gene	processed_transcript	1.04
	Gm18849	Predicted gene, 18849	unprocessed_pseudogene	1.03
	C230029F24Rik	RIKEN cDNA C230029F24 gene	lincRNA	1.02
Civq9	Asb15	Ankyrin repeat and SOCS box-containing 15	protein_coding	1.63
	BC147527	CDNA sequence BC147527	protein_coding	3.53
	Gzmk	Granzyme K	protein_coding	2.83
	Gm15323	Predicted gene 15323	lincRNA	2.28
	4833420G17Rik	RIKEN cDNA 4833420G17 gene	protein_coding	1.96
	Ddx4	DEAD (Asp-Glu-Ala-Asp) box polypeptide 4	protein_coding	1.66
Civq10	Cdc20b	Cell division cycle 20B	protein_coding	1.57
	Gm47914	Predicted gene, 47914	lincRNA	1.52
	3110070M22Rik	RIKEN 3110070M22 gene	protein_coding	1.39
	Gm15327	Predicted gene 15327	processed_transcript	1.16
	4930467J12Rik	RIKEN cDNA 4930467J12 gene	lincRNA	1.07
	Gm15287	Predicted gene 15287	antisense_RNA	1.03
	Vmn2r118	Vomeronasal 2, receptor 118	protein_coding	2.58
	Acsbg2	Acyl-CoA synthetase bubblegum family member 2	protein_coding	1.62
Civq11	Prr22	Proline rich 22	protein_coding	1.49
	Gm16712	Predicted gene, 16712	lincRNA	1.44
	Plin4	Perilipin 4	protein_coding	1.34
	Gm17168	Predicted gene 17168	antisense_RNA	1.13

(B) The table lists all genes showing at least 2-fold differential expression where the WSB allele is higher than CAST allele.

Additional information for all 220 genes is available in Table S3.