Identification of novel neuroprotective loci modulating ischemic stroke volume in a cross between wild-derived inbred mouse strains

Han Kyu Lee*, Samuel J. Widmayer[†], David L. Aylor[†], Douglas A. Marchuk*

* Department of Molecular Genetics and Microbiology, Duke University Medical Center, Durham, North Carolina, USA

[†] Department of Biological Sciences, North Carolina State University, Raleigh, North Carolina, USA

Running title: Novel neuroprotective loci for stroke

Key words: Ischemic stroke, wild-derived mouse strains, Neuroprotection, quantitative

trait mapping

Correspondence should be addressed to:

Douglas A. Marchuk, PhD

213 Research Drive

Duke University Medical Center, BOX 3175

Durham, NC 27710

Phone / Fax: +1-919-684-1945 / +1-919-684-2790

Email: douglas.marchuk@duke.edu

Abstract

To identify genes involved in cerebral infarction we have employed a forward genetic

approach using quantitative trait locus (QTL) mapping for cerebral infarct volume after

middle cerebral artery occlusion, using common inbred mouse strains. This work has

demonstrated that infarct volume is inversely correlated with cerebral collateral vessel

density, since the collateral circulation provides a compensatory shunt for reperfusion of

the ischemic territory. To overcome the limited genetic diversity among domesticated

inbred mouse strains, we have expanded the pool of allelic variation by utilizing the eight

parental strains of the Collaborative Cross. We found that one wild-derived strain,

WSB/EiJ, breaks general rule that collateral vessel density inversely correlates with

infarct volume. WSB/EiJ and another wild-derived strain, CAST/EiJ, show the highest

collateral vessel densities of any inbred strain, but infarct volume of WSB/EiJ is 8.7-fold

larger than CAST/EiJ. QTL mapping between these strains identified 4 new

neuroprotective loci modulating cerebral infarct volume while not affecting collateral

vessel phenotypes. To identify causative variants in genes within the loci, we surveyed

non-synonymous coding SNPs between the strains and also performed RNA sequencing

of brain tissue to determine strain-specific gene expression differences. Combining the

lists of genes harboring coding SNPs predicted to be damaging with genes showing

strain-specific expression differences leaves only 15 genes mapping in any of the four

intervals. The identification of the genes underlying these neuroprotective loci will provide

new understanding of genetic risk factors of ischemic stroke which may provide novel

targets for future therapeutic intervention of human ischemic stroke.

Introduction

Stroke, the sudden death of the brain tissue occurring when the blood flow to the brain is

lost by blockage or rupture, is a leading cause of death in the United States (the fourth)

and the world (the second) (Roger *et al.* 2012; Johnson *et al.* 2016). The majority (over

80%) of strokes are ischemic in origin and results in irreversible death of brain tissue

(infarction). Decades of studies have identified individual risk factors that causes stroke

(eg, smoking, diabetes, obesity, etc.). Furthermore, numerous epidemiologic studies

have estimated that two-thirds of the risk for ischemic stroke is due to genetic factors

(Jamrozik *et al.* 1994; Dichgans 2007). For example, a recent study has reported that a

documented parental ischemic stroke by the age of 65 years was associated with a 3-fold

increase in ischemic stroke risk in offspring (Mozaffarian *et al.* 2008; Mozaffarian *et al.*

2015). However, to date, identification of a significant component of the genetic factors

for ischemic stroke remains elusive.

Animal models of focal cerebral ischemia have been established to investigate the

pathophysiologic events occurring after ischemic stroke. Most stroke models induce

cerebral ischemia within the middle cerebral artery (MCA) territory as most relevant for

thrombo-embolic stroke. MCA occlusion models vary both in the extent of occlusion

(permanent vs. transient occlusion) and the site of occlusion (proximal vs. distal portion

of the vessel). Several reports have determined that permanent occlusion of distal MCA

method produces more restricted and reproducible damage to the cerebral hemisphere

(Majid *et al.* 2000; Lambertsen *et al.* 2002; Carmichael 2005). Furthermore, previous

studies have also demonstrated that different inbred mouse strains show robust

differences in stroke outcomes, providing evidence that the innate response to permanent

focal cerebral ischemia is under strong genetic control (Barone *et al.* 1993; Majid *et al.*

2000; Lambertsen *et al.* 2002; Sugimori *et al.* 2004).

Therefore, we have taken a forward genetic approach (quantitative trait locus - QTL -

mapping) to identify novel genes/pathways (genetic components) involved in modulating

infarct volume in ischemic stroke using the well-established model of permanent

occlusion of distal MCA (pMCAO). Previously, we uncovered several genetic loci that are

involved in regulating ischemic stroke and identified several genes modulating infarct

volume within the genetic loci (Keum and Marchuk 2009; Chu *et al.* 2013; Keum *et al.*

2013; Lee *et al.* 2016; Lee *et al.* 2018). For example, as a major determinant of infarct

volume after pMCAO, we identified a locus on distal chromosome 7 (cerebral infarct

volume QTL 1, *Civq1*) using multiple F2 crosses between inbred mouse strains.

Interestingly, this locus shares the same location on chromosome 7 with a locus

modulating cerebral collateral vessel number (*Canq1*) (Keum and Marchuk 2009; Zhang

et al. 2010; Keum *et al.* 2013; Lee *et al.* 2016). Due to the strong effects of this locus,

infarct volume is inversely correlated with pial collateral vessel density across a broad

range of inbred mouse strains. However, we also found that a small number of inbred

strains break this correlation. One such strain is C3H/HeJ (C3H) that exhibits a similar

number of vessel connections compared to C57BL/6J (B6), but these two strains show

significantly different infarct volumes (Chu *et al.* 2013). Thus, in the common inbred

mouse strains, infarct volume is mainly modulated by collateral circulation, but can also

regulated by a collateral-independent (eg, innate neuroprotection) mechanism. Although

we have identified both collateral-dependent and –independent loci that modulate infarct

volume, genetic variation is limited within the domesticated inbred strains. Therefore, in

1 this study, we utilize eight parental mouse strains of the collaborative cross (CC) mouse

2 to expand genetic variations/differences in the mouse genome and through F2 intercross

3 between two such strains, identify novel genetic loci modulating ischemic stroke via a

4 collateral-independent manner (ie, neuroprotection).

Materials and methods

Animals. All inbred mouse strains were obtained from the Jackson Laboratory (Bar

Harbor, ME) or bred locally from breeding pairs of each strain. Mice (both male and female

animals) were age matched (P21 for collateral vessel perfusion and 12 ± 1 week for

pMCAO) for all experiments. All animal procedures were conducted under protocols

approved by the Duke University IACUC in accordance with NIH guidelines.

Collateral vessel density measurement. As collateral vessel traits are determined by 3

weeks of age and remain constant for many months (Clayton *et al.* 2008), the collateral

vessel phenotype was measured at P21 as previously described (Lee *et al.* 2016; Lee *et*

al. 2018). Mice were anesthetized with ketamine (100 mg/kg) and xylazine (5 mg/kg), and

the ascending thoracic aorta was cannulated. The animals were perfused with freshly

made buffer (1 mg/ml adenosine, 40 g/ml papaverine, and 25 mg/ml heparin in PBS) to

remove the blood. The pial circulation was then exposed after removal of the dorsal

calvarium and adherent dura mater. The cardiac left ventricle was cannulated and a

polyurethane solution with a viscosity sufficient to minimize capillary transit (1:1 resin to

2-butanone, PU4ii, VasQtec) was slowly infused; cerebral circulation was visualized

under a stereomicroscope during infusion. The brain surface was topically rinsed with

10% PBS-buffered formalin and the dye solidified for 20 min. After post-fixation with 10%

PBS-buffered formalin, pial circulation was imaged. All collaterals interconnecting the

anterior- and middle cerebral artery trees of both hemispheres were counted.

Permanent MCAO. Focal cerebral ischemia was induced by direct permanent occlusion

of the distal MCA as previously described (Lee *et al.* 2016; Lee *et al.* 2018). Briefly, adult

mice were anesthetized with ketamine (100 mg/kg) and xylazine (5 mg/kg), and then 0.5%

bupivacaine (5 mg/ml) was also administrated by injection at the incision site. The right

MCA was exposed by a 0.5 cm vertical skin incision midway between the right eye and

ear under a dissecting microscope. After the temporalis muscle was split, a 2-mm burr

hole was made with a high-speed micro drill at the junction of the zygomatic arch and the

squamous bone through the outer surface of the semi-translucent skull. The MCA was

clearly visible at the level of the inferior cerebral vein. The inner layer of the skull was

removed with fine forceps, and the dura was opened with a 32-gauge needle. While

visualizing under an operating microscope, the right MCA was electrocauterized. The

cauterized MCA segment was then transected with microscissors to verify permanent

occlusion. The surgical site was closed with 6-0 sterile nylon sutures. The temperature of

each mouse was maintained at 37°C with a heating pad during the surgery and then was

placed in an animal recovery chamber until the animal was fully recovered from the

anesthetic. Mice were then returned to their cages and allowed free access to food and

water in an air-ventilated room with the ambient temperature set to 25°C.

Infarct volume measurement. Cerebral infarct volumes were measured 24 h after

surgery because the size of the cortical infarct is largest and stable at 24 h after distal

permanent MCA occlusion (Lambertsen *et al.* 2005). Twenty-four hours after pMCAO

surgery, the animals were euthanized, and the brains were carefully removed. The brains

were placed in a brain matrix and sliced into 1 mm coronal sections after being chilled at

-80°C for 4 min to slightly harden the tissue. Each brain slice was placed in 1 well of a 24-

well plate and incubated for 20 min in a solution of 2% 2,3,5-triphenyltetrazolium chloride

(TTC) in PBS at 37°C in the dark. The sections were then washed once with PBS and

fixed with 10% PBS-buffered formalin at 4°C. Then, 24 h after fixation, the caudal face of

each section was scanned using a flatbed color scanner. The scanned images were used

to determine infarct volume (Wexler *et al.* 2002). Image-Pro software (Media Cybernetics)

was used to calculate the infarcted area of each slice by subtracting the infarcted area of

the hemisphere from the non-infarcted area of the hemisphere to minimize error

introduced by edema. The total infarct volume was calculated by summing the individual

slices from each animal.

SNP genotyping. Genomic DNA was isolated from tails of F2 intercross between

CAST/EiJ (CAST) and WSB/EiJ (WSB) mice using DNeasy Tissue kit (Qiagen, Hilden,

Germany). Genome-wide SNP genotyping was performed with a Mouse Universal

Genotyping Array (MUGA, 8K SNPs). Array hybridization including sample preparation

was performed by Neogen/GeneSeek, Inc (Lincoln, NE).

Linkage (QTL) analysis. Genome-wide scans were performed using R/qtl software.

Genotypes from MUGA were prepared for QTL mapping as follows. Total 277 informative

markers for CAST and WSB spaced at approximately 5 cM intervals across the mouse

genome were used for genetic mapping. The significance thresholds for LOD scores were

determined by 1000 permutations using all informative markers. Each QTL was indicated

as a significant when its LOD score exceeded 95% ($p < 0.05$) of the permutation

distribution and QTL intervals exceeded 99% ($p < 0.01$) of the permutation distribution

were considered as candidate intervals. The 95% confidence interval (CI) of each peak

was determined by 1.5-LOD support interval. The physical (Mb) map positions based on

the genomic sequence from the GRCm38/mm10 were calculated using Mouse Map

Converter tool of the Jackson Laboratory (http://cgd.jax.org/mousemapconverter/).

Haplotype analysis. For the intervals of the identified 4 neuroprotective loci (*Civq8* on

chromosome 1, *Civq9* on chromosome 6, *Civq10* on chromosome 13, and *Civq11* on

chromosome 17), SNP data were obtained from the Mouse Phenome Database

(http://phenome.jax.org/). An amino acid change affected detrimental to a protein was

examined by 3 independents *in silico* prediction algorithms, Polymorphism Phenotyping

v2 (PolyPhen-2, http://genetics.bwh.harvard.edu/pph2/index.shtml), Sorting Intolerant

From Tolerant (SIFT, http://sift.jcvi.org), and Protein Variation Effect Analyzer

(PROVEAN, http://provean.jcvi.org).

RNA sequencing analysis. Paired-end, 150 bp sequencing reads were generated from

adult (8 – 12 week) brain tissue mRNA of 6 CAST (3 males and 3 females) and 6 WSB

(3 males and 3 females) mice on the Illumina HiSeq 2500 platform. Adapters for all paired-

end sequencing reads were trimmed using Cutadapt (Martin 2011). Differential gene

expression analysis is occasionally confounded by differences in alignment rate of reads

obtained from strains that are highly diverged from the reference genome. Furthermore,

there are millions of variant sites between the CAST and WSB strains, which would make

these effects substantially pronounced. To alleviate these effects, we performed our

alignment to pseudo-reference genomes, produced by altering the mouse reference

genome (GRCm38.p4) using variant calls for the CAST and WSB strains. Reads from

each strain were aligned to their respective pseudo-reference genome using TopHat (v.

2.1.1) (Trapnell *et al.* 2009) under default settings and re-mapped to reference

coordinates using the Lapels pipeline (Holt *et al.* 2013; Huang *et al.* 2014).

Differential gene expression. Gene expression counts for each sample were obtained

using HTseq (Anders *et al.* 2015) under default settings and subsequently imported into

DESeq2 (Love *et al.* 2014), where gene counts for each mouse were normalized for

differences in sequencing effort and dispersions were calculated for each sample.

Differential expression was determined using a two-sided Wald test comparing two

negative binomial distributions. *p*-values were adjusted using a false-discovery rate of 5%

by Benjamini-Hochberg correction (Benjamin and Hochberg 1995).

Statistics analysis. Results were represented as the mean ± SEM. Significant

differences between datasets were determined using *p*-values *<.05* were considered

significant.

Data availability. Strain used in this study are available through the Jackson Laboratory

catalog. Supplemental data contain one figure (Figure S1) and seven tables (Table S1 –

S7). Figure S1 shows both the number of collateral vessel connections and infarct volume

after pMCAO for all individual F2 animals. Table S1 contains detailed information of all

coding SNPs including *in silico* prediction of three independent algorithms. Table S2

contains detailed information of RNA sequencing analysis within identified

neuroprotective loci. Table S3 contains genotype and phenotype information of 251 F2

(CAST x WSB) animals for QTL mapping. Table S4, S5, S6, and S7 contain genotype

and phenotype information for fine-mapping of *Civq8* on chromosome 1, *Civq9* on

chromosome 6, *Civq10* on chromosome 13, and *Civq11* on chromosome 17, respectively.

RNA sequencing data are available at GEO with the accession number: RNA sequencing

data has been submitted. Supplemental material available at Figshare:

Results

Analysis of pial collateral vessel density and infarct volume after permanent MCAO

for all eight parental strains of Collaborative Cross mouse strain.

Using the eight parental mouse strains of the CC, we first examined the number of

collateral vessel connections between the anterior cerebral artery (ACA) and middle

cerebral artery (MCA) (Figure 1A and B). Strains B6, NOD/ShiLtJ (NOD), NZO/HILtJ

(NZO), and 129S1/SvlmJ (129S1), and a wild-derived strain, PWK/PhJ (PWK) show

similar level of vessel connections (B6 (20.4), NOD (21.0), NZO (21.4), 129S1 (22.4), and

PWK (20.4)), whereas A/J (10.2) showed a lower level. Interestingly, two wild-derived

strains, CAST and WSB, showed the highest number of collateral vessel connections

(CAST (28.8) and WSB (27.3)) that we have ever observed in any inbred strain.

To determine whether collateral vessel density is inversely correlated with infarct volume

in these new strains, we next examined ischemic infarct volume in the permanent

occlusion model. We performed pMCAO and measured infarct volume for the parental

mouse strains (Figure 1C and D). Infarct volume in these parental mouse strains

displayed various levels after ischemic stroke induction; A/J (33.4 mm^3), B6 (8.6 mm^3),

PWK (8.7 mm^3), NOD (12.3 mm^3), NZO (6.8 mm^3), 129S1 (9.0 mm^3), WSB (22.6 mm^3),

and CAST (2.6 mm^3). As expected, most parental mouse strains showed an inverse

correlation between collateral vessel connections and infarct volume. However, one

strain, WSB, breaks this inverse correlation. Although two wild-derived mouse strains,

CAST and WSB, exhibit the highest number of vessel connections we have seen in any

inbred strain (Figure 1B), WSB consistently displayed a dramatically larger infarct volume

(8.7-fold) after pMCAO compared to CAST (22.6 mm^3 vs. 2.6 mm^3) (Figure 1D). This

suggests that, at least in part, WSB may modulate infarct volume after cerebral ischemia

through a collateral-independent mechanism.

Four novel loci contribute infarct volume differences between CAST and WSB.

In order to identify the genetic loci modulating infarct volume between these two strains,

we generated F1 and F2 progeny between CAST and WSB and examined both collateral

vessel density and infarct volume. Consistent with the number of collateral vessel

connections between CAST and WSB, collateral vessel connections in the F1 and F2

generations exhibited a very tight distribution; WSB (27.3), CAST (28.8), F1 (27.7), and

F2 (27.6) (Figure 2A). Moreover, the number of collateral vessel connections in individual

F2 animals showed a similar range of connections (Figure S1A). Thus, genetic variation

within the two strains does not appear to change the collateral vessel phenotype. Next,

using F1 and F2 progeny, we measured infarct volume after pMCAO. In contrast to the

vascular phenotype, infarct volume in the F2 generation was widely distributed, ranging

from 1.1 to 38.9 mm^3 (Figure 2B and Figure S1B) and covering nearly the entire

phenotypic spectrum observed across all inbred strains (Keum and Marchuk 2009; Keum

et al. 2013).

To discover genetic loci that modulate infarct volume we performed genome-wide linkage

analysis. A total of 251 F2 mice (Figure 2B and Supplemental Figure 1B) were used for

genome-wide SNP genotyping using a Mouse Universal Genotyping Array (MUGA, 8K

SNPs). For genome-wide linkage analysis, 277 informative SNP markers spaced at

approximately 5 cM intervals across the mouse genome were selected. From this linkage

mapping, we identified four new linkage peaks that display highly significant linkage to

the infarct volume trait. These were located on chromosomes 1, 6, 13, and 17 (Figure 3).

These new loci were designated as *Civq8* through *Civq11* (Cerebral infarct volume QTL)

based on our previous work that identified seven distinct loci for this trait (Keum and

Marchuk 2009; Chu *et al.* 2013). The novel loci are *Civq8* (chromosome 1, LOD 5.09

(JAX00250952)), *Civq9* (chromosome 6, LOD 7.68 (UNC_rs51474193)), *Civq10*

(chromosome 13, LOD 6.23 (UNC130410312)), and *Civq11* (chromosome 17, LOD 5.45

(UNC170143495)).

Fine-mapping uncovers potential candidate genes in the four new loci.

To better define the linkage intervals, we fine-mapped each locus by adding back

additional SNP markers from the MUGA arrays and extended the interval an additional 1

Mb flanking each arm of 1.5-LOD support interval. Forty-five informative SNP markers

obtained from the MUGA array used for fine-mapping of the *Civq8*, mapping to

chromosome 1. The resulting candidate interval for *Civq8* is from 39 Mb to 70 Mb (Figure

4A and E). For *Civq9*, mapping to chromosome 6, forty-six informative SNP markers were

used and its candidate interval is from 21 Mb to 29 Mb (Figure 4B and E). For *Civq10*,

mapping to chromosome 13, thirty-eight informative SNP markers were used and its

candidate interval is from 108 Mb to 120.4 Mb (Figure 4C and E). For *Civq11*, mapping

to chromosome 17, forty informative SNP markers were used and its candidate interval

is from 54 Mb to 65 Mb (Figure 4D and E).

To identify possible candidate gene(s) modulating infarct volume in these loci, we first

surveyed candidate genes within the candidate interval of each QTLs and then sought

the presence of non-synonymous coding SNPs (hereafter, coding SNP) in these genes.

We identified a total of 330 coding SNPs in 90 coding genes in *Civq8*, 77 coding SNPs in

20 coding genes in *Civq9*, 109 coding SNPs in 34 coding genes in *Civq10*, and 157 coding

genes in 47 coding genes in *Civq11*. Within these intervals, the vast majority of these genes harbor multiple coding SNPs (Figure 4E and Table S1). To determine whether any of these non-synonymous amino acid substitutions might affects protein function, all the coding SNPs were subjected to three independent *in silico* algorithms that predict their functional consequences; SIFT, PolyPhen-2, and PROVEAN. In *Civq8*, 46 genes have a coding SNP (or SNPs) predicted to be damaging by at least one of the prediction algorithms. However, only 11 genes harbor a SNPs predicted to be damaging by all three programs and another 15 harbor damaging SNPs if the threshold is relaxed that only two of the three algorithms predict damaging consequences. In *Civq9*, 7 genes have coding SNPs predicted to be damaging by at least one algorithm, with only one predicted to be damaging by all three programs and another 3 genes by only two algorithms. In *Civq10*, a total 16 genes have coding SNPs predicted to be damaging with only 4 predicted to be damaging by all three programs and another 1 gene by only two algorithms. In *Civq11*, a total 27 genes have coding SNPs predicted to be damaging with only 3 predicted to be damaging by all three programs and another 7 genes by only two algorithms (Table 1 and Table S1). Although genes containing coding SNPs predicted to be damaging by all 3 independents *in silico* algorithms are the best candidate genes, even genes containing coding SNPs predicted to be damaging by only one *in silico* algorithm are still potential candidates. The details of all coding SNPs and functional predictions are listed in Table S1.

Strain-specific differential gene expression discovers candidate genes in the four new loci.

To identify other candidate genes underlying these loci, we sought evidence of strain-

specific differential transcript levels between two wild-derived mouse strains, CAST and

WSB. Strain-specific differences in transcript levels could potentially be caused by

regulatory sequence variation acting in cis, but of any number of potential molecular

mechanisms. We employed RNA sequencing data analysis using adult brain tissues

between CAST and WSB. As shown in Figure 5A, a total 9,563 genes were differentially

expressed between strains but only 220 of these genes are located within one of the four

new QLTs, with 135 in *Civq8*, 10 in *Civq9*, 47 in *Civq10* and 28 in *Civq11*. Assuming that

the most likely candidate genes would show at least a 2-fold difference in strain-specific

expression reduces the list to 47 genes in *Civq8*, 3 genes in *Civq9*, 19 genes in *Civq10*,

and 10 genes in *Civq11* (Figure 4E). All 220 genes are displayed in Table S2 which

includes information on direction of difference, that is, which inbred strain of the pair

exhibits the higher expression.

Upon the assumption genes that exhibit both strain-specific expression differences and

coding SNPs predicted to be damaging would be the most robust candidates, we

compared genes with at least a 2-fold difference in expression between CAST and WSB

within the loci, and asked whether any of these harbored coding SNPs with functional

consequences (Figure 5B). Eight genes meet this combined threshold in *Civq8*, one gene

in *Civq9*, four genes in *Civq10*, and two genes in *Civq11*. As a starting point, these genes

are considered our strongest candidates to modulate infarct volume after ischemic stroke

via a collateral vessel-independent mechanism.

Discussion

Occlusion of the blood vessel in the brain leads to ischemic stroke that results in infarction – irreversible death of the brain tissue. Through decades of research to identify risk factors causing ischemic stroke, our understanding of the molecular events occurring during infarction has dramatically increased. However, although studies of genetic risk factors for stroke susceptibility in the human have been attempted using both family-based linkage (Gretarsdottir *et al.* 2003; Helgadottir *et al.* 2004) and population-based genome-wide association studies (GWAS) (Matarin *et al.* 2007; Matarin *et al.* 2008a; Matarin *et al.* 2008b), the use of these studies to discover druggable targets to treat stroke outcome have been largely disappointing. Furthermore, despite an explosion of the human GWAS for disease phenotypes, no published GWAS for infarct volume in ischemic stroke has been published. Genetic studies in the human related to infarct volume are intrinsically problematic due to uncontrollable variation in the extent and location of the occluded vessel, and especially, variation in the critical time between first recognized symptoms of stroke and medical intervention. Instead, current understanding of the mechanisms underlying infarct damage is primarily based on experimental animal models.

We have employed a forward genetic mapping approach (QTL analysis) to discover novel genes modulating infarction using surgical occlusion mouse model of cerebral ischemia. Through surgical occlusion of the distal middle cerebral artery (pMCAO) in 32 commonly used mouse inbred strains, we have shown that inbred mouse strains demonstrate robust differences in infarct volume (Keum and Marchuk 2009; Keum *et al.* 2013). Exploiting these strain differences, we have mapped several genetic loci (QTLs) that modulate infarct volume in ischemic stroke (Keum and Marchuk 2009; Chu *et al.* 2013; Keum *et al.*

2013). One of these loci located on distal chromosome 7 (cerebral infarct volume QTL 1

(*Civq1*)) is highly significant in regulating infarct volume in multiple pairwise crosses of

inbred mouse strains (Keum and Marchuk 2009; Keum *et al.* 2013). Interestingly, Civq1

is identically overlapped with a locus on chromosome 7 that modulates cerebral collateral

vessel number (collateral artery number QTL 1 (*Canq1*)) (Keum and Marchuk 2009;

Zhang *et al.* 2010; Keum *et al.* 2013; Lee *et al.* 2016). In general, collateral vessels can

serve as a compensatory circulatory shunt when a larger vessel is blocked, enabling

reperfusion of the ischemic cerebral territory. Thus, it is conceivable that cerebral infarct

volume is correlated with the number of collateral vessels in most inbred mouse strains.

Canq1 has recently been shown to be due to variation in the RABEP2 gene (Lucitti *et al.*

2016), encoding a protein that plays a role in endosomal recycling of VEGFR2 (Kofler *et*

al. 2018). This gene also strongly influences the infarct volume phenotype via the

reperfusion mechanism discussed above, although we have identified 3 other genes

mapping within *Civq1*, *Il21r*, *Il4rα*, and *Itgal*, that also modulate infarct volume mainly via

a collateral-independent manner (Keum *et al.* 2013; Lee *et al.* 2016; Lee *et al.* 2018).

Since the collateral vasculature is developmentally established and consequently, may

be difficult to modify therapeutically, we have attempted to identify novel genes

modulating infarct volume in ischemic stroke through a collateral-independent manner (ie.

via neuroprotection). Given the strong influence that the collateral circulation plays in

modulating infarct volume across most inbred mouse strains, we sought strains that

appear to break the inverse correlation between collateral vessel density and infarct

volume. Previously, we found one such strain, C3H, that breaks this inverse correlation.

C3H mice show a similar number of collateral vessel connections compared to the B6

strain, while paradoxically exhibiting a 4-fold increase in infarct volume after pMCAO (Chu

et al. 2013). Genome-wide linkage for infarct volume after pMCAO in F2 progeny of a

cross between B6 and C3H identified a neuroprotective locus, *Civq4*, on chromosome 8

(Chu *et al.* 2013). *Civq4* was identified in a cross between two domesticated inbred mouse

strains that overall exhibit relatively low sequence variation compared to that found within

the entire species. These and other commonly used inbred mouse strains contain 94%

of their genetic background from *Mus musculus domesticus*, 5% from *Mus musculus*

musculus, and less than 1% from *Mus musculus castaneus* (Yang *et al.* 2011). Thus,

genetic mapping using these inbred mouse strains may systemically miss potential

candidate loci that might cause phenotypic variation within the entire species.

Here, to increase the genetic diversity of our mapping approach, we utilized eight founder

inbred mouse strains of the Collaborative Cross mapping population. These strains

include five domesticated inbred strains (A/J, B6, NOD, NZO, and 129S1) and three wild-

derived strains from Mus musculus subspecies (PWK, WSB, and CAST). Specifically,

these wild-derived strains, PWK (*musculus*), WSB (*domesticus*), and CAST (*castaneous*),

are more genetically diverse than the classical inbred strains (Churchill *et al.* 2004;

Threadgill 2004; Aylor *et al.* 2011; Collaborative Cross Consortium 2012). Using all eight

founder strains of the CC mouse, we surveyed both the number of collateral vessel

connections as well as infarct volume after pMCAO. The number of collateral vessel

connection of two wild-derived strains, CAST and WSB, were the highest we have

observed in any inbred mouse strains. Consistent with the role of collateral vessels in

reperfusion of the ischemic territory following pMCAO, CAST mice show the lowest infarct

volume that we have ever measured. However, although CAST and WSB have the largest

number of collateral vessel connections, unlike CAST, upon pMCAO, WSB shows a large

infarct volume. Therefore, we may postulate that this wild-derived WSB strain breaks the

inverse correlation between collateral vessel connections and infarct volume. Supportive

evidence for a non-collateral mechanism is the observation that infarct volume in the F2

animals shows a broad range of values while the collateral vessel connections are nearly

invariant among F2 animals. Based on the ability to reperfuse the ischemic tissue due to

more than sufficient collateral vessel connections, but a much larger infarct volume than

would be predicted, WSB likely contains loci that upon ischemic insult, contribute to

neuronal death. One compelling hypothesis is that most strains contain natural

neuroprotective loci that are defective in WSB. In this F2 cross, we identified the most

significant four novel neuroprotective loci (*Civq8* on chromosome 1, *Civq9* on

chromosome 6, *Civq10* on chromosome 13, and *Civq11* on chromosome 17). Importantly,

these loci identified in an F2 cross between CAST and WSB did not overlap with other

loci that we previously identified with other F2 crosses. More importantly, the *Civq1* /

Canq1 locus, the strongest locus controlling collateral vessel number, is absent in this

cross.

From the linkage analysis for infarct volume, we identified four candidate intervals

regulating infarct volume through a collateral-independent mechanism. In order to identify

candidate genes within the 1.5 LOD-support intervals, we searched for coding SNP

differences between CAST and WSB focusing on genes where the functional

consequence of the variants were predicted using 3 different *in silico* prediction algorithms

(Ng and Henikoff 2003; Adzhubei *et al.* 2010; Choi *et al.* 2012). Coding SNPs predicted

as "damaging" by all 3 algorithms were considered the highest priority candidates, with

full realization that any of the other genes harboring coding SNPs cannot be ignored.

As the genes underlying these four loci might harbor cis-regulatory sequence variation

modulating differences in mRNA levels, we determined strain-specific differential

transcript levels between CAST and WSB using RNA sequencing analysis. Genes

exhibiting a statistically significant, 2-fold or greater difference in transcript levels are

considered our highest priority candidate genes. Finally, to identify a short list of candidate

genes for initial consideration for each of the loci, we sought the intersection of the

predicted functional coding SNPs and those exhibiting strain-specific transcript levels.

Through this association, functional studies can be focused on only 15 genes mapping

within any of the intervals (Figure 5B).

In summary, we identified a wild-derived mouse strain, WSB, that breaks inverse

correlation between a collateral vessel connections and infarct volume after pMCAO. By

selecting two wild-derived mouse strains, CAST and WSB, that contain similarly high

numbers of collateral vessel connections but exhibit large differences in infarct volume

after pMCAO, we identified a strain pair that might uncover novel loci involved in

neuroprotetion. In a cross between these strains, we discovered four novel

neuroprotective genetic loci. Using RNA sequencing of brain tissue and *in silico* analyses

of coding SNPs, we further prioritized the genes mapping within these intervals. The

identification of such neuroprotective genes may provide novel targets for future

therapeutic intervention for human ischemic stroke.

Acknowledgments

The authors thank Dr. Sena Bae for helpful discussion concerning data sorting and

analysis. This work was supported by a grant from NIH grant 5R01HL097281 and the

Foundation Leducq Transatlantic Network of Excellence in Neurovascular Disease (17

CVD 03).

References

Adzhubei, I.A., S. Schmidt, L. Peshkin, V. E. Ramensky, A. Gerasimova, *et al.*, 2010 A method and server for predicting damaging missense mutations. Nat Methods. 7(4):248-9.

Anders, S., P. T. Pyl, and W. Huber, 2015 HTSeq--a Python framework to work with high-throughput sequencing data. Bioinformatics. 31(2):166-9.

Aylor, D. L., W. Valdar, W. Foulds-Mathes, R. J. Buus, R. A. Verdugo, *et al.*, 2011 Genetic analysis of complex traits in the emerging Collaborative Cross. Genome Res. 21(8):1213–22.

Barone, F.C., D. J. Knudsen, A. H. Nelson, G. Z. Feuerstein, and R. N. Willette, 1993 Mouse strain differences in susceptibility to cerebral ischemia are related to cerebral vascular anatomy. J Cereb Blood Flow Metab. 13(4):683-92.

Benjamini, Y., and Y. Hochberg, 1995 Controlling the false discovery rate: a practical and powerful approach to multiple testing. J R Statist Soc B. 57(1): 289-300.

Carmichael, S.T., 2005 Rodent models of focal stroke: size, mechanism, and purpose. NeuroRx. 2(3), 396-409.

Choi, Y., G. E. Sims, S. Murphy, J. R. Miller, and A. P. Chan, 2012 Predicting the Functional Effect of Amino Acid Substitutions and Indels. PLoS One. 7(10):e46688.

Chu, P.L., S. Keum, and D. A. Marchuk, 2013 A novel genetic locus modulates infarct volume independently of the extent of collateral circulation. Physiol Genomics. 45(17):751-63.

Churchill, G. A., D. C. Airey, H. Allayee, J. M. Angel, A. D. Attie *et al.*, 2004 The
Collaborative Cross, a community resource for the genetic analysis of complex traits.
Nat. Genet. 36(11):1133-7.

Clayton, J. A., D. Chalothorn, and J. E. Faber, 2008 Vascular Endothelial Growth Factor-
A Specifies Formation of Native Collaterals and Regulates Collateral Growth in
Ischemia. Circ Res. 103(9):1027-36.

Collaborative Cross Consortium, 2012 The genome architecture of the Collaborative
Cross mouse genetic reference population. Genetics. 190(2):389-401.

Dichgans, M., 2007 Genetics of ischaemic stroke. Lancet Neurol. 6(2):149-61.

Gretarsdottir, S., G. Thorleifsson, S. T. Reynisdottir, A. Manolescu, S. Jonsdottir, *et al.*,
2003 The gene encoding phosphodiesterase 4D confers risk of ischemic stroke. Nat
Genet. 35(2):131-138.

Helgadottir, A., A. Manolescu, G. Thorleifsson, S. Gretarsdottir, H. Jonsdottir, *et al.*, 2004
The gene encoding 5-lipoxygenase activating protein confers risk of myocardial
infarction and stroke. Nat Genet. 36(3):233-9.

Holt, J., S. Huang, L. McMillan, and W. Wang, 2013 Read annotation pipeline for high-
throughput sequencing data, pp. 605-612 in International Conference on
Bioinformatics, Computational Biology and Biomedical Informatics, edited by ACM,
New York.

Huang, S., J. Holt, C. Y. Kao, L. McMillan, and W. Wang, 2014 A novel multi-alignment
pipeline for high-throughput sequencing data. Database (Oxford). bau057.

Jamrozik, K., R. J. Broadhurst, C. S. Anderson, and E. G. Stewart-Wynne. 1994 The role
of lifestyle factors in the etiology of stroke. A population-based case-control study in
Perth, Western Australia. Stroke. 25(1):51-9.

Johnson, W., O. Onuma, M. Owolabi, and S. Sachdev, 2016 Stroke: a global response is
needed. Bull. World Health Organ. 94(9):634-634A.

Keum, S., and D. A. Marchuk, 2009 A locus mapping to mouse chromosome 7 determines
infarct volume in a mouse model of ischemic stroke. Circ Cardiovasc Genet. 2(6):591-
8.

Keum, S., H. K. Lee, P. L. Chu, M. J. Kan, M. N. Huang, *et al.*, 2013 Natural genetic
variation of integrin alpha L (Itgal) modulates ischemic brain injury in stroke. PLoS
Genet. 9(10):e1003807.

Kofler, N., F. Corti, F. Rivera-Molina, Y. Deng, D. Toomre, *et al.*, 2018 The Rab-effector
protein RABEP2 regulates endosomal trafficking to mediate vascular endothelial
growth factor receptor-2 (VEGFR2)-dependent signaling. J Biol Chem. 293(13):4805-
4817.

Lambertsen, K. L., R. Gregersen, and B. Finsen, 2002 Microglial-macrophage synthesis
of tumor necrosis factor after focal cerebral ischemia in mice is strain dependent. J
Cereb Blood Flow Metab. 22(7), 785-97.

Lambertsen, K.L., M. Meldgaard, R. Ladeby, and B. Finsen, 2005 A quantitative study of
microglial-macrophage synthesis of tumor necrosis factor during acute and late focal
cerebral ischemia in mice. J Cereb Blood Flow Metab. 25(1):119-35.

Lee, H.K., S. Keum, H. Sheng, D. S. Warner, D. C. Lo, *et al.*, 2016 Natural allelic variation

of the IL-21 receptor modulates ischemic stroke infarct volume. J Clin Invest.

126(8):2827-38.

Lee, H.K., S. Koh, D. C. Lo, and D. A. Marchuk, 2018 Neuronal IL-4Rα modulates

neuronal apoptosis and cell viability during the acute phases of cerebral ischemia.

FEBS J. 285(15):2785-2798.

Love, M. I., W. Huber, and S. Anders, 2014 Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. Genome Biol. 15(12):550.

Lucitti, J.L., R. Sealock, B. K. Buckley, H. Zhang, L. Xiao, *et al.*, 2016 Variants of Rab

GTPase-Effector Binding Protein-2 Cause Variation in the Collateral Circulation and

Severity of Stroke. Stroke. 47(12):3022-3031.

Majid, A., Y. Y. He, J. M. Gidday, S. S. Kaplan, E. R. Gonzales, *et al.*, 2000 Differences

in vulnerability to permanent focal cerebral ischemia among 3 common mouse strains.

Stroke. 31(11):2707-14.

Martin, M., 2011 Cutadapt removes adapter sequences from high-throughput sequencing

reads. EMBnet.journal 17:10-12.

Matarin, M., W. M. Brown, S. Scholz, J. Simon-Sanchez, H. C. Fung, *et al.*, 2007 A

genome-wide genotyping study in patients with ischaemic stroke: initial analysis and

data release. Lancet Neurol. 6(5):414-20.

Matarin, M., W. M. Brown, A. Singleton, J. A. Hardy, and J. F. Meschia, 2008a Whole
genome analyses suggest ischemic stroke and heart disease share an association
with polymorphisms on chromosome 9p21. Stroke. 39(5):1586-9.

Matarin, M., J. Simon-Sanchez, H. C. Fung, S. Scholz, J. R. Gibbs, *et al.*, 2008b Structural
genomic variation in ischemic stroke. Neurogenetics. 9(2):101-8.

Mozaffarian, D., C. D. Furberg, B. M. Psaty, and D. Siscovick, 2008 Physical activity and
incidence of atrial fibrillation in older adults: the Cardiovascular Health Study.
Circulation. 118(8):800–7.

Mozaffarian, D., E. J. Benjamin, A. S. Go, D. K. Arnett, M. J. Blaha, *et al.*, 2015 Heart
Disease and Stroke Statistics--2015 update: a report from the American Heart
Association. Circulation. 131(4):e29-322.

Ng, P. C., and S. Henikoff, 2003 SIFT: predicting amino acid changes that affect protein
function. Nucleic Acids Res. 31(13):3812–4.

Roger, V.L., A. S. Go, D. M. Lloyd-Jones, E. J. Benjamin, J. D. Berry, *et al.*, 2012 Heart
disease and stroke statistics--2012 update: a report from the American Heart
Association. Circulation. 125(1):e2-e220.

Sugimori, H., H. Yao, H. Ooboshi, S. Ibayashi, and M. Iida, 2004 Krypton laser-induced
photothrombotic distal middle cerebral artery occlusion without craniectomy in mice.
Brain Res Brain Res Protoc. 13(3):189-96.

Threadgill, D. W., 2006 Meeting report for the 4th annual Complex Trait Consortium
meeting: from QTLs to systems genetics. Mamm. Genome. 17(1): 2–4.

Trapnell, C., L. Pachter, and S. L. Salzberg, 2009 TopHat: discovering splice junctions

with RNA-Seq. Bioinformatics. 25(9):1105-11.

Wexler, E.J., E. E. Peters, A. Gonzales, M. L. Gonzales, A. M. Slee, and J. S. Kerr, 2002

An objective procedure for ischemic area evaluation of the stroke intraluminal thread

model in the mouse and rat. J Neurosci Methods. 113(1):51-8.

Yang, H., J. R. Wang, J. P. Didion, R. J. Buus, T. A. Bell, *et al.*, 2011. Subspecific origin

and haplotype diversity in the laboratory mouse. Nat Genet. 43(7): 648–55.

Zhang, H., P. Prabhakar, R. Sealock, and J. E. Faber, 2010 Wide genetic variation in the

native pial collateral circulation is a major determinant of variation in severity of stroke. J

Cereb Blood Flow Metab. 30(5):923-34.

Figure 1

Collateral vessel density and infarct volume after pMCAO in 8 parental strains of

the Collaborative Cross mouse. (A) Representative images of the brains for the 8

parental strains of the CC recombinant inbred mapping panel; A/J, B6, PWK, NOD, NZO,

129S1, WSB, and CAST. For each strain, the area outlined in a red box in the leftmost

image is 3-fold magnified in the image to the right, where the red arrowheads indicate

collateral vessel connections between the ACA and the MCA. Scale bar: 1mm. **(B)** The

graph indicates the average number of collateral vessel connections between ACA and

MCA in the brain. The total number of animals for; A/J, B6, PWK, NOD, NZO, 129S1,

WSB, and CAST were 21, 37, 24, 9, 14, 31, 13, and 32 mice, respectively. Data represent

the mean ± SEM. **(C)** Serial brain sections (1 mm) for each parental strain of the CC

mouse 24 h after pMCAO. The infarct appears as white tissue after 2% TTC staining.

Scale bar: 5 mm. **(D)** The graph shows the infarct volume for each parental strain of the

CC mouse. The total number of animals for; A/J, B6, PWK, NOD, NZO, 129S1, WSB,

and CAST were 19, 32, 27, 9, 10, 32, 18, and 32 animals, respectively. Data represent

the mean ± SEM.

Figure 2

F2 intercross animals between CAST and WSB exhibit a tight distribution of

collateral vessel connections but a wide distribution for pMCAO-induced infarct

volume. (A) The dot plot graph shows the number of collateral vessel connections

between the ACA and MCA for parental strains, WSB and CAST, and their F1 and F2

intercross progeny. The box plot indicates the degree of dispersion and skewness. The

number of animals for WSB, CAST, F1 and F2 was 13, 32, 18, 24 animals, respectively.

Data represent the mean ± SEM. **(B)** The dot and box plot graph show the distribution of

pMCAO-induced infarct volume for WSB, CAST, and their F1 and F2 intercross progeny.

The number of animals for the infarct volume measurements was 18, 32, 14, and 251

animals, respectively. Data represent the mean ± SEM.

Figure 3

Identification of four novel *Civq* loci modulating infarct volume in the F2 intercross

between CAST and WSB. The graph represents the analysis of a genome-wide linkage

scan for infarct volume measured 24 h after pMCAO using 251 F2 progeny (CAST and

WSB). Chromosomes 1 through X are indicated numerically on the x axis. The y axis

shows a logarithm of the odds (LOD) score and the significant levels ($p < 0.05$ and 0.01)

were determined by 1000 permutation tests. Four regions of the genome mapping to

chromosomes 1, 6, 13, and 17 display highly significant linkage to the infarct volume trait

with LOD scores of 5.09, 7.68, 6.23, and 5.45, respectively.

1 **Figure 4**

E

QTL	Chromosome interval (GRCm38/mm10)	Different non-synonymous SNPs between CAST and WSB	Differential gene expressions between CAST and WSB (>2-*fold*)
Civq8	1:39,000,000-70,000,000 (31 Mb)	330 coding SNPs in 90 coding genes	47 genes
Civq9	6:21,000,000-29,000,000 (8 Mb)	77 coding SNPs in 20 coding genes	3 genes
Civq10	13:108,000,000-120,421,639 (22.4 Mb)	109 coding SNPs in 34 coding genes	19 genes
Civq11	17:54,000,000-65,000,000 (11 Mb)	157 coding SNPs in 47 coding genes	10 genes

Fine-mapping and potential candidate genes for the four novel *Civq* loci. (A) The

graph shows the linkage map across chromosome 1 using 45 informative SNP markers,

highlighting the *Civq8* locus. The LOD score at the peak is 5.09 (JAX00250952), and the

1.5-LOD support interval is from 19.627 to 34.627 cM, indicated by red bar on the graph.

(B) The graph shows the linkage map across chromosome 6 using 46 informative SNP

markers, highlighting the *Civq9*. The LOD score at the peak is 7.68 (UNC_rs51474193),

and the 1.5-LOD support interval is from 9.80 to 11.95 cM, indicated by red bar on the

graph. **(C)** The graph shows the linkage map across chromosome 13 using 38 informative

SNP markers, highlighting the *Civq10* locus. The LOD score at the peak is 6.23

(UNC130410312), and the 1.5-LOD support interval is from 61.50 to 67.83 cM, indicated

by red bar on the graph. **(D)** The graph shows the linkage map across chromosome 17

using 40 informative SNP markers, highlighting the *Civq11* locus. The LOD score at the

peak is 5.45 (UNC170143495), and the 1.5-LOD support interval is from 29.50 to 33.10

cM, indicated by red bar on the graph. **(E)** The table provides information for each of the

four loci including confidence intervals, the number of genes harboring coding SNP

differences between CAST and WSB, and the number of genes showing at least 2-fold

expression difference between brain cortex of CAST and WSB.

Figure 5

RNA sequencing data identifies genes showing strain-specific differences in

cerebral cortex gene expression. (A) The volcano plot shows differential gene

expression between CAST and WSB from brain cortex determined by RNA sequencing

analysis. Each dot represents a different gene with the $\log_2$ fold change plotted against

$\log_{10}$ p-value. The total number of significantly differential expressed genes across the

mouse genome is 9,563 genes but only 220 genes map within one of the four QTL

intervals. Differentially-expressed genes in the red box in the leftmost plot are separated

for each interval on the right; green dots indicate 135 genes mapping within *Civq8*, blue

dots indicate 10 genes mapping within *Civq9*, brown dots indicate 47 genes mapping

within *Civq10*, and orange dots indicate 28 genes mapping within *Civq11*. **(B)** For each

locus, this graph identifies genes that show at least a 2-fold difference in expression

between CAST and WSB and also harbor a potentially damaging coding SNP. The bar

goes downward (negative value) for genes with higher gene expression of the CAST

allele and the bar goes upward (positive value) for genes with higher gene expressions

of the WSB allele. The different colors represent how many of the three *in silico* algorithms

used (PolyPhen-2, SIFT, and PROVEAN) predicted that at least one coding SNP in the

gene was damaging.

1 **Table 1**

2 **Candidate genes harboring coding SNPs that are predicted to be damaging by**

3 **three different *in silico* algorithms.**

	Predicted damaging by all 3 *in silico* algorithms			Predicted damaging by only 2 *in silico* algorithms			Predicted damaging by only 1 *in silico* algorithm		
Civq8	Aox2 Casp8 Dytn Nabp1	Aox3 Crygb Il1r1 Nbeal1	Aox4 Dnah7a Mpp4	Aox1 Ercc5 Il18rap Pard3b Timem237	Bmpr2 Gm973 Il1r2 Pms1 Trak2	Ccdc150 Hibch Il1rl1 Sgol2a Wdr75	Boll Clfar D630023F18Rik Il1rl2 Mettl21e Rftn2 Spats2l	Carf Col5a2 Dnah7b Map4k4 Mfsd9 Slc39a10 Zdbf2	Ccdc168 Cyp20a1 Il18r1 Mett21c Plcl1 Spag16
Civq9	Iqub			Hyal4	Pax4	Ptprz1	Cadps2	Cped1	Pot1a
Civq10	Gzma Map3k1	Itga1	Itga2	Mrps30			4833420G17Rik Emb Il31ra Pde4d	Cdc20b Gapt Mcidas Tcstv3	Dhx29 Hcn1 Nnt
Civq11	Adgre1	Arrdc5	Ticam1	Cntnap5c Plin4 Zfp119b	Dennd1c Pspn	Pdzph1 Shd	1700061G19Rik Adgre4 Ebi3 Mpnd Stap2 Zfp119a	4930583I09Rik Chaf1a Fer Plin5 Tnfsf9 Zfp959	Acer1 Clpp Fsd1 Safb2 Vmn2r120

6 Detailed coding SNP information for each gene is available in Table S1.

1 **Table 2**

2 **Differential gene expression between CAST and WSB determined by RNA**

3 **sequencing analysis.**

Chr	Symbol	Description	Gene type	log$_2$ FC (fold change)
	Gm37373	Predicted gene, 37373	TEC	-4.36
	Gm15759	Predicted gene 15759	antisense_RNA	-3.07
	Gm28802	Predicted gene 28802	unprocessed_pseudogene	-2.75
	Als2cr12	Amyotrophic lateral sclerosis 2 chromosome region 12	protein_coding	-2.55
	4930558J18Rik	RIKEN cDNA 4930558J18 gene	lincRNA	-2.31
	Gm37915	Predicted gene, 37915	antisense_RNA	-2.28
	Dnah7a	Dynein, axonemal, heavy chain 7A	protein_coding	-2.24
	Aox2	Aldehyde oxidase 2	protein_coding	-2.23
	Gm26813	Predicted gene, 26813	lincRNA	-1.80
	Gm8241	Predicted gene 8241	processed_pseudogene	-1.72
	Boll	Boule homolog, RNA binding protein	protein_coding	-1.69
1	Gm8419	Predicted gene 8419	processed_pseudogene	-1.65
	Cps1	Carbamoyl-phosphate synthetase 1	protein_coding	-1.64
	C2cd6b	C2 calcium dependent domain containing 6B	lincRNA	-1.60
	E330011M16Rik	RIKEN cDNA E330011M16 gene	TEC*	-1.59
	Gm27607	Predicted gene, 27607	snoRNA	-1.55
	Osgepl1	O-sialoglycoprotein endopeptidase-like 1	protein_coding	-1.52
	Stat4	Signal transducer and activator of transcription 4	protein_coding	-1.46
	Mstn	Myostatin	protein_coding	-1.40
	Gm35801	Predicted gene, 35801	antisense_RNA	-1.28
	Tmem182	Transmembrane protein 182	protein_coding	-1.24
	Gm11600	Predicted gene 11600	processed_pseudogene	-1.24
	Aox3	Aldehyde oxidase 3	protein_coding	-1.05
6	Hyal5	Hyaluronoglucosaminidase 5	protein_coding	-6.24
	Iqub	IQ motif and ubiquitin domain containing	protein_coding	-1.10
	Gm47776	Predicted gene, 47776	processed_pseudogene	-3.67
	Gm21188	Predicted gene, 21188	protein_coding	-3.66
	Mcidas	Multiciliate differentiation and DNA synthesis associated cell cycle protein	protein_coding	-2.10
13	Itga2	Integrin alpha 2	protein_coding	-1.97
	Gm6270	Predicted gene 6270	lincRNA	-1.84
	Gm15290	Predicted gene 15290	lincRNA	-1.69
	Gm15488	Predicted gene 15488	unitary_pseudogene	-1.69
	Gm15285	Predicted gene 15285	processed_pseudogene	-1.25
	Vmn2r120	Vomeronasal 2, receptor 120	protein_coding	-5.23
17	Nudt12os	Nudix (nucleoside diphosphate linked moiety X)-type motif 12, opposite strand	antisense_RNA	-2.98
	Cd70	CD70 antigen	protein_coding	-2.68
	Catsperd	Cation channel sperm associated auxiliary subunit delta	protein_coding	-1.59

4 * To be experimentally confirmed

6 **(A)** The table lists all genes showing at least 2-fold differential expression where the CAST

7 allele is higher than WSB.

Chr	Symbol	Description	Gene type	log$_2$ FC (fold change)
	Rps27a-ps1	Ribosomal protein S27A, pseudogene 1	processed_pseudogene	3.82
	1700066B17Rik	RIKEN cDNA 1700066B17 gene	lincRNA	3.64
	Gm553	Predicted gene 553	lincRNA	3.07
	Gm29018	Predicted gene 29018	antisense_RNA	2.77
	Gm28177	Predicted gene 28177	processed_transcript	2.49
	BC055402	CDNA sequence BC055402	lincRNA	2.19
	Il1rl2	Interleukin 1 receptor-like 2	protein_coding	1.94
	D930019O06Rik	RIKEN cDNA D930019O06	antisense_RNA	1.91
	Sgo2a	Shugoshin 2A	protein_coding	1.78
	Rfx8	Regulatory factor X 8	protein_coding	1.63
	9530026F06Rik	RIKEN cDNA 9530026F06 gene	lincRNA	1.60
1	4933402D24Rik	RIKEN cDNA 4933402D24 gene	protein_coding	1.43
	Dnah7c	Dynein, axonemal, heavy chain 7C	protein_coding	1.39
	Gm15832	Predicted gene 15832	processed_transcript	1.39
	Hibch	3-hydroxyisobutyryl-Coenzyme A hydrolase	protein_coding	1.30
	Myl1	Myosin, light polypeptide 1	protein_coding	1.24
	1700019D03Rik	RIKEN cDNA 1700019D03 gene	protein_coding	1.22
	Gm28826	Predicted gene 28826	lincRNA	1.21
	Il1r2	Interleukin 1 receptor, type II	protein_coding	1.07
	Gm28449	Predicted gene 28449	lincRNA	1.07
	Gm28982	Predicted gene 28982	lincRNA	1.05
	2810408I11Rik	RIKEN cDNA 2810408I11 gene	processed_transcript	1.04
	Gm18849	Predicted gene, 18849	unprocessed_pseudogene	1.03
	C230029F24Rik	RIKEN cDNA C230029F24 gene	lincRNA	1.02
6	Asb15	Ankyrin repeat and SOCS box-containing 15	protein_coding	1.63
	BC147527	CDNA sequence BC147527	protein_coding	3.53
	Gzmk	Granzyme K	protein_coding	2.83
	Gm15323	Predicted gene 15323	lincRNA	2.28
	4833420G17Rik	RIKEN cDNA 4833420G17 gene	protein_coding	1.96
	Ddx4	DEAD (Asp-Glu-Ala-Asp) box polypeptide 4	protein_coding	1.66
13	Cdc20b	Cell division cycle 20B	protein_coding	1.57
	Gm47914	Predicted gene, 47914	lincRNA	1.52
	3110070M22Rik	RIKEN cDNA 3110070M22 gene	protein_coding	1.39
	Gm15327	Predicted gene 15327	processed_transcript	1.16
	4930467J12Rik	RIKEN cDNA 4930467J12 gene	lincRNA	1.07
	Gm15287	Predicted gene 15287	antisense_RNA	1.03
	Vmn2r118	Vomeronasal 2, receptor 118	protein_coding	2.58
	Acsbg2	Acyl-CoA synthetase bubblegum family member 2	protein_coding	1.62
17	Prr22	Proline rich 22	protein_coding	1.49
	Gm16712	Predicted gene, 16712	lincRNA	1.44
	Plin4	Perilipin 4	protein_coding	1.34
	Gm17168	Predicted gene 17168	antisense_RNA	1.13

4   **(B)** The table lists all genes showing at least 2-fold differential expression where the WSB

5   allele is higher than CAST.

6   Additional information for all 220 genes is available in Table S2.