Contingent Convergence: The ability to detect convergent genomic evolution is dependent on demographic similarity

AUTHORS

James R. Whiting[1], Bonnie A. Fraser[1]

[1] *Department of Biosciences, University of Exeter, Geoffrey Pope Building, Exeter, EX4 4QD*

KEYWORDS

Genome scans, population genetics, F_{ST}, D_{XY}, outlier scans, convergence

ABSTRACT

Outlier scans, in which the genome is scanned for signatures of selection, have become a prominent tool in studies of local adaptation, and more recently studies of genetic convergence in natural populations. However, these methods have the potential to be confounded by features of demographic history, which are considerably variable across natural populations. In this study, we use forward-simulations to investigate and illustrate how several measures of genetic differentiation commonly used in outlier scans (F_{ST}, D_{XY} and $\Delta\pi$) are influenced by demographic variation across multiple sampling time points. In a factorial design with 32 treatments, we manipulate the strength of bottlenecks (N of founding individuals), contractions (proportional size of diverging population) and migration rate between an ancestral phenotype population and a diverging phenotype population. Our results illustrate known constraints of individual measures associated with strong contractions and a lack of gene-flow, but notably we demonstrate how relationships between measures are similarly dependent on demography. We find that false-positive signals of convergent evolution (the same simulated genes marked as outliers in independent treatments) are attainable as a product of similar demographic treatment, and that convergent outliers across different measures (particularly F_{ST} and D_{XY}) are achievable with little influence of selection. The data presented act as an illustrative guide of how underappreciated yet quantifiable measures of demographic history influence commonly employed methods for detecting selection.

INTRODUCTION

The advent of modern sequencing technologies has facilitated big-data studies of adaptation and evolution across the genome. These studies often centre around quantifying patterns of variation in genome-wide SNPs, which can be used to highlight regions or genes having experienced selection

relative to the neutral backdrop of the rest of the genome. These analyses, which we refer to here as

outlier scans, have become a common tool in population genetics and have seen great success

across diverse, natural systems in identifying candidate genes associated with the evolution of a

range of adaptive traits (reviewed recently by (1)). In addition, the method of overlapping outlier

scans across independent lineages has been employed extensively in recent studies seeking to test

whether the same regions are involved in independent adaptation events (genetic convergent

evolution) across diverse taxa including birds (2), fish (3–8), insects (9,10), mammals (11), molluscs

(12,13) and plants (14,15).

Following initial excitement in the ability of outlier scans to identify locally adapted genomic regions,

the last 10 years have seen a healthy scepticism develop regarding how indicative genetically

differentiated regions are of natural selection (16). Other genomic processes that illicit

heterogeneity in diversity across the genome, such as background selection (BGS) (17,18), have been

invoked to offer alternative explanations for patterns of genome-wide variation at first attributed to

selection (19). Furthermore, the influence of neutral processes such as genetic drift are similarly

predicted to preclude the informativeness of measures of genetic differentiation if populations are

subject to random differentiation through drift (20).

A major concern arises therefore when applying outlier scans to studies of natural populations, given

the role of often unknown demographic parameters in influencing both BGS and neutral evolution.

For example, the efficacy of selection is dependent on the effective population size (N_e) (21–24),

such that the effects of neutral processes and BGS should be more pronounced in smaller

populations; as has been demonstrated in humans (25) and *Drosophila* (26,27). Additionally,

evidence suggests the effects of BGS are strongest when the interaction between N_e and selection is

intermediate (28). Meanwhile, population sub-division and migration rates influence global

population variance, which has implications for elevating measures of genetic differentiation (29,30)

as sub-populations experience the effects of reduced N_e and increased linkage as discussed above. In

an attempt to mitigate the confounding effects of these processes, some studies advocate the

corroboration of multiple measures of population differentiation and divergence (19,31).

A diverse array of measures of genetic differentiation and divergence have been employed in the

literature, however here we focus on two of the most common measures of relative differentiation

(F_{ST} and π) and a third measure of absolute divergence (D_{XY}) that represents an alternative strategy.

Undeniably, the most commonly used measure is the fixation-index F_{ST} (32,33), which measures the

relative importance of within- and between-population variance. This measure is thus maximised

when genomic regions exhibit the lowest within- and highest between-population variance. F_{ST}

outliers at the right-tail of the distribution are considered candidates for adaptation because they

reflect regions with dramatically different allele frequencies or high substitution rate (large

between-population variance) and/or abnormally low nucleotide variation (low within-population

variance) relative to the rest of the genome. Changes in nucleotide diversity (π) are another

indicator of adaptation, as selection on a beneficial allele limits variation within a population

resulting in selective sweeps (34). Comparisons of the ratio of π between diverging populations

reveal regions under selection as local π is reduced in one population in comparison to the other.

Whilst broadly similar to F_{ST}, changes in π (Δπ) do not discriminate between which copy of a

polymorphism is fixed, such that a substitution between populations is equivalent to a shared non-

polymorphic site. Δπ outliers therefore represent regions of the genome with abnormally reduced π

in either population relative to the rest of the genome. As a measure of absolute genetic divergence,

D_{XY} (35) does not consider the relative frequencies of polymorphisms within populations (19,31). D_{XY}

can be approximated as the likelihood of drawing the same allele from two populations, quantified

as the average number of pairwise differences between sequence comparisons between two

populations. This measure is therefore influenced by ancestral π and the substitution rate, so

outliers highlight regions that are highly variable in the ancestral population (large ancestral π) or

exhibit many substitutions and thus increased sequence divergence.

Because each measure of differentiation quantifies genetic variation between populations in a

slightly different way, each varies in a unique way with demographic variation. Whilst we possess

good predictions for how individual measures are influenced by demography, and subsequently

neutral processes, we know little about how the relationships between measures are affected.

Because different measures are affected differently by demographic history, through its

relationships with neutral processes and BGS (19,36,37), we should expect that their combined

predictive power for regions of the genome experiencing selection is similarly variable across

different demographic histories. In addition, the combining of scans using different measures may

provide unwarranted confidence in results if different measures are susceptible to demography in a

similar fashion.

This complex interplay between selection measures and demography is further exasperated where

studies compare multiple populations to identify convergent genomic evolution. Here measures of

population divergence are used across independent pairs of evolutionary replicates with 'outlier' loci

compared across results (6,14,38). Whilst this strategy has had some success in identifying adaptive

convergent loci, the differential effects of contrasting demographic histories in independent

evolution events, and the subsequent effects on measures of population divergence, have rarely

been considered in how they affect our ability to detect convergent genetic evolution.

We sought to address these concerns by using forward-simulations to investigate the effects of

different demographic histories on the relationships between measures of divergence with

selection. Whilst we focus on the commonly employed method of examining genome-wide SNP

variation between ancestral and derived populations, we should note that other methods are

employed in the literature (e.g. replicated clines, phylogenomics, single population measures e.g.

Tajima's D). We also investigated how demography influenced the overlap of outliers when each

demographic treatment represents an independent evolutionary replicate, as is the case in studies

of convergent genetic evolution. We varied demography through manipulating the number of

founding individuals (bottlenecks), the population size of the derived population (contractions) and

the presence/absence of migration. We simulated the effects of selection on realistic genes

designed from features taken from the guppy genome assembly, a prominent model system for

studies of convergent evolution (6,39). Moreover, we measured genetic divergence at four set time

points through F_{ST}, D_{XY} and $\Delta\pi$ between a founding population and the diverging population to

examine temporal relationships.

Our aim was to investigate the significance of demographic history when employing outlier scans,

and highlight demographic scenarios that are susceptible to 1) Type-I error for individual measures

(outliers under weak/no selection) and 2) Type-III error when employing multiple measures. Type-III

error (accepting a correct result for an incorrect reason) here would occur if common outliers across

measures are attainable irrespective of selection; thus, using evidence from multiple measures to

validate a claim of adaptive significance is prone to error. We sought to answer the following

questions: 1) How do demographic factors influence measures of divergence through time? 2) How

well do measures of divergence identify regions of the genome under strong selection through time

and across demographic factors? 3) How are the different measures of divergence related through

time and across demographic factors? 4) Where do we detect the strongest signals of convergence

(i.e. overlapping outliers using single or multiple measures), and are these consistent with selection?

METHODS

The forward-simulation software *SLiM 3.0* (40) was used to simulate population divergence under

contrasting demographic treatments in a fully factorial design across a founding population of N =

1000 across with a mutation rate based on the guppy genome (41) and scaled 100-fold over three

raw mutation rates ($4.89e^{-5}$, $4.89e^{-6}$, $4.89e^{-7}$) to ensure robustness of results across different, realistic

values of θ ($4N_e\mu$). SLiM employs a classic Wright-Fisher model to simulate populations, in which a

population of diploid hermaphrodites proceeds through generations such that an individual's

contribution towards the next generation is proportional its relative fitness.

Demographic treatments included varying degrees of population contraction relative to the founding

population size (0.01, 0.1, 0.5, 1.0), migration as a proportion of individuals exchanged between

populations (0.0, 0.002) and founding bottlenecks as a number of individuals (2, 20, 100, 1000).

Combining these treatment levels in a fully factorial experimental design generated a total of 32

different independent demographic histories that all genes experienced.

SLiM runs were performed over simulated genes of varying length (in order to ensure robustness

across different genomic elements); with exon number (max = 10), exon lengths, and intron lengths

drawn at random from the guppy genome gene annotation file. Genes were also given 1kb flanking

regions and a divergence selection coefficient (S_σ) drawn at random.

Selection in our model was the fitness consequences incurred through distance from a phenotypic

optimum in a one-dimensional landscape. Intensity of selection was manipulated by modifying the

standard deviation of the normal distribution curve from which the density distribution was

calculated through the "dnorm" function in SLiM. Standard deviations for fitness distributions (S_σ)

were 0.1, 0.5, 1.0, 5.0, and 10.0, with 0.1 representing the steepest fitness peak and 10.0 the

shallowest. All subsequent analyses' standard deviations were 1/x-transformed so that $S_{1/\sigma}$ = 10

represented the strongest selection treatment (Figure 1). Phenotypes were calculated per individual

as the additive phenotypic effects of exonic non-synonymous mutations, which appeared at a rate of

7/3 relative to synonymous mutations (assuming that most mutations in the third base of a codon

do not alter the amino acid). Effect sizes for mutations with phenotypic effects were drawn from a

Gaussian distribution with mean = 0 and σ = 1. The remaining synonymous exon mutations, and

those in introns and flanks, had no effect on fitness. Recombination rate was set at a constant of $1e^{-8}$

across all genomic elements.

Figure 1: Experimental design for simulations. **A)** Selection treatments assigned to genes, illustrated

as the relationship between relative fitness of an individual and its phenotype. Each facet represents

a different fitness landscape modified through editing the standard deviation of the normal

distribution of fitness consequences in the diverging (blue, dashed) population. Facet labels

constitute 1/standard deviation applied. The x-axes represent phenotype as calculated through non-

synonymous mutations and the y-axes represent relative fitness of individuals. The phenotype of the

ancestral population (red, solid) is the same in all treatments (mean = 0, sd = 1), whilst diverging

populations have a diverged phenotypic optimum (mean = 10, sd = S_σ). **B)** Demographic

representation of treatment factors. **C)** Representation of simulation timeline for treatments,

illustrating that all treatments share an ancestral burn-in population before splitting into 32

replicated "ancestral" (solid) and "derived" (dotted) population pairs. Red, dashed lines denote

sampling generations at which F_{ST}, D_{XY} and $\Delta\pi$ are calculated and averaged across the preceding 50

generations. The purpose of this averaging was to achieve a general sense of population

differentiation at four discrete sampling points, such that values represent stable patterns rather

than more stochastic generation to generation variation.

For each simulation, populations were seeded with 1000 individuals and allowed to proceed for a

burn-in period of 5*2*N (10,000) generations to develop stable levels of natural variation under

stabilising selection, adapting the methodology of (36). During this period, burn-in populations

evolved towards a phenotypic optimum of 0, defined as a normal distribution with mean = 0 and σ =

1.0 (Figure 1). Burn-in populations were then subjected to each demographic treatment to simulate

the founding of multiple populations from a shared ancestral state. During this 'divergence period',

the founding population continued to evolve around a phenotypic optimum of 0, whilst the

diverging population's phenotypic optimum was centred around 10 with fitness consequences

defined according to the selection treatment. Individuals also experienced fitness costs associated

with phenotypic proximity to other individuals within the population as a proxy for competition and

to ensure a realistic amount of phenotypic variation persisted within populations. Fitness costs due

to competitive proximity were drawn from a Gaussian distribution with mean = 1, σ = 0.4 and

occurred reciprocally between local individuals with phenotypes with a difference of ≥ 1.2 (3 * 0.4).

Each gene was then run across all demographic treatment levels in parallel. Simulations were

sampled at 0.05*2*N (100), 0.158*2*N (316), 1.581*2*N (3162) and 5*2*N (10,000) generations

(Figure 1C). At each sampling point, F_{ST} (calculated according to (33)), D_{XY} (calculated according to

(35)), and mean heterozygosity (π) of each population were calculated as the average across the

preceding 50 generations. Averaging was performed to gauge a general sense of differentiation

around the four sampling points, such that measures would not be dramatically biased by events

occurring within individual generations. The change in mean heterozygosity between the founder

and divergent population (Δπ) was calculated at each sampling point as the $\log_{10}$-transformed ratio

of founder π to divergent π, such that reduced diversity in the diverging population increases the

value of Δπ. Statistics were calculated over all sites, akin to assessing differentiation over a gene or

chromosomal interval.

214
$$F_{ST} = 1 - \frac{H_S}{H_T}$$

216
$$D_{XY} = \sum_{ij} x_i y_j d_{ij}$$

218
$$\Delta\pi = \log_{10} \frac{\pi_{ancestral}}{\pi_{diverged}}$$

In total, ~1,000 genes were simulated across 32 demographic treatments across three mutation

rates. In addition, a fourth set of simulations were performed in which the phenotypic optimum for

both populations was kept at 0 (Pheno$_{Null}$). This dataset was used to assess whether patterns

associated with selection were driven by phenotypic divergence or variable stabilising selection

within populations. Sets of genes were generated randomly for each mutation rate. Outliers for gene

sets were taken as upper quantiles (either the upper 1% or 5% depending on analysis, see below).

RESULTS

Demography modifies measures of divergence

Bottlenecks, simulated by modifying the number of founding individuals, had little effect on F_{ST} and

D_{XY}, producing minimal variance between treatments over all sampling points (Figure 2A).

Conversely, there was a relatively pronounced effect on $\Delta\pi$ that persisted for several hundred

generations. Genes that were simulated with the most extreme bottlenecks displayed inflated

measures of divergence because of a sharp reduction in π in the diverging population.

Contractions, simulated by modifying the stable number of individuals within the diverging

population, had pronounced and variable effects on all measures. F_{ST} was negatively associated with

contractions, with values becoming inflated with any reduction in diverging population size. This

effect was generally consistent through time, however early sampling points exhibited the largest

variance between the most extreme contraction treatments, whereas later on the strongest

variance was observed between weaker treatments (Figure 2B). A similar effect was observed for

$\Delta\pi$; however, this measure was particularly susceptible to inflation under the most extreme

contraction scenarios, with substantially more elevated values observed between contraction

proportions of 0.01 and 0.1 compared with 0.1 and either 0.5 or 1.0. Whilst this pattern was broadly

consistent through sampling points, variance was most pronounced at the initial sampling point (100

generations). Such observations are unsurprising given that both F_{ST} and $\Delta\pi$ increase with any

process that reduces within-population genetic variance. D_{XY} was largely unaffected by contraction

treatments up until the third sampling point (3162 generations). Following this, in contrast to F_{ST} and

$\Delta\pi$, D_{XY} decreased when diverging population sizes were reduced (Figure 2B).

The inclusion of migration expectedly reduced absolute values of all measures of divergence. For F_{ST}

and D_{XY}, variance between migration treatments increased across the scope of the simulation,

however D_{XY} was generally more resilient to variance induced through migration. $\Delta\pi$ was also

reduced in the presence of migration, although the effect of migration on Δπ was generally

consistent across all generations and did not increase through time, as was the case for F_{ST} and D_{XY}.

Unsurprisingly, there was a consistent trend regardless of demographic treatment for F_{ST} and D_{XY} to

increase generally over time, however this was not the case for Δπ, which was generally constant

through time whilst marginally declining. The majority of these patterns were apparent in the

Pheno$_{Null}$ simulations (Figure S3), however genetic differentiation for all measures was reduced by

the removal of divergent phenotypes when gene flow was present.

Figure 2: Effects of demographic treatments on measures of genetic divergence across all sampling

generations. Point colour and shape denote treatment groups for bottlenecks (**A**), population

contraction (**B**) and migration (**C**). Each point represents values of divergence averaged across all

genes, and all other treatment levels beyond those of the treatment group of interest.

Demography moderates the association between measures of divergence and selection

To assess the effect of demographic treatments on correlations with $S_{1/\sigma}$, we subsetted our data

within each treatment level for each demographic factor (e.g. subsetting contraction groups of 0.01,

0.1, 0.5 and 1.0) and calculated Pearson's correlation coefficients across all treatments with the

treatment level of interest (such that for each population contraction level, 8 treatment groups were

included and averaged over). Bottlenecks had a minimal effect on the correlations observed

between strength of selection and measures of divergence with minimal variance observed between

bottleneck treatments in all comparisons (Figure 3A).

Population contractions had significant effects on relative (F_{ST} and $\Delta\pi$) measures and marginal

effects on absolute (D_{XY}) measures of divergence (Figure 3B). Relative measures in smaller diverging

populations (0.01, 0.1) were less informative, and indeed largely uninformative, of selection than

those observed in treatments with minimal (0.5) or no (1.0) contractions (Table 1). For F_{ST} the

variance in correlation coefficients between contraction treatment groups generally increased

across time whereas variance between treatments for $\Delta\pi$ peaked at 3,162 generations. For D_{XY},

correlations with selection were broadly resistant to contraction-induced variance up until 10,000

generations, in which associations with selection were reduced by stronger contractions (Figure 3B).

Expectedly, the absence of migration largely precluded the ability of relative measures of divergence

to predict strength of selection, with notable variance observed between no migration (M= 0.0) and

minimal migration (M = 0.002) treatments by 3,162 generations for all divergence measures (Figure

3C) (Table 1). Both F_{ST} and D_{XY} variance between migration treatments was greatest at the 10,000

generations sampling point, whereas similar variance was observed between migration treatments

for $\Delta\pi$ at both 3,162 and 10,000 generations. This observation again highlights the significance of

temporal effects on F_{ST} and $\Delta\pi$.

Pheno$_{Null}$ simulations resulted in weaker correlations with $S_{1/\sigma}$ for F_{ST}, whilst D_{XY} became negatively

correlated with $S_{1/\sigma}$ over time (Figure S6B) rather than positively correlated. $\Delta\pi$ became more

positively correlated with $S_{1/\sigma}$ over time for all demographic treatments (Figure S6C). These results

demonstrate that increases in F_{ST} are partly dependent on divergent selection, increases in D_{XY} are

exclusively dependent on divergent selection (and adaptive substitutions), and $\Delta\pi$ increases are less

indicative of divergent selection and more of stabilising selection and within-population variance.

Table 1: Correlation coefficients between measures of divergence and selection, separated according to individual treatment levels across all sampling generations. For each measure, the largest correlation coefficient is highlighted in bold.

Treatment	Level	F_{ST}				D_{XY}				$\Delta\pi$			
		100	316	3,162	10,000	100	316	3,162	10,000	100	316	3,162	10,000
Bottleneck	2	-0.006	0.040	0.129	0.169	0.016	0.019	0.033	0.095	0.006	0.083	0.148	0.072
	20	0.013	0.051	0.140	0.165	0.031	0.026	0.043	0.093	0.010	0.097	0.137	0.071
	100	0.010	0.043	0.147	0.177	0.024	0.022	0.050	0.102	0.009	0.084	0.138	0.074
	1000	0.007	0.040	0.133	0.165	0.020	0.021	0.041	0.097	0.019	0.079	0.149	0.070
Contraction	0.01	0.015	0.019	0.033	0.053	0.030	0.034	0.039	0.052	0.004	0.008	0.006	-0.024
	0.1	0.000	-0.001	0.087	0.157	0.015	0.006	0.020	0.110	0.034	0.040	0.189	0.122
	0.5	0.009	0.078	0.269	0.304	0.025	0.027	0.063	0.147	0.013	0.168	0.378	0.214
	1	0.000	0.136	0.306	0.355	0.022	0.020	0.051	0.138	0.031	0.260	**0.406**	0.245
Migration	0	0.003	0.029	0.074	0.062	0.020	0.014	0.006	-0.015	0.018	0.081	0.094	0.007
	0.002	0.012	0.101	0.335	**0.429**	0.027	0.033	0.092	**0.223**	0.003	0.105	0.241	0.191

Figure 3: Effects of demographic treatments on the relationship between selection ($S_{1/\sigma}$) and
measures of genetic divergence across all sampling generations. Point colour and shape denote
treatment groups for bottlenecks (**A**), population contraction (**B**) and migration (**C**). Each point
represents correlation coefficients averaged across all genes, and all other treatment levels beyond
those of the treatment group of interest.

By examining the correlation coefficients of all 32 unique demographic histories, we can investigate
the combined effect of migration and contraction and directly compare effectiveness of individual
measures across time (Figure 4). F_{ST} consistently outperforms D_{XY} in terms of associating with $S_{1/\sigma}$
under most demographic treatments, particularly when contractions are weak and migration is
present. By 10,000 generations however, the relative dominance of FST appears to subside, with
both F_{ST} and D_{XY} performing as well as each other in the presence of migration (Figure 4). $\Delta\pi$ is
similarly more informative than D_{XY} across sampling generations under most demographic
treatments. Interestingly, at sampling points 316 and 3,162 contractions produce the biggest bias
towards $\Delta\pi$, however by 10,000 generations zero migration produces the largest deviance away
from the y=x relationship (Figure 4). The resilience of $\Delta\pi$ under no-migration treatments is also
apparent in F_{ST} - $\Delta\pi$ comparisons, such that at 316 and 3,162 generations $\Delta\pi$ is more informative

than F_{ST} in the absence of migration. By 10,000 generations however F_{ST} outperforms $\Delta\pi$ under most

scenarios, indicative of the decaying association between $\Delta\pi$ and $S_{1/\sigma}$ discussed previously.

Figure 4: Pairwise comparisons between the correlation coefficients of $S_{1/\sigma}$ with F_{ST}, D_{XY} and $\Delta\pi$

across all sampling points. Each data point represents a unique demographic treatment with points

coloured according to contraction level and shaped according to migration level. Correlation 1 refers

to the first measure listed in the comparison and Correlation 2 to the second. The y=x line is plotted

within each facet to illustrate biases towards one measure. Points below the line are biased towards

Correlation 1, whilst points above the line are biased towards Correlation 2.

Demography moderates the tail end of divergence distributions

Ultimately, outlier tests are concerned with the extreme tail end of the distribution, and as such

correlation coefficients across the whole dataset may fail to capture consistency in the tail ends. To

counteract this, we asked at each sampling point what the relative proportions of outliers above the

99th quantile were per $S_{1/\sigma}$ treatment, and how these proportions varied according to individual

demographic treatments (Figure 5) and collectively (Figure 6) after 10,000 generations (results for

other sampling points and mutation rates Figure S10-20). Each simulated 'genome' consisted of

~1000 genes evenly distributed across the 5 treatment levels of $S_{1/\sigma}$, meaning expected null

proportions of each $S_{1/\sigma}$ value are 0.2. This analysis assumes that if measures of differentiation are

good at highlighting genes under the strongest selection coefficients, genes within the upper 1% of

the distribution should be disproportionately associated with stronger selection ($S_{1/\sigma}$ = 2 or 10). We

found that migration and contraction treatments influenced the relative proportions of $S_{1/\sigma}$

treatments in the upper F_{ST} quantile. Under the weakest contraction treatment, the proportion of

$S_{1/\sigma}$ = 10 outliers increased from 0.19 (contraction = 0.01) to 0.33 (contraction = 1.0) (null

expectation of 0.20 given 5 selection treatments). Migration also dramatically increased the

proportion of genes under the strongest selection from 0.19 to 0.63. The extreme effect of migration

is evident when combining contraction and migration treatments (Figure 6). Whilst weaker

contractions yield a greater proportion of strong selection outliers (moving left to right in the figure),

migration is the dominant factor given that the treatment with the most extreme contraction (0.01)

and migration present has a stronger selection bias than the treatment with no contraction and

migration absent (Figure 6).

The upper quantile of D_{XY} outliers was unaffected by bottlenecks and contractions but was heavily

biased towards $S_{1/\sigma}$ = 10 genes in the presence of migration (Figure 5). Proportions of $S_{1/\sigma}$ = 10

outliers doubled in the presence of migration from 0.27 to 0.54. Proportions across the 16 unique

contraction/migration patterns produced results analogous to F_{ST}, with migration the dominant

factor in biasing proportions towards strong selection genes (Figure 6).

Except for the most extreme population contraction, $\Delta\pi$ outlier proportions were strongly indicative

of genes under strong selection, yielding outlier proportions for $S_{1/\sigma}$ = 10 genes of 0.72 (contraction =

0.1), 0.61 (contraction = 0.5) and 0.58 (contraction = 1.0) at sampling generation 10,000. Effects of

contractions were visible for $\Delta\pi$ outliers by generation 100 at the most extreme treatments (Figure

S10) and clear across all 3 weaker contraction treatments by 3162 generations (Figure S12).

Migration had a more modest effect on outlier proportions, increasing the proportion of genes

under the strongest selection from 0.16 to 0.41 at sampling generation 10,000. This modest effect of

migration was not as persistent through time however, emerging only by generation 3,162 with $S_{1/\sigma}$

=10 proportions of 0.18 (migration = 0) and 0.24 (migration = 0.002). These results are corroborated

across the 16 contraction/migration treatments, as proportions acquired from treatments with

extreme contractions (contraction = 0.01) are relatively uniform with only a small biasing effect of

migration (Figure 6). For all other treatments, outliers are heavily biased towards strong selection

genes.

Taken together, these results highlight that outlier detection using F_{ST} and D_{XY} are heavily dependent

on the presence of migration, and to a lesser extent the absence or minimising of population

contractions for F_{ST}. $\Delta\pi$ outliers appear rapidly and are generally robust in relation to population

contractions up until the most extreme treatment. Whilst migration also influences $\Delta\pi$, the biasing

of outliers towards genes under the strongest selection treatment is less extreme.

Removing divergent selection (Pheno$_{Null}$) moderated outlier proportions in line with expectations

given correlations with $S_{1/\sigma}$ (Figures S21-24). Interestingly, over time both F_{ST} and $\Delta\pi$ outliers were

skewed further towards stronger selection treatments when migration was absent (stronger

patterns of reduced within-population variance) in comparison to simulations with phenotypic

divergence, whilst D_{XY} outliers became biased towards weaker selection treatments in the presence

of gene flow.

394
395

396   **Figure 5:** The effect of demography on representation of genes under different selection treatments

397   as a proportion of outliers above the 99% quantile. Colour of bars within stacks denotes different

398   selection treatments. Vertical facets represent different demographic treatments and horizontal

399   facets show effect per measure of genetic divergence. Results shown are from sampling generation

400   10,000. Results for other sampling points can be found in Figures S10-12.

Figure 6: The combined effects of contractions (proportion relative to founding population size = 1,000) and migration in 16 unique demographic treatments (all combinations of contraction and migration treatments) on representation of genes under different selection treatments as a proportion of outliers above the 99% quantile. Full information regarding demographic treatment is described on the x-axis. Colour of bars within stacks denotes different selection treatments. Horizontal facets show effect per measure of genetic divergence. Results shown are from sampling generation 10,000.

Demography moderates associations between measures of divergence

Given F_{ST}, D_{XY} and $\Delta\pi$ are all measures of population genetic divergence, there is an assumption that positive associations should exist between them. We employed the same analysis as above for correlations with $S_{1/\sigma}$, but instead examined correlations between individual measures. Bottlenecks

had minor effects on the correlations observed between measures of divergence (Figure 7A).

Expectedly, effects were strongest at the first sampling point, with the most extreme bottlenecks

increasing the correlation coefficients between F_{ST} & D_{XY} and D_{XY} & $\Delta\pi$ and decreasing the correlation

coefficient between F_{ST} & $\Delta\pi$. Variation in correlation coefficients was generally minimal however

compared to the effects of other demographic treatments.

F_{ST} & D_{XY} were positively associated irrespective of demography, but smaller population sizes

generally increased the association (Figure 7B) with variance between treatments generally

decreasing through time. Similarly, F_{ST} & $\Delta\pi$ were typically positively associated, however

contractions had a more pronounced effect on the association. Extreme contractions increased the

association between F_{ST} & $\Delta\pi$, with contraction-induced variance increasing across sampling points

(Figure 7B). At 10,000 generations when variance was largest, F_{ST} & $\Delta\pi$ ranged from being generally

well associated under extreme contractions ($R^2 = 0.60$) to unassociated when populations were

equally sized ($R^2 = 0.08$). D_{XY} & $\Delta\pi$ correlations were initially weak early in the simulations, however

increased generally over time, with populations contractions similarly inflating associations between

measures. Interestingly, associations with D_{XY} and both F_{ST} and $\Delta\pi$ appear to peak for weak

contraction treatments at 3,162 generations, suggesting a probable temporal factor within the

simulations driving positive associations with D_{XY} at some point between 316 and 3,162 generations.

Migration induced substantial variance between correlations of divergence measures, with effects

dependent on sampling point (Figure 7C). In the absence of migration, F_{ST} & D_{XY} were strongly

associated at 100 and 316 generations, whilst associations were weaker in migration-present

treatments. However, by 3,162 generations F_{ST} & D_{XY} associations increased substantially in the

presence of migration, and by 10,000 generations the effect of migration had reversed entirely. F_{ST} &

$\Delta\pi$ associations were strong initially, but migration weakened the association over time so

migration-present associations ranged from $R^2 = 0.67$ (100 gens) to $R^2 = 0.14$ (10,000 gens). In

contrast, in the absence of migration positive associations between F_{ST} & $\Delta\pi$ were relatively stable

($R^2 = 0.55 – 0.51$ over the same time period). Interestingly, without migration D_{XY} & $\Delta\pi$ were largely

unassociated, but migration-present treatments induced a negative association between D_{XY} & $\Delta\pi$

over time from $R^2 = -0.04$ (100 gens) to $R^2 = -0.28$ (1000 gens).

Contextualised by our previous demonstrations of associations with selection, these results highlight

that selection can induce positive associations between measures. Namely, F_{ST} & D_{XY} become

strongly associated by generation 10,000 with migration present, which we know is when F_{ST} and D_{XY}

are most strongly linked to selection. Crucially, however this relationship only emerges after several

thousand generations, before which we observe positive associations in migration-absent

treatments when associations with selection are weak for both measures. Extreme contractions also

induce positive associations between F_{ST} & D_{XY} that are likely indicative of false-positives given the

poor association with selection in these treatments. These results highlight that positive associations

between statistics are also achievable under neutrality. It is also interesting to note the decay of

associations with $\Delta\pi$ and both F_{ST} and D_{XY} under demographies in which selection should be

effective. These observations are most likely driven by the substitution rate. Substitution rates are

increased under stronger selection (42) and drive increased F_{ST} and D_{XY} but not $\Delta\pi$.

Figure 7: Effects of demographic treatments on the relationship between measures of genetic

divergence across all sampling generations. Point colour and shape denote treatment groups for

bottlenecks (**A**), population contraction (**B**) and migration (**C**). Each point represents correlation

coefficients averaged across all genes, and all other treatment levels beyond those of the treatment

group of interest.

In Pheno$_{Null}$ simulations the influence of bottleneck treatments and contraction treatments was

broadly unchanged, however the notable emergence of a correlation between measures of F_{ST} and

D_{XY} in simulations with gene flow failed to emerge by generation 10,000 (Figure S27). This

demonstrates that this positive correlation is driven by divergently adaptive allele frequency shifts

and substitutions. In addition, the decay of the relationship between F_{ST} and $\Delta\pi$ in the face of

migration did not occur without divergent selection. Rather, this relationship remained strongly

positive ($R^2 > 0.5$) up to 10,000 generations regardless of migration treatment (Figure S27C).

To examine the overlap of outliers across measures of divergence, we again sub-divided our

treatments into treatment-specific levels and compared the three-way and all pairwise upper 95%

quantile gene overlaps across measures of divergence at sampling point 10,000. Three-way

agreement was generally greater than expected by-chance under most conditions, with the largest

overlap observed under extreme contractions (Figure 8). Our results demonstrate large overlap

between F_{ST} and D_{XY} outliers across all treatments, which was marginally increased by contraction

treatments that favour the effectiveness of selection. This association suggests stronger selection

may improve the positive association between F_{ST} and D_{XY}, however a causative effect is not clear.

Whilst overlap increased by selection-favouring conditions, all treatment levels exhibited above-

expected levels of F_{ST} - D_{XY} overlap, suggesting a strong potential for false-positives. Overlap

between $\Delta\pi$ and both F_{ST} and D_{XY} was substantially less but had contrary associations with

contractions. More extreme contractions tended to increase the overlap between $\Delta\pi$ and F_{ST}

outliers, suggesting a shared risk of false-positives. In contrast, $\Delta\pi$ and D_{XY} outlier overlap tended to

increase under weaker contractions, suggesting a beneficial effect of combining these measures to

remove false-positives. Migration increased outlier overlap between $\Delta\pi$ and both F_{ST} and D_{XY}.

Figure 8: Proportional overlap of 95% outliers across measures of divergence, separated according to treatment levels of bottlenecks (**A**), contractions (**B**) and migration (**C**).

Demography drives signals of convergence irrespective of selection

To assess the convergence of outliers across demographic treatments, we downsampled the dataset such that we retained a random 10% of $S_{1/\sigma} = 10$ and a random 50% of $S_{1/\sigma} = 2$ genes per iteration. This step was taken to try to reduce the proportions of genes under the strongest selection treatment within the dataset from their initial ~20% proportions to a more realistic ~2% and ~10% respectively. Downsampling was bootstrapped 100 times, with the proportional overlap between demographic treatments recorded per iteration and averaged across all runs. Following

502 downsampling, we examined proportional overlap of the two 95% outlier sets with an expected

503 proportional overlap of 0.05 under a null hypothesis.

505 By sampling generation 10,000, proportion of overlap in outlier genes was apparent in F_{ST} results

506 across all demographic treatments that had no migration (Figure 9). Within this cluster, the overlap

507 of outliers above the 95% ranged between 0.07 and 0.32, suggesting a 1.4- to 6.4-fold enrichment

508 above the null expectation. These minimum and maximum values were observed between extremes

509 of contraction (minimum) or when contractions were absent (maximum). Given the poor association

510 between F_{ST} and selection in the absence of migration, convergent F_{ST} outliers here are likely driven

511 by neutral processes and would constitute a false-positive signal. Outside of this cluster, there was

512 also reasonable clustering of results across treatments that included migration and weaker

513 contractions (0.1, 0.5 or 1.0). Here, overlap of outliers ranged between 0.08 and 0.26, representing a

514 1.6- to 5.2-fold enrichment above the expected. In the presence of migration and minimal

515 contractions F_{ST} is a good predictor of selection in our dataset, suggesting this convergence may be

516 driven by inferred selection. Beyond these clusters, overlap between treatments of contrasting

517 migration treatment failed to exceed 0.18. Interestingly, the clustering of F_{ST} outliers in the presence

518 of migration was heavily reduced when divergent selection was removed ($Pheno_{Null}$) but clustering of

519 outliers in the absence of migration was still strong (Figure S39).

Figure 9: The proportional overlap of outliers above the 95% quantile for F_{ST} values, averaged across 100 randomly downsampled datasets for each of the 32 demographic treatments. Colouring of axis labels denotes migration treatment level.

Clustering of D_{XY} outliers followed a broadly similar pattern to F_{ST} with migration treatments dominating the clustering of treatments and inducing proportional overlaps of 0.11-0.34 (2.2- to 6.8-fold enrichment) in the absence of migration (Figure S31). In general, proportional overlap of Δπ outliers was lower than for F_{ST} and D_{XY}, with a maximum overlap observed of 0.21 (fold-enrichment 4.2) (Figure S32). This value was observed in two treatments with minimal contractions (0.5-1.0) and migration present. Generally, however, clustering by demography was weaker in Δπ than for F_{ST} and D_{XY}, corroborating previous results suggesting Δπ may be less susceptible to variation in migration and contraction (with the exception of extreme contraction).

Because migration was the dominant factor in clustering of treatments with convergent overlaps, we

sought to investigate what features of simulated genes drove variance in measures of divergence

with treatments separated by migration factor using a Random Forest approach (Figure 10). For F_{ST},

we observe that the best predictor of variance in the absence of migration is diversity (π) of the

diverging population whilst treatments with migration are most strongly influenced by $S_{1/\sigma}$. This is

consistent with the notion of F_{ST} relying on within-population genetic variance in the absence of

efficient selection. In contrast, D_{XY} in the absence of migration is most strongly driven by gene

features, predominantly the proportion of simulated gene that is coding and by the length of the

simulated gene. With migration, D_{XY} variance is driven by selection and diverging π. Unsurprisingly,

$\Delta\pi$ is also strongly influenced by diverging π (the denominator in its calculation), but we can see a

secondary role for selection in treatments with migration and little influence of gene features.

Removing divergent selection (Pheno$_{Null}$) did not modify the relative importance of any features of

simulations for F_{ST}, but did affect D_{XY} and $\Delta\pi$ (Figure S44). With migration, the relative importance

for diverging population π in explaining D_{XY} variance increased substantially. For $\Delta\pi$, selection was as

important as diverging population π in explaining variance.

553
554
Figure 10: The relative importance of simulated gene features in eliciting variance in measures of

genetic variance. Boxplots denote the relative importance of each explanatory variable across 16

individual treatments with/without migration taken at sampling generation 10,000. Variables

considered are within-population π of the diverging population and gene features including: Number

of Exons (Exon N), Gene Size, and the proportion of gene that is coding (Target Proportion). Also

included are the selection ($S_{1/\sigma}$) imposed on the gene and the generation at which the optimum

phenotype is reached (Pheno Gen).

DISCUSSION

Summary of results

Here, we show that differing demography can have dramatic effects on how measures of population

divergence identify regions of the genome under selection. Effects are also strongly time-dependent.

Using simulated populations, we have demonstrated the relative influences of founding bottlenecks,

population contractions, and migration on three commonly used measures of genetic divergence

(F_{ST}, D_{XY}, $\Delta\pi$), whilst demonstrating the relative usefulness of each measure for informing on

selection under different demographic conditions.

We find that founding bottlenecks have little effect on population divergence measures, with only

minor influence over differences in nucleotide diversity ($\Delta\pi$) shortly after founding. Population

contractions artificially inflate F_{ST} and $\Delta\pi$ but reduce D_{XY} and can erase the relationship of F_{ST} and D_{XY}

with selection under the most extreme contractions. These results can be explained in part due to

the relationship between founding population size and relative (F_{ST} and $\Delta\pi$) and absolute (D_{xy})

divergence measures. Relative measures are, in part, driven by intra-population changes in allele

frequency, which become exaggerated in smaller populations as a product of drift (23,43). As a

consequence, we observe inflated measures of relative divergence as allele frequencies drift in our

smaller diverging population.

D_{xy} in contrast increases under weaker population contractions as a product of the relationship

between the number of segregating sites and the population-level mutation rate ($4N_e\mu$) (44). D_{xy} is a

measure of sequence divergence and is averaged across all sites (although averaging across all

segregating sites is possible), which results in higher D_{XY} as segregating sites are introduced into

either population at rate of $4N_e\mu$. This relationship with the number of segregating sites can be

observed by examining the positive relationships between D_{xy} and π in our diverging population

(Figure S45). The absence of a strong contraction effect in moderating D_{xy} associations with $S_{1/\sigma}$ likely

stems from the lack of allele frequency relevance. Consider for example, two SNPs with minor allele

frequencies of 0/0.5 (SNP 1) and 0.5/0.5 (SNP 2) in the founding/diverging population. Each locus

contributes equally towards D_{xy} (SNP 1 = [0 x 0.5] + [1 x 0.5] = 0.5; SNP 2 = [0.5 x 0.5] + [0.5 x 0.5] =

0.5), whereas the reduction of within-population variance observed for SNP 1 inflates F_{ST} and $\Delta\pi$.

Migration, even at the relatively modest rate of 0.2% employed here, reduced absolute values for all

measures of divergence substantially. However, whilst absolute values were reduced, their

informativeness of selection coefficients increased dramatically. Such a result is expected given the

role of migration and gene-flow in homogenising neutral loci (reducing measures of divergence),

whilst retaining population divergence around adaptive loci (increasing informativeness). This

dynamic is often invoked within the population 'genomic islands of divergence' model to explain

patterns of heterogenous genomic divergence in studies of speciation-with-gene-flow (45,46).

Migration exhibited interesting temporal patterns that are useful for discussing the discrepancies

observed between $\Delta\pi$ and both F_{ST} and D_{xy}. Of the measures considered here, $\Delta\pi$ uniquely

disregards SNP substitutions in its calculation, as fixed substitutions have no influence on intra-

population heterozygosity beyond reducing the heterozygosity of proximate SNPs in the aftermath

of a hard sweep. This characteristic explains why $\Delta\pi$ responds to selection more rapidly than F_{ST} and

D_{xy} and is influenced by migration in a constant manner across time. In addition, a failure to account

for adaptive substitutions in $\Delta\pi$ is the most probable explanation for $\Delta\pi$ informativeness peaking at

3,162 generations and decaying by 10,000 generations. In contrast, the contributions of adaptive

substitutions to F_{ST} and D_{xy} over time increases their predictive power. This temporal observation

has important implications for studies of rapid adaptation on the scale of 10s to 100s of generations,

in which our simulations suggest $\Delta\pi$ may be the most informative measure of divergence, whilst D_{xy}

was largely uninformative until > 3,162 generations.

A novel element to our simulations involved examining the effects of demographic variation on the

associations between measures of divergence themselves. The relative weaknesses of individual

measures of divergences has prompted a recent movement within the literature to employ multiple

measures of divergence to avoid false-positives (19). Our results provide some support for this

recommendation, with outlier overlap across measures generally increasing under conditions that

favour informativeness of selection, suggesting their combined use may be useful in removing false-

positives. Importantly, we also find that F_{ST} and D_{xy} (and to a lesser extent F_{ST} and $\Delta\pi$) highlight

convergent outliers across demographic treatments regardless of the effectiveness of selection.

Cruickshank and Hahn (19) suggested that a failure of F_{ST} to exhibit similar patterns to D_{xy} in

genomic-islands-of-divergence tests highlighted a particular susceptibility of F_{ST} to BGS (although see

(36)). Our results are in line with the notion that D_{xy} is more resistant to BGS given that contraction

treatment levels resulted in minimal variance in D_{xy} (assuming treatment levels of population

contraction are analogous to different rates of BGS across the genome in exhibiting reduced N_e). Our

results add support to the emergence of a positive association between F_{ST} and D_{XY}, but only in the

presence of gene flow between populations and only after several thousand generations, as has

been previously simulated (47). The latter dependence on time is due to the delayed predictive

power of D_{xy}.

However, our results also demonstrate that F_{ST} - D_{xy} associations emerge under demographic

conditions where neither is well correlated with selection, and that this relationship decays over

time in the absence of migration (Figure 7C). No-migration treatments are consistent with Isolation-

By-Distance demographic histories and similarly inter-specific comparisons. With regards to the

latter, empirical observations of negative correlations between F_{ST} and D_{XY} within and between

clades of birds (48,49) and within a radiation of monkeyflowers (50) support the simulated

prediction of an eventual negative relationship between these measures. This negative association

in the absence of effective selection is predictable, given F_{ST} is negatively associated with local π

(increases within-population variance), D_{XY} is positively associated with local π (reduces likelihood of

common sequence) and π is generally well-conserved in stable populations even across long time

periods (51–53).

Similarly, F_{ST} – $\Delta\pi$ outliers were found under most demographic scenarios and marginally increased

under extreme contractions. These results support the notion that whilst the use of multiple

measures may be useful in reducing Type-I error, there is an inherent risk associated with Type-III

error (accepting true-positives for the wrong reason) given the risk of Type-I error under certain

demographic histories.

In comparing our results to a second dataset that lacked divergent selection (Pheno$_{Null}$), we find

consistent support that patterns associated with D_{XY} are strongly dependent on divergent selection;

F_{ST} patterns are attainable with only stabilising selection (but are weaker than when divergent

selection is included); and $\Delta\pi$ patterns are primarily driven by variable stabilising selection and made

weaker by the inclusion of phenotypic divergence. These comparisons are useful in informing

specifically on the role of adaptive allele frequency changes and substitutions in driving patterns of

genetic differentiation. In particular, our observation that positive F_{ST} - D_{XY} correlations emerge only

with effective, divergent selection in the face of gene-flow will be informative in assessing the

usefulness of this combination of statistics in studies of adaptation.

Application of results

We sought to utilise our experimental design to investigate the effects of variable demography in

studies of convergence. Our simulation design, in which an ancestral burn-in population is used to

found 32 independent pairs of ancestral – diverging populations is analogous to the commonly

employed sampling method of examining replicated ecotype population pairs in model systems,

such as various ecotype pairs of the three-spined stickleback, *Gasterosteus aculeatus* (3,4); high/low

predation Trinidadian guppies, *Poecilia reticulata* (6); crab/wave ecotype periwinkles, *Littorina*

saxitilis (12,13,54), and alpine/montane ecotypes of *Heliosperma pusillum* (14). By overlapping

outliers called from one demographic treatment with others, we demonstrate that signals of

convergent genes (proportions above a random null expectation) are attainable for F_{ST} and D_{XY}

across multiple replicates, with clustering predominantly driven by the presence or absence of

migration (with minimal convergence between clusters). Given that within our simulations selection

is expectedly ineffective when migration is absent, convergence of outliers across migration-free

treatments suggests a different subset of genes within our analysis, which are not under strong

selection, are repeatedly reporting the strongest signals of genetic divergence for both F_{ST} and D_{XY}.

In seeking to explain convergence patterns associated with migration, we found that for both F_{ST} and

D_{XY} there were different sources of variance depending on migration treatment. F_{ST} was heavily

influenced by within-population genetic diversity (π) in the absence of migration. This is consistent

with previous theoretical and empirical work highlighting the susceptibility of F_{ST} to processes that

reduce local population diversity disproportionately across the genome, such as heterogeneity in

recombination landscape (22,47), structural variation (55), and gene density (22,56) in the absence

of effective selection. With migration present, and effective selection, we find that convergence is

most probably driven by selection in our simulations. These results therefore demonstrate

convergence within demographically similar pairs, and by extension non-convergence across

demographically dissimilar pairs, is only selection-dependent under certain conditions. Without

migration, D_{XY} convergence was achieved through features of simulated genes, predominantly

length of gene and proportion of gene that was coding. Such an effect may be explained by the

expectation of reduced neutrality of mutations within coding regions, producing a consistent effect

across demographic treatments for genes with increased coding content to exhibit reduced neutral

polymorphism (across both the ancestral and diverging population) and lower D_{XY} as a result. The

secondary effect of gene length is less clear, given that D_{XY} is averaged across all sites, but may stem

from potentially lower levels of linked selection in larger genes (43) or the increased accumulation of

segregating sites in larger genes (given mutation occurs with a fixed probability per site). With

effective selection in the presence of migration, we observe the expected effect of simulated $S_{1/\sigma}$ in

producing convergent D_{XY} outliers. Interestingly however, we also observe a fairly strong effect of

diverging population π that hints towards a susceptibility to BGS even in the presence of effective

selection. This dependence on the efficacy of selection may explain the discrepancies within the

current literature regarding the susceptibility of both F_{ST} (19,29,57) and D_{XY} (36) to BGS.

It is perhaps surprising that population contractions (with the exception of the most extreme factor

level) did not have a dominant influence on our convergence matrix, given the strong effect on

population contractions in reducing the efficacy of selection. However, this finding is consistent with

the increased influence of random drift in populations with reduced N_e, thereby limiting

repeatability of outliers.

In addition, in our analyses we group genes within 32 unique demographic treatments, assuming the

effects of contraction and migration are equivalent for all genes. This may be unrepresentative of

genomes sampled from the wild, in which gene flow and effective population size can vary across

genomic regions through structural variation, variable recombination rate and BGS as previously

discussed. A prominent example of such a mechanism is the well-characterised consequence of

reduced gene-flow within inversions that carry locally adapted alleles. Assuming inversion variants

are fixed between populations, migration across the locus is limited by the prevention of

recombinant haplotypes and resistance to introgression (47,55). Recent genome scan studies have

highlighted convergent outliers within inversions (58–60), confirming theoretical models regarding

the role for chromosomal rearrangements in shielding adaptive haplotypes from introgression

during the adaptation process. Our simulations highlight that measures of divergence across

inversions carrying locally adapted variants should behave according to reduced migration, and

reduced N_e as a consequence, with inflated measures of divergence and differentiation relative to

other genomic regions. This is predicted to lead to their over-representation in genome scan

outliers, thus caution should be taken regarding the adaptive significance of these outliers relative to

absolutely lower values of genetic divergence attained from regions outside of chromosomal

rearrangements.

Limitations of simulations

The factorial design of our study represents a substantial progression in understanding the interplay

between individual features of demographic history and multiple measures of population

divergence. However, like all models it is limited. Computational limitations constrained absolute

population sizes to a maximum of 1,000 individuals and generations to a maximum of 10,000. To

mitigate these constraints, we repeated the analysis over multiple mutation rates to illustrate

patterns over 100-fold variation of θ. In general, most trends were consistent, suggesting that

results should be consistent across taxa of variable effective population size. However certain

patterns are exaggerated, for instance it is clear that the time lag on informativeness of D_{XY} on

selection is shortest when θ is largest, which results in stronger positive associations between F_{ST}

and D_{XY} by generation 10,000. It is well documented that both N_e (61) and mutation rate (μ) (62) are

highly variable across taxa, which suggests that applying knowledge of the relationships between

measures of population differentiation will vary by system.

Furthermore, temporal variation within our simulations is confounded by the time at which major

shifts in the diverging population's phenotype (Figure S46-49), and by extension selective sweeps,

occur. This variation is random with respect to adaptive mutations arising *de novo*, but is also

influenced by demography (Figure S50). For example, treatments that are founded with strong

bottlenecks are less likely to evolve using variation in the founder, increasing dependence on *de*

novo mutations. Additionally, variation in our population contraction parameter modifies the per

population number of new mutations per generation. This is also true for features of simulated

genes, such as absolute length of coding sequence. Predictable temporal variation in the time at

which adaptation is likely to occur is a probable source of variance between measures of

differentiation and is particularly clear for $\Delta\pi$.

Concluding remarks

In conclusion, we have used forward-in-time simulations to perform a factorial experiment in which

we demonstrate the relationships between three measures of genetic differentiation, selection, and

features of demographic history that are commonly variable in natural populations. In agreement

with previous theoretical work, we find the reliance of measures of genetic divergence to indicate

regions under selection are dependent on demography and variable through time, with a notable lag

in D_{XY}. Furthermore, we provide novel comparisons between measures of genetic differentiation

that call into question the perceived wisdom that using multiple measures of genetic differentiation

can rule out false-positives. We also demonstrate that signals of convergent evolution across

independent replicates can be driven by similar demographic histories with minimal influences of

selection. Therefore, we strongly advise those using overlapping outlier scans to carefully consider

the demographic context of their system to avoid false-positives.

Ultimately, the need to quantify and correct for variable demographic histories when interpreting

measures of genetic differentiation and divergence is not new advice. It is however still relatively

uncommon to see demography taken into consideration in published genome scan studies. It is

tempting to assume that replication in study design or analysis in the form of taking multiple

measures of genetic divergence can reduce the risk of attaining false-positives. We hope to

emphasise in this study that this is not the case, as false-positives (i.e. genome scan outliers that are

not associated with regions under natural selection) can be driven by non-random genomic or

common demographic features that cannot be bypassed through replication. Moreover, many of the

patterns we observe are variable through time, such that the relevant pitfalls of analyses will depend

on the age of the populations being considered.

Finally, we hope that researchers interested in studying adaptation in the genomes of natural

populations will find this study informative in aiding interpretations of outlier scans and choosing

appropriate measures suitable to their demographic context.

ACKNOWLEDGEMENTS

783   The authors wish to thank all members of the Fraser lab for insightful discussions on this project, as

784   well as members of the University of Exeter's EB theme and the University of Sussex's EBE group.

785   The authors would also like to thank Dr Kai Zeng for feedback on an earlier copy of this manuscript.

AUTHOR CONTRIBUTIONS

788   JRW and BAF conceived the project, developed the experimental design and contributed towards

789   the writing of the manuscript. JRW wrote and implemented the pipeline for simulations and analysis.

DATA ACCESSIBILITY

792   A full pipeline including all bash, Eidos and R scripts necessary to repeat this analysis can be

793   downloaded from github (https://github.com/JimWhiting91/Contingent_Convergence_Pipeline).

FUNDING INFORMATION

796   This work was funded as part of a European Research Council grant (758382 GUPPYCon) awarded

797   to BAF.

CONFLICTS OF INTEREST

800 The authors declare no conflicts of interest

References

1.   Ahrens CW, Rymer PD, Stow A, Bragg J, Dillon S, Umbers KDL, et al. The search for loci under

selection: trends, biases and progress. Mol Ecol. 2018;27(6):1342–56.

2.   Cooper EA, Uy JAC. Genomic evidence for convergent evolution of a key trait underlying

divergence in island birds. Mol Ecol. 2017;26(14):3760–74.

3.   Hohenlohe PA, Bassham S, Etter PD, Stiffler N, Johnson EA, Cresko WA. Population genomics

of parallel adaptation in threespine stickleback using sequenced RAD tags. PLoS Genet.

2010;6(2).

4.   Jones FC, Chan YF, Schmutz J, Grimwood J, Brady SD, Southwick AM, et al. A genome-wide

SNP genotyping array reveals patterns of global and repeated species-pair divergence in

sticklebacks. Curr Biol. 2012;22(1):83–90.

5.   Rougemont Q, Gagnaire P, Perrier C, Genthon C, Besnard A, Launey S, et al. Inferring the

demographic history underlying parallel genomic divergence among pairs of parasitic and

nonparasitic lamprey ecotypes. Mol Ecol. 2017;26(1):142–62.

6.   Fraser BA, Künstner A, Reznick DN, Dreyer C, Weigel D. Population genomics of natural and

experimental populations of guppies (Poecilia reticulata). Mol Ecol. 2015;24(2):389–408.

7.   Reid NM, Proestou DA, Clark BW, Warren WC, Colbourne JK, Shaw JR, et al. The genomic

landscape of rapid repeated evolutionary adaptation to toxic pollution in wild fish. Science

(80-). 2016;354(6317):1305–8.

8.   Meier JI, Marques DA, Wagner CE, Excoffier L, Seehausen O. Genomics of parallel ecological

speciation in Lake Victoria cichlids. Mol Biol Evol [Internet]. 2018;(April 2018):1–37. Available

from: https://academic.oup.com/mbe/advance-

article/doi/10.1093/molbev/msy051/4958411

9.   Soria-Carrasco V, Gompert Z, Comeault AA, Farkas TE, Parchman TL, Johnston JS, et al. Stick

insect genomes reveal natural selection's role in parallel speciation. Science (80-).

2014;344(6185):738–42.

10.   Van Belleghem SM, Vangestel C, De Wolf K, De Corte Z, Most M, Rastas P, et al. Evolution at

two time frames. PLoS Genet. 2018;

11.   Waterhouse MD, Erb LP, Beever EA, Russello MA. Adaptive population divergence and

directional gene flow across steep elevational gradients in a climate-sensitive mammal. Mol

Ecol. 2018;27(11):2512–28.

12.   Ravinet M, Westram A, Johannesson K, Butlin R, André C, Panova M. Shared and nonshared

genomic divergence in parallel ecotypes of Littorina saxatilis at a local scale. Mol Ecol.

2016;25(1):287–305.

13.   Westram AM, Galindo J, Alm Rosenblad M, Grahame JW, Panova M, Butlin RK. Do the same

genes underlie parallel phenotypic divergence in different Littorina saxatilis populations? Mol

Ecol. 2014;23(18):4603–16.

14.   Trucchi E, Frajman B, Haverkamp THA, Schönswetter P, Paun O. Genomic analyses suggest

parallel ecological divergence in heliosperma pusillum (caryophyllaceae). New Phytol.

2017;216(1):267–78.

15.   Roda F, Ambrose L, Walter GM, Liu HL, Schaul A, Lowe A, et al. Genomic evidence for the

parallel evolution of coastal forms in the *Senecio lautus* complex. Mol Ecol.

2013;22(11):2941–52.

16.   Ellegren H, Wolf JBW. Parallelism in genomic landscapes of differentiation, conserved

genomic features and the role of linked selection. J Evol Biol [Internet]. 2017;30(8):1516–8.

Available from: http://doi.wiley.com/10.1111/jeb.13113

17.   Charlesworth B, Morgan MT, Charlesworth D. The effect of deleterious mutations on neutral

molecular variation. Genetics. 1993;134(4):1289–303.

18.   Bank C, Ewing GB, Ferrer-Admettla A, Foll M, Jensen JD. Thinking too positive? Revisiting

current methods of population genetic selection inference. Trends Genet. 2014;30(12):540–

6.

19.   Cruickshank TE, Hahn MW. Reanalysis suggests that genomic islands of speciation are due to

reduced diversity, not reduced gene flow. Mol Ecol. 2014;23(13):3133–57.

20.   Kimura M. The neutral theory of molecular evolution. Cambridge University Press; 1983.

21.   Charlesworth D, Charlesworth B, Morgan MT. The pattern of neutral molecular variation

under the background selection model. Genetics. 1995;141(4):1619–32.

22.   Cutter AD, Payseur BA. Genomic signatures of selection at linked sites: Unifying the disparity

among species. Nat Rev Genet [Internet]. 2013;14(4):262–74. Available from:

http://dx.doi.org/10.1038/nrg3425

23.   Charlesworth B. Effective population size and patterns of molecular evolution and variation.

Nat Rev Genet. 2009;10(3):195.

24.   Yeaman S, Otto SP. Establishment and maintenance of adaptive genetic divergence under

migration, selection, and drift. Evol Int J Org Evol. 2011;65(7):2123–9.

25.   Torres R, Szpiech ZA, Hernandez RD. Human demographic history has amplified the effects of

background selection across the genome. PLoS Genet. 2018;14(6):1–27.

26.   Sella G, Petrov DA, Przeworski M, Andolfatto P. Pervasive natural selection in the Drosophila

genome? PLoS Genet. 2009;5(6):e1000495.

27.   Charlesworth B. Background selection and patterns of genetic diversity in Drosophila

melanogaster. Genet Res (Camb). 1996;68(2):131–49.

28.   Zeng K. A coalescent model of background selection with recombination, demography and

variation in selection coefficients. Heredity (Edinb). 2013;110(4):363.

29.   Charlesworth B, Nordborg M, Charlesworth D. The effects of local selection, balanced

polymorphism and background selection on equilibrium patterns of genetic diversity in

subdivided populations. Genet Res (Camb). 1997;70(2):155–74.

30.   Hoban S, Kelley JL, Lotterhos KE, Antolin MF, Bradburd G, Lowry DB, et al. Finding the

Genomic Basis of Local Adaptation: Pitfalls, Practical Solutions, and Future Directions. Am Nat

[Internet]. 2016;188(4):379–97. Available from:

http://www.journals.uchicago.edu/doi/10.1086/688018

31.   Charlesworth B. Measures of divergence between populations and the effect of forces that

reduce variability. Mol Biol Evol. 1998;15(5):538–43.

32.   Weir BS, Cockerham CC. Estimating F-Statistics for the Analysis of Population Structure

Author (s): B . S . Weir and C . Clark Cockerham Published by : Society for the Study of

Evolution Stable URL : http://www.jstor.org/stable/2408641 Accessed : 01-08-2016 16 : 07

UTC Your u. Evolution (N Y). 1984;38(6):1358–70.

33.   Hudson RR, Slatkin M, Maddison WP. Estimation of levels of gene flow from DNA sequence

data. Genetics. 1992;132(2):583–9.

34.   Smith JM, Haigh J. The hitch-hiking effect of a favourable gene. Genet Res (Camb).

1974;23(1):23–35.

35.   Nei M. Molecular evolutionary genetics. Columbia university press; 1987.

36.   Matthey-Doret R, Whitlock MC. Background selection and the statistics of population

differentiation: consequences for detecting local adaptation. bioRxiv. 2018;326256.

37.   Whitlock MC, Lotterhos KE. Reliable detection of loci responsible for local adaptation:

inference of a null model through trimming the distribution of F ST. Am Nat.

2015;186(S1):S24–36.

38.   Hohenlohe PA, Bassham S, Currey M, Cresko WA. Extensive linkage disequilibrium and

parallel adaptive divergence across threespine stickleback genomes. Philos Trans R Soc L B

Biol Sci [Internet]. 2012;367(1587):395–408. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/22201169

39. Reznick D, Endler JA. The impact of predation on life history evolution in Trinidadian guppies
(Poecilia reticulata). Evolution (N Y). 1982;36(1):160–77.

40. Haller BC, Messer PW. SLiM 3: Forward genetic simulations beyond the Wright–Fisher model.
Mol Biol Evol. 2019;36(3):632–7.

41. Künstner A, Hoffmann M, Fraser BA, Kottler VA, Sharma E, Weigel D, et al. The genome of the
trinidadian guppy, Poecilia reticulata, and variation in the Guanapo population. PLoS One.
2016;11(12):1–25.

42. Lanfear R, Kokko H, Eyre-Walker A. Population size and the rate of evolution. Trends Ecol
Evol. 2014;29(1):33–41.

43. Ellegren H, Galtier N. Determinants of genetic diversity. Nat Rev Genet. 2016;17(7):422–33.

44. Hartl DL, Clark AG, Clark AG. Principles of population genetics. Vol. 116. Sinauer associates
Sunderland; 1997.

45. Nosil P, Funk DJ, Ortiz-Barrientos D. Divergent selection and heterogeneous genomic
divergence. Mol Ecol. 2009;18(3):375–402.

46. Turner TL, Hahn MW, Nuzhdin S V. Genomic islands of speciation in Anopheles gambiae. PLoS
Biol. 2005;3(9):e285.

47. Ravinet M, Faria R, Butlin RK, Galindo J, Bierne N, Rafajlović M, et al. Interpreting the genomic
landscape of speciation: a road map for finding barriers to gene flow. J Evol Biol.
2017;30(8):1450–77.

48. Vijay N, Weissensteiner M, Burri R, Kawakami T, Ellegren H, Wolf JBW. Genomewide patterns
of variation in genetic diversity are shared among populations, species and higher-order taxa.
Mol Ecol. 2017;26(16):4284–95.

49. Irwin DE, Alcaide M, Delmore KE, Irwin JH, Owens GL. Recurrent selection explains parallel
evolution of genomic regions of high relative but low absolute differentiation in a ring
species. Mol Ecol. 2016;25(18):4488–507.

50. Stankowski S, Chase MA, Fuiten AM, Ralph PL, Streisfeld MA. The tempo of linked selection:
Rapid emergence of a heterogeneous genomic landscape during a radiation of
monkeyflowers [Internet]. bioRxiv. 2018. 342352 p. Available from:
https://www.biorxiv.org/content/early/2018/06/21/342352

51. Romiguier J, Gayral P, Ballenghien M, Bernard A, Cahais V, Chenuil A, et al. Comparative
population genomics in animals uncovers the determinants of genetic diversity. Nature.
2014;515(7526):261.

52. Van Doren BM, Campagna L, Helm B, Illera JC, Lovette IJ, Liedvogel M. Correlated patterns of
genetic diversity and differentiation across an avian family. Mol Ecol. 2017;26(15):3982–97.

53.   Dutoit L, Vijay N, Mugal CF, Bossu CM, Burri R, Wolf J, et al. Covariation in levels of nucleotide

diversity in homologous regions of the avian genome long after completion of lineage sorting.

Proc R Soc B. 2017;284(1849):20162756.

54.   Kess T, Galindo J, Boulding EG. Genomic divergence between Spanish Littorina saxatilis

ecotypes unravels limited admixture and extensive parallelism associated with population

history. Int J Bus Innov Res. 2018;17(3):8311–27.

55.   Kirkpatrick M, Barton N. Chromosome inversions, local adaptation and speciation. Genetics.

2006;173(1):419–34.

56.   Enard D, Messer PW, Petrov DA. Genome-wide signals of positive selection in human

evolution. Genome Res. 2014;

57.   Noor MAF, Bennett SM. Islands of speciation or mirages in the desert? Examining the role of

restricted recombination in maintaining species. Heredity (Edinb). 2009;103(6):439.

58.   Jones FC, Grabherr MG, Chan YF, Russell P, Mauceli E, Johnson J, et al. The genomic basis of

adaptive evolution in threespine sticklebacks. Nature [Internet]. 2012;484(7392):55–61.

Available from: http://www.nature.com/doifinder/10.1038/nature10944

59.   Nishikawa H, Iijima T, Kajitani R, Yamaguchi J, Ando T, Suzuki Y, et al. A genetic mechanism for

female-limited Batesian mimicry in Papilio butterfly. Nat Genet. 2015;47(4):405.

60.   Morales HE, Faria R, Johannesson K, Larsson T, Panova M, Westram AM, et al. Genomic

architecture of parallel ecological divergence: beyond a single environmental contrast.

bioRxiv. 2018;447854.

61.   Frankham R. Effective population size/adult population size ratios in wildlife: a review. Genet

Res (Camb). 1995;66(2):95–107.

62.   Hodgkinson A, Eyre-Walker A. Variation in the mutation rate across mammalian genomes.

Nat Rev Genet. 2011;12(11):756.