Environmental light is required for maintenance of long-term memory

in *Drosophila*

Show Inami[1], Shoma Sato[1], Shu Kondo[2], Hiromu Tanimoto[3], Toshihiro Kitamoto[4, 5], and

Takaomi Sakai[1, *]

[1]Department of Biological Sciences, Tokyo Metropolitan University, Tokyo, Japan

[2]Genetic Strains Research Center, National Institute of Genetics, Shizuoka, Japan

[3]Tohoku University Graduate School of Life Sciences, Sendai, Japan

[4]Department of Anesthesia and Pharmacology, Carver College of Medicine, University of Iowa,

Iowa, USA

[5]Interdisciplinary Programs in Genetics and Neuroscience, University of Iowa, Iowa, USA

*For correspondence: sakai-takaomi@tmu.ac.jp

Abstract

Long-term memory (LTM) is stored as functional modifications of relevant neural circuits in the brain. However, it remains poorly understood how the consolidated memory is maintained for a long period. Here, we show that in *Drosophila*, universally available environmental light plays a critical role in LTM maintenance by reactivating the memory modulator cAMP response element-binding protein (CREB) via neuropeptide signaling. LTM is impaired when flies are kept in constant darkness during the memory maintenance phase. Because light activates the brain neurons expressing the neuropeptide Pigment-dispersing factor (Pdf) through the brain photoreceptors, we examined the possible involvement of Pdf in LTM maintenance and found that temporal knockdown of Pdf during the memory maintenance phase impairs LTM in light–dark cycles. Furthermore, we demonstrated that the Pdf receptor (Pdfr) and CREB are required in the memory center for LTM maintenance and Pdfr regulates light-dependent transcription via CREB.

Introduction

A newly formed memory is initially labile, but under certain circumstances, it is consolidated into a more stable long-term memory (LTM). Previous studies using various animal species have shown that activation of specific transcription factors, such as the cAMP response element-binding protein (CREB), and the corresponding *de novo* protein synthesis are essential for memory consolidation (Alberini, 2009; Kandel, 2012; Lee et al., 2008; Yin and Tully, 1996). Once consolidated, LTM requires continual maintenance for its long-term storage and subsequent recall, because memory traces gradually decay owing to molecular turnover and cellular reorganization. Similarly to memory consolidation, transcriptional activation and *de novo* protein synthesis are required for LTM maintenance (Bekinschtein et al., 2007; Fioriti et al., 2015; Hirano et al., 2016; Majumdar et al., 2012). For example, the maintenance of hippocampal long-term potentiation (LTP) and spatial memory in mice is dependent on the prion-like translational regulator CPEB3 (Fioriti et al., 2015). Similarly, LTM maintenance in *Drosophila* also requires Orb2, a *Drosophila* homolog of CPEB (Majumdar et al., 2012). Furthermore, transcriptional regulations through CREB and its coactivator CRTC play crucial roles in LTM maintenance in the *Drosophila* memory center, namely, the mushroom bodies (MBs) (Hirano et al., 2016), suggesting that memory consolidation and maintenance share some of the same molecular mechanisms (Bekinschtein et al., 2007). An obvious and important question is how transcriptional activation and the following protein synthesis are triggered during the memory maintenance phase. Unlike transcriptional and translation activation involved in memory consolidation, those associated with LTM maintenance cannot be directly controlled by the stimuli that induce memory formation, because there is a significant time separation between stimulus-induced memory formation and the memory maintenance process, and LTM should be continually maintained. However, the molecular and cellular underpinnings of active LTM maintenance that is transcription- and translation-dependent still remain elusive.

Earth's rotation generates the daily cycle of day and night, and the rhythmic light–dark (LD) cycles have a significant impact on animal behavior and physiology. In animals, light is not only essential for acquiring information for image-forming vision in nature but also acts as a powerful modulator of brain functions such as circadian entrainment, hormone secretion, sleep-wake cycles, mood, and cognitive functions (Altimus et al., 2008; Crocker and Sehgal, 2010; Fernandez et al., 2018; LeGates et al., 2012; Vandewalle et al., 2009). Using the diurnal fruitfly *Drosophila melanogaster,* here, we found that LTM was severely impaired in flies kept in constant darkness (DD) after memory consolidation. Thus, LTM maintenance is light-dependent. In *Drosophila*, light activates photoreceptors in the brain neurons

expressing the Pigment-dispersing factor (Pdf), a neuropeptide, and increases their spontaneous firing rate (Fogle et al., 2011; Ni et al., 2017; Sheeba et al., 2008). We found that the brain photoreceptors in Pdf-expressing neurons (hereafter, Pdf neurons) and the Pdf receptor (Pdfr) in the *Drosophila* memory center are essential for light-driven LTM maintenance. We further showed that light activates CREB-dependent transcription in MBs and that Pdfr in MBs regulates light-dependent CREB transcription. Overall, our results demonstrate for the first time that environmental light, which is available daily to all animals under normal conditions, plays a critical role in LTM maintenance by reactivating the evolutionarily conserved memory modulator CREB via neuropeptide signaling.

Results

Light is essential for LTM maintenance

To investigate whether lighting conditions affect LTM in *Drosophila*, the courtship conditioning assay was carried out (Griffith and Ejima, 2009; Keleman et al., 2012; Sakai et al., 2004; Siegel and Hall, 1979). In this assay, males receive stressors from nonreceptive mated females (e.g., sexual rejection) to block successful mating (conditioning) (Lee et al., 2017), and memory is subsequently observed as experience-dependent courtship suppression toward virgin females. One hour conditioning generates short-term memory (STM), which persists for at least 8 h, whereas 7 h conditioning induces LTM, which persists for at least 5 days (d) (Ishimoto et al., 2009; Sakai et al., 2012; Sakai et al., 2004). The courtship activity of naïve and conditioned males was quantified using a courtship index (CI); subsequently, a memory index (MI) was calculated to quantify courtship memory (see Materials and methods). When males were conditioned for 7 h in light or darkness and the conditioned males were subsequently kept under LD cycle**s** until the test, they showed lower courtship activity on d 5 (five days after conditioning) than naïve males, and there was no significant difference in MI between these flies [Figure 1A; (1) and (2)], indicating that conditioning in darkness has no adverse effects on LTM. However, when flies were conditioned in light and then kept in DD after the conditioning and before the test, LTM was severely attenuated [Figure 1A; (3)]. DD for two consecutive days after conditioning was sufficient to impair LTM [Figure 1A; (6)], but not DD for only one day [Figure 1A; (4) and (5)], indicating that flies cannot maintain their LTM when DD lasts more than two days. In constant light (LL), the *Drosophila* circadian clock does not work normally, and flies show arrhythmic locomotor activity (Qiu and Hardin, 1996). When flies were kept in LL after conditioning, their LTM was intact [Figure 1A; (7)], as previously reported (Sakai et al., 2004). Furthermore, the LTM of several clock mutants except for *period* (*per*)

mutants is intact (Sakai et al., 2004). Thus, light input, but not the circadian clock, is necessary for LTM

maintenance.

Sleep plays an important role in the consolidation of *Drosophila* courtship memory (Donlea et al.,

2011). However, it remains unclear whether an abnormal sleep phenotype is attributable to disturbed

LTM maintenance of flies kept in DD. To examine whether light conditions affect *Drosophila* sleep (see

Materials and methods), sleep amount was measured in flies kept in LD and DD (Figure 2A). The mean

amount of daytime sleep of flies kept in DD was lower than that of flies kept in LD (Figures 2A–C).

However, when flies in LD were slightly deprived of sleep to adjust the amount of daytime sleep to the

level of that in DD (Figures 2D–F), their LTM maintenance was not attenuated (Figure 2G). Thus, LTM

impairment induced by DD does not simply result from the reduced amount of sleep.

Expression of brain photoreceptors in Pdf neurons is necessary for LTM maintenance in LD

Because light directly activates Pdf neurons through circadian photoreceptors [Rhodopsin 7 (Rh7) and

Cryptochrome (Cry)] (Fogle et al., 2011; Ni et al., 2017; Sheeba et al., 2008), Pdf neurons may regulate

light-driven LTM maintenance. To examine whether Rh7 and Cry are required for LTM, *Rh7* and *cry* null

mutant flies ($Rh7^1$ and cry^{01}, respectively) were used. Since memory on d 1 (one day after conditioning)

was intact in $Rh7^1$ and cry^{01} flies after 7 h conditioning (Figure 1B), *Rh7* and *cry* are dispensable for

memory consolidation. Unlike the memory on d 1, $Rh7^1$ and cry^{01} flies showed memory impairment on d

5 (Figure 1C), suggesting that Rh7 and Cry are required for keeping LTM more than one day. We next

sought to determine the temporal requirement of brain photoreceptors for LTM maintenance. For this

purpose, we performed temporal knockdown of *Rh7* or *cry* in Pdf neurons using the TARGET system

(McGuire et al., 2003) and RNA interference (RNAi) technology. The effectiveness of *Rh7* and *cry* RNAi

was confirmed by qRT-PCR (Figures 3A and B). To knockdown *Rh7* during memory consolidation,

maintenance or test phase, the temperature was raised to 30 °C for 24 h during three experimental periods

(Figures 1D–G): starting at 24 h before the end of conditioning, 48–72 h after conditioning (memory

maintenance phase), and 24 h before the test initiation. When flies were kept for 24 h before the test

initiation at the restrictive temperature (RT), *Rh7* should remain suppressed during the 10 min test. In *Rh7*

knockdown flies, LTM was impaired only when the temperature was shifted during the memory

maintenance phase (Figure 1F). Temporal induction of *cry* RNAi in Pdf neurons did not affect LTM

(Figures 1D–G). This is probably because temporal *cry* knockdown was insufficient to impaire LTM

since nonconditional knockdown of *cry* in Pdf neurons (*Pdf*-GAL4 / UAS-*cry* RNAi) resulted in severe

memory impairment on d 5, as was observed in *cry^{01}* flies (Figure 1H).

In *Rh7* knockdown flies, the 48–72 h RT exposure resulted in impaired LTM, indicating that *Rh7*

expression is required during the maintenance phase. However, the 24 h RT exposure immediately before

the test did not affect LTM. When flies were exposed to RT for 48–72 h, the targeted genes should remain

knocked down even after flies are returned to the permissive temperature (PT). As a result, functions of

the target genes should be suppressed for over 24 h during the maintenance phase. In contrast, when flies

were exposed to RT for 24 h immediately before the test, the genes should be suppressed during the

maintenance phase for 24 h or less. As shown in Figure 1A, the length of time in DD is the critical factor

for LTM defect: 1-d DD had no effect whereas 2-d DD led to LTM impairment. Thus, it is most likely

that only long-term (>24 h) suppression of the light-Pdf pathway results in impaired LTM. These results

support the conclusion that brain photoreceptors are necessary for LTM maintenance and Pdf neurons

regulate light-driven LTM maintenance.

***Pdf* expression is critical for LTM maintenance in LD**

To demonstrate that the Pdf signaling pathway regulates light-driven LTM maintenance, we performed

temporal knockdown of *Pdf* in Pdf neurons (Figures 4A–D). The effectiveness of *Pdf* RNAi was

confirmed by qRT-PCR (Figure 3C) and immunostaining using an anti-Pdf antibody (Figures 3F and G).

LTM was impaired only when *Pdf* was knocked down during the memory maintenance phase (Figure 4C),

indicating that *Pdf* expression is required for LTM maintenance. Since memory on d 1 was intact in *Pdf^{01}*

null mutant flies after 7 h conditioning (Figure 4E), Pdf is dispensable for memory consolidation. Unlike

the memory on d 1, *Pdf^{01}* mutant flies showed memory impairment on d 5, which was rescued by *Pdf*

expression (Figure 4F), indicating that Pdf plays a critical role in LTM maintenance.

We next used two additional GAL4 drivers, c929 and *R18F07* to further investigate neuronal cell

types involved in Pdf-mediated LTM maintenance. Pdf neurons form two clusters, small lateral ventral

neurons (s-LNvs) and large lateral ventral neurons (l-LNvs) (Dubowy and Sehgal, 2017). c929 drives the

expression of GAL4 in peptidergic neurons including l-LNvs but not s-LNvs (Shimada et al., 2016;

Taghert et al., 2001), and *R18F07* drives GAL4 expression in all s-LNvs and only weakly in one of the

l-LNvs (Figure 4G). *Pdf* knockdown in c929-positive neurons impaired LTM, but not that in

R18F07-positive neurons (Figure 4H). Taken together, it is most likely that Pdf expression in l-LNvs is

essential for LTM maintenance.

Activity of Pdf neurons regulates light-dependent LTM maintenance

We next examined whether the temporal activation of Pdf neurons induced by the temperature-sensitive

cation channel TrpA1 can compensate for the DD‐dependent LTM impairment (Figures 5A–D). When

Pdf neurons were activated in *Pdf*-GAL4/UAS-*TrpA1* flies within each subjective day or night in DD for

2 d after conditioning, LTM was maintained for 5 d in *Pdf*-GAL4/UAS-*TrpA1* flies (Figures 5C and D).

Under the same temperature-shift conditions, GAL4 and UAS control flies still showed LTM impairment

(Figures 5C and D). Thus, this finding indicates that the activation of Pdf neurons during either a

subjective day or night is sufficient to rescue the LTM impairment.

Consistently, the electrical silencing of Pdf neurons during the memory maintenance phase

attenuated LTM in LD (Figures 5E and F). Furthermore, l-LNv-specific electrical silencing (see Materials

and methods) also impaired LTM (Figures 5G and H), indicating that Pdf in l-LNvs plays a predominant

role in LTM maintenance. Taken together, these results support the conclusion that the light-inducible

activation of l-LNvs is essential for LTM maintenance.

Pdfr in MB α/β neurons regulates LTM maintenance

Because Pdfr activation increases cAMP levels (Hyun et al., 2005; Mertens et al., 2005) and MBs are the

critical brain structures for learning and memory (Davis, 2011; Dubnau and Chiang, 2013), it is possible

that Pdf activates Pdfr in MBs and drives LTM maintenance. However, previous studies using

Pdfr-GAL4 lines, in which GAL4 is fused to the upstream regulatory sequences of *Pdfr*, or an anti-Pdfr

antibody did not indicate Pdfr expression in MBs (Im and Taghert, 2010; Mertens et al., 2005). In

contrast, a recent study using RNA sequencing revealed that *Pdfr* is expressed in MBs (Crocker et al.,

2016). In this study, we generated a GAL4 knock-in fly strain, *Pdfr::2A::GAL4*, in which a transgene

containing the 2A peptide sequence and GAL4 was inserted immediately upstream of the stop codon of

Pdfr (Figures 6A and B). In *Pdfr::2A::GAL4* flies, GAL4 was expected to be expressed in the cells

endogenously expressing *Pdfr*. mCherry::NLS signals driven by *Pdfr::2A::GAL4* were detected in

Per-expressing clock neurons (Figure 6C). Furthermore, reporter gene expression driven by

Pdfr::2A::GAL4 partially colocalized with that driven by MB-LexA (Figure 7A). Several Kenyon cells

with the colocalization of *Pdfr::2A::GAL4* and MB-LexA were identified [5 ± 0.3 (mean ± SEM) cells in

one hemisphere, $N = 6$]. In the MB-LexA line, LexA is specifically expressed in all lobes (Figure 7A).

However, considering that LexA-positive neurons in this line are only a subset of all MB neurons, the

number of Kenyon cells with the colocalization of *Pdfr::2A::GAL4* and MB-LexA should be only a part

of the all Pdfr-positive MB neurons. In contrast to the *Pdfr::2A::GAL4* line, when the previously

generated *Pdfr*-GAL4 line was used, we did not detect the colocalization of *Pdfr*-GAL4 and MB-LexA in

the Kenyon cells (Figure 7B). To examine whether Pdfr is expressed in the MB α/β core region, we used

another LexA line (*R58F02*-LexA) with LexA expression in the MB α/β core region (Figure 7C). When

this line was used, the Kenyon cells with the colocalization of *Pdfr::2A::GAL4* and *R58F02*-LexA were

also detected (4 ± 1.0 cells in one hemisphere; *N* = 5). However, no colocalization of GAL4 and LexA

was detected (*N* = 5) when the *R19B03*-LexA line with weak LexA expression in the MB α/β core region

was used (Figure 7D). These findings indicate that Pdfr is indeed expressed in MB α/β core region.

To examine whether the Pdfr is required for LTM, we performed knockdown of *Pdfr* in

Pdfr-expressing neurons. The effectiveness of *Pdfr* RNAi was confirmed by qRT-PCR (Figure 3D, E).

LTM was attenuated when *Pdfr* was knocked down in *Pdfr::2A::GAL4*-positive neurons (Figure 8A).

Furthermore, *Pdfr*-null mutant flies (*Pdfr^{5304}*) also showed severe LTM impairment (Figure 8B), which

was rescued by *Pdfr* expression when an MB-GAL4 line, OK107, was used (Figure 8B). However, when

the previously generated *Pdfr*-GAL4 line was used, the impairment was not rescued (Figure 8B). Using

an anti-Per antibody, we confirmed that GAL4-positive neurons in OK107 did not include Per-positive

clock neurons (Figure 9A). Thus, Pdfr expression in MB neurons is essential for LTM maintenance.

LTM was also attenuated when *Pdfr* was knocked down in MB α/β or γ neurons (Figures 8C and D).

In contrast, targeted expression of *Pdfr* RNAi in Pdf neurons, GAL4-positive neurons in the previously

generated *Pdfr*-GAL4 line, and MB α'/β' neurons did not affect LTM (Figures 8C and D). The temporal

knockdown of *Pdfr* in MB γ neurons during conditioning or the memory maintenance phase did not affect

LTM (Figures 8E–G). However, temporal *Pdfr* knockdown in MB α/β neurons during the memory

maintenance impaired LTM (Figure 8G). In the MB α/β GAL4 line used in this experiment,

GAL4-positive neurons did not include Per-positive clock cells (Figure 9B). Thus, Pdfr in MB α/β

neurons is likely involved in intracellular signaling necessary for LTM maintenance; however, it remains

unclear whether Pdfr in MB γ neurons plays an acute physiological role in LTM.

CREB activity in MB α/β neurons is necessary for LTM maintenance

A *Drosophila* homolog of CREB (CrebB) is required for the consolidation and maintenance of olfactory

memory (Hirano et al., 2016; Yin and Tully, 1996). Our previous studies demonstrated that the

consolidation of courtship memory is also regulated by CrebB (Ishimoto et al., 2009; Sakai et al., 2004).

Pdf released in response to light may activate Pdfr, increase cAMP levels, and subsequently enhance

CrebB transcription in MB α/β neurons. Consistent with this idea, LTM was attenuated when a repressor

isoform of CrebB (CrebB-B; also known as dCREB2-b) was expressed in MB α/β neurons during

conditioning or the memory maintenance phase (Figures 10A–C), indicating that CrebB expression in

MB α/β neurons is required for both the consolidation and maintenance of LTM. On the other hand,

CrebB expression in MB γ neurons during conditioning, but not the maintenance phase, attenuated LTM

(Figures 10A–C), suggesting that CrebB in MB γ neurons is required only for memory consolidation.

Pdfr is essential for light-dependent CrebB activity in MB α/β neurons

To determine whether light and Pdfr activate CrebB transcription in MBs, we used the

CRE>mCherry::STOP>*luciferase* (*luc*) reporter, which is a *luc*-based reporter gene under the control of

CrebB-binding sites (CRE) (Tanenhaus et al., 2012). Using this reporter construct, in which the

mCherry::STOP sequence is flipped out in the presence of the GAL4-induced recombinase Flippase

(FLP), we can measure the MB-specific transcriptional activity of CrebB with MB-GAL4. The induction

of CrebB-B driven by a heat-shock promoter severely attenuated the CRE-dependent Luc activity in MBs

(Figure 11A), demonstrating that this reporter can reliably be used as an indicator of CrebB activity.

When naïve flies were kept in LD, robust CrebB activity was detected in MB α/β neurons, but not in MB

γ neurons (Figure 11B). When they were kept in DD for 1 d, CrebB activity in MB α/β neurons was

reduced by approximately 50% compared with that in control flies (Figure 11B). On the second day of

DD, it was further reduced (Figure 11B), indicating that CrebB activity in MB α/β neurons is

light-dependent. *Pdfr* knockdown induced an approximately 50% reduction in CrebB activity, and CrebB

activity was severely attenuated in MB α/β neurons with the *Pdfr* null mutant background (Figure 11B).

Furthermore, activation of Pdf neurons by TrpA1 increased CrebB activity in MB α/β neurons (Figure

11C). Thus, the Pdf/Pdfr signaling pathway is essential for light-driven CrebB activity in MB α/β neurons.

Next, the effect of 7 h conditioning on CrebB activity was examined. CrebB activity in MB α/β and γ

neurons increased immediately after 7 h conditioning (Figure 11D). This result is consistent with a

previous report (Ishimoto et al., 2009). However, no conditioning-dependent increase in CrebB activity

was observed during the memory maintenance phase (Figure 11E).

Discussion

In nocturnal mice, an ultradian LD cycle condition induces spatial learning defects, although such

an aberrant light condition does not impair the molecular clock or sleep amount (LeGates et al., 2012).

The ultradian LD cycle also attenuates hippocampal long-term potentiation (Fernandez et al., 2018). In

contrast, a short pulse of white light during the night before learning enhances long-lasting fear memory

through the activation of hippocampal p21-activated kinase 1 (Shan et al., 2015). In humans, who are

naturally diurnal, prior exposure to orange light promotes working memory, indicating that light also

modulates human cognitive brain functions (Chellappa et al., 2014). Thus, regardless of the nocturnal or

diurnal nature of animals, lighting conditions can positively or negatively modify the acquisition or

consolidation of memories (Cajochen et al., 2011; Chellappa et al., 2014; LeGates et al., 2012; Shan et al.,

2015). However, to the best of our knowledge, it has never been studied whether environmental light

affects LTM maintenance. In this study using *Drosophila*, we demonstrated for the first time that

environmental light is an essential factor for the appropriate maintenance of LTM.

Here we found that the brain photoreceptors Cry and Rh7, which are essential for light-dependent

activation of Pdf neurons (Fogle et al., 2011; Ni et al., 2017; Sheeba et al., 2008), play important roles in

LTM maintenance. *Drosophila* has three light-sensing organs: the compound eyes, ocelli, and Hofbauer–

Buchner (H–B) eyelets. Besides the brain photoreceptors, compound eyes also play key roles in light

entrainment of the circadian clock (Helfrich-Förster et al., 2002), and l-LNvs receive visual information

via the compound eyes (Muraro and Ceriani, 2015). Furthermore, the H–B eyelets are also important for

circadian photoreception, and axons of H–B eyelet photoreceptors project to the circadian pacemaker

neurons including l-LNvs (Li et al., 2018). Although it remains unclear whether visual information via the

compound eyes or H–B eyelets is also important for LTM maintenance, light inputs from photoreceptors

in *Drosophila* light-sensing organs may also partially contribute to light-driven LTM maintenance.

The *Drosophila* Pdf neuropeptide regulates various biological phenomena such as circadian

behavioral rhythms, light-driven arousal, geotactic behavior, rival-induced prolonged mating, and sex

pheromone biosynthesis (Helfrich-Förster et al., 2000; Kim et al., 2013; Krupp et al., 2013; Mertens et al.,

2005; Renn et al., 1999). In the present study, we found that *Drosophila* has a light-driven memory

maintenance system, and that the Pdf signaling pathway regulates this system. Pdf expression was

essential for LTM maintenance in LD (Figure 4). LTM was impaired when flies were kept in DD for 2 d

after 7 h conditioning (Figure 1), and the activation of Pdf neurons was sufficient to rescue the LTM

impairment induced by DD (Figure 5). Moreover, the electrical silencing of Pdf neurons impaired LTM in

LD (Figure 5). Considering that light activates Pdf neurons through brain photoreceptors (Fogle et al.,

2011; Ni et al., 2017; Sheeba et al., 2008) and Rh7 and Cry in Pdf neurons are necessary for light-driven

LTM maintenance (Figure 1), it is most likely that light-inducible Pdf released from Pdf neurons regulates

LTM maintenance. Our study using a GAL4 knock-in fly, *Pdfr::2A::GAL4*, revealed that Pdfr is

expressed in MB neurons (Figure 7). We further confirmed that Pdfr in MB α/β neurons is necessary for

LTM maintenance (Figure 8G) and Pdfr expression in MB α/β neurons is necessary for the light-driven

transcription through CrebB, which is essential for LTM maintenance (Figures 10 and 11). Taken together,

our study shows that this light-driven transcription system via the Pdf/Pdfr/CrebB pathway in MB α/β

neurons regulates LTM maintenance in *Drosophila*.

The targeted expression of a CrebB repressor in MB α/β neurons during 7 h conditioning impaired

LTM, as was observed in that in MB γ neurons (Figure 10), indicating that CrebB-dependent transcription

in both MB α/β and γ neurons is necessary for memory consolidation. In addition, the targeted expression

of a CrebB repressor in MB α/β neurons during the memory maintenance phase also impaired LTM

(Figure 10), whereas, that in MB γ neurons did not (Figure 10). It has been reported that MB γ neurons

are necessary for 1 d memory of courtship conditioning (Kruttner et al., 2015). Thus, consolidated

courtship memory may last in MB γ neurons only for 1–2 d at most, whereas longer-lasting LTM seems

to be established and stored in MB α/β neurons. In contrast to the MB γ neurons, Pdfr expression and

CrebB activity in MB α/β neurons are essential for LTM maintenance, and the light-dependent activation

of CrebB transcription was evident in MB α/β neurons, indicating that 5 d memory is maintained in MB

α/β neurons through the Pdf/Pdfr/CrebB pathway in LD. Thus, long-lasting LTM (> 1 d memory) seems

to be established and stored in MB α/β neurons.

When flies were kept in DD for two consecutive days, LTM was impaired (Figure 1) and CrebB

activity in MB α/β neurons was severely attenuated (Figure 11). However, DD for only 1 day was not

sufficient to impair LTM (Figure 1) but it induced 50% reduction in CrebB activity (Figure 11). Thus,

when CrebB activity in MB α/β neurons is severely attenuated, LTM maintenance may break down. Can

Pdfr be the only factor that regulates CrebB activity in MB α/β neurons? Pdf/Pdfr signaling should play a

role in the regulation of CrebB activity in MB α/β neurons because a *Pdfr* null mutation severely

attenuated CrebB activity (Figure 11). However, it remains possible that signaling pathways other than

the Pdf/Pdfr signaling pathway also partially contribute to CrebB activity in MB α/β neurons.

In contrast to LTM maintenance, LTM formation occurred in darkness (Figure 1). This finding is

consistent with the previous report that STM after 1 h conditioning under a dim red light, which blocks

visual input, is intact (Joiner and Griffith, 1997). Although we found that CrebB activity during

conditioning was necessary for memory consolidation (Figure 10), *Pdfr* in MBs was not (Figure 8).

Considering that flies were able to establish LTM when they were conditioned in darkness (Figure 1), it is

likely that light-independent CrebB transcription without using Pdfr in MBs plays an important role in

memory consolidation (Figure 12A). Unlike memory consolidation, LTM maintenance requires a

light-driven transcription system via the Pdf/Pdfr/CrebB pathway in MB α/β neurons (Figure 12B). In

naïve males, however, light can also increase the CrebB transcription activity in MB α/β neurons (Figure

11). In naïve males, it is most unlikely that this light-driven transcription system provides proteins that are

required for LTM maintenance. Thus, it may play a role in innate brain functions other than LTM

maintenance. Assuming that this hypothesis is true, how are proteins that are required for LTM

maintenance provided in only conditioned males? Although it remains unclear, repetitive exposure to

stressors using mated females during courtship conditioning may trigger the change in the target genes of

CrebB in the light-driven transcription system, and such a system in MB α/β neurons may provide gene

products that are required for maintaining consolidated LTM in a courtship conditioning-dependent

manner.

In nature, animals learn much from their experience throughout the day. Through their experience,

LTM should be formed and maintained for a long period. If environmental light, which is available daily

to all animals in nature, can be used for transcriptional activation in the brain, such a light-driven

transcription system is considered reasonable and effective for continually providing *de novo* protein

synthesis required for LTM maintenance. When diurnal Nile grass rats were housed under dim LD cycles,

24 h spatial memory is impaired, and this lighting condition inhibits the expression of brain-derived

neurotrophic factors and dendritic spine density in the hippocampus (Soler et al., 2018). Although it is

unknown whether the rapid forgetting in the Nile grass rats under dim LD cycles results from the

inhibition of *de novo* protein synthesis required for LTM maintenance, light-dependent *de novo* protein

synthesis in the memory center may be conserved in many animal species. As is observed in *Drosophila*,

in mammals, repeated exposure to stressors also induces a long-lasting reduction of male sexual

motivation (Hawley et al., 2011; Hawley et al., 2013). It is thus interesting to examine whether

environmental light and the evolutionarily conserved memory modulator CREB also play critical roles in

the maintenance of such depressed sexual motivation in mammals including humans.

Materials and methods

Fly stocks

All flies were raised on glucose-yeast-cornmeal medium in 12:12 LD cycles at 25.0 ± 0.5 °C (45–60%

relative humidity). Virgin males and females were collected without anesthesia within 8 h after eclosion.

Fly stocks used for this study are as follows: wild-type Canton-S (CS), Pdf^{01} (BL26654),

$Pdfr^{5304}$(BL33068), cry^{01}(Provided by Dr. Taishi Yoshii), $Rh7^{1}$ (BL76022), Pdf-GAL4 (BL6900), c929

(BL25373), $R14F03$ (BL48648), $R18F07$ (BL47876), OK107 (BL854), MB-LexA (provided by Dr.

Minoru Saitoe), $R58F02$-LexA (Gift from Dr. Yoshinori Aso), $R19B03$-LexA (BL52537), $R61G12$-LexA

(BL52685), $R41C10$ (BL50121), $R55D03$ (BL47656), $R19B03$ (BL49830), c305a (BL30829),

$nSyb$-GAL4 (BL51941), UAS-FLP (BL4539), UAS-$Kir2.1::eGFP$ (BL6596), UAS-$TrpA1$ (BL26263),

UAS-$mCD8::GFP$ (BL5137), UAS-$mCherry::NLS$ (BL38424), UAS-$mCD8::RFP$ (BL32218), UAS-cry

RNAi (NIG3772R-3), UAS-$Rh7$ RNAi (VDRC109688), UAS-Pdf RNAi (VDRC4380), UAS-$Pdfr$ RNAi

(VDRC42724), UAS-$CrebB$-B (also known as UAS-$dCREB2$-b, BL7219), tub-GAL80^{ts} (BL7017),

LexAop2-$FLPL$ (BL55820), LexAop2-$mCD8::GFP$ (BL32203), LexAop-$TrpA1$ (provided from Dr. Mani

Ramaswami), hs-$CrebB$-B (also known as hs-$dCREB2$-b) (Sakai et al., 2004; Yin et al., 1994),

$Pdfr$-GAL4 (provided by Dr. Leslie C. Griffith), UAS-Pdf (provided by Dr. Taishi Yoshii), UAS-$Pdfr$

(provided from Dr. Taishi Yoshii), UAS-luc RNAi (provided by Dr. Kanae Ando),

UAS>STOP>$Kir2.1::eGFP$ (provided by Dr. David J. Anderson), and CRE>mCherry::STOP>luc

(provided by Dr. Jerry C. P. Yin). $Pdfr::2A::GAL4$ was generated by the targeted integration of a

2A-GAL4 cassette (Diao and White, 2012) into the $Pdfr$ locus, such that it replaces the stop codon of the

RA isoform of the $Pdfr$ transcripts. The 2A-GAL4 cassette contained a $3xP3$-RFP transgene (Horn et al.,

2000) flanked by loxP sites as a selection marker, which was removed by Cre expression after the

transgenic lines were established. All lines for behavior experiments except for cry^{01}, Pdf^{01}, $Pdfr^{5304}$, $Rh7^{1}$,

UAS-$Pdfr$, UAS-luc RNAi, LexAop2-$FLPL$, and $Pdfr::2A::GAL4$ were outcrossed for at least five

generations to $white^{1118}$ flies with the Canton-S (CS) genetic background.

Courtship conditioning assay

The courtship conditioning assay was carried out as previously described (Sakai et al., 2004) with some

modifications. Unreceptive mated females were prepared as "trainers" one or two days before they were

used for courtship conditioning. In this conditioning, a virgin CS female and a male (3–6 d old) were

placed in an acrylic courtship chamber (15 mm in diameter $\times$ 3 mm in depth) for copulation. For LTM, a

3–5 d old male was placed with a mated female (4–7 d old) in a conditioning chamber (15 mm in

diameter, 5 mm in depth) containing food for 7 h either with (conditioned) or without (naïve) a single

premated female (7 h conditioning). If males remated with mated females during conditioning, we

discarded such males after conditioning. After 7 h conditioning, only flies showing courtship behaviors

toward the mated female but not copulating successfully were transferred to a glass tube with food (12

mm in diameter × 75 mm in depth) and kept in isolation for 1 or 5 days until the test. The test was

performed using a freeze-killed virgin female in a test chamber (15 mm in diameter, 3 mm in depth). All

procedures in the experiments were carried out at 25 ± 1.0 °C (45–60% relative humidity) except for the

temperature shift experiments. Male courtship behaviors of individual flies were calculated manually as a

courtship index (CI), defined as the percentage of time spent in performing courtship behaviors during a

given observation period (10 min). We first measured CI in conditioned and naïve males ($CI_{Conditioned}$ and

$CI_{Naïve}$), and then mean $CI_{Naïve}$ was calculated. To quantify courtship memory as previously reported (Lee

et al., 2017), memory index (MI) was calculated using the following formula: MI = (mean $CI_{Naïve}$ - mean

$CI_{Conditioned}$) / mean $CI_{Naïve}$. In the statistical analysis of CI, the Kolmogorov–Smirnov test was used to

determine whether the data are normally distributed. When the data were not distributed normally, we

carried out the log transformation of the data. When the basic data or transformed data are normally

distributed, Student's *t*-test was used for comparisons. When the basic data and transformed data are not

distributed normally, we used the Mann–Whitney *U* test for comparisons. The computer software IBM

SPSS Statistics 22 (IBM Japan, Ltd.) was used for these tests. In the statistical analysis of MI, the

permutation test with 10000 random permutations was used. The free statistical package R was used for

these tests (Koemans et al., 2017).

Lighting conditions in courtship conditioning

When courtship conditioning was performed in a dark place, conditioning chambers were placed in a

temperature-regulated (25.0 ± 0.5 °C) light-tight incubator (MIR-254, Sanyo Electric Co., Ltd.). To check

whether mating occurs during conditioning, we observed the flies for 10 sec every 30 min by opening the

incubator door. To determine whether lighting conditions affect memory maintenance, conditioned and

naïve flies were kept in a light-tight incubator (MIR-254, Sanyo Electric Co., Ltd.) for one, two, or four

days after 7 h conditioning (see Figure 1A).

Temporal activation of Pdf neurons

The temperature-sensitive cation channel TrpA1 was used to activate Pdf neurons (Hamada et al., 2008).

For the activation of Pdf neurons, during two days of DD (the second and third days after 7 h

conditioning), *Pdf*-GAL4/UAS-*TrpA1* flies were kept at 30 °C for 8 h within each subjective day

[circadian time (CT) 0–8] or night (CT 12–20). However, DD-dependent LTM impairment was not

improved. Since it is possible that activation of Pdf neurons under this condition is insufficient for LTM

maintenance, the flies were kept at 34 °C for 8 h within each subjective day (CT 0–8) or night (CT 12–20).

UAS-*TrpA1*/+ and *Pdf*-GAL4/+ flies were used as the control.

Temporal gene expression using TARGET system

The *tub*-GAL80^{ts} transgene used in the TARGET system (McGuire et al., 2003) encodes a ubiquitously

expressed, temperature-sensitive GAL4 repressor that is active at the permissive temperature (PT, 25 °C)

but not at the restrictive temperature (RT, 30 or 32 °C). By using UAS-*Rh7* RNAi, UAS-*cry* RNAi,

UAS-*Pdf* RNAi, or UAS-*Pdfr* RNAi combined with the TARGET system, we knocked down *Rh7*, *cry*,

Pdf, or *Pdfr* in GAL4-positive neurons at RT (30 °C), but not at PT. In these experiments, we shifted PT

to RT and vice versa during the three experimental phases (Figures 8E–G): 24 h before the end of

conditioning, 48–72 h after conditioning, and 24 h before the initiation of the test. Furthermore, to drive

the expression of a UAS-*CrebB-B* construct in MBs during a specific temporal phase, the TARGET

system was also used. In *CrebB-B* experiments, we shifted PT to RT (30 °C) and vice versa during the

two experimental phases (Figures 10A–C): 10 h before the end of conditioning, 48–72 h after

conditioning. To drive the expression of a UAS-*Kir2.1* construct in Pdf neurons during the memory

maintenance phase using the TARGET system, flies were kept at RT (32°C) for 48–72 h after

conditioning. Subsequently, they were kept at PT.

l-LNv-specific electrical silencing

To assay whether l-LNv-specific electrical silencing affects LTM, two binary gene expression systems

(GAL4/UAS and LexA/LexAop) combined with flippase (FLP/FRT) were used. Using this system, the

specific target gene is expressed in GAL4- and LexA-coexpressing neurons. *R61G12*-LexA and

R14F03-GAL4 lines were used in the experiments. We confirmed that *Kir2.1::eGFP* is expressed in all

l-LNvs but not s-LNvs (Figure 5H). Thus, in *R61G12*-LexA/LexAop-*FLPL*;

UAS>STOP>*Kir2.1::eGFP/R14F03* flies, according to the l-LNvs-specific expression of Kir2.1, only

l-LNvs will be electrically silenced.

Real-time quantitative reverse transcription PCR (qRT-PCR)

A PicoPure RNA Isolation Kit (KIT0204, Thermo Fisher Scientific) was used for collecting total RNA

from three whole brains in each genotype. cDNA was synthesized by the reverse transcription reaction

using a QuantiTect Reverse Transcription Kit (#205311, QIAGEN). Real-time quantitative PCR was

carried out using the THUNDERBIRD SYBR qPCR Mix (QPS-201, TOYOBO) and a Chromo 4

Detector (CFB-3240, MJ Research). The primer sequences (custom-made by Eurofins Genomics) used

for qRT-PCR were as follows: *Pdf*-Forward, 5'-ATCGGGATCTCCTCGACTGG-3'; *Pdf*-Reverse,

5'-ATGGGCCCAAGGAGTTCTCG-3'; *Pdfr*-Forward, 5'- CGTCTCATGCAGCAGATGGG-3';

Pdfr-Reverse, 5'-TAAGGGCGAACAGGGAGAGG -3'; *Rh7*-Forward,

5'-GCAAGGACCACGACAAGCAC-3'; *Rh7*-Reverse, 5'- GCTCAGGTCGTACTTGTCCC-3';

cry-Forward, 5'-CTCGCCGATAAGGATCAGGG-3'; *cry*-Reverse,

5'-ATGTCCTGCAACGAGTCCAG-3'; *rp49*-Forward, 5'-AAGATCGTGAAGAAGCGCAC-3';

rp49-Reverse, 5'-TGTGCACCAGGAACTTCTTG-3'. The expression level of each mRNA was

normalized by that of *rp49* mRNA. The average of the normalized mRNA expression level in control flies

was calculated using data from 4–6 independent assays. The expression level of each mRNA was

normalized by that of *rp49* mRNA. The average of the normalized mRNA expression level in control flies

was calculated using data from 4–6 independent assays. The mean (± SEM) ratio was calculated using

data from 4–6 independent assays. The Kolmogorov–Smirnov test was used to estimate whether the data

are normally distributed. When the basic data or log-transformed data were normally distributed,

Student's *t*-test or one-way ANOVA followed by post-hoc analysis using Scheffe's test was used. When

the basic data and log-transformed data were not distributed normally, the Kruskal–Wallis test followed

by the rank-sum test for multiple pairwise comparisons was used. The computer software IBM SPSS

Statistics 22 (IBM Japan, Ltd.) was used for these tests.

Luciferase assay

To test whether light or Pdfr activation promotes CrebB activity in MBs, the Luciferase (Luc) reporter

was used (Tanenhaus et al., 2012). Luc should be expressed in MBs in the combination of

CRE>mCherry::STOP>*luc* reporter, MB-GAL4, and UAS-*FLP*. Three MB-GAL4 lines (*R41C10*,

R55D03, and *R19B03*) were used in these experiments. *In vitro* Luc activity was measured using a

Luciferase Assay System (E1501, Promega). Three adult male heads were collected into a 1.5 ml

Eppendorf tube and homogenized in 50 µl of Glo Lysis Buffer (E266A, Promega) at Zeitgeber time (ZT)

0–2. After centrifugation, 10 µl of the resulting supernatant and 50 µl of Luciferin solution were used to

analyze Luc activity. The luminescence of each sample was measured using a luminometer (GloMax

20/20, Promega) and normalized to total protein concentration using a Protein Assay Kit (#5000006,

Bio-Rad). The UAS-*FLP* transgene used in this study displayed leaky expression of Luc (211–279

relative luminescence unit (RLU/mg). Thus, first, in the control line

(UAS-*FLP*/CRE>mCherry::STOP>*luc*), the mean normalized luminescence (mean $L_{control}$) was calculated.

CrebB activity index (CAI) (RLU/mg) was defined as the difference between the "mean $L_{control}$" and the

measured luminescence in each sample in each genotype, and finally, we calculated the mean CAI. To

examine whether CrebB activity increases after 7 h conditioning, we measured the CAI in each sample

using conditioned or naïve flies ($CAI_{Conditioned}$ and $CAI_{Naïve}$), and then mean $CAI_{Naïve}$ was calculated.

Finally, we calculated the conditioning-dependent CrebB activity using the following formula:

conditioning-dependent CrebB activity = ($CAI_{Conditioned}$ - mean $CAI_{Naïve}$) / mean $CAI_{Naïve}$. To examine

whether induction of the CrebB-B inhibits CrebB activity, we used UAS-*FLP*/*hs-CrebB-B*;

CRE>mCherry::STOP>*luc*/*R19B03* flies. For the heat-shock treatment, 3–4 d old male flies were grouped

as 20 flies per a food vial. Flies were heat-shocked at 32°C for three days. Luc activity was measured

immediately after the heat-shock treatment at ZT 0–2. To examine whether activation of Pdf neurons

increases CrebB activity in the MBs, we used UAS-*FLP*/*R61G12*-LexA; CRE>mCherry::STOP>*luc*

LexAop-*TrpA1*/*R41C10* flies. For the activation of Pdf neurons by TrpA1, the 3–5 d old males were

grouped as 20 flies per a food vial. Flies were heat-shocked at 32°C during two days of DD (the second

and third days after transfer). Luc activity was measured at ZT 0–2 after flies were returned to normal LD

cycle. The Kolmogorov–Smirnov test was used to estimate whether the data are normally distributed.

Because the basic data were normally distributed, Student's *t*-test was used for comparisons of two means,

and one-way ANOVA followed by post-hoc analysis using Schefe's test was carried out for multiple

pairwise comparisons. The computer software IBM SPSS Statistics 22 (IBM Japan, Ltd.) was used for

these tests.

Immunohistochemistry

Immunohistochemistry was performed as previously described (Mabuchi et al., 2016; Shimada et al.,

2016). For Pdf staining, brains were stained with a mouse anti-Pdf antibody (PDF C7-s, Developmental

Studies Hybridoma Bank at the University of Iowa, 1:200), and Alexa Fluor 568 anti-mouse IgG (A11004,

Thermo Fisher Scientific) was used as the secondary antibody (1:1000). For GFP staining, brains were

stained with a rabbit anti-GFP antibody (A11122, Thermo Fisher Scientific, 1:200), and Alexa Fluor 488

anti-rabbit IgG (A11008, Thermo Fisher Scientific, 1:1000) was used as the secondary antibody. For Per

staining, brains were stained with a rabbit anti-Per antibody (sc-33742, Santa Cruz Biotechnology, 1:250),

and Alexa Fluor 568 anti-rabbit IgG (A11011, Thermo Fisher Scientific) was used as the secondary

antibody (1:1000). Fluorescence signals were observed under a confocal microscope [C2$^+$ (Nikon) or

LSM710 (Zeiss)].

Quantitative analysis of Pdf immunoreactivity in l-LNvs

To examine whether temporal knockdown of *Pdf* in l-LNvs by the TARGET system inhibits Pdf

immunoreactivity, *Pdf*-GAL4/UAS-*Pdf* RNAi; +/*tub*-GAL80^{ts} flies were used. *Pdf*-GAL4/+;

+/*tub*-GAL80^{ts} flies were used as the control. After eclosion, all flies were kept for 3–6 days at PT, and

then the temperature was shifted to RT at ZT 8. After 24 h, the temperature was shifted again to PT.

Subsequently, we performed brain dissection for antibody staining 1 h after the temperature shift to PT

(ZT 9). A confocal image stack of the brain hemisphere containing l-LNvs was Z-projected into several

sequential sections. Z-sections were collected at 1 μm intervals. The signal intensity of Pdf

immunoreactivity was quantified in a manually set region of interest of the cell body region in each l-LNv

using the NIS elements Ar (Nikon). In all samples, image data were acquired under identical conditions.

When the basic data or log-transformed data were distributed normally, Student's *t*-test was used. When

they were not normally distributed, we performed the Mann–Whitney *U* test.

Sleep analysis

Single male flies (2–3 d old) were introduced into glass tubes (3 mm in diameter × 75 mm in length)

containing fly food, and the glass tubes were set in a MB5 MultiBeam Activity Monitor (Trikinetics) to

monitor the locomotor activity of individual flies. In this system, 17 independent infrared beams per glass

tube were used to detect fly movement. When a fly repositions from one beam to the next, it was counted

as one beam-crossing. Flies were acclimated in the glass tubes for 3 days in LD cycles at 25 °C before

measuring sleep amount. Locomotor activity data were collected at 1-min intervals for 5 days and

analyzed with a Microsoft Excel-based program as previously described (Kume et al., 2005). Sleep was

defined as behavioral inactivity for 5 min or more (Huber et al., 2004). Total sleep amount during the day

or night was analyzed as previously described (Shimada et al., 2016).

The Kolmogorov–Smirnov test was used to examine whether the data are normally distributed. For

comparisons of two groups, when the basic data or log-transformed data were normally distributed,

Student's *t*-test was used. When they were not normally distributed, we performed the Mann–Whitney *U*

test. In multiple group analysis of data, basic data and log-transformed data were not distributed normally.

Thus, we performed nonparametric ANOVA (Kruskal–Wallis test) followed by the rank-sum test for

multiple pairwise comparisons. The computer software IBM SPSS Statistics 22 (IBM Japan, Ltd.) was

used for these tests.

Sleep deprivation

To deprive flies of sleep, a MB5 MultiBeam Activity Monitor with glass tubes containing single naïve or

conditioned flies was horizontally shaken using a shaker (NJ-022NS, Nissin). The shaker was placed

inside a temperature-regulated incubator (MIR-254, Sanyo Electric Co., Ltd.). The shaking speed was set

to 200 rpm. On the 2nd and 3rd day after 7 h conditioning, the MB5 MultiBeam Activity Monitor was

shaken for 20 sec per 3 min during only the daytime. All experiments were carried out in 12:12 LD cycles

at 25.0 ± 0.5 °C.

Acknowledgments

We thank Yuto Kurata and Yushun Sato for technical assistance. We also thank Toshiro Aigaki for

helpful discussions. This work was supported by JSPS KAKENHI Grants.

References

References

Alberini, C.M. (2009). Transcription factors in long-term memory and synaptic plasticity. Physiol Rev *89*,

121—145.

Altimus, C.M., Guler, A.D., Villa, K.L., McNeill, D.S., Legates, T.A., and Hattar, S. (2008). Rods-cones

and melanopsin detect light and dark to modulate sleep independent of image formation. Proc Natl

Acad Sci U S A *105*, 19998–20003.

Bekinschtein, P., Cammarota, M., Igaz, L.M., Bevilaqua, L.R., Izquierdo, I., and Medina, J.H. (2007).

Persistence of long-term memory storage requires a late protein synthesis- and BDNF-dependent

phase in the hippocampus. Neuron *53*, 261–277.

Cajochen, C., Frey, S., Anders, D., Spati, J., Bues, M., Pross, A., Mager, R., Wirz-Justice, A., and Stefani,

O. (2011). Evening exposure to a light-emitting diodes (LED)-backlit computer screen affects

circadian physiology and cognitive performance. J Appl Physiol (1985) *110*, 1432–1438.

Chellappa, S.L., Ly, J.Q., Meyer, C., Balteau, E., Degueldre, C., Luxen, A., Phillips, C., Cooper, H.M.,

and Vandewalle, G. (2014). Photic memory for executive brain responses. Proc Natl Acad Sci U S A

111, 6087–6091.

Crocker, A., Guan, X.J., Murphy, C.T., and Murthy, M. (2016). Cell-type-specific transcriptome analysis

in the *Drosophila* mushroom body reveals memory-related changes in gene expression. Cell Rep *15*,

1580–1596.

Crocker, A., and Sehgal, A. (2010). Genetic analysis of sleep. Genes Dev *24*, 1220–1235.

Davis, R.L. (2011). Traces of *Drosophila* memory. Neuron *70*, 8–19.

Diao, F., and White, B.H. (2012). A novel approach for directing transgene expression in *Drosophila*:

T2A-Gal4 in-frame fusion. Genetics *190*, 1139–1144.

Donlea, J.M., Thimgan, M.S., Suzuki, Y., Gottschalk, L., and Shaw, P.J. (2011). Inducing sleep by remote

control facilitates memory consolidation in *Drosophila*. Science *332*, 1571–1576.

Dubnau, J., and Chiang, A.S. (2013). Systems memory consolidation in *Drosophila*. Curr Opin Neurobiol

23, 84–91.

Dubowy, C., and Sehgal, A. (2017). Circadian rhythms and sleep in *Drosophila melanogaster*. Genetics

205, 1373–1397.

Fernandez, D.C., Fogerson, P.M., Lazzerini Ospri L., Thomsen, M.B., Layne, R.M., Severin, D., Zhan, J.,

Singer, J.H., Kirkwood, A., Zhao, H., Berson, D.M., and Hattar, S. (2018). Light affects mood and

learning through distinct retina-brain pathways. Cell *175*, 71–84.

Fioriti, L., Myers, C., Huang, Y.Y., Li, X., Stephan, J.S., Trifilieff, P., Colnaghi, L., Kosmidis, S., Drisaldi,

B., Pavlopoulos, E., and Kandel, E.R. (2015). The persistence of hippocampal-based memory

requires protein synthesis mediated by the prion-like protein CPEB3. Neuron *86*, 1433–1448.

Fogle, K.J., Parson, K.G., Dahm, N.A., and Holmes, T.C. (2011). CRYPTOCHROME is a blue-light

sensor that regulates neuronal firing rate. Science *331*, 1409–1413.

Griffith, L.C., and Ejima, A. (2009). Courtship learning in *Drosophila melanogaster*: diverse plasticity of

a reproductive behavior. Learn Mem *16*, 743–750.

Hamada, F.N., Rosenzweig, M., Kang, K., Pulver, S.R., Ghezzi, A., Jegla, T.J., and Garrity, P.A. (2008).

An internal thermal sensor controlling temperature preference in *Drosophila*. Nature *454*, 217–220.

Hawley, W., Grissom, E., Keskitalo, L., Hastings, T., and Dohanich, G. (2011). Sexual motivation and

anxiety-like behaviors of male rats after exposure to a trauma followed by situational reminders.

Physiol Behav *102*, 181–187.

Hawley, W.R., Grissom, E.M., Belkin, M.N., James, T.F., and Dohanich, G.P. (2013). Decreased sexual

motivation and heightened anxiety in male Long-Evans rats are correlated with the memory for a

traumatic event. Arch Sex Behav *42*, 659–668.

Helfrich-Förster, C., Edwards, T., Yasuyama, K., Wisotzki, B., Schneuwly, S., Stanewsky, R.,

Meinertzhagen, I.A., Hofbauer, A. (2002). The extraretinal eyelet of *Drosophila*: development,

ultrastructure, and putative circadian function. J Neurosci *22*, 9255–9266.

Helfrich-Förster, C., Tauber, M., Park, J.H., Muhlig-Versen, M., Schneuwly, S., and Hofbauer, A. (2000).

Ectopic expression of the neuropeptide pigment-dispersing factor alters behavioral rhythms in

Drosophila melanogaster. J Neurosci *20*, 3339–3353.

Hirano, Y., Ihara, K., Masuda, T., Yamamoto, T., Iwata, I., Takahashi, A., Awata, H., Nakamura, N.,

Takakura, M., Suzuki, Y.*, et al.* (2016). Shifting transcriptional machinery is required for long-term

memory maintenance and modification in *Drosophila* mushroom bodies. Nat Commun *7*, 13471.

Horn, C., Jaunich, B., and Wimmer, E.A. (2000). Highly sensitive, fluorescent transformation marker for

Drosophila transgenesis. Dev Genes Evol *210*, 623–629.

Huber, R., Hill, S.L., Holladay, C., Biesiadecki, M., Tononi, G., and Cirelli, C. (2004). Sleep homeostasis

in *Drosophila melanogaster*. Sleep *27*, 628–639.

Hyun, S., Lee, Y., Hong, S.T., Bang, S., Paik, D., Kang, J., Shin, J., Lee, J., Jeon, K., Hwang, S.*, et al.*

(2005). *Drosophila* GPCR Han is a receptor for the circadian clock neuropeptide PDF. Neuron *48*,

267–278.

Im, S.H., and Taghert, P.H. (2010). PDF receptor expression reveals direct interactions between circadian

oscillators in *Drosophila*. J Comp Neurol *518*, 1925–1945.

Ishimoto, H., Sakai, T., and Kitamoto, T. (2009). Ecdysone signaling regulates the formation of long-term

courtship memory in adult *Drosophila melanogaster*. Proc Natl Acad Sci U S A *106*, 6381–6386.

Joiner, M.L.A., and Griffith, L.C. (1997). CaM kinase II and visual input modulate memory formation in

the neuronal circuit controlling courtship conditioning. J Neurosci *17*, 9384–9391.

Kandel, E.R. (2012). The molecular biology of memory: cAMP, PKA, CRE, CREB-1, CREB-2, and

CPEB. Mol Brain *5*, 14.

Keleman, K., Vrontou, E., Kruttner, S., Yu, J.Y., Kurtovic-Kozaric, A., and Dickson, B.J. (2012).

Dopamine neurons modulate pheromone responses in *Drosophila* courtship learning. Nature *489*,

145–149.

Kim, W.J., Jan, L.Y., and Jan, Y.N. (2013). A PDF/NPF neuropeptide signaling circuitry of male

Drosophila melanogaster controls rival-induced prolonged mating. Neuron *80*, 1190–1205.

Koemans, T.S., Oppitz, C., Donders, R.A.T., van Bokhoven, H., Schenck, A., Keleman, K., and Kramer,

J.M. (2017). *Drosophila* courtship conditioning as a measure of learning and memory. J Vis Exp. 124,

55808.

Krupp, J.J., Billeter, J.C., Wong, A., Choi, C., Nitabach, M.N., and Levine, J.D. (2013).

Pigment-dispersing factor modulates pheromone production in clock cells that influence mating in

Drosophila. Neuron *79*, 54–68.

Kruttner, S., Traunmuller, L., Dag, U., Jandrasits, K., Stepien, B., Iyer, N., Fradkin, L.G., Noordermeer,

J.N., Mensh, B.D., and Keleman, K. (2015). Synaptic Orb2A bridges memory acquisition and late

memory consolidation in *Drosophila*. Cell Rep *11*, 1953–1965.

Kume, K., Kume, S., Park, S.K., Hirsh, J., and Jackson, F.R. (2005). Dopamine is a regulator of arousal in

the fruit fly. J Neurosci *25*, 7377–7384.

Lee, S.S., Ding, Y., Karapetians, N., Rivera-Perez, C., Noriega, F.G., and Adams, M.E. (2017). Hormonal

signaling cascade during an early-adult critical period required for courtship memory retention in

Drosophila. Curr Biol *27*, 2798–2809 e2793.

Lee, Y.S., Bailey, C.H., Kandel, E.R., and Kaang, B.K. (2008). Transcriptional regulation of long-term

memory in the marine snail *Aplysia*. Mol Brain *1*, 3.

LeGates, T.A., Altimus, C.M., Wang, H., Lee, H.K., Yang, S., Zhao, H., Kirkwood, A., Weber, E.T., and

Hattar, S. (2012). Aberrant light directly impairs mood and learning through melanopsin-expressing

neurons. Nature *491*, 594–598.

Li, M.T., Cao, L.H., Xiao, N., Tang, M., Deng, B., Yang, T., Yoshii, T., and Luo, D.G. (2018)

Hub-organized parallel circuits of central circadian pacemaker neurons for visual photoentrainment

in *Drosophila*. Nat Commun *9*, 4247.

Mabuchi, I., Shimada, N., Sato, S., Ienaga, K., Inami, S., and Sakai, T. (2016). Mushroom body signaling

is required for locomotor activity rhythms in *Drosophila*. Neurosci Res *111*, 25–33.

Majumdar, A., Cesario, W.C., White-Grindley, E., Jiang, H., Ren, F., Khan, M.R., Li, L., Choi, E.M.,

Kannan, K., Guo, F*., et al.* (2012). Critical role of amyloid-like oligomers of *Drosophila* Orb2 in the

persistence of memory. Cell *148*, 515–529.

McGuire, S.E., Le, P.T., Osborn, A.J., Matsumoto, K., and Davis, R.L. (2003). Spatiotemporal rescue of

memory dysfunction in *Drosophila*. Science *302*, 1765–1768.

Mertens, I., Vandingenen, A., Johnson, E.C., Shafer, O.T., Li, W., Trigg, J.S., De Loof, A., Schoofs, L.,

and Taghert, P.H. (2005). PDF receptor signaling in *Drosophila* contributes to both circadian and

geotactic behaviors. Neuron *48*, 213–219.

Muraro, N.I., and Ceriani, M.F. (2015) Acetylcholine from visual circuits modulates the activity of

arousal neurons in *Drosophila*. J Neurosci *35*, 16315–16327.

Ni, J.D., Baik, L.S., Holmes, T.C., and Montell, C. (2017). A rhodopsin in the brain functions in circadian

photoentrainment in *Drosophila*. Nature *545*, 340–344.

Qiu, J., and Hardin, P.E. (1996). *per* mRNA cycling is locked to lights-off under photoperiodic conditions

that support circadian feedback loop function. Mol Cell Biol *16*, 4182–4188.

Renn, S.C., Park, J.H., Rosbash, M., Hall, J.C., and Taghert, P.H. (1999). A *pdf* neuropeptide gene

mutation and ablation of PDF neurons each cause severe abnormalities of behavioral circadian

rhythms in *Drosophila*. Cell *99*, 791–802.

Sakai, T., Sato, S., Ishimoto, H., and Kitamoto, T. (2012). Significance of the centrally expressed TRP

channel *painless* in *Drosophila* courtship memory. Learn Mem *20*, 34–40.

Sakai, T., Tamura, T., Kitamoto, T., and Kidokoro, Y. (2004). A clock gene, *period*, plays a key role in

long-term memory formation in *Drosophila*. Proc Natl Acad Sci U S A *101*, 16058–16063.

Shan, L.L., Guo, H., Song, N.N., Jia, Z.P., Hu, X.T., Huang, J.F., Ding, Y.Q., Richter-Levin, G., Zhou,

Q.X., and Xu, L. (2015). Light exposure before learning improves memory consolidation at night.

Sci Rep *5*, 15578.

Sheeba, V., Gu, H., Sharma, V.K., O'Dowd, D.K., and Holmes, T.C. (2008). Circadian- and

light-dependent regulation of resting membrane potential and spontaneous action potential firing of

Drosophila circadian pacemaker neurons. J Neurophysiol *99*, 976–988.

Shimada, N., Inami, S., Sato, S., Kitamoto, T., and Sakai, T. (2016). Modulation of light-driven arousal by

LIM-homeodomain transcription factor Apterous in large PDF-positive lateral neurons of the

Drosophila brain. Sci Rep *6*, 37255.

Siegel, R.W., and Hall, J.C. (1979). Conditioned responses in courtship behavior of normal and mutant

Drosophila. Proc Natl Acad Sci U S A *76*, 3430–3434.

Soler, J.E., Robison, A.J., Núñez, A.A., Yan, L. (2018). Light modulates hippocampal function and spatial

learning in a diurnal rodent species: A study using male nile grass rat (*Arvicanthis niloticus*).

Hippocampus *28*, 189–200.

Taghert, P.H., Hewes, R.S., Park, J.H., O'Brien, M.A., Han, M., Peck, M.E. (2001). Multiple amidated

neuropeptides are required for normal circadian locomotor rhythms in *Drosophila*. J Neurosci *21*,

6673–6686.

Tanenhaus, A.K., Zhang, J., and Yin, J.C. (2012). *In vivo* circadian oscillation of dCREB2 and

NF-kappaB activity in the *Drosophila* nervous system. PLoS One *7*, e45130.

Vandewalle, G., Maquet, P., and Dijk, D.J. (2009). Light as a modulator of cognitive brain function.

Trends Cogn Sci *13*, 429–438.

Yin, J. C., Wallach, J. S., Del Vecchio, M., Wilder, E. L., Zhou, H., Quinn, W. G., and Tully, T. (1994).

Induction of a dominant negative CREB transgene specifically blocks long-term memory in

Drosophila. Cell *79*, 49–58.

Yin, J.C., and Tully, T. (1996). CREB and the formation of long-term memory. Curr Opin Neurobiol *6*,

264–268.

    Figure legends

Figure 1.   Light and brain photoreceptors are essential for LTM maintenance

Box-and-whisker plots for a set of CI data show 20th, 25th, 75th, and 80th centiles. In the

box-and-whisker plots, the black square in each box indicates the mean, the line in each box is drawn at

the median, white boxes show naïve males, and gray boxes show conditioned males. The Mann–Whitney

U test was used for comparisons of CI. The permutation test with 10000 random permutations was used

for comparisons of MI among experimental conditions or genotypes. *, $P < 0.05$; **, $P < 0.01$; ***, $P <$

0.001; NS, not significant; *N*, sample size in each box. **(A)** LTM was measured under various lighting

conditions. Wild-type males were used in the experiments. Males were tested on d 5 after 7 h

conditioning. A schematic drawing of lighting conditions in courtship conditioning is shown on the left.

The white box indicates day (light) and the black box indicates night (dark). The gray box indicates that

experiments were conducted in darkness during the daytime. For MI, asterisks indicate a comparison

between the control (1) and test groups. **(B)** Memory on d 1 in wild-type (WT), cry^{01}, and $Rh7^1$ flies. **(C)**

Memory on d 5 in WT, cry^{01}, and $Rh7^1$ flies. **(D–G)** Temporal knockdown of *Rh7* or *cry* in Pdf neurons

using the TARGET system. The upper panel shows a schematic drawing of temperature shift experiments

using the temperature-sensitive GAL4 repressor *tub*-GAL80^{ts}. All flies were raised at 25°C. PT,

permissive temperature (25°C); RT, restrictive temperature (30°C). **(D)** All experiments were performed

at PT (25°C). **(E)** Flies were kept at RT for 24 h before the end of conditioning. **(F)** Flies were kept at RT

for 48–72 h after 7 h conditioning. **(G)** Flies were kept at RT for 24 h before the test. **(H)** *Pdf*-GAL4/+,

UAS-*cry* RNAi/+, and *Pdf*- GAL4/UAS-*cry* RNAi flies were used. For MI, the asterisk shows a

comparison between F_1 flies and GAL4 control, and the hashmark shows a comparison between F_1 flies

and UAS control. ###, $P < 0.001$

Figure 2.   Total sleep amount in darkness decreases in comparison with that in light.

(A–F) Sleep amount was measured for 5 d using wild-type males. Data are presented as mean ± SEM.

(A) Continuous sleep amounts of control (black line) and experimental flies (magenta line). In control

flies, sleep amount was measured in LD. In experimental flies, sleep amount was measured for two days

in LD, which was subsequently shifted to DD for two days, and then back to LD. $N = 31$ in black line; N

$= 30$ in magenta line. **(B)** Total daytime sleep amounts in control (black bars) and experimental flies

(magenta bars). *, $P < 0.05$; ***, $P < 0.001$; NS, not significant. $N = 31$ for black bars; $N = 30$ for

magenta bars. Student's *t*-test or the Mann–Whitney *U* test was used for comparisons. **(C)** Total nighttime

sleep amounts in control (black bars) and experimental flies (magenta bars). $N = 31$ for black bars; $N = 30$

for magenta bars. Student's t-test was used for comparisons. NS, not significant. **(D)** Continuous sleep

amounts in naïve (deep blue line) and conditioned flies (orange line). $N = 30$ for deep blue line; $N = 31$

for orange line. Flies were deprived of sleep during the daytime on the 2nd and 3rd days after 7 h

conditioning. SD, sleep deprivation. **(E, F)** Nonparametric ANOVA (Kruskal–Wallis test) followed by

post-hoc analysis using the rank-sum test was carried out for multiple pairwise comparisons. Bars with

the same letter indicate values that are not significantly different ($P > 0.05$). $N = 30$ for deep blue bars; N

$= 31$ for orange bars. **(E)** Total daytime sleep amounts in naïve (deep blue bars) and conditioned flies

(orange bars). **(F)** Total nighttime sleep amounts in naïve (deep blue bars) and conditioned flies (orange

bars). **(G)** Memory on d 5 was measured using sleep-deprived flies. Control flies were kept in LD without

sleep deprivation. In the box-and-whisker plots, the white boxes indicate naïve males and the gray boxes

indicate conditioned males. The Mann–Whitney U test was used for comparison of CI. The permutation

test with 10000 random permutations was used for comparisons of MI among genotypes. *, $P < 0.05$; **,

$P < 0.01$; N, sample size in each box.

Figure 3.   Real-time qRT-PCR analysis and quantitative analysis of Pdf immunoreactivity.

(A) Analysis of *Rh7* mRNA expression level. The pan-neuronal driver *nSyb*-GAL4 was used for the

induction of *Rh7* RNAi. Mean ± SEM was calculated from five replicates. **(B)** Analysis of *cry* mRNA

expression level. *nSyb*-GAL4 was used for the induction of *cry* RNAi. Mean ± SEM was calculated from

four replicates. **(A, B)** One-way ANOVA followed by post-hoc analysis using Scheffe's test was carried

out for multiple pairwise comparisons. *, $P < 0.05$; NS, not significant. **(C)** Analysis of *Pdf* mRNA

expression level. *Pdf*-GAL4 was used for the induction of *Pdf* RNAi. Mean ± SEM was calculated from

five to six replicates. Nonparametric ANOVA (Kruskal–Wallis test) followed by the rank-sum test for

multiple pairwise comparisons was used. *, $P < 0.05$; NS, not significant. **(D, E)** Analysis of *Pdfr* mRNA

expression level. *nSyb*-GAL4 (D) or MB-GAL4 OK107 (E) was used. Mean ± SEM was calculated from

six (D) or four (E) replicates. **(D)** One-way ANOVA followed by post-hoc analysis using Scheffe's test

was carried out for multiple pairwise comparisons. *, $P < 0.05$; NS, not significant. **(E)** F_1 flies between

UAS-*Pdfr* RNAi and MB-GAL4 (OK107) lines were used. Student's t-test was used for comparisons

between F_1 and GAL4 control flies. *, $P < 0.05$. **(F, G)** Quantitative analysis of Pdf immunoreactivity in

l-LNvs. Adult brains were dissected at ZT 9. In the box-and-whisker plots, white boxes indicate

Pdf-GAL4/+; *tub*-GAL80^{ts}/+ flies and gray boxes indicate *Pdf*-GAL4/ UAS-*Pdf* RNAi; *tub*-GAL80^{ts}/+

flies. **(F)** All experiments were carried out at PT (25°C). $N = 26$ for white box; $N = 27$ for gray box.

Student's *t*-test was used for statistical analysis. NS, not significant. **(G)** Flies were kept at RT (30°C) for

24 h. $N = 50$ for white box; $N = 38$ for gray box. The Mann–Whitney *U* test was used for statistical

analysis. ***, $P < 0.001$.

Figure 4.   Pdf expression is essential for LTM maintenance

(A–F, H) Males were tested on d 1 or d 5 after 7 h conditioning. In the box-and-whisker plots, the white

boxes indicate naïve males and the gray boxes indicate conditioned males. The Mann–Whitney *U* test was

used for comparisons of CI. The permutation test with 10000 random permutations was used for

comparisons of MI. *****, $P < 0.05$; ******, $P < 0.01$; *******, $P < 0.001$; ##, $P < 0.01$; NS, not significant; *N*, sample

size in each box. **(A)** All experiments were performed at PT (25°C). **(B)** Flies were kept at RT for 24 h

before the end of conditioning. **(C)** Flies were kept at RT for 48–72 h after 7 h conditioning. **(D)** Flies

were kept at RT for 24 h before the test. **(A–D)** memory on d 5 in *Pdf*-GAL4 / UAS-*Pdf* RNAi;

tub-GAL80^{ts}/+ flies and control flies (*Pdf*-GAL4 /+; *tub*-GAL80^{ts}/+). **(E)** Memory on d 1 in UAS-*Pdf* /

Pdf-GAL4; *Pdf*^{01}/*Pdf*^{01} flies and control flies (UAS-*Pdf* /+; *Pdf*^{01}/*Pdf*^{01}). **(F)** Memory on d 5 in UAS-*Pdf* /

Pdf-GAL4; *Pdf*^{01}/*Pdf*^{01} flies and control flies. **(G)** Confocal section images at Pdf neuron level of the

adult brain. Triangles, l-LNvs; arrows, s-LNvs. Scale bars represent 50 μm. UAS-*mCherry::NLS/R18F07*

flies were used. Magenta, mCherry::NLS; green, Pdf. **(H)** Knockdown of *Pdf* in l-LNvs and/or s-LNvs

using three GAL4 drivers (*Pdf*-GAL4, c929, and *R18F07*). For MI, asterisks indicate a comparison

between F_1 and GAL4 control flies, and hashmarks indicate a statistical comparison between F_1 and UAS

control flies.

Figure 5.   Activity of Pdf neurons is required for LTM maintenance

(A–D) Temporal activation of Pdf neurons compensates for DD-dependent LTM impairment.

Pdf-GAL4/+, UAS-*TrpA1*/+, and *Pdf*-GAL4/UAS-*TrpA1* males were used. For MI, asterisks indicate a

comparison between F_1 (GAL4/UAS) and GAL4 control flies, and hash marks indicate a statistical

comparison between F_1 and UAS control flies. **(A)** All experiments were performed at PT. **(B)** On the 2nd

and 3rd days after 7 h conditioning, flies were kept in DD. **(C)** On the 2nd and 3rd days after 7 h

conditioning, flies were kept at 34°C during the period between CT 0 and CT 8. **(D)** On the 2nd and 3rd

days after 7 h conditioning, flies were kept at 34°C during the period between CT 12 and CT 20. **(E, F)**

UAS-*Kir2.1*/*Pdf*-GAL4; *tub*-GAL80^{ts}/+ flies were used. UAS-*Kir2.1*/*tub*-GAL80^{ts} flies were used as the

control. **(E)** All experiments were carried out at PT (25°C). **(F)** Flies were kept at RT (32°C) for 48–72 h

after 7 h conditioning. **(G)** l-LNv-specific silencing impairs LTM. *R61G12*-LexA/LexAop-*FLPL*;

UAS>STOP>*Kir2.1*/*R14F03* flies were used. For MI, asterisks indicate a comparison between

l-LNv-silenced flies and the LexA/LexAop control (*R61G12*-LexA/LexAop-*FLPL*), and the hashmark

indicates a comparison between l-LNv-silenced flies and the GAL4/UAS control

(UAS>STOP>*Kir2.1*/*R14F03*). **(A–G)** The Mann–Whitney U test was used for comparisons of CI. The

permutation test with 10000 random permutations was used for comparisons of MI. *****, $P < 0.05$; ******, $P <$

0.01; *******, $P < 0.001$; #, $P < 0.05$; ##, $P < 0.01$; NS, not significant; N, sample size in each box. **(H)**

Confocal section images at the level of Pdf neurons of the adult brain. Scale bars represent 10 μm.

Triangles, l-LNvs; arrows, s-LNvs.

Figure 6. GAL4-expressing neurons in the adult brain in *Pdfr::2A::GAL4* flies

(A, B) *Pdfr::2A::GAL4*; UAS-*mCD8::GFP*/+ flies were used for visualizing GAL4-positive cells. Brains

were stained with an anti-GFP antibody. Stacked confocal images of the adult brain showing an anterior

(A) or posterior view (B). Scale bars, 100 μm. **(C)** *Pdfr::2A::GAL4*; *UAS-mCherry::NLS*/+ flies were

used for visualizing GAL4-positive cells. Adult brains were dissected at ZT 0–2, in which flies were

entrained in more than 3 LD cycles. All brains were stained with an anti-Per antibody. Magenta,

mCherry::NLS; green, Per. Partially stacked or sectional images of specific clusters of clock neurons

(DN1s, LNds, l-LNvs, and s-LNvs). Rectangles, DN1s; triangles, LNds or l-LNvs; arrows, s-LNvs. Scale

bars, 50 (DN1s) or 10 μm (LNds, l-LNvs, and s-LNvs).

Figure 7. Pdfr::2A::GAL4-positive MB neurons

(A) Confocal image of whole brain and sectional images at the level of Kenyon cells in MBs.

Pdfr::2A::GAL4; UAS-*mCherry::NLS*/+ ; MB-LexA LexAop-*mCD8::GFP*/+ flies were used to visualize

Pdfr-positive neurons (magenta, mCherry::NLS) and MB neurons (green, mCD8::GFP). The small

window located in the upper left in each picture shows an enlarged image of a single cell enclosed in a

small white square. Scale bars, 5 (enlarged images), 50 (Kenyon cells), or 100 μm (whole brains). **(B)**

UAS-*mCherry::NLS*/ *Pdfr*-GAL4*;* MB-LexA LexAop-*mCD8::GFP/*+ flies were used. **(C)**

Pdfr::2A::GAL4; UAS-*mCherry::NLS*/ LexAop-*mCD8::GFP* ; *R58F02*-LexA/+ flies were used.

Triangles show the colocalization of *Pdfr::2A::GAL4* and *R58F02*-LexA. **(D)** *Pdfr::2A::GAL4*;

UAS-*mCherry::NLS*/ LexAop-*mCD8::GFP* ; *R19B03*-LexA/+ flies were used. (**A–D**) Brains were

stained with an anti-GFP antibody.

Figure 8.   Pdfr expression in MB α/β neurons is essential for LTM maintenance

The Mann–Whitney U test was used for comparisons of CI. The permutation test with 10000 random

permutations was used for comparisons of MI among genotypes. *, $P < 0.05$; **, $P < 0.01$; ***, $P <$

0.001; #, $P < 0.05$; ##, $P < 0.01$; NS, not significant; N, sample size in each box. (**A**) Memory on d 5 in

Pdfr::2A::GAL4/UAS-*luc* RNAi (control) and *Pdfr::2A::GAL4*/UAS-*Pdfr* RNAi flies. (**B**) Memory on d

5 in *Pdfr^{5304}*; *Pdfr*-GAL4/+; UAS-*Pdfr* /+ and *Pdfr^{5304}*; UAS-*Pdfr* /+; OK107/+ flies. (**C**) *Pdfr* RNAi was

driven by *Pdf*-GAL4, *Pdfr* -GAL4, MB α/β -GAL4 (*R41C10*), MB γ-GAL4 (*R55D03*), and MB

α'/β'-GAL4 (c305a). For MI, asterisks indicate a comparison between F_1 and GAL4 control flies, and

hash marks indicate a statistical comparison between F_1 and UAS control flies. (**D**) Stacked confocal

images of the adult brain. *R41C10*/ UAS-*mCD8::GFP*, *R55D03*/UAS-*mCD8::GFP*, and

c305a/UAS-*mCD8::RFP* flies were used. Scale bars represent 100 μm. (**E–G**) Temporal knockdown of

Pdfr using MB-GAL4 lines. All flies were raised at 25°C. The temperature was shifted during two

experimental periods. (**E**) All experiments were performed at PT (25°C). (**F**) Flies were kept at RT

(30°C) for 24 h before the end of conditioning. (**G**) Flies were kept at RT for 48–72 h after 7 h

conditioning.

Figure 9.   GAL4-expressing neurons in the adult brain in OK107 and *R41C10*.

(**A, B**) Brains were stained with an anti-Per antibody and anti-GFP antibody. Magenta, Per; green,

mCD8::GFP. Adult brains were dissected at ZT 0–2, in which flies were entrained in more than 3 LD

cycles. Partially stacked images of specific clusters of clock neurons (DN1s, LNds, and LNvs). Stacked

confocal images showing DN1s, LNds or LNvs. Triangles, LNds or l-LNvs; arrows, s-LNvs. Scale bars,

50 μm (DN1s) or 10 μm (LNds and LNvs). (**A**) OK107/ UAS-*mCD8::GFP* flies were used. (**B**) *R41C10*/

UAS-*mCD8::GFP* flies were used.

Figure 10.   Temporal expression of CrebB repressor in MB neurons

(**A–C**) The Mann–Whitney U test was used for comparisons of CI. The permutation test with 10000

random permutations was used for comparisons of MI. For MI, asterisks indicate a comparison between

the UAS control and test groups. *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$; NS, not significant; N, sample

size in each box. All flies were raised at 25°C. The temperature was shifted during two experimental

periods. **(A)** All experiments were performed at PT (25°C). **(B)** Flies were kept at RT (30°C) for 10 h

before the end of conditioning. **(C)** Flies were kept at RT for 48–72 h after 7 h conditioning.

Figure 11.   **Light-dependent transcriptional activity of CrebB via Pdfr in MB α/β neurons**

(A) Using UAS-*FLP*/*hs-CrebB-B*; CRE>mCherry::STOP>*luc*/*R19B03* and UAS-*FLP*/+;

CRE>mCherry::STOP>*luc*/*R19B03* flies, we examined whether induction of the CrebB repressor

CrebB-B inhibits CrebB activity. $N = 9$–10 for each bar. In heat-shock treatment, males (3–4 d old) were

kept at 32°C for three days. Luc activity was measured immediately after the heat-shock treatment. Error

bars indicate SEM. Student's *t*-test was used. ***, $P < 0.001$. **(B)** CrebB activity in MB α/β or γ neurons

was measured in LD or DD using MB α/β-GAL4 (*R41C10*) or MB γ-GAL4 (*R55D03*). Samples were

prepared between ZT 0 and ZT 2. N= 8–13 in each bar. Error bars indicate SEM. One-way ANOVA

followed by post-hoc analysis using Scheffe's test was used. Bars with the same letter indicate values that

are not significantly different ($P > 0.05$). **(C)** UAS-*FLP*/*R61G12*-LexA; CRE>mCherry::STOP>*luc*

LexAop-*TrpA1*/*R41C10* and UAS-*FLP*/*R61G12*-LexA; CRE>mCherry::STOP>*luc* /*R41C10* flies were

used. $N = 5$–6 for each bar. For activation of Pdf neurons by TrpA1, males (3–5 d old) were kept at 32°C

during two days of DD. Error bars indicate SEM. Student's *t*-test was used. **, $P < 0.01$. **(D, E)**

Conditioning-dependent CrebB activity in MB α/β or γ neurons was measured immediately after 7 h

conditioning **(D)** and on the 2nd day after 7 h conditioning **(E)**. Student's *t*-test was used. ∗, $P < 0.05$; NS,

not significant. N= 6–9 in each bar.

Figure 12.   **A possible model of light-driven memory maintenance system.**

(A) During courtship conditioning, males receive stressors from mated females, and CrebB-dependent

transcription activity increases. Consequently, memory is consolidated. **(B)** During the memory

maintenance phase, light directly activates l-LNvs, and then Pdf is released. The released Pdf activates

Pdfr in MB α/β neurons, and promotes CrebB-dependent transcription required for LTM maintenance.

A
Anterior

Pdfr::2A::GAL4 / UAS-*mCD8::GFP*

B
Posterior

C

| | DN1s | LNds Section 1 | LNds Section 2 | l-LNvs | s-LNvs |

Pdfr::2A::GAL4 / UAS-*mCherry::NLS*

Section 1

GFP | mCherry::NLS | Merge | MB-LexA

Section 2

Section 3

B GFP | mCherry::NLS | Merge

C GFP | mCherry::NLS | Merge | *R58F02*-LexA

D GFP | mCherry::NLS | Merge | *R19B03*-LexA

Figure 8

A

| DN1s | LNds | LNvs |

Per

GFP

Merge

UAS-*mCD8::GFP* / OK107

B

| DN1s | LNds | LNvs |

Per

GFP

Merge

UAS-*mCD8::GFP* / *R41C10*

UAS-*CrebB-B*	+	+	+	+		+	+	+	+		+	+	+	+
tub-GAL80ts	+	+	+	+		+	+	+	+		+	+	+	+
R41C10 (α/β)		+					+					+		
R55D03 (γ)			+					+					+	
c305a (α'/β')				+					+					+

A

Stressors

♀ ♂

CREB
CRE

Consolidation

B

Light

♂

Pdf

CREB
CRE

Maintenance