An approximate full-likelihood method for inferring selection and allele frequency trajectories from DNA sequence data

Aaron J. Stern[1,*], Peter R. Wilton[2], Rasmus Nielsen[2,3]

1 Graduate Group in Computation Biology, University of California, Berkeley, Berkeley, CA 94702

2 Department of Integrative Biology, University of California, Berkeley, Berkeley, CA 94702

3 Department of Statistics, University of California, Berkeley, Berkeley, CA 94702

*Please address correspondence to `ajstern@berkeley.edu`.

Abstract

Most current methods for detecting natural selection from DNA sequence data are limited in that they are either based on summary statistics or a composite likelihood, and as a consequence, do not make full use of the information available in DNA sequence data. We here present a new importance sampling approach for approximating the full likelihood function for the selection coefficient. The method treats the ancestral recombination graph (ARG) as a latent variable that is integrated out using previously published Markov Chain Monte Carlo (MCMC) methods. The method can be used for detecting selection, estimating selection coefficients, testing models of changes in the strength of selection, estimating the time of the start of a selective sweep, and for inferring the allele frequency trajectory of a selected or neutral allele. We perform extensive simulations to evaluate the method and show that it uniformly improves power to detect selection compared to current popular methods such as nSL and SDS, under various demographic models and can provide reliable inferences of allele frequency trajectories under many conditions. We also explore the potential of our method to detect extremely recent changes in the strength of selection. We use the method to infer

the past allele frequency trajectory for a lactase persistence SNP (*MCM6*) in Europeans. 17

We also study a set of 11 pigmentation-associated variants. Several genes show evidence 18

of strong selection particularly within the last 5,000 years, including *ASIP*, *KITLG*, and 19

TYR. However, selection on *OCA2/HERC2* seems to be much older and, in contrast to 20

previous claims, we find no evidence of selection on *TYRP1*. 21

Author summary 22

Current methods to study natural selection using modern population genomic data are 23

limited in their power and flexibility. Here, we present a new method to infer natural 24

selection that builds on recent methodological advances in estimating genome-wide 25

genealogies. By using importance sampling we are able to efficiently estimate the 26

likelihood function of the selection coefficient. We show our method improves power to 27

test for selection over competing methods across a diverse range of scenarios, and also 28

accurately infers the selection coefficient. We also demonstrate a novel capability of our 29

model, using it to infer the allele's frequency over time. We validate these results with a 30

study of a lactase persistence SNP in Europeans, and also study a set of 11 31

pigmentation-associated variants. 32

Introduction

Direct observation of the change in allele frequency over time (the allele frequency
trajectory) allows one to make powerful inferences regarding whether selection acted on
the allele [1, 2]. However, outside of certain contexts such as experimental evolution of
viruses or bacteria [3–6] or analyses of ancient DNA samples [7, 8], in most cases such
direct observations of allele frequencies at multiple points in the history of a population
are unavailable. Instead, selection must be inferred from contemporary, modern data. A
wide variety of methods have been developed to detect selection based on patterns
observed from modern DNA sequences (e.g. [9–11]).

The hitch-hiking effect provides a key signature of selection in modern
datasets [12, 13]. Hitch-hiking causes aberrations in the spatial pattern of genetic
diversity, including the site frequency spectrum (SFS) [14, 15] and the pattern of
haplotype homozygosity [9]. Methods designed to detect these aberrations are
particularly useful in the setting where a single population is surveyed, and the only
information available is variation within this single population.

The most familiar methods for detecting selection are based on linear functions of
the SFS, such as Tajima's D, Fu and Li's D, or Fay and Wu's H [14, 16, 17]. An
advantage of SFS-based methods is that they do not require the data to be phased.
However, these methods have several limitations: they tend to confound selection with
other non-equilibrium conditions, such as a fluctuating population size [10, 18]; they are
not suitable for estimating parameters such as the value of the selection coefficient s;
significance can usually only be established using an empirical null distribution; and
crucially, these methods do not incorporate any features of the haplotype structure.

To make fuller use of information provided by phased sequence data, a number of
methods have incorporated summary statistics based on haplotype structure. In a broad
sense, these methods are based on calculations of haplotype similarity in a window
around some core site of interest [9]. Several methods have adapted this general concept
to specifically detect ongoing selection [11, 19, 20]. More recently, [21] showed that the
density of singletons surrounding a focal SNP can be a powerful signal of extremely
recent selection in large cohorts. In addition to recent and ongoing selection, it has been
demonstrated that these methods have compelling advantages to detecting selection

from standing variation [20–22]. However, these methods share the major limitation of SFS-based method in that they are not suitable for parametric inference and it is unclear how to establish significance without use of an empirical null model.

Recently, supervised machine learning methods have been proposed as an alternative to traditional summary-statistic based methods (see e.g., [23]). Standard machine learning techniques applied to population genomic data afford some major advantages over methods based on summary-statistics: standard techniques can produce accurate classifiers based on summaries of the data that live in much higher-dimensional space than the aforementioned summary statistics, and these techniques often encompass a wide space of classification functions that are often non-linear (see e.g. [24, 25]). Some studies have demonstrated these methods can have improved robustness to demographic model mis-specification [22, 26]. Although these methods can potentially detect complex patterns left by selection, they demand training on large data sets which typically are simulated using models that may not accurately correspond to the empirical data.

In contrast to the aforementioned methods, one might aim to develop a full likelihood method which would take into account the full data set, rather than merely summary statistics. A common strategy for obtaining the full likelihood has been to find the distribution of the genealogy under selection. For example, Krone and Neuhauser described the distribution of the coalescence tree of a locus under weak selection and no recombination [27]. Alternatively, one can describe how the genealogy depends on the trajectory of the derived allele (first described by [28]), and in turn how the trajectory depends on selection. To this end, Coop and Griffiths [29] developed a sampling method for approximating the full likelihood of the selection coefficient. Their method uses sampling to marginalize out two layers of latent variables: the allele frequency trajectory and the genealogy of the locus. To estimate the likelihood function, they perform random sampling of both the trajectory, and the genealogy conditioned on the trajectory. Unfortunately, selection likelihood methods that consider the both coalescence and recombination are generally considered computationally intractable.

Composite likelihood methods (see e.g. [10, 30]) are able to approximate the likelihood function using tractable expressions for the frequency distribution of a neutral site linked to the selected site [15, 31]. These methods approximate the joint distribution of frequencies observed at linked sites as the product of their marginals. These

approaches can be applied to test for selection, and estimate the strength of selection. The approximations made by composite likelihood methods are more accurate under strong selection (arguably beyond the strength of most recent selection in humans), and thus have less power to detect weak selection — although to some extent low power to detect weak selection is a natural outcome of any selection method.

Approximate Bayesian computation (ABC) and rejection sampling methods approximate the likelihood function by simulation. One advantage over the composite likelihood approach is that ABC can capture dependencies between linked neutral sites. For example, methods have been used to jointly infer the strength and timing of selection acting on a locus and determine whether a sweep occurred from a *de novo vs* standing variant [32–35]. However, a major disadvantage of such approaches is that the amount of simulation necessary to obtain an accurate estimate grows dramatically with the dimensionality of the model parameters. There is an additional tradeoff between information utilized from the data and computational burden; as the number of the summary statistics used increases, the number of simulations required to approximate the likelihood at a fixed parameter value also increases (for a discussion, see e.g. [36]).

The method we present in this paper draws inspiration from the Coop & Griffiths method [29], and has several key similarities: our method produces a likelihood; involves integrating out the allele frequency trajectory and genealogy, i.e., the aforementioned two hidden layers; and both methods account for selection by modeling how allele frequency changes depend on selection. However, there are several key differences between this method and our approach: while Coop and Griffiths assume no recombination of the locus, our method is based on the coalescent with recombination (i.e. the ancestral recombination graph or ARG) [37]. Also, whereas Coop & Griffiths simulate random trajectories, we use dynamic programming algorithms similar to those used in Hidden Markov Models (HMMs) to completely marginalize the latent trajectory. The hidden states represent allele frequencies and the emission probabilities are coalescence probabilities. While the framework is not a traditional HMM because the process is time-inhomogenous and the emission space changes with time, the similarities with traditional HMMs are, nonetheless, so significant that we will refer to this as an HMM. Lastly, our method uses a novel importance sampling scheme that allows us to sample ARGs assuming a neutral prior, and find the likelihood function at arbitrary

values of s; this drastically reduces the amount of ARG sampling necessary. ₁₂₈

Furthermore, the new method is, to our knowledge, the first that is capable of inferring the allele frequency trajectories for models with recombination and selection using only modern data. We are able to accomplish this task using the aforementioned Markovian structure of both coalescence and the trajectory, forming a HMM over these two hidden states and solving for the posterior marginals of each hidden allele frequency state over time. Recently, Edge & Coop proposed a method to reconstruct changes to polygenic scores over time via such estimates of the local trees, but their method is not suitable for estimating allele frequency changes or selection at individual loci [38].

Materials and methods

Overview

We begin with an overview of our method for jointly inferring selection and the allele frequency trajectory, which we summarize in Fig. 1. Our method begins with input in the form of phased SNP data from a linked genomic region (Fig. 1A), although technically, it is also possible to use unphased data, and sample possible phasings. While the method generalizes to arbitrary sample size, we recommend using $n = 25 - 100$ diploid individuals when using ARGweaver as done in this study, as ARGweaver runtime increases quadratically with sample size. The method also generalizes to arbitrarily long regions, although we recommend using regions of $10^2 - 10^3$kb, roughly the size of many LD blocks in the human genome [39].

Next, we sample the genealogy at the selected site from its posterior distribution, assuming a neutral model (Fig. 1B). By sampling over this distribution, we marginalize out the hidden coalescence events, the first of two latent variables or "hidden layers" in our model. Specifically, we sample the full ancestral recombination graph (ARG) of the input haplotypes. The ARG is a graph that summarizes all of the common ancestry and recombination events that have occurred within the sample. We sample ARGs rather than gene trees in order to account for recombination, and to incorporate information from sites in long-range linkage disequilibrium with the selected site. Then we extract the genealogy at the site of interest (the "local tree") and from here on, this is the only

component of the ARG that goes into our subsequent calculations. To perform ARG ₁₅₇ sampling, we choose to use ARGweaver [37], which is the only currently available ₁₅₈ method to sample the posterior ARG. In practice, it is possible and straightforward to ₁₅₉ adapt this method to other ARG inference methods designed for larger samples, but ₁₆₀ sampling the posterior yields beneficial statistical properties (see "Importance Sampling" ₁₆₁ under Materials and Methods). ₁₆₂

Then, for each local tree we have sampled, we form a hidden Markov model (HMM, ₁₆₃ Fig. 1C) indexed in time according to the discretization chosen for ARGweaver; in this ₁₆₄ HMM, observed states are coalescence times in this local tree and hidden states are the ₁₆₅ selected allele's frequency trajectory over time (i.e., the second hidden layer of our ₁₆₆ overall model). We use a discrete-time model of the coalescent process to match the ₁₆₇ model used by ARGweaver, so that the length of the HMM is of manageable, finite ₁₆₈ length. Emission probabilities (i.e., coalescence probabilities) depend both on the allele ₁₆₉ frequency and the current coalescent state. Hence, the model is time-inhomogenous as ₁₇₀ the coalescent state changes through time. However, the dependence structure is ₁₇₁ otherwise identical to traditional HMMs and all the usual dynamic programming ₁₇₂ algorithms apply. The transition probabilities of allele frequencies depend on the ₁₇₃ selection coefficient s, the parameter we are ultimately interested in estimating. ₁₇₄ Marginalizing out the allele frequency trajectory from the HMM yields the probability ₁₇₅ of the sample local tree as a function of s. To obtain the likelihood function of s, we ₁₇₆ perform importance sampling over all sample trees, reweighting their coalescent ₁₇₇ probabilities and summing them up. This approach allows us to use trees sampled ₁₇₈ exclusively under a prior of selective neutrality ($s = 0$) to calculate the likelihood ₁₇₉ function at arbitrary values of s. In other words, this approach allows us to minimize ₁₈₀ the amount of ARG sampling necessary to estimate the likelihood function, which is ₁₈₁ notable because ARG sampling is generally the most computationally intensive step of ₁₈₂ our method. ₁₈₃

Finally, we can analyze the results to test for selection or estimate the selection ₁₈₄ coefficient (Fig. 1D). Additionally, we show that we can decode the HMMs depicted in ₁₈₅ Fig. 1C and use them to obtain a posterior estimate of the allele frequency trajectory ₁₈₆ (Fig. 1E). ₁₈₇

A glossary to accompany the following derivations and description of the method is ₁₈₈

Fig 1. A: To apply our method for inferring selection, we begin by sampling the posterior ARG of a set of recombining chromosomes. B: For each sample ARG, we extract local trees at the site of interest (blue). C: For each sample local tree, we run an HMM to calculate the likelihood of selection, marginalizing out the hidden allele frequency trajectory based on coalescence in the sample tree. We later use the recursions performed in this step to calculate the posterior allele frequency trajectory. D: An example of the estimated likelihood function for an allele under neutrality (top) and selection (bottom). E: An example of the inferred allele frequency trajectory compared to the ground truth trajectory under neutrality (top) and selection (bottom). Both (D) and (E) are inferred from data simulated under a European demographic model with $n = 50$ haplotypes, conditioning on the derived allele segregating at 75% in the present day. with $s = 0$ and $s = 0.003$, respectively.

Coalescent model for a site under selection 190

First, let us consider how the distribution of the local tree T at a site under selection 191
depends on the frequency trajectory of an allele at that site. We assume that the tree is 192
labeled, i.e. we know which branches subtend each allele. We also assume the tree to be 193
compatible with the infinite sites assumption, i.e. that there is at most one mutation 194
event that has occurred at the focal site, and thus the site is bi-allelic. We model the 195
likelihood of the tree using a structured coalescent; moving backwards in time from the 196
time of sampling until the time of the mutation, lineages can only coalesce with other 197
lineages that subtend the same allele, and the coalescence rate within the derived and 198
ancestral classes depends on both the derived allele frequency $X(t)$ and the effective 199
population size $N(t)$, both indexed by the time $t \geq 0$ in coalescent units before the 200
present day. Proceeding back in time, lineages coalesce freely after the time of mutation, 201
and the coalescence rate depends only on $N(t)$. In the rest of this section we treat the 202
trajectory $X(t)$ as known, but in practice the trajectory is hidden and highly stochastic; 203
in a later section we develop a hidden Markov model to efficiently integrate out $X(t)$. 204

We use a discrete-time model of the coalescent employed also by ARGweaver [37]. 205
That is, we only observe the coalescent process at a set of K discrete timepoints 206
$\{t_1, \ldots, t_K\}$, and also make the additional assumption that all lineages must coalesce by 207
t_K. (Typically t_K is set to $\sim 100 \times N_e$, implying coalescence would be extremely unlikely 208
to occur after t_K, and hence this assumption is very reasonable.) Henceforth, using this 209
discretization we also discretize X and N; we assume $X(t) = X_i$ for $t \in (t_i, t_{i+1}]$, and 210
$N(t) = N_i$ for $t \in (t_i, t_{i+1}]$. 211

We use C to track the number of lineages remaining at these timepoints leading back 212
into the past; as long as we keep track of the number of lineages belonging to each of 213
the allelic classes, by exchangeability of lineages within an allelic class, we can model 214
the likelihood function in the usual way, as independent of the topology given the 215
waiting times. Hence, we define three simultaneous, related processes 216
$C = (C^{\mathrm{der}}, C^{\mathrm{anc}}, C^{\mathrm{mix}})$. The processes C^{der} and C^{anc} refer to coalescence within the 217
derived and ancestral classes during the time going back from the time of sampling to 218

the time of the mutation. The mixed process C^{mix} refers to coalescence going backwards from the time of the mutation. We call it the mixed process because it includes un-coalesced lineages from C^{anc}, as well as the lineage ancestral to all derived lineages. Assuming the infinite sites model, C^{mix} will have one additional lineage relative to C^{anc} at the time of the mutation, and will eventually reach $C^{\mathrm{anc}} = C^{\mathrm{mix}}$ once that lineage coalesces with one of the other lineages in the ancestral class. In Fig. 2 we illustrate the lines-of-descent process in the these three classes.

We model the probability of transitioning from $C_i \to C_{i+1}$ lineages during some time interval $[t_i, t_{i+1}]$ using a simple variation of Tavare's formula for the exact distribution of the number of lines of descent remaining after t generations [40]. We use Tavare's formula in order to model the coalescent at discrete timepoints, allowing multiple coalescences at each epoch.

We write the probability of C given the trajectory X (note this is distinct from the full likelihood of s) as

$$\mathbb{P}(C \mid X, N) = \prod_{i=0}^{K-1} \mathbb{P}(C_{i+1} \mid C_i, X_i, N_i) \tag{1}$$

More precisely, in terms of the derived, ancestral, and mixed processes,

$$\mathbb{P}(C \mid X, N) = \prod_{i=0}^{i^*-1} \mathbb{P}(C_{i+1}^{\mathrm{der}} \mid C_i^{\mathrm{der}}, X_i, N_i)\, \mathbb{P}(C_{i+1}^{\mathrm{anc}} \mid C_i^{\mathrm{anc}}, X_i, N_i) \times \prod_{i=i^*}^{K-1} \mathbb{P}(C_{i+1}^{\mathrm{mix}} \mid C_i^{\mathrm{mix}}, N_i) \tag{2}$$

where $i^* := \max\{i : X_i > 0\}$ denotes the index of the epoch during which the allele arose via mutation. Naturally, the mixed process—which we only keep track of while the derived allele is nonexistent—does not depend on X. We can write the transition probabilities using a variation of Tavare's formula for the transition probabilities of the number of lines of descent [40]; in place of the effective population size, we substitute the size of a allelic class z^{class} in order to reflect the coalescence rate within an allelic class:

Time	0	1	2	3	4	5
C^{der}	4	3	2	1	1	1
C^{anc}	3	3	2	2	2	1
C^{mix}	4	4	3	3	2	1

Fig 2. Top: Coalescence conditioned on the allele frequency trajectory (dashed blue line). Blue lineages subtend the derived allele, whereas black lineages do not. Black lineages belong to the ancestral class while the derived allele has $X_t > 0$, and they belong to the mixed class while $X_t = 0$. Bottom: the numbers of derived, ancestral, and mixed lineages at each time point. Numbers with unshaded cells factor into the likelihood calculation, whereas numbers with shaded cells do not.

$$\mathbb{P}(C_{i+1}^{\text{class}} = b \mid C_i^{\text{class}} = a, Z_i = z_i^{\text{class}}) \quad (3)$$

$$= \sum_{k=b}^{a} \left\{ \exp\left(\frac{-\binom{k}{2}}{2z_i}(t_{i+1} - t_i) \right) \frac{(2k-1)(-1)^{k-b}}{b!(k-b)!(k+b-1)} \prod_{l=0}^{k-1} \frac{(b+l)(a-l)}{(a+l)} \right\} \quad (4)$$

where

$$z_i^{\text{class}} = \begin{cases} N_i X_i & : \quad \text{class} = \text{der} \\ N_i(1 - X_i) & : \quad \text{class} = \text{anc} \\ N_i & : \quad \text{class} = \text{mix} \end{cases} \tag{5}$$

We note that this formula is known to be computationally unstable for large values of C, large values of N, and/or small values of $\Delta t_i = t_{i+1} - t_i$; under such conditions, the asymptotic distribution of $C_{i+1}^{\text{class}} \mid C_i^{\text{class}} = a$ (where a is, e.g., the number of derived lines of descent present at t_i) takes on a normal distribution [41]:

$$C_{i+1}^{\text{class}} \mid C_i^{\text{class}} = a \sim \mathcal{N}(\mu(\Delta t), \sigma^2(\Delta t)) \tag{6}$$

where

$$\mu(\Delta t) = \frac{2\eta}{\Delta \tau} \tag{7}$$

and

$$\sigma^2(\Delta t) = 2\eta/\Delta\tau(\eta + \beta)^2(1 + \eta/(\eta + \beta) - \eta/\alpha - \eta/(\alpha + \beta) - 2\eta)\beta^{-2} \tag{8}$$

and

$$\Delta\tau = \Delta t/(2z_i^{\text{class}}) \tag{9}$$

where $\alpha = a\Delta\tau/2$, $\beta = -\Delta\tau/2$, and $\eta = \alpha\beta/[\alpha(e^\beta - 1) + \beta e^b]$ [41]. In practice, for samples of $n = 50$ haplotypes under constant $N_e = 10^4$, we find this approximation is unnecessary; however, for the same sample size under a European demographic model, which exhibits very large recent N_e, we find it necessary to use this approximation during the roughly 10^3 generations preceding the present day, prior to which Δt is sufficiently large that we change over to Tavare's exact formula [42].

Marginalizing the hidden allele frequency states

In the previous sections we showed how we obtain $\mathbb{P}(C \mid X)$ and $\mathbb{P}(X \mid s)$. Here we illustrate how to model coalescence in the two allelic classes when the trajectory X is as a random latent variable. The probability of C given s is thus

$$\mathbb{P}(C \mid s) = \sum_{x \in \mathcal{X}} \mathbb{P}(C \mid X = x)\mathbb{P}(X = x \mid s). \qquad (10)$$

Naively, this involves a prohibitively large sum over d^{K-1} terms in $\mathcal{X}$, the space of possible trajectories. But due to the conditional independence of the likelihood, we can calculate the likelihood much faster using a recursion similar to the forward and backward algorithms commonly deployed on HMMs; we show the derivations of these recursions in S1 Text. In essence, this algorithm works by iteratively marginalizing out the allele frequency in each epoch; the backward recursion marginalizes frequency during earlier epochs first and later epochs last, and the forward recursions marginalizes in reverse order. At each epoch i the recursions yield $b_i(x_i)$ and $f_i(x_i)$; these correspond to $\mathbb{P}(C_{1:i} \mid X_i, N_{i-1})$ and $\mathbb{P}(C_{i+1:K-1}, X_i \mid X_i, N_i)$, where $C_{a:b} = C_a, C_{a+1}, \ldots, C_b$.

Consequently, the two recursions can be used together to obtain the the posterior probability of the allele frequency during the ith epoch X_i,

$$\mathbb{P}(X_i = x_i \mid C, s) = \frac{b_i(x_i)f_i(x_i)}{\sum_{x_i'} b_i(x_i')f_i(x_i')} \qquad (11)$$

which gives the posterior marginal of X_i using the familiar forward-backward algorithm.

Importance sampling to estimate the likelihood function

The above formulas pertain immediately only to the case in which the local tree is observed directly and without noise. In practical settings, the local tree is hidden to us and we must integrate over the space of possible local trees using sampling methods. Here we describe a novel importance sampling method to reweight posterior samples of the ARG to approximate the likelihood function of selection. Although we use s to express the argument of the likelihood function, we use this as shorthand for estimating

the likelihood function of arbitrarily complex parameters; for example, one could 277
estimate the selection coefficient s, as well as the time of selection's onset, t_s, before 278
which the allele behaved neutrally. 279

We are given haplotype data D representing n haplotypes with l sites that are fixed 280
for the derived allele. We wish to use D to infer the maximum-likelihood value of s for 281
some site $k \in \{1, 2, \ldots, l\}$, where l is the number of sites in the region, assuming that all 282
other sites are selectively neutral (i.e. $s_j = 0 \ \forall j \in \{1, 2, \ldots, k-1, k+1, \ldots, l\}$). In 283
other words, we restrict ourselves to testing simple hypotheses of the form "site k has 284
selection coefficient s_k and all of its flanking sites are selectively neutral." 285

The likelihood of s under the data can be expressed as the expected value of the 286
likelihood of the ARG $\mathcal{G}$ given the data D, with respect to the distribution of $\mathcal{G}$ given s: 287

$$\mathrm{L}(s) = \mathbb{E}_{\mathcal{G}|s}[\mathbb{P}(D \mid \mathcal{G}, s)] \tag{12}$$

At this stage, we introduce G, the discrete-time approximation of $\mathcal{G}$ (discussed in 288
more detail by [37]), and we assume 289

$$\mathrm{L}(s) = \mathbb{E}_{G|s}[\mathbb{P}(D \mid G, s)] \tag{13}$$

By importance sampling, we are able to express the expectation over an alternative 290
distribution $q(G)$, as long as $\mathbb{P}(G, D \mid s) > 0 \Rightarrow q(G) > 0$. Notice that this implies we 291
can conduct sampling under $q(G)$ once, and reweight these samples for arbitrary values 292
of s without having to conduct additional sampling. In other words, approximating $L(s)$ 293
using importance sampling does not require sampling under each value of s at which 294
you want to approximate $L(s)$. 295

In this paper we specifically consider the importance sampling proposal distribution 296
$q(G) = \mathbb{P}(G \mid D, s = 0)$, which corresponds to the posterior ARG assuming a neutral 297
model. Later, we evaluate the performance of the estimator using the Markov chain 298
Monte Carlo method ARGweaver, which samples from the posterior [37]. One can 299
obtain the importance sampling estimate of the full likelihood $L(s)$ by expressing Eq. 13 300

as an expectation over a different distribution, i.e. the posterior distribution of the

ARG (assuming selective neutrality):

$$L(s) = \mathbb{E}_{G|s}[\mathbb{P}(D \mid G, s)] = \mathbb{E}_{G|D,s=0}\left[\frac{\mathbb{P}(D \mid G, s)\mathbb{P}(G \mid s)}{\mathbb{P}(G \mid D, s = 0)}\right] \tag{14}$$

We can express Eq. 14 using the Monte Carlo approximation

$$\widehat{L}(s) = \frac{1}{M}\sum_{m=1}^{M}\mathbb{P}(D \mid G^{(m)}, s)\frac{\mathbb{P}(G^{(m)} \mid s)}{\mathbb{P}(G^{(m)} \mid D, s = 0)} \to L(s) \tag{15}$$

where $G^{(m)} \sim \mathbb{P}(G \mid D, s = 0)$, $m = 1, 2, \ldots, M$, and "$\to$", here and in the following, means that the left-hand side converges almost surely to the right-hand side as M goes to infinity, assuming that a Law of Large Numbers for ergodic processes holds (the Birkhoff–Khinchin theorem).

Hence, if we sample genealogies from the posterior under selective neutrality, that is, $G^{(m)} \sim \mathbb{P}(G \mid D, s = 0)$, $m = 1, 2, \ldots, M$ (where M is the number of ARGs sampled), then the right-hand side of Eq. 15 can be used as a Monte Carlo estimator of the likelihood function. However, in practice this estimator is highly unstable. However, a more stable estimator of the likelihood ratio $\frac{L(s)}{L(s=0)}$ can be derived. We can divide through Eq.14 by $L(s = 0) = \mathbb{P}(D \mid s = 0)$ to get

$$\frac{\mathrm{L}(s)}{\mathrm{L}(s = 0)} = \mathbb{E}_{G|D,s=0}\left[\frac{\mathbb{P}(D, G \mid s)}{\mathbb{P}(D, G \mid s = 0)}\right] \tag{16}$$

Because we assume the data are conditionally independent of selection given the full ARG, we can simplify this as

$$\frac{\mathrm{L}(s)}{\mathrm{L}(s = 0)} = \mathbb{E}_{G|D,s=0}\left[\frac{\mathbb{P}(D \mid G)\mathbb{P}(G \mid s)}{\mathbb{P}(D \mid G)\mathbb{P}(G \mid s = 0)}\right] = \mathbb{E}_{G|D,s=0}\left[\frac{\mathbb{P}(G \mid s)}{\mathbb{P}(G \mid s = 0)}\right] \tag{17}$$

A key development in our method is that although we sample the ARG of the entire sequence, we only calculate likelihoods using the marginal tree at the selected site,

which we will call G_k. First, let us define $G_{\backslash k}$ as the rest of the ARG omitting the local tree at site k, G_k. Consequently, G is equivalent to $(G_k, G_{\backslash k})$.

We make a key assumption that, for differing sweep parameters s and s'

$$\mathbb{P}(G_{\backslash k} \mid G_k, s) \approx \mathbb{P}(G_{\backslash k} \mid G_k, s') \tag{18}$$

That is, we assume that $G_{\backslash k}$ is approximately conditionally independent of s given the marginal tree at the selected site, G_k. Thus, we can reduce Eq. 17 to

$$
\begin{aligned}
\frac{\mathrm{L}(s)}{\mathrm{L}(s=0)} &= \mathbb{E}_{G|D,s=0}\left[\frac{\mathbb{P}(G \mid s)}{\mathbb{P}(G \mid s=0)}\right] \\
&= \mathbb{E}_{G|D,s=0}\left[\frac{\mathbb{P}(G_{\backslash k} \mid G_k, s)}{\mathbb{P}(G_{\backslash k} \mid G_k, s=0)} \frac{\mathbb{P}(G_k \mid s)}{\mathbb{P}(G_k \mid s=0)}\right] \\
&\approx \mathbb{E}_{G|D,s=0}\left[\frac{\mathbb{P}(G_k \mid s)}{\mathbb{P}(G_k \mid s=0)}\right]
\end{aligned}
$$

which suggests the following importance sampling estimator using genealogies sampled from ARGweaver will converge almost surely to a close approximation to the likelihood ratio:

$$\widehat{\mathrm{LR}}(s) = \frac{1}{M}\sum_{m=1}^{M}\frac{\mathbb{P}(G_k^{(m)} \mid s)}{\mathbb{P}(G_k^{(m)} \mid s=0)} \rightarrow \mathbb{E}_{G|D,s=0}\left[\frac{\mathbb{P}(G_k \mid s)}{\mathbb{P}(G_k \mid s=0)}\right] \approx \frac{\mathrm{L}(s)}{\mathrm{L}(s=0)} \tag{19}$$

where $G^{(m)} \sim \mathbb{P}(G|D, s=0)$ for $m = 1, 2, \ldots, M$.

Finally, due to exchangeability of lineages within the derived and ancestral allelic classes, we can assume

$$\mathbb{P}(G_k \mid s) \propto \mathbb{P}(C_k \mid s) \Rightarrow \widehat{\mathrm{LR}}(s) = \frac{1}{M}\sum_{m=1}^{M}\Omega^{(m)}(s) \tag{20}$$

where

$$\Omega^{(m)}(s) := \frac{\mathbb{P}(C_k^{(m)} \mid s)}{\mathbb{P}(C_k^{(m)} \mid s=0)} \tag{21}$$

denotes the summand of the importance sampling estimator. That is, the topology

within allelic classes is not important, and instead we need only the lines of descent $\quad$ 331

process within each class. $\qquad$ 332

We can maximize the likelihood ratio over different values of s to obtain the $\qquad$ 333

maximum-likelihood estimate of s $\qquad$ 334

$$\hat{s} = \operatorname{argmax}_s \widehat{\mathrm{LR}}(s) \tag{22}$$

Finally, we show in S1 Text that an importance sampling estimate of $\pi(x_i \mid D, s)$, $\quad$ 335

the posterior marginal of the allele frequency at timepoint i, X_i, is given by $\qquad$ 336

$$\hat{\pi}(x_i \mid D, s) := \frac{\sum_{m=1}^{M} \mathbb{P}(X_i \mid C_k^{(m)}, s)\Omega^{(m)}(s)}{\sum_{m=1}^{M} \Omega^{(m)}(s)} \tag{23}$$

where in the summand we use the posterior marginal established in Eq. 11. In practice, $\quad$ 337

we fix $s = \hat{s}$. A concern is, therefore, that this estimator does not take uncertainty in $\quad$ 338

the estimate of s into account. This problem can be addressed by using a Bayesian $\qquad$ 339

approach, which we demonstrate briefly in S1 Text. $\qquad$ 340

The method is implemented in package named `CLUES`, available for download at $\quad$ 341

`https://github.com/35ajstern/clues`, with accompanying documentation $\qquad$ 342

currently provided in S1 Text. In this paper and the currect software release we assume $\quad$ 343

positive directional selection with an additive effect on fitness, our method can be easily $\quad$ 344

extended to general dominance relationships as well as negative selection. $\qquad$ 345

Simulations $\qquad$ 346

To evaluate the power of `CLUES` to determine whether a site has been subject to $\qquad$ 347

selection, we simulated a dataset of $n = 25$ diploid individuals under two different $\qquad$ 348

demographic models; (1) a model of constant effective population size ($N = 10^4$), and $\quad$ 349

(2) a model of European (CEU) demography [43]. We performed both sets of $\qquad$ 350

simulations using the program `discoal` [44]. We set $\mu = 2r = 2.5 \times 10^{-8}$ mut/bp/gen, $\quad$ 351

$L = 1 \times 10^5$ bp or 2×10^5 bp for the constant-size and CEU models, respectively, and $\quad$ 352

simulated conditional on a variety of present-day frequencies and selection coefficients, $\quad$ 353

the latter of which we ranged from weak to strong values. Under each condition, we $\qquad$ 354

simulated 100 independent iterations. We also sampled 1 ancient haplotype; because $_{355}$ ARGweaver, which we used subsequently to sample the posterior ARG, does not $_{356}$ incorporate any information about ancestral/derived states, it is best practice to add an $_{357}$ ancient individual or outgroup to help polarize the the alleles. In practical settings $_{358}$ where the ancestral state is unknown, ARGWeaver accomodates specification of missing $_{359}$ data on the ancient haplotype. For the constant-size and CEU models, we used ancient $_{360}$ sampling dates of 2×10^4 and 1.6×10^4 generations before present, respectively. $_{361}$ Because `discoal` can only simulate piecewise-constant population sizes, we specified $_{362}$ population sizes to take on the value of their harmonic mean over the epoch, calculated $_{363}$ from the original CEU model. Commands to run simulations of trajectories, local trees, $_{364}$ and haplotypes are described in S1 Text. $_{365}$

Importantly, we conditioned simulations on the site of interest segregating at a $_{366}$ particular frequency in the present day. Hence, when we considered the power to $_{367}$ discriminate between neutral and selected alleles, we controlled the present-day $_{368}$ frequency to be equal in both of these cases. Avoiding this step would otherwise $_{369}$ upwardly bias estimates of the statistical power, due simply to the tendency for selected $_{370}$ alleles to segregate at higher frequencies than neutral alleles [45]. (If the allele frequency $_{371}$ in itself is also of interest, this part of the likelihood could trivially be added at a later $_{372}$ stage, by simply using the stationary distribution of the allele frequency; see "Allele $_{373}$ frequency transition probabilities" in S1 Text.) We then simulate the allele frequency $_{374}$ backwards in time, from the present-day frequency, until the allele reaches a frequency $_{375}$ of 0. Simulators such as `discoal` achieve this by using the conditional Wright-Fisher $_{376}$ diffusion (see e.g. [46]). In the case where effective population size changes over time, $_{377}$ running conditional simulations requires additional considerations because the $_{378}$ probability of a mutation entering the population scales approximately linearly with $_{379}$ population size. Naively sampling the trajectory backwards in time will therefore $_{380}$ produce a bias, unless trajectories where the mutation occurs while N_e is low are $_{381}$ somehow penalized. Thus, approaches such as reweighting sample trajectories using $_{382}$ importance sampling have been used to correct this bias [47]. The program `discoal` $_{383}$ implements a similar bias-correcting scheme using rejection sampling that rejects $_{384}$ trajectories where the mutation occurs while N_e is low with higher probability than $_{385}$ trajectories where the mutation occurs while N_e is high. $_{386}$

Next, we inferred the posterior ARG given the sequence data we simulated using ARGweaver [37]. This method works by proposing adjustments to an initial ARG, and randomly accepting or rejecting these proposals based on calculations of the prior probability of the proposed ARG, as well as its likelihood given the sequence data. Because the prior probability is based on the effective population size, we specified the same effective population size in the prior as we used to generate the sequence data. We found it important to adjust the proposal mechanism of ARGweaver; specifically, we adjusted resample window size and the number of resamples per window to achieve an acceptance rate of about 30-70%. In total, we sampled 3×10^3 ARGs for each simulation, discarding the first 1×10^3 as a burn-in period, and subsequently thinning the remaining samples to reduce the computational burden of downstream analyses; we used a thinning rate of 100 samples, resulting in $M = 20$ approximately independent samples. Reducing the thinning rate would increase accuracy and convergence of the inference at the cost of additional computation to calculate the likelihood of each additional sample tree. Commands to conduct ARG-sampling and local tree extraction are described in S1 Text.

Using utilities in the ARGweaver package, we extracted local trees at the selected site (at the center of the locus) from these sample ARGs. We then analyze this final set of trees using CLUES. We also analyzed the same sequence data using nSL, H_{12}, and Tajima's D [14, 19, 48]. The nSL method is essentially equivalent to iHS [11], except nSL does not require specifying a genetic map; despite this, these methods have been shown to have very similar statistical power with a slight advantage of nSL under some conditions. H_{12} is a method to calculate haplotype homozygosity merging the two most common haplogroups; thus, it is a test for selection that is robust to the origin of a sweep, i.e. whether it is hard or soft. Tajima's D is a site frequency spectrum-based statistic which is sensitive to skews in the frequency distribution of linked alleles caused by hitchhiking on the partially swept selected allele. We used scripts provided by [22] to calculate D and H_{12}, using a window size of 100kb centered on the selected site. We compare testing for selection under these methods by comparing their power curves under both the constant N_e and CEU demography models (Figs. 3,4).

We also conducted a similar simulation study for detecting recent selection starting 100 generations ago. We simulated under the same CEU demographic model as

previously described, but instead sample $n = 50$ diploids. We conducted ARG sampling and thinning as previously described, but in our analysis of the sample trees using CLUES, we calculated the likelihood for models of selection where $s = 0$ up until 100 generations ago, and $s \geq 0$ from that point until the present day. This sweep from standing variation (SSV) model differs from the hard sweep model we used previously, which assumes s is constant throughout history. Instead of optimizing the likelihood function only with respect to s, we optimized with respect to two parameters, s and t_s, jointly; here t_s represents the time of the onset of selection.

Results

Testing for selection

We found that across all scenarios, CLUES matches or exceeds the statistical power of the other methods evaluated (Figs. 3,4). As expected, all methods had highest power under large values of both the selection coefficient and the derived allele frequency (Fig 3I). Under these conditions, CLUES had 100% power at the 1% significance threshhold; the next most powerful method, nSL, had 68% power at the same significance level. CLUES also demonstrated improvement in power under weak selection; as the selection coefficient was decreased, nSL retained about 20% power when $s = 0.003$ and <5% power when $s = 0.001$, and Tajima's D and H_{12} retained <5% power under both $s = 0.001, 0.003$ (Fig 3G,H). By contrast, CLUES retained approximately 45% and 90% power under $s = 0.001, 0.003$, respectively. We conclude that CLUES has high power across a wide regime of selection strengths, and has notably improved power over standard methods under weaker values of s.

We also considered the effect of present-day allele frequency on statistical power. Previous studies have shown a strong dependence of power on current allele frequency, with methods such as nSl and iHs having highest power at allele frequencies in the 70-90% range (see e.g. [11]). We tested for selection at alleles ranging in present day frequency from 25% to 75%, and while CLUES showed the expected pattern of increasing power with frequency, it also improved on the performance of other methods at lower frequencies. For example, under strong selection ($s = 0.01$), the power of CLUES changed

from 100% to 90% to 85% as the frequency is decreased from 75% to 50% to 25% (Fig. 3C,F,I). By contrast, the power of the next most powerful method, H_{12}, dropped from approximately 65% to 45% to 15% (Fig. 3C,F,I). Under moderate selection ($s = 0.003$), these effects were even more drastic, with the power of CLUES and nSL (the next most powerful method in this regime) changing from 90% to 60% to 50% and 20% to 5% to <5%, respectively. We conclude that CLUES has high power compared to standard methods across a wide range of allele frequencies, with the most major improvements in performance occurring when the derived allele is at lower frequencies (<50%). We found that using the approximation due to Griffiths (Eq. 6, [41]) decreased power of CLUES by increasing variability of the null distribution of the likelihood ratios. Hence, for testing under nonequilibrium demography we used the exact lines-of-descent probabilities (Eq. ??). By contrast, as we will later show, we found the approximation given by Eq. 6 for $t \in [0, 1000]$ to improve estimation of allele frequency trajectories under this demographic model.

We also considered the same testing procedure under non-equilibrium demography, simulating under the previously described model of CEU demography (Fig. 34). We found in general reduced power to detect selection under this regime relative to the constant population size regime (Fig. 4I, cf. Fig. 3I), consistent with the well-known confounding of expanding population size with selection [10]. Nonetheless, CLUES demonstrated improved power relative to the competing methods across a wide range of selection coefficients (Fig. 4C,F,I), as well as across a wide range of derived allele frequencies (Fig. 4G,H,I).

Estimating selection coefficients

Using the simulations from the previous section to study statistical power in testing for selection, we used our estimate of the likelihood surface for s to estimate the value of the selection coefficient via maximum likelihood (see Eq. 22). We obtained selection coefficient estimates under importance sampling using ARGweaver (Fig. 5), as well as selection coefficient estimates based on the true local tree observed directly (S1 Fig). Generally, the estimates are approximately unbiased. For example, the mean estimates of $s = 0, 1 \times 10^{-3}, 3 \times 10^{-3}, 1 \times 10^{-2}$ were approximately

Fig 3. ROC curves illustrating performance of tests between selection and neutrality. Rows correspond to simulations conditioned on the same present-day allele frequency, and columns correspond to simulations with the same value of s. Simulations were performed under a model of constant effective population size ($N_e = 10^4$) using a locus of 100kb, $n = 25$ diploid individuals and $\mu = 2.5 \times 10^{-8}$ mut/bp/gen, $r = 1.25 \times 10^{-8}$ recombinations/bp/gen.

$\bar{\hat{s}} = 1.9 \times 10^{-4}, 9.6 \times 10^{-4}, 3.2 \times 10^{-3}, 1.3 \times 10^{-2}$ when the present day frequency was fixed to 75% (Fig. 5A). Relative to inference when the true tree is observed, we found that the importance sampling estimates had increased variance, reflecting uncertainty in the tree. For example, we saw increased variability in the importance sampling vs. true tree estimates under constant population size (Fig. 5A vs. S1 FigA), as well as under CEU demography (Fig. 5B vs. S1 FigB). This pattern is consistent with the additional uncertainty in s when the local tree is not observed directly. Notably, we found that importance sampling under a model of CEU demography yields estimates with a slight bias towards lower values of s, especially under strong selection (e.g. $s = 0.01$).

Fig 4. ROC curves illustrating performance of tests between selection and neutrality. Rows correspond to simulations conditioned on the same present-day allele frequency, and columns correspond to simulations with the same value of s. Simulations were performed under a model of European demography using a locus of 200kb, $n = 25$ diploid individuals and $\mu = 2.5 \times 10^{-8}$ mut/bp/gen, $r = 1.25 \times 10^{-8}$ recombinations/bp/gen.

Fig 5. Inference of selection coefficients of varying strength using importance sampling method based on ARGweaver local trees. A: Constant population size. B: Tennessen CEU model. Marker color denotes present-day allele frequency (25/50/75% correspond to yellow/red/purple, respectively).

Inferring allele frequency trajectories

Using the same simulations and importance sampling estimates we obtained in the 488
previous sections, we decoded the hidden Markov model (HMM) described in the 489
section Materials & Methods. Specifically, we take $\hat{s}$, the maximum likelihood estimate 490
of s, and plug it into the posterior marginal (Eq. 11) to obtain a probabilistic estimate 491
of the allele frequency during a particular epoch; we do this independently for each 492
epoch in our discrete-time model. To get a point estimate, we choose to use the 493
posterior marginal mean; i.e., for each epoch, we choose the mean of the posterior 494
marginal distribution. We illustrate the accuracy of these allele frequency trajectory 495
estimates assuming the true local tree is observed and under importance sampling when 496
the true tree is unknown in Fig. 6. We find that estimates of the allele frequency 497
trajectory are generally unbiased for both true trees (Fig. 6 A,B) and importance 498
sampling (Fig. 6 C,D), with increased variance in the trajectory estimates in the 499
importance sampling setting. We also illustrated variability in true vs. inferred 500
trajectories controlling for s (S6 Fig, here setting $s = 0$). 501

Whereas inference tended to be relatively accurate for high-frequency alleles (Fig. 6 502
B,D), when the derived allele was simulated conditioned on lower frequencies (e.g. 25%, 503
Fig. 6 A,C), estimates tend to be downwardly biased. We tracked this bias to a lack of 504
convergence in ARGweaver; specifically, we found that across different demographic 505
scenarios and selection coefficients, ARGweaver can drastically overestimate the 506
occurrence of very recent coalescences (in our case, in the last 100 generations; see S5 507
Fig). Under constant population size, we see a nearly 7-fold excess in the number of 508
recent coalescences inferred by ARGweaver. Naturally, this bias will affect estimates for 509
low-frequency alleles more strongly, as fewer lineages subtend the derived allele, and 510
thus a larger proportion of them are susceptible to this bias. 511

Because recombination rates vary substantially throughout the genomes of humans 512
and other organisms, we also evaluated the accuracy of the estimates assuming $\mu = r$, 513
larger than the $\mu = 2\rho$ setting we used in the other simulations, and estimation 514
accuracy to be robust to this increase in recombination rate (S2 Fig). 515

We also examined trajectory inference under non-equilibrium demography; i.e., the 516
aforementioned model of CEU demography (S3 Fig). Under the CEU model, we found 517

trajectory estimates to have increased variance under importance sampling vs. true ^{518}
trees, but also a slight downward bias in estimating the selection coefficient under ^{519}
strong selection (i.e. $s = 0.01$; see Fig. 5B, S3 Fig D). As this bias does not occur under ^{520}
the true trees (S1 Fig B, S3 Fig B), we inspected the posterior trees sampled by ^{521}
ARGweaver for patterns consistent with this bias. We found that under this ^{522}
demographic model in particular, ARGweaver tends to under-sample trees with short ^{523}
times to most recent common ancestor (TMRCAs; see S4 Fig). For reference, nearly ^{524}
60% of runs under constant N_e contained even a single sample tree that had a TMRCA ^{525}
less than or equal to that of the true TMRCA (S4 Fig A). By comparison, under ^{526}
$s = 0.01$ and CEU demography, only 11% of ARGweaver runs met this criterion (S4 Fig ^{527}
B). Some bias is to be expected, as trees were sampled under a posterior distribution ^{528}
that assumes selective neutrality; however, these results suggest that, if ARGweaver is ^{529}
sampling from the true posterior assuming selective neutrality, then importance ^{530}
sampling estimates (of the selection coefficient, for example) will at least have much ^{531}
higher variance under the CEU model than under constant population size. ^{532}

We further investigated whether uncertainty in s due to importance sampling ^{533}
variance drove the downward bias when estimating strong selection (Fig. 5B and S3 Fig ^{534}
D). First, we obtained importance sampling estimates of the trajectory fixing s to its ^{535}
true value (S7 Fig A). If uncertainty in s were the cause of the bias, then fixing the true ^{536}
value of s ought to correct for bias due to uncertainty. While we observe less bias in the ^{537}
estimates when fixing the true value of s, the bias is not totally eliminated. We observe ^{538}
a similar reduction in the bias of estimates under neutrality when we fix $s = 0$ (see S6 ^{539}
Fig B,E,H, vs. S6 Fig C,F,I). Thus, we conclude the bias is due to a lack of convergence ^{540}
in ARGweaver, which appears to be exacerbated in settings where strong selection is ^{541}
combined with non-equilibrium demography. ^{542}

We also investigated whether incorporating uncertainty in the estimate of s, rather ^{543}
than fixing $s = \hat{s}$, would improve the accuracy of trajectory inference. One strategy for ^{544}
modeling uncertainty in s is to apply a prior distribution to s. We found that ^{545}
marginalizing out s with respect to its posterior distribution (assuming a uniform prior ^{546}
on s) did not have a noticeable effect on inference for large values of s (S7 Fig B). This ^{547}
result is concordant with our observation that for large values of s, the likelihood ^{548}
surface peaks so strongly that the posterior remains tightly concentrated around the ^{549}

MLE $\hat{s}$. Hence, applying a prior distribution to s does not appear to be an adequate strategy to model uncertainty in s.

Fig 6. Allele frequency trajectories inferred from true trees (top row) and ARGweaver local trees (bottom row). Stepwise trajectories are inferred (vertical bars denote 25th-75th percentiles), dashed trajectories are the ground truth. Columns correspond to different initial allele frequencies (left: 25%, center: 50%, right: 75%), and rows/colors correspond to different selection coefficients. For each condition we show 25 randomly selected simulations and their corresponding inferences. All data are simulated under a model of constant effective population size ($N_e = 10^4$).

Inferring extremely recent selection

We applied our likelihood model of a sweep from a standing variant (SSV) to two types of datasets: selection from a standing variant starting 100 generations ago and selection with constant s (including $s = 0$), both described in 'Simulations' under Materials and Methods. We inferred trajectories under the best case scenario where the true trees are observed (Fig. 7A,B). We found that overall the method inferred the trajectory, as well as the strength and timing of selection, with highest accuracy when selection is strong (e.g. $s = 0.03$ in Fig. 7A,B). However, we found that as s took on smaller values ($s = 0.01$), many combinations of s and t_s had very similar likelihood (Fig. 7B), and thus estimates of s, t_s, and the allele frequency trajectory tended to be noisier than under very strong selection (Figs. 7A,B). Adding the extra parameter t_s did not cause

overfitting when inferring the trajectories of hard sweeps (Fig. 7A). We also found good
power to distinguish between hard vs. soft sweeps (i.e. sweeps from a standing variant),
as apparent in the trajectories inferred in Fig. 7A. We calculated statistical power to
test for a hard sweep using the statistic $\max_{s,t_s}\{L(s,t_s)\}\big/\max_{s}\{L(s,t_s=\infty)\}$; intuitively,
this statistic is the ratio of the highest likelihood under any model with a SSV ($t_s \neq \infty$)
to the highest likelihood of any hard sweep ($t_s = \infty$). At the 1% significance level we
found 60% and 100% power to distinguish soft vs. hard sweeps with $s = 0.01, 0.03$,
respectively.

We also performed importance sampling using ARGweaver and evaluated the power
of the importance sampling estimates to detect recent selection vs. neutrality (Fig. 7C).
Instead of comparing our method to nSL, which is not designed to detect signals of
extremely recent selection, we compared to Singleton Density Score (SDS; [21]), as well
as H_{12} and Tajima's D. We found that for lower values of s, all methods had generally
low power. Although CLUES exhibited fairly high power (44%) to detect very strong
recent selection ($s = 0.03$) —even outperforming SDS—we found that H_{12} has about
the same power (45%) in this particular case. The lower power ($<5\%$) of SDS is
consistent with the fact that the method was explicitly designed to have high power for
large datasets ($n > 1000$ for selection coefficients of this magnitude). Although we
demonstrate that CLUES has substantial power to detect extremely recent selection, we
found that importance sampling point estimates of s, t_s, and the trajectory were highly
vulnerable to biases in the distribution sampled by ARGweaver (S5 Fig). Specifically,
we found that across various demographic and selection conditions, ARGweaver samples
trees with substantially more recent coalescent events than in the true trees.
Specifically, under the European demographic model with the settings used here to
study recent selection, we find ARGweaver samples about a 4-fold excess of recent
coalescent events (S5 Fig B). Clearly, this bias would produce a false signature of recent
selection under neutral conditions. Thus, we did not further explore importance
sampling estimates of s and the trajectory under the recent selection model. We
conclude that potential ARG-sampling methods that avoid this bias will improve upon
power to detect recent selection, as well as point estimates of the strength, timing of
selection, and the allele frequency trajectory.

Fig 7. (A) Trajectories inferred from true trees under both hard sweeps and recent selection on a standing variant (i.e. soft sweeps) when both s and time of selection onset are unknown. (B) The log-likelihood surface for joint inference of s and onset of selection, averaged over 100 simulations, taking the true tree as observed. (C) ROC curves (using importance sampling) illustrating performance of tests between selection from a standing variant where onset of selection occurs 100 generations ago. We condition on a present day frequency of 50%.

Fine-mapping the selected site

Strongly selected alleles tend to have high levels of LD to linked neutral sites, and thus many methods to detect selection are limited in their ability to determine the exact site

under selection. To assess whether the likelihood ratio statistic produced by CLUES can be used to fine-map the selected site, we ran CLUES at linked neutral sites in a locus centered on a site under positive selection. Simulations were identical to those used in the simulation study under constant $N_e = 10^4$, with a present-day selected allele frequency of 75%, $s = 0.01$, and 100 independent simulations. We chose sites with the maximal squared correlation coefficient r^2 to the selected allele, such that r^2 did not exceed a threshold value, and vary that threshold from 0.50 to 0.99 (Fig. 8). We found that when the true tree is observed (or sampled with high accuracy), the likelihood ratio statistic identifies the selected site correctly in a head-to-head test with the neutral linked site with 85% accuracy even when $r^2 \leq 0.99$; this quantity reaches 100% for $r^2 \leq 0.50$ (Fig. 8A). When the likelihood ratio is estimated using importance sampling via ARGweaver, the accuracy declines to about 50% and 85%, respectively (Fig. 8A). Because the exact causal site may not be known in many studies, we also investigated how the estimate of the selection coefficient, s, depends on r^2 between the site analyzed and the site under selection. We estimate s given the true tree, and find that, on average, estimates of s decline with r^2, such that for $r^2 \leq 0.50$, the mean estimate of s at these neutral sites is less than 20% of the true value of s at the causal site (Fig. 8B).

Effects of background selection

We also assessed the effects of linked selection on inference of selection at a particular site. In particular, we consider the effects of background selection (BGS) on inference of positive selection on a linked site (S8 Fig). We performed forward simulations in SLiM 3 [49], assuming a model of constant $N_e = 10^3$, a locus of length 1Mb, $n = 25$ diploid individuals and $\mu = 2.5 \times 10^{-8}$ mut/bp/gen, $r = 1.25 \times 10^{-8}$ recombinations/bp/gen. We let mutations be neutral with probability 90% and (negatively) selected with probability 10%. We simulated under deleterious effects of $s = 0, -0.001, -0.003$ and -0.01, performing 100 independent replicates under each case. To simulate selection at a focal allele, 100 generations prior to the present day, we choose a random neutral allele conditional on its frequency falling in the interval [0.005,0.015] and its position falling in the interval $[4 \times 10^5, 6 \times 10^5]$bp (to ensure it is somewhat centered in the 1Mb region), and then endow it with a selection coefficient of $s = 0.15$. Prior to this timepoint, we

Fig 8. (A) Probability of correctly identifying the selected site in a head-to-head test with a linked neutral site. Vertical bars represent 95% CIs estimated by plug-in bootstrap. (B) Estimates of the selection coefficient at the causal vs. linked neutral sites, given the true tree. Mean estimates are represented by blue hash marks.

perform a burn-in phase of 19900 generations with only neutral and deleterious   627

mutation. From the sampled haplotypes, we perform importance sampling using   628

ARGweaver to estimate the likelihood of the selection coefficient.   629

We find that our method is quite robust to BGS; e.g., the median estimate of the selection coefficient is approximately unbiased (mean $\hat{s} = 0.13$) as the strength of BGS is increased (S8 Fig). Also, regardless of BGS strength there is nearly perfect power to detect positive selection when comparing to neutral simulations (S8 Fig). We note that the strength of selection on the beneficial allele investigated here is somewhat strong; for weaker selection on a beneficial allele, inference with sites under BGS may be a more significant determinant of the beneficial allele's trajectory (see e.g. [50]). We also note that our simulations assume a model fo equilibrium demography, but under non-equilbrium conditions (e.g., rapid population size expansion) BGS has a magnified effect on neutral diversity, which may further bias estimates of selection [51].

Effects of demographic model misspecification

To explore the effects of demographic model misspecification on inference of selection, we ran CLUES on datasets simulated under a model of European demography (described earlier in Methods), using a mismatched model of constant $N_e = 10^4$ to calculate the likelihood, and compare them to calculations under the true demographic model (S9 Fig). We report likelihood ratios (S9 Fig A,C) as well as estimates of the selection coefficient s (S9 Fig B,D) both given the true tree (S9 Fig A,B) and approximated via importance sampling using ARGweaver (S9 Fig C,D). We find that statistical power to detect selection, especially strong selection, is not substantially impeded by model misspecification (S9 Fig A,C). We do, however, find upward bias in estimates of the selection coefficient when s is 0 or close to 0 (S9 Fig B,D).

Analysis of a lactase persistence SNP

To assess performance of CLUES on empirical data, we applied our method to study selection acting on the SNP rs4988235 in the *MCM6 gene*, known to regulate the neighboring *LCT* gene and affect the lactase persistence trait. The derived allele (A) current segregates at approximately 72% in the 1000 Genomes Phase 3 reference panel (British in England and Scotland, henceforth GBR; see S10 Fig A). We conducted sampling in ARGweaver assuming a model of European demography [43], using a 300kbp region centered around the focal SNP and polarizing alleles using the genomes

of three ancient individuals (Altai Neandertal, Denisova, and Vindija 659
Neandertal [52–54]). We sampled $M = 200$ ARGs, extracted local trees using tools in 660
the ARGweaver package, and conducted importance sampling to estimate likelihood 661
surfaces and trajectories using CLUES. 662

We found very strong evidence for selection on rs4988235 ($s = 0.0161$, 663
$\log \mathrm{LR} = 131.82$). The trajectory as well as the value of the selection coefficient inferred 664
by CLUES are consistent with previous estimates of the trajectory and $s = 0.018$ due to 665
Mathieson and Mathieson (2018), illustrated in Fig. 9 [55]. Their method incorporates 666
genomic times series spanning thousands of generations using an HMM-based approach, 667
where hidden states are population-wide allele frequencies, observed states are 668
genotypes of sampled ancient individuals, and transition probabilities are governed by 669
the selection coefficient. Our approach, by contrast, does not utilize any 670
ancient/timecourse data except for the 3 aforementioned ancient individuals, which we 671
use to simply polarize the derived and ancestral states of each allele. 672

Fig 9. Comparison of inferred allele frequency trajectories for a sweep at rs4988235 (MCM6) in GBR under an ancient DNA (aDNA) based method vs. CLUES, which only uses contemporary modern data. Black curve is the posterior median allele frequency, whereas gray areas are a 95% posterior interval. The red surface is the posterior of the frequency trajectory within Steppe ancestry conditioned on an ancient DNA time series, adapted from [55].

Analysis of *EDAR* 673

Another canonical example of a common SNP under selection in humans in rs3827760 674
(see e.g. [56]). This SNP codes for a V→A change that is present at high frequency in 675
East Asian populations (e.g. 94% in Han Chinese [CHB] in the 1000 Genomes 676

database), intermediate-high frequency in Central and South America, and low 677

frequency in other geographical regions S10 Fig. This variant is associated with a 678

number of traits, including tooth shape and hair straightness [57, 58]. To estimate 679

selection on this SNP, we conducted sampling in ARGweaver assuming a model of East 680

Asian demography [59], using a 300kbp region centered around the focal SNP and 681

polarizing alleles using the genomes of three ancient individuals (Altai Neandertal, 682

Denisova, and Vindija Neandertal [52–54]). We sampled $M = 200$ ARGs, extracted 683

local trees using tools in the ARGweaver package, and conducted importance sampling 684

to estimate likelihood surfaces and trajectories using CLUES. We estimate that rs3827760 685

has undergone selection with $s = 0.0047$, corresponding to an allele age of roughly 45kya 686

(S11 Fig). Our estimates are in stark contrast to some previous estimates obtained 687

using ABC methods, which estimate > 30-fold stronger selection on this SNP, and an 688

allele age of 1.4-6.9kya [32]. Our results are consistent with ancient DNA evidence, 689

which suggest the derived allele to have originated prior to 7.5kya [8]. 690

Analysis of pigmentation alleles 691

Using the same GBR panel from 1000 Genomes Phase 3, we analyzed a set of SNPs 692

associated with pigmentation-related traits, some of which were previously identified as 693

likely targets of recent selection [21]. We conducted sampling in ARGweaver assuming a 694

model of European demography, using a 300kbp region centered around the focal SNP 695

and sampling $M = 200$ approximately iid ARGs. We ran CLUES and estimated likelihood 696

surfaces and allele frequency trajectories for these SNPs (Fig. 10). We found significant 697

concordance between the SDS values and our likelihood ratio statistics paired for each 698

SNP ($p = 1.7 \times 10^{-3}$, Spearman one-sided) [21]. We also illustrated the geographical 699

distribution of these SNPs among diverse populations (S12 Fig) using GGV [60]. 700

We found several signals of very strong selection acting on rs619865 (*ASIP*, $s \approx 0.10$, 701

Fig. 10I), rs12821256 (*KITLG*, $s \approx 0.016$, Fig. 10H), and rs1393350 (*TYR*, $s \approx 0.011$, 702

Fig. 10J); these SNPs are significantly associated with freckling, blonde hair color, and 703

freckling and blue/green eye color, respectively [61–63]. Interestingly, these SNPs all 704

demonstrated a signal of selection mostly concentrated in the last $\sim$5 kya. The 705

geographical distribution of the frequency of these SNPs shows that the derived version 706

of these variants are mostly concentrated in European populations, with minimal   707
sharing with populations located in Africa and Asia (S12 Fig I,H,J). For example, *TYR*   708
and *KITLG* segregate at a frequency ~20% in several European populations and have a   709
frequency close to 0% in African and East Asian populations (S12 Fig J). These three   710
SNPs are the only ones in this set of SNPs which have a frequency of nearly 0% across   711
the African populations surveyed, with the exception of OCA2/HERC2 (S12 Fig   712
A,H,I,J), consistent with our evidence for recent selection at these loci. The frequencies   713
of these variants in GBR ranges from ~10-20%; by contrast, the only other variant in   714
this set with comparable frequency in GBR (13%), rs35264875 (*TPCN2*), we find   715
inconclusive evidence of selection (Fig. 10F), consistent with its comparably even   716
geographical distribution relative to the aforementioned SNPs at *ASIP, KITLG*, and   717
TYR (S12 Fig F).   718

At rs12896399 (*SLC24A4*, Fig. 10B), a SNP identified to be significantly associated   719
with hair color [62], we found strong evidence for moderate selection ($s \approx 0.005$). This   720
result is consistent with a previous analysis that suggested positive selection acted on   721
this allele in Out-of-Africa (OoA) populations, based on its high allele differentiation   722
relative to a YRI panel, and low haplotype diversity within CEU individuals [63]. Our   723
results, paired with the apparent low levels of differentiation between European and   724
Asian populations relative to differentiation between OoA populations and African   725
populations at this locus (S12 Fig B) are consistent with our estimate that selection   726
acted on *SLC24A4* as early as ~30 kya, during the OoA bottleneck as inferred   727
by [43,59].   728

Notably, we find moderate evidence for selection on rs12913832 (*OCA2/HERC2*,   729
Fig. 10 A, S13 Fig), a SNP previously shown to be causal for blue-brown eye color [64]   730
and significantly associated with hair color [62]. This gene exhibits abberantly high   731
differentiation across populations [65], consistent with a model of local adaptation of eye   732
color. Compared to previous estimates based on ancient DNA samples [66], we estimate   733
substantially weaker selection acting on this gene ($s \approx 0.002$ vs. $s \approx 0.04$), and we find   734
no evidence to support a recent increase in selection acting on this SNP (i.e., our   735
method found a hard sweep to have higher likelihood than a SSV). Our estimate of   736
moderate selection and lack of a recent change in the selection coefficient imply that   737
selection on OCA2/HERC2 began at least ~50 kya, roughly the time of the start of the   738

OoA bottleneck estimated by [43,59]. Our analysis suggests that selection on ⁷³⁹

OCA2/HERC2 may have begun much earlier than previously suggested [66]. We also ⁷⁴⁰

note that the aforementioned rs12913832 (*OCA2/HERC2*)—as well as rs2153271 ⁷⁴¹

(*BNC2*), a SNP which is significantly associated with freckling (Fig. 10 G)—occur in ⁷⁴²

high-frequency archaic haplotypes [67]. While our method is not explicitly designed to ⁷⁴³

control for population structure between archaic and modern human lineages, we do ⁷⁴⁴

find moderate evidence for selection on both of these SNPs. ⁷⁴⁵

One surprising result is that we found no signal of selection acting at rs13289810 ⁷⁴⁶

(*TYRP1*, $s \approx 0$, Fig. 10E). In Europeans, *TYRP1* is associated with hair and eye ⁷⁴⁷

pigmentation [68–71]. Some analyses of European populations have indicated evidence ⁷⁴⁸

for positive selection on *TYRP1* [56,63,69]. Our results temper these claims, and ⁷⁴⁹

appear consistent with the fairly even geographical distribution of rs13289810 frequency ⁷⁵⁰

across European, African, and Middle Eastern populations (S12 Fig E). ⁷⁵¹

Fig 10. Allele frequencies trajectories inferred for 11 pigmentation-associated SNPs in GBR.

Discussion

We have developed an approach to use modern population genomic data to approximate the full likelihood of selection acting on a locus. We use this approach to test for and estimate the strength and timing of selection, as well as estimate the full allele frequency trajectory. The method is effective across a span of selection coefficients ($s = 0 - 0.01$), derived allele frequencies ($f = 25\% - 75\%$), and under multiple demographic models.

Our method draws on previously published methods to estimate the ancestral recombination graph (ARG). We chose to use ARGweaver because it is the only currently available method for sampling the posterior of the ARG; as shown in our derivation of the importance sampling estimates, we rely on sampling from the posterior in order to make rigorous guarantees regarding convergence and consistency of our estimators. Intuitively, it is important to model the uncertainty in the local tree in order to marginalize out this latent variable. We showed that estimates of the selection coefficient and the trajectory are generally accurate, barring scenarios where importance sampling is inefficient, or ARGweaver produces a bias in the inferred trees. In light of these biases, under certain conditions—primarily when the derived allele is at low frequencies ($\leq 25\%$)—importance sampling using ARGweaver trees has limited power to detect selection.

Another important limitation of ARGweaver is its computational cost; in order to study selection on short timescales, large sample sizes are necessary, often on the order of thousands of individuals [21]. The runtime of ARGweaver grows dramatically with increasing sample size; not only does the cost of the individual sampling steps increase with sample size, but also so does the size of the state space, necessitating more samples be taken in order to achieve convergence to the stationary distribution.

However, we see potential to make use of recent advances in inference of local trees in order to further advance approximate full-likelihood methods to infer selection (see e.g., [72–75]; it is worth noting that some of these methods, such as [75], do not infer the ARG in a strict sense, but rather the sequence of local trees along a recombining locus). A major benefit of these methods is that they are far more scalable than ARGweaver, and hence offer more potential to study selection on short, punctuated timescales. However, they also possess several limitations: Firstly, several of these

methods only infer topologies, rather than branch lengths [73, 74]. While it is possible to infer branch lengths condition on topology estimates, it is unclear how accurate these estimates would be. By contrast, methods that infer branch lengths along with topology entail a slight tradeoff in their scalability [72, 75]. Another limitation of these methods it that they only yield a point estimate of the local tree, rather than estimating uncertainty in the tree. Nonetheless, it may be feasible to quantify uncertainty in the local tree using a jackknife approach where the local tree is inferred over random subsets of the individuals.

It may also be possible to make use of recent advances in inferring pairwise coalescence times (e.g., [76]) to build an approximation to the full likelihood. Recently, Albers & McVean proposed a composite likelihood method to estimate allele age by "sandwiching" the age using identity-by-descent tracts at the site of interest [77]. However, their method does not extend to inferring how the allele frequency changed over time, and does not explicitly model selection.

Currently our method assumes correct knowledge of the demographic history. The effects of latent or mis-specified population structure on inference of selection are well known (e.g., [78]), but in future work one might try to determine the exact effects of mis-specification of effective population size on both inferring the local tree, and inferring selection conditional on the local tree. One approach to dealing with this is to extend the importance sampling approach we use to correct for selection to additionally correct for demography, when ARG sampling is performed under a mis-specified demographic model.

Furthermore, many aspects of our model of selection (e.g. coalescence, allele frequency transitions) assume a panmictic population. To extend our model to more complex demographic models and/or linked selection (i.e., allowing multiple sites to be subject to selection) would entail drastically increased computational cost (e.g., marginalizing allele frequencies corresponding to each population, rather than the allele frequency in a single population). Using a deterministic approximation of the allele frequency trajectory would circumvent this issue, but it would also raise new issues, such as how to model allele frequencies when $s = 0$.

Despite its limitations, the method presented here provides the first close approximation to a full likelihood function for the selection coefficient under simple

models. As demonstrated by our simulations, full likelihood methods have the potential to greatly improve power to detect selection and estimate the strength of selection under a variety of conditions. It also provides a rigorous and accurate method for estimating allele frequency trajectories, and is the first to achieve so using modern data. As methods for inferring ARGs improve in the future, so too will the derived methods for detecting and quantifying selection and inferring allele frequency changes.

Acknowledgments

We thank Melissa Hubisz and Andrew Kern for extensive help with the software packages ARGweaver and Discoal, respectively. We also thank Graham Coop, Michael Edge, Yun Song, as well as Vladimir Shchur and other members of the Nielsen Lab, for helpful discussions.

Funding

RN was supported by (NIH) R01GM116044.

Author contributions

AJS: Conceptualization, formal analysis, investigation, methodology, software, validation, visualization, writing (original draft), writing (review and editing); PRW: Software, writing (original draft); RN: Conceptualization, methodology, supervision, writing (original draft), writing (review and editing).

Potential competing interests

We have no competing interests to report.

References

1. Watterson GA. Testing Selection at a Single Locus. Biometrics. 1982;38(2):323–331.

2. Mathieson I, McVean G. Estimating selection coefficients in spatially structured populations from time series data of allele frequencies. Genetics. 2013;193(3):973–984.

3. Williamson EG, Slatkin M. Using Maximum Likelihood to Estimate Population Size From Temporal Changes in Allele Frequencies. 1999;.

4. Bollback JP, York TL, Nielsen R. Estimation of 2Nes from temporal allele frequency data. Genetics. 2008;179(1):497–502. doi:10.1534/genetics.107.085019.

5. Lang GI, Rice DP, Hickman MJ, Sodergren E, Weinstock GM, Botstein D, et al. Pervasive genetic hitchhiking and clonal interference in forty evolving yeast populations. Nature. 2013;500(7464):571.

6. Good BH, McDonald MJ, Barrick JE, Lenski RE, Desai MM. The dynamics of molecular evolution over 60,000 generations. Nature. 2017;551(7678):45.

7. Lazaridis I, Patterson N, Mittnik *et al* Alissa. Ancient human genomes suggest three ancestral populations for present-day Europeans. Nature. 2014;513(7518):409–13. doi:10.1038/nature13673.

8. Mathieson I, Lazaridis I, Rohland N, Mallick S, Patterson N, Roodenberg SA, et al. Genome-wide patterns of selection in. Nature. 2015;528(7583):499–503. doi:10.1038/nature16152.

9. Sabeti PC, Reich DE, Higgins JM, Levine HZP, Richter DJ, Schaffner SF, et al. Detecting recent positive selection in the human genome from haplotype structure. 2002;419(October). doi:10.1038/nature01027.1.

10. Nielsen R, Williamson S, Kim Y, Hubisz MJ, Clark AG, Bustamante C. Genomic scans for selective sweeps using SNP data. 2005; p. 1566–1575. doi:10.1101/gr.4252305.

11. Voight BF, Kudaravalli S, Wen X, Pritchard JK. A map of recent positive selection in the human genome. PLoS biology. 2006;4(3):e72. doi:10.1371/journal.pbio.0040072.

12. Smith JM, Haigh J. The hitch-hiking effect of a favourable gene. Genetics Research. 1974;23(1):23–35.

13. Kaplan NL, Hudson R, Langley C. The" hitchhiking effect" revisited. Genetics. 1989;123(4):887–899.

14. Tajima F. Statistical method for testing the neutral mutation hypothesis by DNA polymorphism. Genetics. 1989;123(3):585–595.

15. Stephan W, Wiehe TH, Lenz MW. The effect of strongly selected substitutions on neutral polymorphism: analytical results based on diffusion theory. Theoretical Population Biology. 1992;41(2):237–254.

16. Fu YX, Li WH. Statistical tests of neutrality of mutations. Genetics. 1993;133(3):693–709.

17. Fay JC, Wu Ci. Hitchhiking Under Positive Darwinian Selection. 2000;.

18. Teshima KM, Coop G, Przeworski M. How reliable are empirical genomic scans for selective sweeps ? 2006;(773):702–712. doi:10.1101/gr.5105206.702.

19. Ferrer-Admetlla A, Liang M, Korneliussen T, Nielsen R. On detecting incomplete soft or hard selective sweeps using haplotype structure. Molecular biology and evolution. 2014;31(5):1275–1291.

20. Garud NR, Messer PW, Buzbas EO, Petrov Da. Recent selective sweeps in North American Drosophila melanogaster show signatures of soft sweeps. PLoS genetics. 2015;11(2):e1005004. doi:10.1371/journal.pgen.1005004.

21. Field Y, Boyle EA, Telis N, Gao Z, Gaulton KJ. Detection of human adaptation during the past 2 , 000 years. 2016; p. 1–18.

22. Schrider DR, Kern AD. S/HIC: Robust Identification of Soft and Hard Sweeps Using Machine Learning. PLoS genetics. 2016;12(3):e1005928. doi:10.1371/journal.pgen.1005928.

23. Schrider DR, Kern AD. Supervised Machine Learning for Population Genetics: A New Paradigm. Trends in Genetics. 2018;.

24. Lin K, Li H, Schlötterer C, Futschik A. Distinguishing positive selection from neutral evolution: boosting the performance of summary statistics. Genetics. 2011;187(1):229–244.

25. Ronen R, Udpa N, Halperin E, Bafna V. Learning natural selection from the site frequency spectrum. Genetics. 2013;195(1):181–193.

26. Sheehan S, Song YS. Deep learning for population genetic inference. PLoS computational biology. 2016;12(3):e1004845.

27. Krone SM, Neuhauser C. Ancestral processes with selection. Theoretical population biology. 1997;51(3):210–237.

28. Kaplan NL, Darden T, Hudson RR. The Coalescent Process in Models With Selection. 1988;829(2):819–829.

29. Coop G, Griffiths RC. Ancestral inference on gene trees under selection. Theoretical population biology. 2004;66(3):219–32. doi:10.1016/j.tpb.2004.06.006.

30. Vy HMT, Kim Y. A composite-likelihood method for detecting incomplete selective sweep from population genomic data. Genetics. 2015;200(2):633–649.

31. Kim Y, Stephan W. Detecting a Local Signature of Genetic Hitchhiking Along a Recombining Chromosome. 2002;777(February):765–777.

32. Peter BM, Huerta-Sanchez E, Nielsen R. Distinguishing between selective sweeps from standing variation and from a de novo mutation. PLoS genetics. 2012;8(10):e1003011.

33. Ormond L, Foll M, Ewing GB, Pfeifer SP, Jensen JD. Inferring the age of a fixed beneficial allele. Molecular ecology. 2016;25(1):157–169.

34. Ilardo MA, Moltke I, Korneliussen TS, Cheng J, Stern AJ, Racimo F, et al. Physiological and genetic adaptations to diving in sea nomads. Cell. 2018;173(3):569–580.

35. Corl A, Bi K, Luke C, Challa AS, Stern AJ, Sinervo B, et al. The genetic basis of adaptation following plastic changes in coloration in a novel environment. Current Biology. 2018;28(18):2970–2977.

36. Sugden LA, Atkinson EG, Fischer AP, Rong S, Henn BM, Ramachandran S. Localization of adaptive variants in human genomes using averaged one-dependence estimation. Nature communications. 2018;9(1):703.

37. Rasmussen MD, Hubisz MJ, Gronau I, Siepel A. Genome-wide inference of ancestral recombination graphs. PLoS genetics. 2014;10(5):e1004342. doi:10.1371/journal.pgen.1004342.

38. Edge MD, Coop G. Reconstructing the history of polygenic scores using coalescent trees. Genetics. 2019;211(1):235–262.

39. Berisa T, Pickrell JK. Approximately independent linkage disequilibrium blocks in human populations. Bioinformatics. 2016;32(2):283.

40. Tavaré S. Line-of-descent and genealogical processes, and their applications in population genetics models. Theoretical population biology. 1984;26(2):119–164.

41. Griffiths R. Asymptotic line-of-descent distributions. Journal of Mathematical Biology. 1984;21(1):67–75.

42. Jewett EM, Rosenberg NA. Theory and applications of a deterministic approximation to the coalescent model. Theoretical population biology. 2014;93:14–29.

43. Tennessen JA, Bigham AW, O'connor TD, Fu W, Kenny EE, Gravel S, et al. Evolution and functional impact of rare coding variation from deep sequencing of human exomes. science. 2012;337(6090):64–69.

44. Kern AD, Schrider DR. Discoal: flexible coalescent simulations with selection. Bioinformatics. 2016;32(24):3839.

45. Wright S. The distribution of gene frequencies under irreversible mutation. Proceedings of the National Academy of Sciences. 1938;24(7):253–259.

46. Przeworski M. The signature of positive selection at randomly chosen loci. Genetics. 2002;160(3):1179–1189.

47. Slatkin M. Simulating genealogies of selected alleles in a population of variable size. Genetics Research. 2001;78(1):49–57.

48. Garud NR, Messer PW, Buzbas EO, Petrov DA. Recent selective sweeps in North American Drosophila melanogaster show signatures of soft sweeps. PLoS Genet. 2015;11(2):e1005004.

49. Haller BC, Messer PW. SLiM 3: forward genetic simulations beyond the Wright–Fisher model. Molecular biology and evolution. 2019;36(3):632–637.

50. Barton NH. Linkage and the limits to natural selection. Genetics. 1995;140(2):821–841.

51. Torres R, Szpiech ZA, Hernandez RD. Human demographic history has amplified the effects of background selection across the genome. PLoS genetics. 2018;14(6):e1007387.

52. Meyer M, Kircher M, Gansauge MT, Li H, Racimo F, Mallick S, et al. A high-coverage genome sequence from an archaic Denisovan individual. Science. 2012;338(6104):222–226.

53. Prüfer K, Racimo F, Patterson N, Jay F, Sankararaman S, Sawyer S, et al. The complete genome sequence of a Neanderthal from the Altai Mountains. Nature. 2014;505(7481):43.

54. Prüfer K, de Filippo C, Grote S, Mafessoni F, Korlević P, Hajdinjak M, et al. A high-coverage Neandertal genome from Vindija Cave in Croatia. Science. 2017;358(6363):655–658.

55. Mathieson S, Mathieson I. FADS1 and the Timing of Human Adaptation to Agriculture. Molecular Biology and Evolution. 2018;35(12):2957–2970. doi:10.1093/molbev/msy180.

56. Grossman SR, Shylakhter I, Karlsson EK, Byrne EH, Morales S, Frieden G, et al. A composite of multiple signals distinguishes causal variants in regions of positive selection. Science. 2010;327(5967):883–886.

57. Kimura R, Yamaguchi T, Takeda M, Kondo O, Toma T, Haneji K, et al. A common variation in EDAR is a genetic determinant of shovel-shaped incisors. The American Journal of Human Genetics. 2009;85(4):528–535.

58. Wu S, Tan J, Yang Y, Peng Q, Zhang M, Li J, et al. Genome-wide scans reveal variants at EDAR predominantly affecting hair straightness in Han Chinese and Uyghur populations. Human genetics. 2016;135(11):1279–1286.

59. Gravel S, Henn BM, Gutenkunst RN, Indap AR, Marth GT, Clark AG, et al. Demographic history and rare allele sharing among human populations. Proceedings of the National Academy of Sciences of the United States of America. 2011;108(29):11983–8. doi:10.1073/pnas.1019276108.

60. Marcus JH, Novembre J. Visualizing the geography of genetic variants. Bioinformatics. 2017;33(4):594–595.

61. Eriksson N, Macpherson JM, Tung JY, Hon LS, Naughton B, Saxonov S, et al. Web-based, participant-driven studies yield novel genetic associations for common traits. PLoS genetics. 2010;6(6):e1000993.

62. Han J, Kraft P, Nan H, Guo Q, Chen C, Qureshi A, et al. A genome-wide association study identifies novel alleles associated with hair color and skin pigmentation. PLoS genetics. 2008;4(5):e1000074.

63. Sulem P, Gudbjartsson DF, Stacey SN, Helgason A, Rafnar T, Magnusson KP, et al. Genetic determinants of hair, eye and skin pigmentation in Europeans. Nature genetics. 2007;39(12):1443.

64. Sturm RA, Duffy DL, Zhao ZZ, Leite FP, Stark MS, Hayward NK, et al. A single SNP in an evolutionary conserved region within intron 86 of the HERC2 gene determines human blue-brown eye color. The American Journal of Human Genetics. 2008;82(2):424–431.

65. Huerta-Sánchez E, Jin X, Bianba Z, Peter BM, Vinckenbosch N, Liang Y, et al. Altitude adaptation in Tibetans caused by introgression of Denisovan-like DNA. Nature. 2014;512(7513):194.

66. Wilde S, Timpson A, Kirsanow K, Kaiser E, Kayser M, Unterländer M, et al. Direct evidence for positive selection of skin, hair, and eye pigmentation in Europeans during the last 5,000 y. Proceedings of the National Academy of Sciences. 2014;111(13):4832–4837.

67. Gittelman RM, Schraiber JG, Vernot B, Mikacenic C, Wurfel MM, Akey JM. Archaic hominin admixture facilitated adaptation to out-of-Africa environments. Current Biology. 2016;26(24):3375–3382.

68. Frudakis T, Thomas M, Gaskin Z, Venkateswarlu K, Chandra KS, Ginjupalli S, et al. Sequences associated with human iris pigmentation. Genetics. 2003;165(4):2071–2083.

69. Sulem P, Gudbjartsson DF, Stacey SN, Helgason A, Rafnar T, Jakobsdottir M, et al. Two newly identified genetic determinants of pigmentation in Europeans. Nature genetics. 2008;40(7):835.

70. Liu F, Wollstein A, Hysi PG, Ankra-Badu GA, Spector TD, Park D, et al. Digital quantification of human eye color highlights genetic association of three new loci. PLoS genetics. 2010;6(5):e1000934.

71. Kenny EE, Timpson NJ, Sikora M, Yee MC, Moreno-Estrada A, Eng C, et al. Melanesian blond hair is caused by an amino acid change in TYRP1. Science. 2012;336(6081):554–554.

72. Mirzaei S, Wu Y. RENT+: an improved method for inferring local genealogical trees from haplotypes with recombination. Bioinformatics. 2016;33(7):1021–1030.

73. Kelleher J, Wong Y, Albers P, Wohns AW, McVean G. Inferring the ancestry of everyone. BioRxiv. 2018; p. 458067.

74. Shchur V, Ziganurova L, Durbin R. Fast and scalable genome-wide inference of local tree topologies from large number of haplotypes based on tree consistent PBWT data structure. bioRxiv. 2019; p. 542035.

75. Speidel L, Forest M, Shi S, Myers S. A method for genome-wide genealogy estimation for thousands of samples. BioRxiv. 2019; p. 550558.

76. Palamara PF, Terhorst J, Song YS, Price AL. High-throughput inference of pairwise coalescence times identifies signals of selection and enriched disease heritability. bioRxiv. 2018; p. 276931.

77. Albers PK, McVean G. Dating genomic variants and shared ancestry in population-scale sequencing data. bioRxiv. 2018;doi:10.1101/416610.

78. Galtier N, Depaulis F, Barton NH. Detecting bottlenecks and selective sweeps from DNA sequence polymorphism. Genetics. 2000;155(2):981–987.

Supporting information

S1 Fig.   Selection coefficients inferred directly from the true local trees.
Left: constant population size ($N_e = 10^4$). Right: Tennessen CEU demographic model.
Marker color denotes present-day allele frequency (25/50/75% correspond to
yellow/red/purple, respectively).

**S2 Fig.   Allele frequency trajectories inferred from ARGweaver local trees
when** $\mu = r$**.** We set $\mu = 2.5 \times 10^{-8}$ mut/bp/gen, $r = 2.5 \times 10^{-8}$
recombinations/bp/gen and fix the present day allele frequency to $X_0 = 50\%$ Stepwise
trajectories are inferred, dashed trajectories are the ground truth. Vertical bars denote
the 25-75th percentile range of estimates. For each condition we show 20 randomly
selected simulations and their corresponding inferences. All data simulated under a
demographic model with constant size $N = 10^4$.

S3 Fig.   Inferring allele frequency trajectories under CEU demography.
Trajectories were inferred from true local trees (top row) and importance sampling on
ARGweaver local trees (bottom row). Columns correspond to different present-day
allele frequencies (left: 50%, right: 75%). For each condition we show 20 randomly
selected simulations (dashed, translucent lines) and their corresponding inferences
(piecewise constant curves; dots and vertical bars indicate the median and 25-75
percentiles of estimates, respectively). The gray box indicates the timing of the
bottleneck, occurring approximately 920-2040 generations ago. Simulations were done
under the European demographic model described in Methods and Materials using a
locus of 200kb, $n = 25$ diploid individuals and $\mu = 2.5 \times 10^{-8}$ mut/bp/gen,
$r = 1.25 \times 10^{-8}$ recombinations/bp/gen.

S4 Fig.   ARGweaver proposes less accurate trees under non-equilibrium   862

demography. Left: constant $N_e = 10^4$. Right: Tennessen CEU demographic model.   863

We found in Fig. S2 Fig that importance sampling using ARGweaver tends to   864

underestimate the selection coefficient under a model of CEU demography. To   865

demonstrate that the proposal distribution for sampling the local tree is the source of   866

this bias, we use TMRCA of the local tree as a heuristic the locus's selection coefficient.   867

For the sake of argument, we postulate that as one decreases the TMRCA of a local   868

tree, the maximum-likelihood estimate of the selection coefficient strictly descreases. If   869

so, then if the minimum value of the sampled TMRCAs is greater than the true   870

TMRCA, then this instance of the importance sampling estimate will underestimate the   871

selection coefficient. Hence, one can measure importance sampling efficiency by looking   872

at the probability that the minimum value of the sampled TMRCAs is less than the   873

true TMRCA. This is shown graphically by the proportion of points that fall in the red   874

upper triangle. The selection coefficients $s = 0, 0.001, 0.003, 0.01$ are indicated by   875

purple, blue, green, and orange, respectively. Simulations were done under the   876

European demographic model described in Methods and Materials using a locus of   877

200kb, $n = 25$ diploid individuals and $\mu = 2.5 \times 10^{-8}$ mut/bp/gen, $r = 1.25 \times 10^{-8}$ 878

recombinations/bp/gen. We fixed the present-day allele frequency to be 75%.   879

S5 Fig.   ARGweaver infers an excess of recent coalescences. As a diagnostic

for the trees outputted by ARGweaver, we compared the amount of coalescence in the

sample trees vs. the local trees. Let $a_{i,s}$ be the number of coalescent events during

epoch i in the sth replicate of the simulation. We calculate

$$e_i = \frac{\frac{1}{M} \sum_{s=1}^{S} \sum_{m=1}^{M} a_{i,s}^{(m)}}{\sum_{s=1}^{S} a_{i,s}}$$

as an estimate of the "fold excess" of coalescence, where $a_{i,s}^{(m)}$ denotes the mth   881

ARGweaver sample of $a_{i,s}$. Notice that if the sample trees closely approximate the true   882

tree, then $e_i \approx 0$. The dashed line indicates no excess, i.e., no bias in the estimates. We   883

find that ARGweaver can have a nearly $4\times$ excess of inferred coalescence events in the   884

most recent epochs (e.g. [0,100] generations ago). Here we simulated recent selection   885

starting 100 generations ago under a model of European demography with $n = 50$ 886

diploid individuals, a physical length of 200kb, and $\mu = 2.5 \times 10^{-8}$ mut/bp/gen,   887

$r = 1.25 \times 10^{-8}$ mut/bp/gen. We condition on the variant having a present-day   888

frequency of 50%.   889

S6 Fig. Performance of trajectory inference across replicates. We illustrated inferred vs. true trajectories, holding selection coefficient constant across the replicates. In this case, we chose $s = 0$ in order to maximize the amount of variability in the trajectories of the replicates. Rows correspond to simulations conditioned on different present-day allele frequencies (0.25: A-C,0.50: D-F,0.75: G-I). Left column (A,D,G): trajectories inferred using the true local tree. Middle column (B,E,H): trajectories inferred using importance sampling, fixing the selection coefficient to the ground truth ($s = 0$). Right column (C,F,I): trajectories inferred under importance sampling and estimating s. Simulations were done under the constant size model described in Methods and Materials using a locus of 100kb, $n = 25$ diploid individuals and $\mu = 2.5 \times 10^{-8}$ mut/bp/gen, $r = 1.25 \times 10^{-8}$ recombinations/bp/gen.

S7 Fig. Effect of uncertainty in s on trajectory inference. A: trajectories inferred using importance sampling (i.e. Eq. 23), fixing $\hat{s} = s$. B: trajectories inferred using importance sampling and integrating over a uniform prior on s (see S1 Text for formulae). Simulations were done under the European demographic model described in Methods and Materials using a locus of 200kb, $n = 25$ diploid individuals and $\mu = 2.5 \times 10^{-8}$ mut/bp/gen, $r = 1.25 \times 10^{-8}$ recombinations/bp/gen.

S8 Fig.   Effect of background selection on inference of selection. Estimates   910
of the selection coefficient of a selected allele ($s = 0.1$) linked to sites under background   911
selection of varying strength. Simulations were done using forward simulations assuming   912
constant $N_e = 10^3$ using a locus of 1Mb, $n = 25$ diploid individuals and $\mu = 2.5 \times 10^{-8}$ 913
mut/bp/gen, $r = 1.25 \times 10^{-8}$ recombinations/bp/gen. Mean estimates are represented   914
by blue hash marks.   915

True trees

Importance sampling

S9 Fig. Effect of model misspecification on inference of selection.

Likelihood ratios (A,C) and selection coefficient estimates (B,D) calculated given the true tree (A,B) and via importance sampling using ARGweaver (C,D). Blue violin plots represent estimates using the correct (European) demographic model, whereas orange plots represent estimates using a model of constant $N_e = 10^4$. Simulations were done under the European demographic model described in Methods and Materials using a locus of 200kb, $n = 25$ diploid individuals and $\mu = 2.5 \times 10^{-8}$ mut/bp/gen,

$$r = 1.25 \times 10^{-8} \text{ recombinations/bp/gen.}$$

(A) LCT (rs4988235)

(B) EDAR (rs3827760)

S10 Fig.   Geographical distribution of pigmentation SNPs. Population-wide   926
allele frequencies of (A) rs4988235 (MCM6) and (B) rs3827760 (EDAR) plotted   927
geographically using GGV.   928

S11 Fig.   Allele frequency trajectory of rs3827760 (EDAR) in CHB.   930
Frequency trajectory inferred using importance sampling under a model of CHB   931
demography.   932

S12 Fig. Geographical distribution of pigmentation SNPs. Population-wide allele frequencies of pigmentation SNPs from Fig. 10 plotted geographically using GGV.

S13 Fig.   Allele frequency trajectory estimate of rs12913832
(OCA2/HERC2). The same trajectory estimate as in Fig. 10A with x-axis limits
extended to illustrate earlier history of the allele.

S1 Text. Appendix and commands to reproduce simulations and analyses.