In vitro Evaluation of Antifungal Drug Combinations against

Multidrug-resistant *Candida auris* isolates from New York Outbreak

Brittany O'Brien[1], Sudha Chaturvedi[1,2], Vishnu Chaturvedi[1,2]*

Mycology Laboratory[1], Wadsworth Center New York State Department of Health, and

Department of Biomedical Sciences[2], University at Albany School of Public Health,

Albany, NY, USA

Running title: *C. auris* antifungal combination

*Corresponding author: Mycology Laboratory, Wadsworth Center, New York State

Department of Health, 120 New Scotland Ave., Albany, NY 12208, USA

Tel: +1-518-474-4177; FAX: +1-518-486-7971; E-mail: Vishnu.Chaturvedi@health.ny.gov

orcid.org/0000-0002-3922-9676

Abstract

Since 2016, New York hospitals and healthcare facilities have faced an unprecedented

outbreak of pathogenic yeast *Candida auris*. We tested over one thousand *C. auris*

isolates from affected facilities and found high-resistance to fluconazole (FLC, MIC_{50}

>256 mg/L), and variable resistance to other antifungal drugs. Therefore, we evaluated

if two-drug combinations are effective *in vitro* against multidrug-resistant *C. auris*. Broth

micro-dilution (BMD) plates were custom-designed, and quality controlled by TREK

Diagnostic System. We used MIC_{100} endpoints for the drug combination readings as

reported earlier for the intra- and inter-laboratory agreements against *Candida* species

and *Aspergillus fumigatus*. The study results were derived from 12,960 MIC_{100} readings,

for fifteen *C. auris* isolates tested against 864 possible two-drug antifungal combinations

for nine antifungal drugs. Flucytosine (5FC) at 1.0 mg/L potentiated the most successful

combinations with other drugs. Micafungin (MFG), Anidulafungin (AFG), Caspofungin

(CAS) at individual concentrations of 0.25 mg/L in combination with 5FC (1.0 mg/L)

yielded MIC_{100} for 14, 13, and 12 of 15 *C. auris* test isolates. AMB / 5FC (0.25/1.0 mg/L)

yielded MIC_{100} for 13 isolates. None of the combinations were effective for *C. auris* 18-1,

which tested resistant against FLC and 5FC, except POS/5FC (0.12/1.0 mg/L). The

simplified two-drug combination susceptibility test format would permit laboratories to

provide clinicians and public health experts with additional data to deal with

multidrug-resistant *C. auris*.

Introduction

Candida auris is an emerging multidrug-resistant yeast, causing invasive healthcare-associated infection with high mortality worldwide (1). It was described in 2009 as a new yeast species from the ear discharge of a Japanese patient (2). In the ensuing years, many reports of multidrug-resistant C. auris infections came from South Asia, South Africa, South America and travel-related cases from other parts of the world (3-6). The largest, localized, uninterrupted C. auris outbreak in the US healthcare facilities continues to afflict the New York metro areas (2013-2019); C. auris caused 23 deaths among the first 51 clinical case-patients in New York (7). In response, the New York State Department of Health (NYSDOH) laboratory scientists and epidemiologists developed enhanced protocols for the unprecedented surveillance and testing for C. auris (8). To date, over 15,000 clinical samples from 150 facilities were processed in the NYSDOH laboratory. The CLSI broth microdilution method (BMD) was used to test nearly one thousand C. auris isolates, which revealed high-resistance to fluconazole (MIC_{50} >256 mg/L), and variable resistance to other broad-spectrum antifungal drugs (Chaturvedi et al. unpublished data). Therefore, we investigated the efficacy of two-drug combinations against a representative set of fifteen drug-resistant C. auris.

Materials and Methods

Isolates–*C. auris* isolates were initially processed on Sabouraud dulcitol agar at 40℃,

and single colonies used to confirm identify with MALDI-TOF-MS and ITS -D1/D2

sequencing as described earlier. The isolates were stored at -80℃ in 20% sterile glycerol.

Drug combinations–Fluconazole (FLC) was not tested in combinations because of wide-

spread resistance in our isolate collection. Nine drugs were selected for the combination

testing: amphotericin B (AMB), anidulafungin (AFG), caspofungin (CAS), micafungin

(MFG), flucytosine (5FC), isavuconazole (ISA), posaconazole (POS), voriconazole (VRC)

and itraconazole (ITC). An advisory from the Centers for Diseases Control and

Prevention was used for C. auris antifungal susceptibility testing and interpretation

(https://www.cdc.gov/fungal/candida-auris/c-auris-antifungal.html).

Combination testing–Broadly, we followed a method described earlier from a multi-

laboratory evaluation of antifungal combination testing (9). Two fixed concentrations of

each drug were tested, a value conforming to Cmax in the product insert, and a

concentration considered resistant for a majority of pathogenic yeasts. The drug

combinations were dispensed in a total volume of 100 μl. The 96-well microtiter plates

were custom-manufactured and tested for sterility by TREK Diagnostic System, Part of

Thermo Fisher Scientific, Oakwood Village, OH, USA. They were transferred under dry

ice to NYSDOH Mycology Laboratory and stored at -80℃until used. The schematic of

plate design is provided in supplementary file 1. We also prepared two-fold dilutions of

each drug combination by adding 100 μl sterile RPMI 1640 to the initial well and

transferring 100 μl of 200 μl mix into micro-well of a new 96-well microtiter plate. This

dilution scheme provided 1/2, 1/4, 1/8, 1/16, and 1/32 concentrations of each drug in the

initial well. The inoculation, incubation, and MIC_{100} end-point reading, and *Candida*

krusei ATCC was the quality control strain, were as published earlier from a multi-

laboratory evaluation of antifungal combination testing (9). Briefly, *C. auris* cultures

were streaked on Sabouraud dextrose agar for 24 hours at 35℃, cells suspended in

sterile water to absorbance (A_{530}) 0.08-0.1 absorbance as measured with a Mettler Toledo

UV5 bio spectrophotometer; 20 μl of the cell suspension was added to the 11 ml of

RPMI 1640 broth tube. One-hundred-microliter of the suspension in RPMI 1640 was

added to each well of the 96-well microtiter plate with the drug combinations. The

plates were incubated at 35℃ for 48-hrs, and MIC_{100} reading was obtained with an

illuminated convex mirror plate reader. Twenty microliters of the drug-fungal

suspension from each well, after MIC_{100} determination, was transferred to the well of a

fresh 96-well microtiter plate that contained 180 μl of sterile RPMI-1640 to set-up the

sterility testing. We incubated the sterility plates at 35℃ for 48-hrs, and determined the

minimum fungicidal concentration (MFC) from each well with no visible growth. In

preliminary experiments, a good correlation was observed between MFC values

obtained from microbroth plates and Sabrouad dextrose agar (details not shown).

Results

Among the test group, *C. auris* 16-1 was susceptible to all antifungals tested

while the remaining fourteen isolates showed resistance to FLC (Table 1). The

distribution of resistance in remaining *C. auris* was triazole resistance (2 isolates),

echinocandin resistance (6 isolates), and AMB resistance (4 isolates) (Table 1). Only *C.*

auris 18-1 and 18-13 showed resistance to 5FC. The combination test results were

derived from 12,960 MIC_{100} readings, for fifteen *C. auris* isolates tested against 864 two-

drug antifungal combinations for nine antifungal drugs (Supplementary Table 1).

Flucytosine (5FC) at 1.0 mg/L potentiated the most successful combinations with other

drugs (Table 2). MFG, AFG, and CAS at individual concentrations of 0.25 mg/L

combined well with 5FC (1.0 mg/L) to yield MIC_{100} for 14, 13, and 12 out of 15 *C. auris*

isolates tested. AMB / 5FC (0.25/1.0 mg/L) yielded MIC_{100} for 13 isolates. Triazoles POS,

ISA and VRC also combined well with 5FC (0.25/2.0 mg/L) to yield MIC_{100} for 12, 13,

and 13 isolates, respectively. The summary results for the remaining combinations are

presented in supplementary table 2. Remarkably, MIC_{100} readings were obtained for all

selected combinations. *C. auris* 18-1, resistant to FLC and 5FC initially, also tested

resistant to all antifungal combinations except POS/5FC (0.12/1.0 mg/L) (Table 2).

Discussion

The clinical and public health microbiology community is at high alert due to the alarming rise in the incidence and spread of drug-resistant *C. auris* globally (10). An aggressive therapeutic intervention is indicated for invasive *C. auris* infections associated with high mortality(7, 11). The results of the current study fill a gap in the published literature, as there is no report on the efficacy of antifungal combinations against *C. auris*. Our findings suggested that *C. auris* with various resistance patterns are susceptible to low dose combinations of existing drugs. Several of the effective combinations of antifungals exerted fungicidal activities, which was an additional indicator of the drug combination efficacy. The earlier multi-laboratory evaluation showed that MIC_{100} endpoints provide a reliable method for antifungal combination testing vis-a-vis ΣFIC (the summation of fractional inhibitory concentration) index (9). We further optimized the method by using fixed drug concentrations instead of the checkerboard dilutions, as described by other authors in earlier publications (12, 13). As almost all NY isolates belonged to the South Asian clade, it remains to be determined by other investigators if the resistant *C. auris* strains from South Africa and South America clades respond similarly to the antifungal combinations. Further clinical relevance of the findings reported in the study could be evaluated by pharmacokinetic/pharmacodynamic studies and testing in the animal models (14, 15).

In conclusion, a limited set of two drug-combinations of antifungals were efficacious *in*

vitro against fifteen multidrug-resistant *C. auris* strains.

The simplified two-drug combination susceptibility test format would permit

laboratories to provide clinicians and public health experts with additional data to deal

with multidrug-resistant *C. auris*.

Acknowledgments

We are grateful to the members of the New York State Healthcare Epidemiology

and Infection Control Program, *Candida auris* Investigation Workgroup, and the staff

members from various hospitals and long-term-care facilities for assistance with

surveillance samples.

Funding

This publication was supported in part by Cooperative Agreement number

NU50CK000516, funded by the Centers for Disease Control and Prevention. Its contents

are solely the responsibility of the authors and do not necessarily represent the official

views of the Centers for Disease Control and Prevention or the Department of Health

and Human Services.

Transparency declaration

B. O'Brien performed the experiments and tabulated and analyzed the data, S.

Chaturvedi supervised single drug testing and interpretation, helped interpreting

combination testing and critiqued draft manuscript, and V. Chaturvedi conceived and

designed the study, interpreted data, and wrote the manuscript.

Supplementary Data

Supplementary File 1-Design of 96-well microtiter plates for the antifungal combination

testing

Supplementary Table 1–Master file for all antifungal combination test results

Supplementary Table 2–Summary of less effective antifungal combinations against

Candida auris

168
169

170 Table 1. Broth microdilution determination of antifungal MIC$_{50}$ (mg/L) and interpretative

171 criteria (S, I, R) for the test isolates of *Candida auris*

C. auris	Source	Antifungal drug									
Susceptible		FLC	VOR	ITC	ISA	POS	AFG	CAS	MFG	AMB	5FC
16-1	Ear	8	0.25	0.5	0.25	0.25	0.12	0.12	0.12	0.5	0.5
		S	S	S	S	S	S	S	S	S	S
17-12	Urine	32	0.5	0.06	0.015	0.03	8	8	8	1	0.032
		S	S	S	S	S	R	R	R	S	S
18-14	NA	>256	1	0.25	0.5	0.03	0.06	0.015	0.06	1	0.047
		R	S	S	S	S	S	S	S	S	S
18-15	Blood	>256	1	0.25	0.5	0.03	0.12	0.03	0.06	1.5	0.064
		R	S	S	S	S	S	S	S	S	S
17-1	Blood	>256	2	0.5	0.5	0.25	4	4	4	1	0.094
		R	S	S	S	S	R	R	R	S	S
17-13	Urine	>256	1	0.25	0.5	0.03	0.06	0.03	0.06	3	0.032
		R	S	S	S	S	S	S	S	R	S
17-14	Groin swab	>256	2	0.5	0.5	0.12	4	2	4	0.75	0.032
		R	S	S	S	S	R	R	R	S	S
17-15	Skin rash swab	>256	2	0.5	0.5	0.25	4	2	4	0.75	0.032
		R	S	S	S	S	R	R	R	S	S
17-16	Urine	>256	2	0.5	0.5	0.12	4	2	4	1	0.023
		R	S	S	S	S	R	R	R	S	S
17-17	Nares swab	>256	4	0.5	0.5	0.06	0.25	0.03	0.06	3	0.064
		R	R	S	S	S	S	S	S	R	S
17-18	Nares swab	>256	2	0.5	1	0.12	0.25	0.06	0.06	3	0.064
		R	S	S	S	S	S	S	S	R	S
18-1	Wound swab	128	0.5	0.12	0.12	0.03	0.25	0.03	0.12	1	>32
		R	S	S	S	S	S	S	S	S	R
18-2	Ascites fluid	>256	2	1	1	0.25	8	2	4	1.5	0.064
		R	S	R	R?	S	R	R	R	S	S
18-5	Urine	>256	2	0.25	0.5	0.06	0.12	0.03	0.12	2	0.094
		R	S	S	S	S	S	S	S	R	S
18-13	Rectal swab	256	0.5	0.12	0.12	0.03	0.12	0.03	0.12	1	>32
		R	S	S	S	S	S	S	S	S	R

173
174 NA not available

Table 2. Most promising antifungal combinations for *Candida auris*

Combi-nation	End-reading	*Candida auris*														
		16-1	17-1	17-12	17-13	17-14	17-15	17-16	17-17	17-18	18-1	18-2	18-5	18-13	18-14	18-15
MFG + 5FC	MIC$_{100}$	0.12*/1.0	0.12/1.0	0.06/0.5	0.12/1.0	0.12/1.0	0.12/1.0	0.12/1.0	0.12/1.0	0.25/2.0	NA	0.12/1.0	0.12/1.0	0.12/1.0	0.12/1.0	0.12/1.0
AMB + 5FC	MIC$_{100}$	0.25/1.0	0.25/1.0	0.25/1.0	0.25/1.0	0.25/1.0	0.25/1.0	0.25/1.0	0.25/1.0	0.25/1.0	NA	0.25/1.0	0.25/1.0	NA	0.25/1.0	0.25/1.0
AFG + 5FC	MIC$_{100}$	0.12 /1.0	0.12/1.0	0.12/1.0	0.12/1.0	0.12/1.0	0.25/2.0	0.12/1.0	0.12/1.0	0.25/2.0	NA	0.12/1.0	0.12/1.0	NA	0.12/1.0	0.12/1.0
CAS + 5FC	MIC$_{100}$	0.12/1.0	0.12/1.0	0.12/1.0	0.12/1.0	0.12/1.0	0.12/1.0	0.25/2.0	0.12/1.0	NA	NA	0.12/1.0	0.25/2.0	NA	0.12/1.0	0.12/1.0
POS + 5FC	MIC$_{100}$	0.12/1.0	0.12/1.0	0.12/1.0	NA	NA	0.12/1.0	0.12/1.0	0.12/1.0	0.12/1.0	0.12/1.0	0.12/1.0	0.12/1.0	NA	0.12/1.0	0.25/2.0
ISA + 5FC	MIC$_{100}$	0.12/1.0	0.12/1.0	0.12/1.0	0.12/1.0	0.12/1.0	0.12/1.0	0.12/1.0	0.12/1.0	0.12/1.0	NA	0.12/1.0	0.12/1.0	NA	0.12/1.0	0.25/2.0
ITC + 5FC	MIC$_{100}$	0.03/0.5	0.06/1.0	0.06/1.0	0.06/1.0	NA	0.06/1.0	0.06/1.0	0.06/1.0	0.06/1.0	NA	0.06/1.0	0.06/1.0	NA	0.06/1.0	0.12/2.0
VOR + 5FC	MIC$_{100}$	0.25/1.0	0.25/1.0	0.125/0.5	0.25/1.0	0.25/1.0	0.25/1.0	0.25/1.0	0.25/1.0	0.25/1.0	NA	0.25/1.0	0.25/1.0	NA	0.25/1.0	0.5/2.0

* mg/L

NA Not available

References

1.   Lockhart SR, Etienne KA, Vallabhaneni S, Farooqi J, Chowdhary A, Govender NP,
Colombo AL, Calvo B, Cuomo CA, Desjardins CA. 2017. Simultaneous Emergence of
Multidrug-Resistant *Candida auris* on 3 Continents Confirmed by Whole-Genome
Sequencing and Epidemiological Analyses. Clinical Infectious Diseases 64:134-140.

2.   Satoh K, Makimura K, Hasumi Y, Nishiyama Y, Uchida K, Yamaguchi H. 2009. *Candida*
auris sp. nov., a novel ascomycetous yeast isolated from the external ear canal of an
inpatient in a Japanese hospital. Microbiology and immunology 53:41-44.

3.   Vallabhaneni S, Kallen A, Tsay S, Chow N, Welsh R, Kerins J, Kemble SK, Pacilli M, Black
SR, Landon E, Ridgway J, Palmore TN, Zelzany A, Adams EH, Quinn M, Chaturvedi S,
Greenko J, Fernandez R, Southwick K, Furuya EY, Calfee DP, Hamula C, Patel G, Barrett P,
Lafaro P, Berkow EL, Moulton-Meissner H, Noble-Wang J, Fagan RP, Jackson BR,
Lockhart SR, Litvintseva AP, Chiller TM. 2017. Investigation of the First Seven Reported
Cases of *Candida auris*, a Globally Emerging Invasive, Multidrug-Resistant Fungus-United
States, May 2013-August 2016. Am J Transplant 17:296-299.

4.   Govender NP, Magobo RE, Mpembe R, Mhlanga M, Matlapeng P, Corcoran C, Govind C,
Lowman W, Senekal M, Thomas J. 2018. *Candida auris* in South Africa, 2012-2016.
Emerg Infect Dis 24:2036-2040.

5.   Chowdhary A, Kumar VA, Sharma C, Prakash A, Agarwal K, Babu R, Dinesh K, Karim S,
Singh S, Hagen F. 2014. Multidrug-resistant endemic clonal strain of *Candida auris* in
India. European journal of clinical microbiology & infectious diseases 33:919-926.

6.   Calvo B, Melo ASA, Perozo-Mena A, Hernandez M, Francisco EC, Hagen F, Meis JF,
Colombo AL. 2016. First report of *Candida auris* in America: Clinical and microbiological
aspects of 18 episodes of candidemia. Journal of Infection 73:369-374.

7.   Adams E, Quinn M, Tsay S, Poirot E, Chaturvedi S, Southwick K, Greenko J, Fernandez R,
Kallen A, Vallabhaneni S, Haley V, Hutton B, Blog D, Lutterloh E, Zucker H, Workgroup.
Cal. 2018. *Candida auris* in Healthcare Facilities, New York, USA, 2013-2017. Emerging
Infectious Diseases 24:1816-1824.

8.   Leach L, Zhu Y, Chaturvedi S. 2018. Development and Validation of a Real-Time PCR
Assay for Rapid Detection of *Candida auris* from Surveillance Samples. J Clin Microbiol
56.

9.   Ren P, Luo M, Lin S, Ghannoum MA, Isham N, Diekema DJ, Pfaller MA, Messer S,
Lockhart SR, Iqbal N, Chaturvedi V. 2015. Multilaboratory testing of antifungal drug
combinations against *Candida* species and *Aspergillus fumigatus*: utility of 100 percent
inhibition as the endpoint. Antimicrob Agents Chemother 59:1759-66.

10. Tsay S, Kallen A, Jackson BR, Chiller TM, Vallabhaneni S. 2018. Approach to the
Investigation and Management of Patients With *Candida auris*, an Emerging Multidrug-
Resistant Yeast. Clin Infect Dis 66:306-311.

11. Ruiz-Gaitan A, Moret AM, Tasias-Pitarch M, Aleixandre-Lopez AI, Martinez-Morel H,
Calabuig E, Salavert-Lleti M, Ramirez P, Lopez-Hontangas JL, Hagen F, Meis JF, Mollar-
Maseres J, Peman J. 2018. An outbreak due to *Candida auris* with prolonged
colonisation and candidaemia in a tertiary care European hospital. Mycoses 61:498-505.

12. Pfaller MA, Barry AL, Fuchs PC, Gerlach EH, Hardy DJ, McLaughlin JC. 1993. Comparison
of fixed concentration and fixed ratio options for dilution susceptibility testing of gram-

negative bacilli to ampicillin and ampicillin/sulbactam. Eur J Clin Microbiol Infect Dis
12:356-62.
13. Bradford PA, Huband MD, Stone GG. 2018. A Systematic Approach to the Selection of
the Appropriate Avibactam Concentration for Use with Ceftazidime in Broth
Microdilution Susceptibility Testing. Antimicrobial Agents and Chemotherapy
62:e00223-18.
14. Lepak AJ, Andes DR. 2015. Antifungal Pharmacokinetics and Pharmacodynamics. Cold
Spring Harbor Perspectives in Medicine 5.
15. Hager CL, Larkin EL, Long L, Zohra Abidi F, Shaw KJ, Ghannoum MA. 2018. *In Vitro* and *In*
Vivo Evaluation of the Antifungal Activity of APX001A/APX001 against *Candida auris*.
Antimicrobial Agents and Chemotherapy 62:e02319-17.
236
237