Title: Ambient temperature effects on stress-induced hyperthermia in Svalbard ptarmigan

Andreas Nord[1, 2, 3,*] and Lars P. Folkow[2]

[1] Department of Biology, Section for Evolutionary Ecology, Lund University, SE-223 62

Lund, Sweden

[2] Department of Arctic and Marine Biology, University of Tromsø – the Arctic University of

Norway, NO-9037 Tromsø, Norway

[3] Scottish Centre for Ecology and the Natural Environment, University of Glasgow,

Rowardennan G63 0AW, United Kingdom

* Corresponding author: andreas.nord@biol.lu.se

Running head: Stress-induced hyperthermia in the world's northernmost land bird

Keywords: Arctic, body temperature, handling, stress, stress-induced hyperthermia,

thermoregulation

Summary statement: Stress-induced hyperthermia (SIH) is displayed also by the high-arctic

Svalbard ptarmigan, suggesting that this response in birds is regulated as in other endotherms.

However, environmental temperature, body size and insulation must be taken into

consideration when evaluating thermal responses to acute stressors in birds.

**ABSTRACT**

Stress-induced hyperthermia (SIH) is commonly observed during handling in homeotherms.

However, in birds, handling in cold environments typically elicits hypothermia. It is unclear

whether this indicates that SIH is differently regulated in this taxon or if it is due to size,

because body temperatures changes during handling in low temperature have only been

measured in small birds ≤ 0.03 kg (that are more likely to suffer high heat loss when handled).

We have, therefore, studied thermal responses to handling stress in the intermediate-sized

(0.5-1.0 kg) Svalbard ptarmigan (*Lagopus muta hyperborea*) in 0°C and -20°C, in winter and

spring. Handling caused elevated core body temperature, and peripheral vasoconstriction that

reduced back skin temperature. Core temperature increased less and back skin temperature

decreased more in -20°C than in 0°C, probably because of higher heat loss rate at the lower

temperature. Responses were qualitatively consistent between seasons, despite higher body

condition/insulation in winter and dramatic seasonal changes in photoperiod, possibly

affecting stress responsiveness. Our study supports the notion that SIH is a general

thermoregulatory reaction to acute stressors in endotherms, but also suggests that body size

and thermal environment should be taken into account when evaluating this response in birds.

## INTRODUCTION

Stress-induced hyperthermia (SIH), is a ubiquitous feature of the body's response to acute stressors, such as restraint or an altered social context, in mammals and birds (e.g. Briese and Cabanac, 1991; Cabanac and Briese, 1992; Carere and Van Oers, 2004; Gray et al., 2008; Korhonen et al., 2000; Meyer et al., 2008). It is believed that the increase in core body temperature ($T_c$) during SIH represents a sympathetically mediated elevation of the hypothalamic set point, i.e. an 'active hyperthermia' resembling pathogen-induced fevers (Briese and Cabanac, 1991; Kluger et al., 1987; Oka et al., 2001; Vinkers et al., 2009). Hence, SIH is sometimes referred to as 'stress fever' or 'psychogenic fever' (e.g. IUPS Thermal Commission, 2003), although several studies suggest it employs different neural pathways (Gray et al., 2008; Vinkers et al., 2009). Thermal responses leading to elevated $T_c$ during SIH include cutaneous vasoconstriction and shivering thermogenesis (e.g. Briese and Cabanac, 1991; Herborn et al. 2015; Jerem et al., 2015; Kluger et al., 1987; Oka et al., 2001). The diversion of peripheral blood flow to the core, together with stress-induced tachycardia and increased ventilation rate (Cabanac and Aizawa, 2000; Cabanac and Guillemette, 2001; Greenacre and Lusby, 2004; Mans et al., 2012), probably prepares the animal for escape or interaction (i.e., a "fight or flight" response), at the same time as centralization of the blood pool and increased blood clotting function could minimize blood loss in the event of injury (Cannon 1915).

Because SIH probably reflects set point change it is predicted that, for a given stressor, $T_c$ changes should be independent of the ambient temperature ($T_a$) under which stress is perceived. This prediction is supported by laboratory studies of rodents (e.g. Briese and Cabanac, 1991; Kluger et al., 1987; Long et al., 1990). However, studies of birds have revealed remarkable variation in the thermal responses to restraint or handling at different $T_a$'s. Specifically, handling in thermoneutral conditions sometimes invokes SIH (Cabanac and Aizawa, 2000; Cabanac and Guillemette, 2001; Carere and Van Oers, 2004; Herborn et al., 2015), sometimes hypothermia (Maggini et al., 2018; Møller, 2010), and sometimes seem to leave $T_c$ unaltered (Lewden et al., 2017), whereas handling below thermoneutrality consistently seems to elicit hypothermia without any co-occurring signs of shock such as lack of muscle tonus (Andreasson et al., 2019; Lewden et al., 2017; Udvardy, 1955;).

Cooling rather than warming of the core need not mean that birds have differentially regulated stress responses to handling. Rather, because the cooling phenomenon seems confined to

small (< 0.03 kg) species with inherently higher thermal conductance, it is likely that stress-
associated *hypothermia* results from changes to the rate and avenues of heat transfer caused
by the handling *per se*. For example, a small bird enclosed by a hand is subject to a substantial
increase in the proportion of surface area amenable for conductive heat transfer, suffers
reduced insulation when the plumage is compressed, and probably conducts heat over a
considerably steeper thermal gradient (as a consequence of the handling-induced reduction in
plumage depth). This explanation remains speculative, though, because body temperature
responses to handling in larger, better insulated, birds have never been recorded in low $T_a$
(e.g. Cabanac and Aizawa, 2000; Cabanac and Guillemette, 2001; Herborn et al., 2015). In
addition, there are no continuous data from birds on the relationship between temperature
changes in the periphery and the core during stress, which further complicates our
understanding of the regulatory processes involved in situations where 'atypical' thermal
responses are observed.  Thus, additional data from large and well-insulated birds measured in
low $T_a$ are required to better understand why this taxon displays such variable stress-induced
body temperature responses.
90
We measured the thermal responses to handling in the Svalbard ptarmigan (*Lagopus muta*
*hyperborea* Sundevall) (Fig. 1). This 0.5 to 1.0 kg bird is endemic to the high-arctic Svalbard
archipelago (76-81°N), where it is exposed to $T_a$ ranging ca. -40 to +20°C and a photoperiod
varying 24 h over the course of the year. Its thermal conductance varies accordingly, being the
lowest in winter when it is dark and cold and birds are in prime body condition, and the
highest in summer when birds are at their leanest and have moulted into a less insulating
plumage (Mortensen and Blix 1986; Nord & Folkow 2018). These properties make the
Svalbard ptarmigan a good model for studying how body temperatures of a relatively large
bird respond to handling, and if these responses differ in different thermal environments and
in line with variation in insulation. Accordingly, we measured temperature changes both
peripherally (back and head skin; assumed to reflect changes in cutaneous circulation) and in
the body core (to assess whether birds responded with hypo- or hyperthermia) in Svalbard
ptarmigan subjected to handling stress at thermoneutrality (0°C) and far below
thermoneutrality (-20°C) (Mortensen and Blix, 1986), both in winter when insulation peaked,
and in spring when insulation was declining. Measurement of both deep and peripheral
temperatures, and at times of year when thermal conductance is notably different (Mortensen
and Blix, 1986; Nord and Folkow, 2018), was expected to provide information on the
thermoregulatory processes involved in acute stress in birds. If the normal thermal response to

stress in birds is an increase in $T_c$ (i.e., SIH), and the reduced $T_c$ during handling in smaller

species of birds is, in fact, a consequence of increased heat loss rate (e.g. from plumage

disturbances or increased conductive cooling; cf. Andreasson et al., 2019; Lewden et al.,

2017) rather than representing a differentially regulated response, we predicted that the larger

and well-insulated Svalbard ptarmigan would show an elevation of $T_c$ that was qualitatively

similar at thermoneutral (0°C) vs. very low (-20°C) $T_a$'s, and that this would be preceded by

reduced peripheral temperature (reflecting cutaneous vasoconstriction). Because the Svalbard

ptarmigan has low thermal conductance compared to other species, even at its leanest in

summer (Mortensen & Blix 1986), we also predict that any seasonal effect on the thermal

responses to handling should be minor.

**MATERIALS AND METHODS**

**Birds and housing**

Twelve male Svalbard ptarmigan were used. Six were captured as chicks near Longyearbyen,

Svalbard (78°13' N, 15°38' E) 16 to 17 months before the start of the experiment, and the

other six were captive bred at the University of Tromsø – the Arctic University of Norway. At

the time of the study, all females in the facility were part of the breeding population that is

exempt from experimental work year round. However, in Svalbard ptarmigan, both sexes

show the same seasonal responses and are of similar size (Mortensen et al., 1983), suggesting

that stress responsiveness should be qualitatively similar in males and females. All birds were

maintained indoors in thermoneutrality under simulated Longyearbyen photoperiod.   The

morphological and physiological changes associated with winter acclimation do not differ

between captive and wild-caught Svalbard ptarmigan as long as the birds are maintained in

natural photoperiod (Lindgård and Stokkan, 1989; Stokkan et al., 1986). Housing conditions

followed Nord and Folkow (2018).

**Measurement of body temperature and handling protocol**

All birds were physically mature (i.e., older than 1 year). Studies were undertaken under polar

night conditions in early winter (DD) (07 December 2015 to 17 January 2016), when body

condition peaked (body mass: 1064.2 ± 33.0 g (s.d.); dissectible fat: 261.5 ± 16.7 g) and in

spring (03 May to 03 June 2016) in continuous light (LL) when body condition was declining

(body mass: 828.1 ± 18.1 g; dissectible fat: 142.0 ± 9.2 g). Data were collected at ambient

temperatures ($T_a$) 0°C ($T_a$ = -0.1 ± 0.6°C (s.d.), which is within the thermoneutral zone, and at

-20°C (-19.9 ± 0.5°C), which is far below thermoneutrality in Svalbard ptarmigan

independent of season (Mortensen and Blix, 1986; Nord and Folkow, 2018). We used 11 birds

in winter, of which 10 were measured in 0°C and five also in -20°C. One additional bird was

measured only in -20°C. Seven of the 11 winter birds, and one additional bird, were measured

in spring (i.e., 0°C: $n$ = 8 birds; -20°C: $n$ = 5 birds, of which 4 had been measured in -20°C

also in winter). In winter, 7 birds were first measured in 0°C, and 4 birds were first measured

in -20°C. In spring, all birds were first measured in 0°C and then in -20°C. Repeated

measurement within seasons were spaced 27 ± 3 (s.d.) days (range 24 to 31 days). Birds were

monitored as part of a study that is not reported here for 21 days after each measurement

session. For this reason, sampling in both $T_a$'s in winter could not be achieved within the

period of constant darkness. In spring, measurement in both $T_a$'s was not always possible,

since some individuals were allocated to the on-site breeding program by the end of May.

Starting at 10:06 am ± 30 min (s.d.) (GMT +1), birds were weighed and fitted with 36-gauge

type T (copper-constantan) thermocouples (Omega Engineering, Stamford, CT, USA) for

measurement of $T_c$ (70 mm into the colon), back skin temperature ($T_{back}$) between the wings,

and head skin temperature ($T_{head}$) on the scalp following Nord & Folkow (2018), within 10-15

min of collection from the cage. Skin thermocouples were attached using a small amount of

cyanoacrylate glue (Loctite® Power Easy Gel, Henkel, Düsseldorf, Germany) and taking care

not to cover the thermocouple junction. This glue does not cause any skin damage or irritation

in Svalbard ptarmigan (AN, LPF, pers. obs.). Thermocouples were calibrated at 0°C (Ice point

drywell model 5115) and 40°C (High precision bath model 6025, both Fluke Calibration,

American Fork, UT, USA) prior to use.

Once thermocouples were attached and before the handling experiment, we placed the bird in

a 33.6 l transparent acrylic glass metabolic chamber (that was ventilated with ambient air at

5.1 ± 0.3 l min$^{-1}$) inside a climatic chamber (model 24/50 DU, Weiss Technik, Giessen,

Germany) for measurement of resting metabolic rate and body temperatures during 1 h 44

min ± 13 min (s.d.), as part of a different study. Both metabolic rate and body temperatures

stabilized at lower levels within 30 min after instrumentation. After completion of metabolic

measurements, we opened the metabolic chamber, side-pinned (terminology *sensu* Herborn et

al., 2015) the bird, and administered an immune challenge (100 µl 1 mg kg$^{-1}$ intramuscular

LPS), also as part of the other study that is not reported here. Instead, we here report the

changes in body temperatures that were recorded during and after the handling stress that was induced thereby. This stressor lasted, on average, $3.2 \pm 0.6$ min (s.d.) (range: 2.5 to 5.3 min) and did not differ in length between seasons or $T_a$'s (season $\times$ $T_a$, season, $T_a$: all $P \geq 0.3$). Thus, the order of events during measurements were: capture and instrumentation (10-15 min); equilibration and baseline data collection at relevant $T_a$ (104 min); handling stress (3 min) and post-stressor data collection (20 min). Because the metabolic chamber was fully open when birds were handled, metabolic rate could not be measured during, and in the 20 min period following, stress exposure.

During winter measurements, the climatic chamber was always dark (except for illumination with dim red light, << 1 lx, to allow video inspection) to simulate polar night. In spring, the chamber was always fully illuminated by full spectrum white light bulbs. All but three birds had been subjected to a similar handling- and measurement protocol on 2 to 8 instances in a previous study (mean $\pm$ s.d.: $6 \pm 2$) (for details on this protocol, see Nord and Folkow, 2018). Data were recorded at 10 samples/s, and were digitized from raw signals using a ML796 PowerLab/16SP A-D converter (ADInstruments, Sydney, Australia).

Ethical approval for experimental procedures and reuse of birds was issued by the Norwegian Food Safety Authority (permit no. 6640). Live capture and import of Svalbard ptarmigan chicks was under permissions issued by the Governor of Svalbard (permit no. 2014/00290-2 a.522-01), the Norwegian Environment Agency (permit 2018/7288) and the Norwegian Food Safety Authority (permit no. 2014/150134).

**Data analyses**

We only used data from periods where the birds were at full rest (standing or walking but not moving vigorously, or perched with ptiloerection) immediately before handling. This criterion was met in 24 of 31 instances (winter, 0°C: 8 of 10; winter, -20°C: 5 of 6; spring, 0°C: 7 of 8; spring, -20°C: 3 of 5). We also dismissed data from $T_{head}$ thermocouples that fell off or broke (winter, -20°C: 1; spring, 0°C: 2; spring, -20°C: 1). $T_c$ and $T_{back}$ data were complete.

Pre-handling data were collected during 2 min immediately before handling (i.e., $\geq$ 1.5 h after the start of body temperature measurement). Handling and post-handling data were then collected during 20 min after the end of this stressor, i.e., a period sufficient to encompass the body temperature response to acute stress in other gallinaceous birds (Herborn et al., 2015),

but not long enough to include any thermal or metabolic effects of LPS (Marais et al., 2011).

We binned data in 5 s (i.e., 50 samples) averages, partitioned in pre-handling, handling, and

post-handling, periods. We then calculated: 'response amplitude' as the maximum or

minimum body temperature attained during handling; 'duration of response' as the first time

point where a 30 s running mean for each body temperature had returned to, or intersected,

pre-handling values once the stressor had been removed; and 'response magnitude' as mean

body temperature change (relative to pre-handling values) over the duration of the response. If

a given body temperature had not returned to pre-handling levels by the end of the focal

period, we set 'duration of response' to 20 min (i.e. the length of the post-handling

observation period), and calculated 'response magnitude' based on this period.

Statistics were done using R 3.4.3 for Windows (R Core Team, 2018). We analysed all

response variables in tissue-specific mixed effect models fitted with maximum likelihood

(lmer function in the lme4 package) (Bates et al., 2015), with season and $T_a$ and their

interaction as factors, and 'bird id' as a random intercept to account for repeated

measurements. We did not include body mass or body condition in any models, because both

parameters varied strongly between, but only very little within, seasons (above). These

metrics, therefore, conveyed largely the same statistical information as 'season', so testing for

any of the former in presence of the latter was not warranted. Final models were derived by

sequentially excluding terms with the highest $P$-values based on likelihood ratio tests for the

original and alternative models, starting with the interactions, until only significant ($P \leq 0.05$)

variables remained. We then refitted the final model using restricted maximum likelihood

(Zuur et al., 2009) and calculated predicted means ± s.e.m. using the emmeans package

(Lenth, 2019).

**RESULTS**

Parameter estimates and test statistics are reported in Table S1.

All $T_c$ responses to handling were positive and attenuated in -20°C compared to in 0°C.

Accordingly, birds reached 0.19 ± 0.06°C greater maximum $T_c$ in 0°C ($\Delta T_c = 0.52 \pm 0.04$°C)

than in -20°C ($\Delta T_c = 0.32 \pm 0.05$°C) ($P = 0.009$, Fig. 2). Response duration, i.e. the time

taken for $T_c$ to return to pre-handling levels for at least 30 s once the stressor had been

removed, was longer in 0°C (13.4 ± 1.1 min) than in -20°C (5.6 ± 1.4 min) by 7.8 ± 1.8 min

(+138%) ($P < 0.001$, Fig. 2). The response magnitude, i.e. the mean deviation from pre-

handling $T_c$ during the response, was larger when birds were measured in thermoneutrality

($0.29 \pm 0.03°C$) than when they were measured in -20°C ($0.17 \pm 0.04°C$), by $0.12 \pm 0.04°C$ ($P$

$= 0.016$). The modifying effect of $T_a$ was always uniform across seasons (season $\times$ $T_a$: all $P \geq$

0.22), and there was no difference in mean responses between winter and spring ($P \geq 0.43$ in

all cases; Table S1).

$T_{back}$ decreased on average during handling. However, one bird showed increased $T_{back}$ in three

out of four measurement sessions, and two additional birds showed positive response when

measured in thermoneutrality in spring (Fig 1). In all but one of these cases, positive

responses were preceded by a slight $T_{back}$ decrease (Fig. 2B). The strongest $\Delta T_{back}$ response

was $1.38 \pm 0.44°C$ below pre-handling levels. There was a tendency for a stronger maximum

response at -20°C ($-2.03 \pm 0.77°C$) than at 0°C ($-1.01 \pm 0.53°C$) ($P = 0.092$; Fig. 2). Birds had

recovered this decrease within $10.5 \pm 1.3$ min, but tended to take longer to do so at -20°C

($12.0 \pm 2.2$ min) than at 0°C ($7.2 \pm 1.8$ min) ($P = 0.098$; Fig. 2). Average response magnitude

was uniform, at $0.45 \pm 0.16°C$ below pre-handling $T_{back}$, across $T_a$'s. Neither season nor the

season-by-$T_a$ interaction affected the $T_{back}$ response to handling (all $P \geq 0.12$; Table S1).

The maximum $\Delta T_{head}$ response was positive on average ($+0.67 \pm 0.35°C$) (Fig. 2). This

increase subsided within $7.5 \pm 0.4$ min after removal of the stressor (Fig. 2). The response

magnitude index showed that $\Delta T_{head}$ was elevated, on average by $0.45 \pm 0.16°C$, during this

time. Season, $T_a$, and their interaction, did not affect any $T_{head}$ metric (all $P \geq 0.1$; Table S1).

**DISCUSSION**

Handling of Svalbard ptarmigan was associated with increased $T_c$ and $T_{head}$ and decreased

$T_{back}$ that lasted, on average, 6.6 to 9.7 min after the stressor had been removed (Fig. 2). The

maximum $T_c$ response was small but distinct and within the range of those recorded over

similar time periods in other bird species of comparable body sizes but at higher $T_a$

(Bittencourt et al., 2015; Cabanac and Aizawa, 2000; Cabanac and Guillemette, 2001; Gray et

al., 2008). Both the maximum increase in $T_c$ and the $T_c$ response magnitude were more

pronounced in 0°C than in -20°C (Fig. 2; Table S1). This probably reflects that low $T_a$ blunted

the rate of increase in $T_c$. Thus, since the stressor was of fixed duration and probably too short

to allow the birds to reach a new set-point, $T_c$ did not have time to increase as much in -20°C

as it did in 0°C. $T_{back}$, moreover, tended to decrease about twice as much in -20°C than in 0°C

(Fig. 2). This could suggest that a stronger peripheral vasoconstrictor response was employed

to elevate $T_c$ when $T_a$ was lower, but more likely reflects that the drop in skin temperature

when cutaneous blood flow was diminished in the cold, was larger in the lower $T_a$ due to a

more rapid heat loss rate. Thus, the difference in response strength in the two $T_a$'s most likely

reflected that higher heat loss rate in the cold slowed and blunted the $T_c$ response, and

increased $T_{back}$ change, relative to changes in the milder $T_a$.

In some cases (5 of 24), birds responded to handling by vasodilation of back skin (reflected by

increased $T_{back}$; Fig. 2), but this was preceded by an initial $T_{back}$ decrease in all but one of

these instances. This suggests that also those few birds that displayed positive $T_{back}$ change at

least initially met the stressful stimulus with the expected cutaneous vasoconstrictor response,

but then, for some reason, rapidly switched to vasodilation. Our data do not allow us to

determine the proximate explanation for this result, which could be related to skin

thermocouple placement in relation to vascular structures, or that the thermal state of these

particular birds caused them to revert to heat dissipation because a new set-point temperature

was reached more rapidly. In this latter context, we noted that pre-handling $T_{back}$ was higher

(by 1.05°C) in these birds.

$T_{head}$ on the scalp increased in response to handling, but only after a short-lasting drop that

implied rapid vasoconstriction (Fig. 2). Thus, cutaneous $T_{head}$ initially followed predictions for

peripheral temperature change during acute stress, but rapidly reverted to the opposite

response. The temperature increase was probably a combined effect of increased delivery of

internal heat to the head in conjunction with the rise in $T_c$, and increased blood flow to the

brain in response to the stressor (Hasler et al., 2007; Wang et al., 2005) that together

outweighed any reduced supply of heat due to constriction of head skin vasculature. Increased

blood delivery to the brain likely prompted higher attentiveness and increased cognitive

ability that would aid the animal in decision-making in a threatening situation.

Physiologically, increased temperature in the poorly insulated head (cf. Nord and Folkow,

2018), lasting several minutes after removal of the stressor, probably contributed to reversal

of the $T_c$ response due to its effects on central thermoreceptors, even if the ultimate

explanation for $T_{head}$ change in our study was not related to heat dissipation as such. It,

therefore, appears that this response, together with birds' general heat loss rate, was sufficient

to regain thermal balance in the relatively low $T_a$'s applied in this study. This could also

explain why we mostly did not record any corresponding compensatory peripheral
temperature increase in back skin as $T_c$ returned to normothermia (Fig. 2B). This is in contrast
to previous studies of the body temperature responses to handling conducted in considerably
milder $T_a$, e.g. in the cold-tolerant common eider (*Somateria mollissima*) (Cabanac and
Guillemette, 2001).
318
**CONCLUSIONS**
The Svalbard ptarmigan, like other medium-sized to large birds, displayed a rapid cutaneous
vasoconstrictor response and an increase in $T_c$ during handling, indicative of SIH. These
responses did not vary qualitatively between seasons despite higher overall heat loss rate in
spring (Nord and Folkow, 2018) that should allow more rapid return to pre-handling $T_c$.
Moreover, even when insulation was close to its annual minimum in late spring, handling in
$T_a$ far below thermoneutrality did not cause hypothermic $T_c$. This is in stark contrast to data
from the considerably smaller cold-tolerant black-capped chickadee (*Poecile atricapillus*)
(Lewden et al., 2017) and great tit (*Parus major*) (Andreasson et al., 2019), in which handling
in the cold results in $T_c$ dropping several degrees C below normothermic values. Thus, our
data from cold exposure in the substantially larger Svalbard ptarmigan support the argument
that stress-related *hypothermia* in small birds handled below thermoneutrality probably does
not reflect a regulated process but more likely is due to increased heat loss rate following
mechanical distortion of insulation during handling, or reduced heat production if handling
elicits tonic immobility (cf. Hohtola, 1981). Yet, thermal responses to handling in our birds
were still modified by $T_a$, particularly in the core but to some extent also peripherally on the
back. Hence, our study suggests that when interpreting the body temperature responses to
handling, one must take into account both body size of the model and the thermal
environment in which measurements are undertaken.
338
339

340  **ACKNOWLEDGEMENTS**
341  Hans Lian, Hans Arne Solvang and Renate Thorvaldsen excellently cared for birds outside
342  experimental periods. Comments from Katherine Herborn and Paul Jerem improved a
343  previous version of the manuscript.

344
345
**COMPETING INTERESTS**

We declare no conflicting or financial interests.

**FUNDING**

AN was supported by the Swedish Research Council (grant no. 637-2013-7442), the Carl

Trygger Foundation for Scientific Research (grant no. 14:347), the Birgit and Hellmuth Hertz

Foundation / The Royal Physiographic Society of Lund (grant no. 2017-39034) and the

Längman Cultural Foundation.

**AUTHOR CONTRIBUTION**

AN conceived the idea, AN and LPF developed methodology and AN collected the data. AN

analysed the data and wrote the first draft, which was critically edited by LPF. Both authors

approved the final version of the manuscript and agreed to be accountable for all contents. AN

administered the project and acquired the funding.

**DATA AVAILABILITY**

Data are deposited in figshare (doi: 10.6084/m9.figshare.8080934).

**REFERENCES**

**Andreasson, F. A., Nord, A. and Nilsson, J.-Å.** (2019). Body temperature responses of great

tits *Parus major* to handling in the cold. *Ibis*, in press.

**Bates, D., Maechler, M., Bolker, B. and Walker, S.** (2015). Fitting linear mixed-effects

models using lme4. *J. Stat. Soft.* **67**, 1-48.

**Bittencourt, M. d. A., Melleu, F. F. and Marino-Neto, J.** (2015). Stress-induced core

temperature changes in pigeons (*Columba livia*). *Physiol. Behav.* **139**, 449-458.

**Briese, E. and Cabanac, M.** (1991). Stress hyperthermia: Physiological arguments that it is a

fever. *Physiol. Behav.* **49**, 1153-1157.

**Cabanac, A. and Briese, E.** (1992). Handling elevates the colonic temperature of mice.

*Physiol. Behav.* **51**, 95-98.

**Cabanac, A. J. and Guillemette, M.** (2001). Temperature and heart rate as stress indicators

of handled common eider. *Physiol. Behav.* **74**, 475-479.

**Cabanac, M. and Aizawa, S.** (2000). Fever and tachycardia in a bird (*Gallus domesticus*)
after simple handling. *Physiol. Behav.* **69**, 541-545.

**Cannon, W. B.** (1915). *Bodily changes in pain, hunger, fear and rage*. New York: D.
Appleton and Company.

**Carere, C. and Van Oers, K.** (2004). Shy and bold great tits (*Parus major*): body
temperature and breath rate in response to handling stress. *Physiol. Behav.* **82**, 905-912.

**Gray, D. A., Maloney, S. K. and Kamerman, P. R.** (2008). Restraint increases afebrile body
temperature but attenuates fever in Pekin ducks (*Anas platyrhynchos*). *Am. J. Physiol.* **294**,
R1666-R1671.

**Greenacre, C. B. and Lusby, A. L.** (2004). Physiologic responses of Amazon parrots
(*Amazona* species) to manual restraint. *J. Avian Med. Sur.* **18**, 19-22.

**Hasler, G., Fromm, S., Alvarez, R. P., Luckenbaugh, D. A., Drevets, W. C. and Grillon,**
**C.** (2007). Cerebral blood flow in immediate and sustained anxiety. *J. Neurosci.* **27**, 6313-
6319.

**Herborn, K. A., Graves, J. L., Jerem, P., Evans, N. P., Nager, R., McCafferty, D. J. and**
**McKeegan, D. E. F.** (2015). Skin temperature reveals the intensity of acute stress. *Physiol.*
*Behav.* **152**, 225-230.

**Hohtola, E.** (1981). Tonic immobility and shivering in birds: Evolutionary implications.
*Physiol. Behav.* **27,** 475-480.

**IUPS Thermal Commission.** (2003). Glossary of terms for thermal physiology. *J. Therm.*
*Biol.* **28**, 75-106.

**Jerem, P., Herborn, K. A., McCafferty, D., McKeegan, D. and Nager, R.** (2015). Thermal
imaging to study stress non-invasively in unrestrained birds. *J. Vis. Exp.* (105), e53184.

**Kluger, M. J., O'Reilly, B., Shope, T. R. and Vander, A. J.** (1987). Further evidence that
stress hyperthermia is a fever. *Physiol. Behav.* **39**, 763-766.

**Korhonen, H., Hansen, S. W., Malmkvist, J. and Houbak, B.** (2000). Effect of capture,
immobilization and handling on rectal temperatures of confident and fearful male mink. *J.*
*Anim. Breed. Genet.* **117**, 337-345.

**Lenth, R. V.** (2019). emmeans: Estimated marginal means, aka least-squares means. R
package version 1.3.3. https://CRAN.R-project.org/package=emmeans

**Lewden, A., Nord, A., Petit, M. and Vézina, F.** (2017). Body temperature responses to
handling stress in wintering Black-capped Chickadees (*Poecile atricapillus* L.). *Physiol.*
*Behav.* **179**, 49-54.

**Lindgård, K. and Stokkan, K. A.** (1989). Daylength control of cood intake and body weight
in Svalbard ptarmigan *Lagopus mutus hyperboreus*. *Orn. Scand.* **20**, 176-180.

**Long, N. C., Vander, A. J. and Kluger, M. J.** (1990). Stress-induced rise of body
temperature in rats is the same in warm and cool environments. *Physiol. Behav.* **47**, 773-
775.

**Maggini, I., Tahamtani, F. M., Cardinale, M., Fusani, L. and Carere, C.** (2018). Body
temperature upon mist-netting procedures in three species of migratory songbirds at a
stopover site: implications for welfare. *Anim. Welf.* **27**, 93-101.

**Mans, C., Guzman, D. S.-M., Lahner, L. L., Paul-Murphy, J. and Sladky, K. K.** (2012).
Sedation and physiologic response to manual restraint after intranasal administration of
midazolam in Hispaniolan Amazon parrots (*Amazona ventralis*). *J. Avian Med. Sur.* **26**,
130-139.

**Marais, M., Maloney, S. K. and Gray, D. A.** (2011). The metabolic cost of fever in Pekin
ducks. *J. Therm. Biol.* **36**, 116-120.

**Meyer, L. C. R., Fick, L., Matthee, A., Mitchell, D. and Fuller, A.** (2008). Hyperthermia in
captured impala (*Aepyceros melampus*) a fright not flight response. *J. Wildl. Dis.* **44**, 404-
416.

**Mortensen, A. and Blix, A. S.** (1986). Seasonal changes in resting metabolic rate and mass-
specific conductance in Svalbard ptarmigan, Norwegian rock ptarmigan and Norwegian
willow ptarmigan. *Orn. Scand.* **17**, 8-13.

**Mortensen, A., Unander, S., Kolstad, M. And Blix, A. S.** (1983). Seasonal changes in body
composition and crop content of Spitzbergen ptarmigan *Lagopus mutus hyperboreus*. *Orn.*
*Scand.* **14**, 144-148.

**Møller, A. P.** (2010). Body temperature and fever in a free-living bird. *Comp. Biochem.*
*Physiol. B* **156**, 68-74.

**Nord, A. and Folkow, L. P.** (2018). Seasonal variation in the thermal responses to changing
environmental temperature in the world's northernmost land bird. *J. Exp. Biol.* **221**,
jeb171124.

**Oka, T., Oka, K. and Hori, T.** (2001). Mechanisms and mediators of psychological stress-
induced rise in core temperature. *Psychosom. Med.* **63**, 476-486.

**R Core Team.** (2018). R: A language and environment for statistical computing. Vienna,
Austria: R Foundation for Statistical Computing.

**Stokkan, K. A., Mortensen, A. and Blix, A. S.** (1986). Food intake, feeding rhythm, and
body mass regulation in Svalbard rock ptarmigan. *Am. J. Physiol.* **251**, R264-R267.

**Udvardy, M. D. F.** (1955). Body temperature of parids in the Arctic winter. *Orn. Fenn.* **32**,
101-107.

**Wang, J., Rao, H., Wetmore, G. S., Furlan, P. M., Korczykowski, M., Dinges, D. F. and**
**Detre, J. A.** (2005). Perfusion functional MRI reveals cerebral blood flow pattern under
psychological stress. *Proc. Natl. Acad. Sci. U.S.A.* **102**, 17804-17809.

**Vinkers, C. H., Groenink, L., van Bogaert, M. J. V., Westphal, K. G. C., Kalkman, C. J.,**
**van Oorschot, R., Oosting, R. S., Olivier, B. and Korte, S. M.** (2009). Stress-induced
hyperthermia and infection-induced fever: Two of a kind? *Physiol. Behav.* **98**, 37-43.

**Zuur, A. F., Ieno, E. N., Walker, N. J., Saveliev, A. A. and Smith, G. M.** (2009). *Mixed*
*effects models and extensions in ecology with R.* New York: Springer.

FIGURE LEGENDS

**Fig. 1. Male Svalbard ptarmigan in winter plumage overlooking the slopes of the**
**Arctowski mountain, Spitsbergen, Svalbard (78°12' N, 16°17' E) in early spring.** Photo ©
Andreas Nord

**Fig. 2. Mean ± s.e.m. changes in cloacal temperature ($T_c$ – orange), back skin**
**temperature ($T_{back}$ – red or green), and head skin temperature ($T_{head}$– blue) during**
**handling in Svalbard ptarmigan.** Panels show the response over: (A) the 22 min
observation period relative to pre-handling temperature; (B) the response during the handling
event ± 1 min relative to pre-handling temperature; and (C) mean ± s.e.m. absolute body
temperatures during the observation period. Data were collected in thermoneutrality (0°C) and
below thermoneutrality (-20°C) in continuous darkness (DD) in winter (-20°C: $n = 6$; 0°C: $n =$
10) and continuous light (LL) in spring (-20°C: $n = 5$; 0°C: $n = 8$). The vertical grey bars
show periods of handling. $T_{back}$ changes are presented separately for birds showing negative
(red) and positive (green) averages responses, of which the latter occurred in three birds in
three different measurement conditions. Data were binned in 30 s (A, C) or 15 s (B) intervals
before plotting. Metrics extracted from the depicted thermal responses, and ditto analyses, are
described in Materials & Methods.

B

C

Winter (DD) −20 °C | Winter (DD) 0 °C | Spring (LL) −20 °C | Spring (LL) 0 °C

Tissue temperature (°C)

Time after start of handling (min)