Modern clinical *Mycobacterium tuberculosis* strains leverage type I IFN pathway for a

pro-inflammatory response in the host

Deepthi Shankaran[1,2#], Prabhakar A[1, 2#], Ankur Bothra[3], Sheetal Gandotra[1,2], Vivek Rao[1,2*]

1-CSIR- Institute of genomics and Integrative Biology, New Delhi-110025, India.

2- Academy of Scientific and Innovative Research (AcSIR), ACSIR- Human Resource

Development Centre, (CSIR-HRDC) Campus, Uttar Pradesh- 201 002, India.

3- LPD/MPS, NIAID-NIH, Bethesda, MD 20892, USA. – Current address

#- Equal contribution

*Corresponding author: Tel: +91 11 29879229, E-mail: vivek.rao@igib.in

Running title: IFN1 induction is Mtb strain dependent

Key words: Mycobacterium tuberculosis/ clinical lineages/ Type I IFN/ macrophage

ABSTRACT:

Host phagocytes respond to infections by innate defense mechanisms through metabolic

shuffling in order to restrict the invading pathogen. However, this very plasticity of the host

provides an ideal platform for pathogen mediated manipulation. By employing the macrophage

model of Mtb infection, we identify an important strategy employed by modern clinical lineages

in regulating the host immune–metabolism axis. The potent ability of these strains to specifically

elicit a strong and early macrophage type I IFN response (in contrast to the protracted response

to ancient Mtb), was dependent on an increased ability to localize in acidified phagosomes; this

higher transit via acidified compartments is important for stimulation of the DNA dependent

signaling in infected macrophages. The augmented IFN signaling provided a positive regulatory

loop for enhanced expression of the cellular oxysterol-CH25H which in turn facilitated higher

levels of IL6 in macrophages infected with the modern Mtb strains. Requirement of type I IFN

signaling in mycobacterial intracellular growth highlights another unique ability of Mtb to

manipulate host cell physiology and proinflammatory responses.

Abbreviations: IFN- Interferon, BAF A- bafilomycin A, MDM- monocyte derived macrophages,

Mtb- *Mycobacterium tuberculosis*, BCG- *Bacille Calmette Guerin*, CH25H- Cholesterol 25-

Hydroxylase, lyso[+]- lysotracker positive, L1, L3- Lineage 1,3.

Significance Statement:

Co-evolution with humans has enabled the development of novel adaptive mechanisms for

survival in host specific environments in the human TB pathogen-Mtb. We present one such

mechanism of modern Mtb strains harnessing the type I IFN immune axis to regulate the host

pro-inflammatory response. Our results highlight the use of host intracellular endosomal transit

as a mechanism by these strains to ensure a strong type I IFN response in macrophages. We

also demonstrate the ability of Mtb to regulate macrophage cholesterol metabolism in order to

fine tune the host innate responses. These findings lay the foundation of the future development

of a host axis directed intervention strategy against this pathogen.

Introduction:

A long-standing association with the human population has been critical to the

development of specific immune modulatory mechanisms by *Mycobacterium tuberculosis* (Mtb)

for efficient survival within the host. With the advent of high throughput genome analysis, Mtb

strains have been differentiated into 7 distinct groups that can be classified into phylogenetically

ancient (restricted geographic penetration) and modern (widespread) lineages. Analysis of

strain specific differences in their virulence and disease manifestation has led to identification of

possible molecular correlates of virulence (*1-3*).

The initial contact of Mtb with macrophages activates the cGAS-STING-nucleic acid

signaling pathway leading to type I IFN expression (*4, 5*). This response is mediated by host

factors like ROS induced mitochondrial damage and pathogen associated factors like ESX-1

mediated phagosomal escape and mycobacterial cyclic di-nucleotides (*6-10*). A detrimental

pathogenic effect has been associated with Mtb induced type I IFN signaling in regulating the

expression of cytokines such as TNFα, IL12 and IL10 in macrophages (*11-13*). Moreover, type I

IFNs have been shown to alter the IFN γ dependent activation of macrophages and consequent

mycobacterial growth (*14*). However, another study has also suggested a beneficial role for the

mycobacteria induced type I IFN signaling in IL12 production and antigen presentation by

macrophages (*15*).

Clinical Mtb lineages vary significantly in their inflammatory response induction and

immune modulatory functions (*16-18*). Given the duality of the outcome of type I IFN induction by Mtb in macrophages, mechanisms underlying strain specific type I IFN is still unclear. We demonstrate that the higher type I IFN induction (IFNβ) by modern lineages (L3, L4 and L2) is dependent on bacterial DNA presentation to the cGAS- STING signaling pathway as a result of an early and increased bacterial trafficking via the endo-lysosomal pathway. Consistent with this augmented IFNβ and enhanced expression of the IFN stimulated, immunomodulatory (pro inflammatory) oxysterol- CH25H, significantly higher levels of IL6 were observed in macrophages infected with the modern Mtb strains. This study provides evidence for a novel mechanism of the plasticity of modern Mtb by coupling type I IFN with macrophage pro-inflammatory response. Understanding the physiological consequence of this phenomenon would provide important insights into the successful adaptation of modern Mtb to the host environment.

Results:

Mtb lineages induce differential type I IFN response in macrophages.

The type I IFN response of macrophages has generated interest as a pro- pathogenic response induced by Mtb in macrophages (*19-21*). The THP1-Dual cell line can be used to report on the induction of Type I IFN as it expresses a secreted luciferase reporter gene under the control of an ISG54 minimal promoter in conjunction with five IFN-stimulated response elements. Detectable levels of secreted luciferase as a read out for IFN activation was observed in a temporal manner in response to Mtb Erdman at a MOI of 5 with 10-15 folds activity by 24h and 48h (Fig. 1A).

Further, we also observed a distinct pattern of response in these macrophages infected with clinical Mtb strains of different phylogenetic lineages. Macrophages infected with the modern Mtb lineages (L2- HN878, L3-N4, N24; L4-H_{37}Rv, Erdman, H_{37}Ra) displayed 8-20 folds higher luminescence activity at 24h of infection contrasting with the lower (~2.5 folds) or complete absence of activity in macrophages infected with the ancient lineage I (T83, N72, N73) Mtb and BCG strains, respectively (Fig. 1B, Fig. S1). This higher activation potential of L3 Mtb was maintained across different infective doses (Fig. 1C). While L3 Mtb strains (N4 and N24) could

activate this response even at a low MOI of 1 (4.6 folds) by 24h of infection, these levels were

higher than the average levels of IFN response induction (~3 folds) by L1 Mtb (N72 and N73) at

MOI-10.

Gene expression profiles further validated the temporal and differential IFNβ expression profiles

of Mtb infected macrophages, with the L3 Mtb strain-N24 beginning to induce IFNβ expression

as early as 1h with a gradual increase (~30 folds by 12h of infection) that was sustained even

by 24h of infection (~16 folds) (Fig. 1D). In contrast, the L1 strain (N73) induced IFNβ at a

significantly later time point in infection (6-12h) and at 5-7-fold lower levels even by 24h of

infection.

This pattern of an attenuated IFN response by L1 Mtb was also evident in primary human

MDMs and the mouse macrophage line RAW264.7. Consistently, L3 Mtb induced significantly

higher (4-10 folds) IFN response than L1 Mtb at any point in infection (Fig. 1E, F). The induced

IFN response required active infection as macrophages infected with non-viable Mtb failed to

initiate this response (Fig. 1G). To verify if L1 Mtb actively suppressed type I IFN response, we

employed a co-infection assay of N24 and N73 in different ratios (Fig.1H). At all doses, N24

induced significantly higher response than N73, however, in mixed infections, presence of four-

fold excess N73 (N73:N24::4:1) could not diminish the response induced by N24. Consequently,

IFN levels in mixed infections were comparable to levels in macrophages infected with N24

alone, even at the overall higher MOI of infection of 10. Subsequently, in a trans-activation/

suppression assay, again N73, did not suppress poly IC mediated IFN response (Fig. 1I). In

most cases, response to poly IC was clearly augmented in conditions of co-stimulation with

either N24 or by N73.

Fig.1

Fig.1: Modern lineages have evolved to induce a significantly higher type I IFN response

A-C) Relative luciferase activity in naïve and stimulated THP1 dual macrophages: Infected with Mtb

Erdman at a MOI-5 (A), or with Mtb strains belonging to different lineages [L1-4] at a MOI-5 (B), 2

independent isolates of each lineage are represented as open and closed circles, T83 Vietnamese L1

subtype Mtb is shown as inverted triangle, or with Mtb L1 [grey bars; 1-N72, 2-N73] and L3 [black bars;

3- N4, 4-N24] at different MOIs (C). At the indicated time points, culture supernatants were harvested and

tested for luciferase expression. Values represent average ratios $\pm$ SE of RLU in infected and control

samples from triplicate wells of two independent experiments [N=2].

D-F) Relative expression in macrophages infected with N24 [L3] or N73 [L1] at a MOI-5. D) IFNβ

expression at different time intervals post infection, E) IFNβ, IFIT2 and IP10 mRNA in primary MDMs from

four separate individuals (symbols) and F) expression of IFNβ and IFIT2 mRNA in RAW macrophages.

The average relative values $\pm$ SE (with respect to GAPDH) in infected and control samples from triplicate

wells of multiple replicate experiments are represented.

G-I) Relative luciferase activity in naïve and stimulated THP1 dual macrophages: Infected with live or heat

killed (Δ) Mtb Erdman at MOI-5 (G) or with N24, N73 or in different combinations (H) or stimulated with

poly IC alone or in combination with Mtb at a MOI of 5 (I).

IFN inducing capability correlates with induction of a pro-inflammatory response via

expression of CH25H in macrophages.

To test if L1 Mtb strains were attenuated in overall macrophage activation, we tested gene

expression and cytokine profiles of RAW264.7 macrophages infected with L1 and L3 Mtb. Both

TNFα and IL6 patterns were strikingly similar to type I IFN, with lower levels in the case of L1

than the L3 infected RAW264.7 macrophages after 24 hours (Fig. 2A). This attenuated IL6 and

TNF expression was also observed in bone marrow-derived macrophages from C57BL/6 mice

with L1 Mtb inducing minimal amounts of the cytokines (Fig. 2B). L3 Mtb induced significantly

higher levels of IL6 in a temporal manner attaining peak values by 48h of infection as opposed

to minimal levels of cytokines even by 72h of L1 Mtb infection (Fig. 2C). A similar pattern of low

IL6 induction by L1 Mtb in human monocyte derived macrophages of 4 healthy individuals at 6h

and 24h of infection suggested a universal mechanism of regulation in the macrophage

response pathway by this mycobacterial strain (Fig. 2D).

To check if the regulation was at the transcript level, we compared gene expression by qPCR.

In macrophages infected with N24, ~30 fold higher *il6* expression was observed than in N73

infected cells as early as 6h *p.i.* While the expression declined at a later time point (24h) in

these macrophages, the overall levels were significantly higher than in the N73 infected cells

(Fig. 2E).

Both strains were capable of inducing *il6* by 6h; N24 by 34 folds and N73 by 7 folds. While we

did not observe any change in the levels of *il6* expression in the N73 infected ΔSTING

macrophages, the differential higher response (~5x) seen in the case of N24 infection was lost

(Fig. 2F). Although, *il6* levels were lower at 24h of infection with either Mtb strain, loss of STING

abrogated expression in macrophages suggestive of an inherent regulation of IL6 by IFN1 in

macrophages. More importantly, loss of STING did not significantly affect the TNF inducing

capacity of Mtb in macrophages. Recent evidence points to a positive regulatory circuit for IL6

via cGAS-STING regulated expression of CH25H, responsible for the synthesis of the pro

inflammatory oxysterol, 25 hydroxy-cholesterol (*22, 23*).   Examination of genes involved in

cholesterol biosynthesis revealed a temporal but distinct pattern of expression in L3 and L1 Mtb

infected BMDM (Fig. S2). Most of the genes involved in the cholesterol biosynthetic pathway

were either up-regulated or similar in L1 infected macrophages after 24h of infection. In

contrast, these genes were inhibited in response to infection with L3 Mtb. Surprisingly, in

contrast with the overall pathway, the expression of CH25H was significantly upregulated in the

L3 infected macrophages (60-70-fold higher than the uninfected control) as opposed to a much

lower expression of this gene (3-4 folds) in the L1 infected macrophages, by 6h of infection (Fig.

2G). This difference was, however, lower at 24h, wherein a sharp decline in L3 (23 folds) and

an increase in L1 infected macrophages (10-12 folds) was observed. Given the increased

CH25H expression and requirement of type I IFN for enhanced IL6 expression, we tested if

STING signaling contributed to the early increase of CH25H in L3 Mtb infected macrophages.

The complete loss of CH25H expression in ∆STING macrophages substantiates our hypothesis

of a regulatory role for type I IFN in host oxysterol mediated IL6 response in modern Mtb

infected macrophages (Fig. 2H).

Fig. 2

Fig.2: IL6 response in Mtb infected macrophages primarily depends on STING mediated CH25H

expression.

A-C) Levels of cytokine secreted from RAW264.7 macrophages (A), BMDMs at 24h (B) or at different

time intervals (C) following infection with two L3 or L1 Mtb strains. Values represent mean cytokine

concentration in pg/ml $\pm$ SEM of triplicate assays (N=3).

(D) Levels of IL6 in Mtb infected MDMs from four individuals at 6h and 24h *p.i.* Values represent mean

cytokine concentration in pg/ml in the culture supernatant of triplicate wells for each individual.

E, F) IL6 and TNF gene expression profiles in BMDMs (E) and RAW264.7 macrophages (F) following

infection with N24 (L3) or N73 (L1) Mtb strains. The average relative IL6 values $\pm$ SE (with respect to

GAPDH) at 6h and 24h post infection in triplicate assays is shown.

G, H) Expression of CH25H in BMDMs (G) and RAW macrophages (H) following infection with Mtb

strains at 6h and 24h *p.i.* at MOI-5. The average relative IL6 values $\pm$ SE (with respect to GAPDH) at 6h

and 24h *p.i.* in triplicate assays of (N=2/3) is shown.

L1 Mtb are delayed in inducing type I IFN response in macrophages despite the presence

of an active ligand.

Previous studies have implicated an important role of the Mtb ESX-1 secretion system in cytosolic egress of ligands and activation of the cytosolic surveillance pathway resulting in type I IFN activation (*5, 10*). However, we did not observe any difference between the IFN inducing abilities of $H_{37}Ra$ or $H_{37}Rv$ despite the reported differences in the levels of the Esx-1 proteins (*24*). Further, comparable expression of ESAT6 in our clinical Mtb strains (L1 and L3) (Fig. S3A) argued for an esx-1 independent mechanism of Type 1 IFN induction.

We also did not observe any difference in mitoSOX staining of macrophages infected with either Mtb strain, contrasting with the previously identified mode of mitochondrial ROS mediated type I IFN signaling (*10*) hinting at alternative early events associated with recognition of bacterial ligands (Fig. S3B).

Having ascertained that N73 did not suppress IFN induction, it was logical to assume lack/ lower quantities of the active ligand as an underlying cause for the deferred IFN response by L1 Mtb. In accordance with previous reports, Mtb induced type I IFN response of macrophages was completely dependent on DNA mediated signaling. Wt RAW-luc (ISG) cells also portrayed a differential type I IFN response to Mtb lineages, with 5-10-fold higher IFN by L3 Mtb in comparison to L1 Mtb strains. Only the loss of cGAS, STING and TRIF (~60% decrease), associated with DNA mediated signaling as opposed to RNA sensing, diminished the type I IFN response to L3 Mtb (Fig. 3A, S4A). Lipofectamine based transfection of DNA results in delivery of DNA to the endosomal system, and eventually to the cytosol via formation of multiple transient pores in the endosomal membrane (*25*). As cytosolic sensors such as STING are essential for eliciting this response in case of Mtb infection (*10, 26, 27*), we argued that delivery of mycobacterial DNA by lipofectamine may also be able to elicit this response. In fact, with as low as 125 ng of genomic DNA from Mtb, macrophages responded with IFN expression that was again abolished in cells stimulated with DNAse treated DNA (Fig. 3B). Surprisingly, genomic DNAs from N73 as well as BCG were similar to Erdman and N24 in their IFN inducing capacities (Fig. 3C). Given the presence of an active molecule (DNA) even in Mtb strains with attenuated IFN induction, it was reasonable to assume the lack of DNA presentation as the basis for lower IFN. A preliminary evaluation hinted at a temporal IFN response to Mtb Erdman

extract (cell lysates) when delivered to THP1 cells via transfection, with the response increasing

from ~10 folds at 3h to ~75 folds by 6h and then declining to ~10-fold expression by 24h of

stimulation with different doses (Fig. 3D). Even though very low amounts of the extract (1.6 ng)

was sufficient to induce ten-fold IFN by 6h (Fig. S4B), the complete loss of IFN only in extracts

subjected to DNAse digestion argued for the pivotal role of DNA sensing in Mtb induced type I

IFN in macrophages and its inaccessibility to the IFN signaling machinery of macrophages (Fig.

3E). Further proof of effective ligand masking by L1 Mtb and BCG came from our observation

that delivery of BCG with lipofectamine was sufficient to enhance the Type I IFN response by 6

folds compared to BCG infection without lipofectamine (Fig. 3F).

Fig. 3

Fig.3: Strains with suppressed type I IFN response are well equipped with the active IFN inducing

ligand.

A)   Relative luciferase activity in Wt- ISG or mutant RAW macrophages uninfected or infected with

Mtb strains belonging to L3 / L1 lineages at a MOI of 5 for 24h. Values represent average ratios ± SE

of RLU in infected and control culture supernatants from triplicate wells of three independent

experiments [N=3].

B-F) Relative luciferase activity in naïve and stimulated THP1 dual macrophages: following

transfection with genomic DNA (B), or genomic DNA of mycobacteria (C) or crude Mtb Erdman crude

lysate (D, E), or BCG transfection (F). H, P, R, D-represent treatment of extract with Heat, proteinase

K, RNase and DNase, respectively.   Values represent average ratios $\pm$ SE of RLU in infected and

control culture supernatants from triplicate wells of multiple experiments [N=2, 3].

231   Early localization to acidified compartments is important for Mtb induced type I IFN

232   induction

The early induction of type I IFN in L3 infected macrophages advocated for molecular events

that ensue immediately after infection as causative. Further, analysis of THP1 macrophages by

TEM revealed the differential localization of the Mtb clinical isolates in distinct intracellular

compartments- N24 localized to compact double membranous vesicles at 24h in contrast to N73

that were confined in larger less defined vesicular structures with intracellular particles (Fig. 4A).

In support of this observation, we observed that Mtb N24 localized to lysotracker positive (lyso+)

vesicles at significantly higher numbers than N73 at 24h *p.i.* (Fig. 4B).   This differential

localization to lyso+ vesicles between N24 and N73 was also observed in RAW264.7

macrophages. In a temporal analysis of Mtb transit via the endosomal pathway, differential

localization between Mtb strains arose as early as 1h of infection with more than two-fold N24 in

the lyso+ vesicles as compared to N73 (Fig. 4C). Despite a gradual decrease of N24 in lyso+

positive vesicles with time, consistently 2-fold higher numbers of N24 was seen throughout the

experiment duration.

We have previously demonstrated that L1 strains harbor a mutation in papA2, rendering them

incapable of sulfolipid synthesis (*28*). A dominant role for the Mtb sulfolipids has also been

demonstrated in modulation of the cellular phagosome maturation and acidification (*29-31*). We

questioned whether the lack of sulfolipids renders Mtb incapable of inducing the type I IFN

response. To test this hypothesis, we deleted *mmpL8*, coding for an exporter of mature

sulfolipids to the bacterial cell surface and measured induction of the Type I IFN response in the

reporter cells. There was no loss of IFN inducing ability of Mtb in the absence of *mmpL8* (Fig.

S5), supporting an alternative reason for the differential induction of the type I IFN response.

Another finding supporting the lack of sulfolipid not to be responsible for this phenotype is the

inability of the Vietnamese strain of L1 lineage to induce significant Type I IFN response despite

expressing sulfolipids in the membrane (Fig. 1B).

Fig. 4

Fig.4: Early and sustained localization of the high IFN inducing Mtb strains with acidified vesicles.

A)   Representative TEM images (2) of Mtb N24 and N73 in THP1 macrophages at 24h post infection.

While N24 is found in compact vesicles, N73 localizes in enlarged vesicles (black arrow) with cellular

debris.

B-E) Localization of Mtb N24 and N73 in acidified vesicles. Representative images of macrophages

stained with lysotracker red and infected with GFP expressing N24/ N73 - THP1 (B) at 24h or RAW 264. 7

(C) at 1h, 3h, 6hand 24 h post infection is depicted. The extent of co- localization of Mtb with the lyso+

vesicles in THP1, 24h and RAW264.7 at 1h, 3h, 6h and 24h is represented graphically. Each dot

represents a field of image from 2 independent experiments (> 100 infected cells).

IFN inducing capacity of Mtb relies on efficient acidification of the phagosome.

Given the high numbers of N24 localization to lyso$^+$ compartments, and comparable intracellular

growth rates of N24 and N73 (data not shown), it was reasonable to assume an enhanced

propensity of N24 to be better equipped for survival in this subcellular niche. The ability of Mtb

to counter phagosomal-lysosomal fusion and acidification events is well recognized (*32, 33*). To

understand if phagosomal acidification was important for type I IFN induction, we treated

macrophages with an established phagocytosis activation signal (LPS+ IFNγ) or phagosomal

acidification blockers- bafilomycin A and chloroquine prior to infection with Mtb. Activation of

macrophages with IFNγ and LPS enhanced the presence of Mtb in lyso$^+$ vesicles (Fig. 5A).

Importantly, the numbers of lyso+ N73 Mtb increased to comparable levels as N24 Mtb by 6h of

infection (Fig. 5B). Consequently, a 10-fold higher type I IFN response was observed in N73

infected IFNγ+ LPS pretreated macrophages leading to loss of the differential IFN response

between strains (Fig. 5C and inset panel). While, addition of IFNγ and LPS significantly boosted

type I IFN induction even in N24 infected macrophages, treatment with phagosome acidification

blockers like bafilomycin A or chloroquine on the contrary, stunted (~5 and 2 folds, respectively)

the IFN response in these cells (Fig. 5D) emphasizing the importance of phagosomal trafficking

and acidification in Mtb mediated IFN induction in macrophages.

Fig. 5

Abrogation of DNA signaling arrests mycobacterial growth in macrophages

In an effort to test the physiological relevance of type I IFN, we analyzed the ability of Mtb N24

to grow in Wt ISG and macrophages deficient in STING or TRIF. Over 3 days of culture, Wt

macrophages supported Mtb growth (Fig. 6). In contrast, despite, similar levels of uptake in the

deficient macrophages, Mtb N24 growth was restricted to initial levels at all times

(bacteriostasis) strongly arguing for an important role for the early type I IFN signaling in

intracellular growth of Mtb.

Fig. 6

Fig.6: Growth of Mtb N24 in RAW reporter macrophages. Macrophages were infected with Mtb at an MOI of 5 for 6h and the intracellular bacterial numbers were enumerated by dilution plating. Average bacterial numbers / well $\pm$ SEM in WT (ISG- solid squares) or ΔSTING (open triangle) or ΔTRIF (open circle) are represented graphically at 0, 1, 2 and 3 days *p.i.*

The working model

Clinical Mtb strains differ in their intracellular trafficking routes to arrive at or avoid acidified compartments; modern lineages traffic to arrive early into acidified compartments of host macrophages. The reduced movement of ancient Mtb strains likely hinders the access of Mtb DNA to cytosolic DNA sensors, thereby limiting the nucleic acid response pathway and leading to weaker type I IFN response. The higher activation of type I IFN by the modern strains manifests as a substantial upregulation of the pro-inflammatory cytokine IL6 via induction of cholesterol 25- hydroxylase in a positive feedforward loop and is essential for bacterial growth in macrophages (Fig. 6).

Fig. 7

Fig.7: A putative working model of modern Mtb induced IFN and subsequent IL6 expression via the STING-CH25H pathway.

DISCUSSION

The grand success of Mtb as a human pathogen stems from its extreme plasticity to suit the host cell niche, modulate the host response and survive multiple intracellular stresses. The immediate response of host phagocytes upon contact is the activation of inflammatory responses aimed at controlling the growth of the invading pathogen. Typically the type I IFN response is directed against viral infections, but recent studies have highlighted the macrophage expression of type I IFNβ in response to intracellular (pathogenic) and extracellular bacterial (protective) infections (*34-39*). We observed that modern Mtb strains are proficient in inducing this response than the evolutionarily older strains of Mtb. We also observed a strong correlation between the STING mediated IFN induction and high levels of IL6 response in macrophages, arguing for a novel mechanism of macrophage response regulation by the modern Mtb strains. Recent evidence also connects the type I IFN induced cholesterol 25-hydroxylase (CH25H) with the expression of pro-inflammatory cytokines like IL6 in macrophages (*22, 23, 40-42*). Given the lower expression of genes for cholesterol biosynthesis in the late stage of infection with N24 Mtb in line with the earlier report (*43*), it was surprising that expression levels of CH25H were on the contrary increased, signifying its importance in bridging type I IFN and IL6 expression in macrophages. Moreover, the complete loss of CH25H expression in the STING deficient macrophages following N24 infection highlights this novel regulatory loop of type I IFN- CH25H- IL6 regulation in modern Mtb infections. Overall, we define a novel pathway of pro-inflammatory cytokine expression in Mtb induced macrophages involving the early innate type I IFN response and active modulation of the host cell oxysterol metabolism. Our results identify a novel link between known pathogenicity of modern lineages (*16*) and their ability to mount a feedforward pro-inflammatory response via the Type I IFN axis. As demonstrated previously, loss of STING/ cGAS completely abrogated Mtb type I IFN response in macrophages implicating cellular DNA signaling as a predominant mechanism of

this response. Given the early induction of type I IFN and the absence of this response despite

the presence of active ligand, we argued that a novel mechanism that actively concealed the

DNA from the cellular nucleic acid signaling mechanism was responsible for the lower IFN

induction by the attenuated ancient Mtb strains. Interestingly, loss of TRIF also resulted in

absence of IFN response in N24 infected macrophages suggesting a role for RNA signaling in

this response. Surprisingly, we did not observe loss of IFN induction by cellular extracts treated

with RNAse. Recent reports have identified a TRIF dependent activation of STING in the DNA

induced IFN response of macrophages (*44*); our results demonstrate a dominant role for this

signaling pathway in Mtb mediated type I IFN response.

Two separate observations supported our hypothesis that intracellular localization early in

infection would be primarily accountable for this differential IFN signaling by Mtb strains:

Induction of IFN response by 1) transfected BCG as opposed to direct infection and 2) absolute

requirement of DNA to be transfected. Bacterial lysis by the phagosome-lysosome pathway has

been implicated in several of the innate immune response pathways in phagocytes (*45, 46*).

Our results provide insights into the importance of transit of Mtb strains via an acidified

compartment as an important determinant of IFN induction in macrophages. Based on the

presence of a double membrane compartment, we speculate this to be an autophagosome like

compartment. While on the one hand only live bacteria were able to stimulate the macrophage

type I IFN response, dead Mtb despite their well-documented higher rates of trafficking to

acidified P-L compartments were unable to induce this response in macrophages, pointing at

the ability of modern Mtb strains to utilize its early localization to an acidified compartment for

active presentation of DNA (in contrast to dead bacteria that fail in this presentation) to the

cytosolic signaling complex. Additionally, the loss of Mtb mediated IFN response in Baf A/

chloroquine treated macrophages and a significant increase by LPS+IFNγ pretreatment reveals

an important role of phagosomal acidification in type I IFN induction.

While sulfolipids have been demonstrated in promoting phagosomal acidification and L1 strains

lack sulfolipids, our results do not purport an active role for this lipid in the type I IFN response.

Despite the presence of mature sulfolipids in the L1 subtype Mtb strains T83 (*28*), induction of

type I IFN was relatively low in THP1 macrophages. Moreover, while differences in sulfolipid

expression between $H_{37}Ra$ and $H_{37}Rv$ is well established (*47, 48*), there was no discernable

alteration of IFN expression between the two Mtb strains. The complete lack of change in IFN

induction ability of a sulfolipid deficient Erdman Mtb (ΔM8) further absolves the critical role of

sulfolipid in type I IFN induction by Mtb (Fig. S5).

In summary, we identify a novel means for modern strains of Mtb to mount a pathogenic

immune response via their early passage through an acidified endosomal compartment. This

passage allows presentation of their DNA to cytosolic sensors, which then triggers a cholesterol

driven feed forward loop of pro-inflammatory responses. An understanding of these early

trafficking events in the future will help identify means to counter this pathogenic immune

response triggered by the modern Mtb strains.

Material and Methods

Bacterial Strains and Growth Conditions- Mycobacterial and *E. coli* strains were cultured as

per standard protocols with the appropriate supplements and selection antibiotics as required*.*

The M6 promoter fragment was amplified as described earlier (*49*) and cloned upstream of GFP

in the mycobacterial expression vector- pMV261 and electroplated into mycobacteria.

Expression of ESAT6 in the culture filtrate and secreted Mtb cultures was performed by

immunoblotting with anti ESAT6 monoclonal antibody HYB76-8 as described earlier (*3*). The

*mmpL*8 null mutant (ΔM8) was created by homologous recombination using specialized

transduction and recombineering as described earlier (*50*). Transducing phages generate by

recombineering mediated generation of phages were used for transduction of Wt Erdman Mtb

cultures and hyg^R colonies were screened for deletion by qPCR and checking expression of

sulfolipid in apolar lipid fractions by TLC as per recommended protocols (*51*).

Macrophage culture and maintenance- All the cell lines used were tested for mycoplasma

contamination at regular intervals and cultured according to the manufacturer's

recommendations. THP1 cells were grown in HiglutaXL RPMI-1640 (Himedia laboratories,

India) with 10% FBS and differentiated by using PMA as described earlier (*50*). Human

monocyte derived macrophages (MDMs) were isolated from 20ml as described earlier (*49*). For

primary mouse macrophages, bone marrow cells were harvested from the hind limbs of C57Bl6

mice and differentiated into macrophages in 20% L- cell supernatant for 7 days as described

earlier (*52*).

Preparation of Mtb fractions for stimulation of macrophages- Mtb crude extracts were

prepared from logarithmic liquid cultures with similar bacterial densities for the different strains.

Cells were lysed with a mixture of 0.1 µm and 0.7 µm zirconia beads in PBS using a bead

beater (Biospec India). The resultant lysates were cleared off the cell debris by high speed

centrifugation and sterilized by filtration and used to transfect THP1 dual cells using

Lipofectamine LTX (Invitrogen, USA) reagent according to the recommended protocol. Genomic

DNA was isolated according to the manufacturer's recommendations (Nucleopore gDNA kit)

and used for transfection. Enzymatic treatment of lysates was performed as: 1) Proteinase K- at

55^0C for 1h, 2) DNase, RNase at 37^0C for 1h and thermal denaturation at 96^0C for 30min.

Infection of macrophages with Mtb- Macrophage cultures were seeded at the requisite

density and infected with single cell suspensions (SCS) as described earlier (*50*). At different

time intervals, the cells/ RNA/ supernatants were harvested and used for analysis. Bacterial

numbers were counted from individual wells by standard dilution plating of the lysate.

Macrophages infected with fluorescent Mtb strains were used for lysosomal localization by using

100nm Lysotracker–Red DND-99 (Invitrogen, USA) according to manufacturer's

recommendations. Macrophages transfected with mito-DsRed (kind gift from Dr. Sowmya Sinha

Roy) was used for analysis of mitochondrial architecture. The cells were fixed in 4%

paraformaldehyde for 30 min, mounted on slides with Prolong diamond anti fade (Invitrogen,

USA) and subjected to microscopy in the Leica TCS SP8 Confocal Microscope (Leica

Microsystems, USA). For estimating mitochondrial ROS by Mitosox staining, macrophages were

washed thrice with PBS and stained with 5µM Mitosox (Invitrogen, USA) in HBSS for 30 min at

37^0C, fixed and subjected to high content microscopy in a IN Cell Analyzer 6000(GE healthcare

life sciences, USA).

Transmission Electron Microscopy- THP1 cells were infected with mycobacteria at a MOI of

5 for 24h, fixed in 2.5% gluteraldehyde and 4% paraformaldehyde, dehydrated in graded series

of alcohol and embedded in Epon 812 resin. Ultrathin sections were cut and stained with uranyl

acetate and lead citrate and imaged by using Tecnai G2 20 twin (FEI) transmission electron

microscope (FEI, Thermo Fisher Scientific Inc, USA).

Statistical Analysis

423 All experiments were performed in multiple biological replicates (N=2 or 3). Statistical analysis

424 was done by using Student's t-test and corresponds to a *, **, *** for p-values<0.05, <0.01and

425 <0.001, respectively.

Ethics Statement

427 The protocol for isolation of human blood PBMCs was approved by the Institutional Human

428 ethics committee (Ref no: CSIR-IGIB/IHEC/2017-18 Dt. 08.02.2018). Blood was collected from

429 people with their informed consent as per the Institutional guidelines**.**

References

430
1.   I. Comas, M. Coscolla, T. Luo, S. Borrell, K. E. Holt, M. Kato-Maeda, J. Parkhill, B. Malla, S. Berg,
G. Thwaites, D. Yeboah-Manu, G. Bothamley, J. Mei, L. Wei, S. Bentley, S. R. Harris, S. Niemann,
R. Diel, A. Aseffa, Q. Gao, D. Young, S. Gagneux, Out-of-Africa migration and Neolithic
coexpansion of Mycobacterium tuberculosis with modern humans. *Nat Genet* **45**, 1176-1182
(2013); published online EpubOct (10.1038/ng.2744

ng.2744 [pii]).
2.   M. Pareek, J. Evans, J. Innes, G. Smith, S. Hingley-Wilson, K. E. Lougheed, S. Sridhar, M. Dedicoat,
P. Hawkey, A. Lalvani, Ethnicity and mycobacterial lineage as determinants of tuberculosis
disease phenotype. *Thorax* **68**, 221-229 (2013); published online EpubMar (10.1136/thoraxjnl-
2012-201824

thoraxjnl-2012-201824 [pii]).
3.   V. Rao, N. Dhar, A. K. Tyagi, Modulation of host immune responses by overexpression of
immunodominant antigens of Mycobacterium tuberculosis in bacille Calmette-Guerin. *Scand J*
Immunol **58**, 449-461 (2003); published online EpubOct (1321 [pii]).
4.   S. A. Stanley, J. E. Johndrow, P. Manzanillo, J. S. Cox, The Type I IFN response to infection with
Mycobacterium tuberculosis requires ESX-1-mediated secretion and contributes to
pathogenesis. *Journal of immunology* **178**, 3143-3152 (2007); published online EpubMar 01 (
5.   P. S. Manzanillo, M. U. Shiloh, D. A. Portnoy, J. S. Cox, Mycobacterium tuberculosis activates the
DNA-dependent cytosolic surveillance pathway within macrophages. *Cell host & microbe* **11**,
469-480 (2012); published online EpubMay 17 (10.1016/j.chom.2012.03.007

S1931-3128(12)00125-4 [pii]).
6.   A. C. Collins, H. Cai, T. Li, L. H. Franco, X. D. Li, V. R. Nair, C. R. Scharn, C. E. Stamm, B. Levine, Z. J.
Chen, M. U. Shiloh, Cyclic GMP-AMP Synthase Is an Innate Immune DNA Sensor for
Mycobacterium tuberculosis. *Cell host & microbe* **17**, 820-828 (2015); published online EpubJun
10 (10.1016/j.chom.2015.05.005).
7.   A. K. Pandey, Y. Yang, Z. Jiang, S. M. Fortune, F. Coulombe, M. A. Behr, K. A. Fitzgerald, C. M.
Sassetti, M. A. Kelliher, NOD2, RIP2 and IRF5 play a critical role in the type I interferon response

to Mycobacterium tuberculosis. *PLoS pathogens* **5**, e1000500 (2009); published online EpubJul
(10.1371/journal.ppat.1000500).

8. P. S. Redford, K. D. Mayer-Barber, F. W. McNab, E. Stavropoulos, A. Wack, A. Sher, A. O'Garra,
Influenza A virus impairs control of Mycobacterium tuberculosis coinfection through a type I
interferon receptor-dependent pathway. *J Infect Dis* **209**, 270-274 (2014); published online
EpubJan 15 (10.1093/infdis/jit424

jit424 [pii]).

9. B. Dey, R. J. Dey, L. S. Cheung, S. Pokkali, H. Guo, J. H. Lee, W. R. Bishai, A bacterial cyclic
dinucleotide activates the cytosolic surveillance pathway and mediates innate resistance to
tuberculosis. *Nat Med* **21**, 401-406 (2015); published online EpubApr (10.1038/nm.3813

nm.3813 [pii]).

10. K. E. Wiens, J. D. Ernst, The Mechanism for Type I Interferon Induction by Mycobacterium
tuberculosis is Bacterial Strain-Dependent. *PLoS pathogens* **12**, e1005809 (2016); published
online EpubAug (10.1371/journal.ppat.1005809).

11. R. A. de Paus, A. van Wengen, I. Schmidt, M. Visser, E. M. Verdegaal, J. T. van Dissel, E. van de
Vosse, Inhibition of the type I immune responses of human monocytes by IFN-alpha and IFN-
beta. *Cytokine* **61**, 645-655 (2013); published online EpubFeb (10.1016/j.cyto.2012.12.005

S1043-4666(12)00815-0 [pii]).

12. K. D. Mayer-Barber, B. B. Andrade, S. D. Oland, E. P. Amaral, D. L. Barber, J. Gonzales, S. C.
Derrick, R. Shi, N. P. Kumar, W. Wei, X. Yuan, G. Zhang, Y. Cai, S. Babu, M. Catalfamo, A. M.
Salazar, L. E. Via, C. E. Barry, 3rd, A. Sher, Host-directed therapy of tuberculosis based on
interleukin-1 and type I interferon crosstalk. *Nature* **511**, 99-103 (2014); published online
EpubJul 3 (10.1038/nature13489

nature13489 [pii]).

13. F. W. McNab, J. Ewbank, A. Howes, L. Moreira-Teixeira, A. Martirosyan, N. Ghilardi, M. Saraiva,
A. O'Garra, Type I IFN induces IL-10 production in an IL-27-independent manner and blocks
responsiveness to IFN-gamma for production of IL-12 and bacterial killing in Mycobacterium
tuberculosis-infected macrophages. *Journal of immunology* **193**, 3600-3612 (2014); published
online EpubOct 1 (10.4049/jimmunol.1401088

jimmunol.1401088 [pii]).

14. R. M. Teles, T. G. Graeber, S. R. Krutzik, D. Montoya, M. Schenk, D. J. Lee, E. Komisopoulou, K.
Kelly-Scumpia, R. Chun, S. S. Iyer, E. N. Sarno, T. H. Rea, M. Hewison, J. S. Adams, S. J. Popper, D.
A. Relman, S. Stenger, B. R. Bloom, G. Cheng, R. L. Modlin, Type I interferon suppresses type II
interferon-triggered human anti-mycobacterial responses. *Science* **339**, 1448-1453 (2013);
published online EpubMar 22 (10.1126/science.1233665).

15. E. Giacomini, M. E. Remoli, V. Gafa, M. Pardini, L. Fattorini, E. M. Coccia, IFN-beta improves BCG
immunogenicity by acting on DC maturation. *J Leukoc Biol* **85**, 462-468 (2009); published online
EpubMar (10.1189/jlb.0908583

jlb.0908583 [pii]).

16. D. Portevin, S. Gagneux, I. Comas, D. Young, Human macrophage responses to clinical isolates
from the Mycobacterium tuberculosis complex discriminate between ancient and modern
lineages. *PLoS pathogens* **7**, e1001307 (2011); published online EpubMar
(10.1371/journal.ppat.1001307).

17. G. Weiss, U. E. Schaible, Macrophage defense mechanisms against intracellular bacteria.
Immunological reviews **264**, 182-203 (2015); published online EpubMar (10.1111/imr.12266).

18. C. Manca, L. Tsenova, S. Freeman, A. K. Barczak, M. Tovey, P. J. Murray, C. Barry, G. Kaplan,
Hypervirulent M. tuberculosis W/Beijing strains upregulate type I IFNs and increase expression
of negative regulators of the Jak-Stat pathway. *Journal of interferon & cytokine research : the*

506   *official journal of the International Society for Interferon and Cytokine Research* **25**, 694-701
507   (2005); published online EpubNov (10.1089/jir.2005.25.694).

508  19.  C. H. Liu, H. Liu, B. Ge, Innate immunity in tuberculosis: host defense vs pathogen evasion. *Cell*
509   *Mol Immunol*, (2017); published online EpubSep 11 (10.1038/cmi.2017.88

510 cmi201788 [pii]).

511  20.  L. Petit-Jentreau, L. Tailleux, J. L. Coombes, Purinergic Signaling: A Common Path in the
512   Macrophage Response against Mycobacterium tuberculosis and Toxoplasma gondii. *Front Cell*
513   *Infect Microbiol* **7**, 347 (2017)10.3389/fcimb.2017.00347).

514  21.  A. Dorhoi, V. Yeremeev, G. Nouailles, J. Weiner, 3rd, S. Jorg, E. Heinemann, D. Oberbeck-Muller,
515   J. K. Knaul, A. Vogelzang, S. T. Reece, K. Hahnke, H. J. Mollenkopf, V. Brinkmann, S. H. Kaufmann,
516   Type I IFN signaling triggers immunopathology in tuberculosis-susceptible mice by modulating
517   lung phagocyte dynamics. *European journal of immunology* **44**, 2380-2393 (2014); published
518   online EpubAug (10.1002/eji.201344219).

519  22.  S. Y. Bah, P. Dickinson, T. Forster, B. Kampmann, P. Ghazal, Immune oxysterols: Role in
520   mycobacterial infection and inflammation. *J Steroid Biochem Mol Biol* **169**, 152-163 (2017);
521   published online EpubMay (S0960-0760(16)30111-X [pii]

522 10.1016/j.jsbmb.2016.04.015).

523  23.  E. S. Gold, A. H. Diercks, I. Podolsky, R. L. Podyminogin, P. S. Askovich, P. M. Treuting, A. Aderem,
524   25-Hydroxycholesterol acts as an amplifier of inflammatory signaling. *Proceedings of the*
525   *National Academy of Sciences of the United States of America* **111**, 10666-10671 (2014);
526   published online EpubJul 22 (10.1073/pnas.1404271111

527 1404271111 [pii]).

528  24.  S. M. Pinto, R. Verma, J. Advani, O. Chatterjee, A. H. Patil, S. Kapoor, Y. Subbannayya, R. Raja, S.
529   Gandotra, T. S. K. Prasad, Integrated Multi-Omic Analysis of Mycobacterium tuberculosis H37Ra
530   Redefines Virulence Attributes. *Frontiers in microbiology* **9**, 1314
531   (2018)10.3389/fmicb.2018.01314).

532  25.  Z. ur Rehman, D. Hoekstra, I. S. Zuhorn, Mechanism of polyplex- and lipoplex-mediated delivery
533   of nucleic acids: real-time visualization of transient membrane destabilization without
534   endosomal lysis. *ACS Nano* **7**, 3767-3777 (2013); published online EpubMay 28
535   (10.1021/nn3049494).

536  26.  Y. Cheng, J. S. Schorey, Mycobacterium tuberculosis-induced IFN-beta production requires
537   cytosolic DNA and RNA sensing pathways. *J Exp Med* **215**, 2919-2935 (2018); published online
538   EpubNov 5 (10.1084/jem.20180508

539 jem.20180508 [pii]).

540  27.  L. H. Yamashiro, S. C. Oliveira, A. Bafica, Innate immune sensing of nucleic acids from
541   mycobacteria. *Microbes Infect* **16**, 991-997 (2014); published online EpubDec
542   (10.1016/j.micinf.2014.09.006

543 S1286-4579(14)00213-5 [pii]).

544  28.  V. Panchal, N. Jatana, A. Malik, B. Taneja, R. Pal, A. Bhatt, G. S. Besra, L. Thukral, S. Chaudhary, V.
545   Rao, A novel mutation alters the stability of PapA2 resulting in the complete abrogation of
546   sulfolipids in clinical mycobacterial strains. *Faseb Bioadvances* **1**, 306-319 (2019).

547  29.  M. B. Goren, P. D'Arcy Hart, M. R. Young, J. A. Armstrong, Prevention of phagosome-lysosome
548   fusion in cultured macrophages by sulfatides of Mycobacterium tuberculosis. *Proc Natl Acad Sci*
549   *U S A* **73**, 2510-2514 (1976); published online EpubJul (10.1073/pnas.73.7.2510).

550  30.  C. Passemar, A. Arbues, W. Malaga, I. Mercier, F. Moreau, L. Lepourry, O. Neyrolles, C. Guilhot,
551   C. Astarie-Dequeker, Multiple deletions in the polyketide synthase gene repertoire of
552   Mycobacterium tuberculosis reveal functional overlap of cell envelope lipids in host-pathogen
553   interactions. *Cell Microbiol* **16**, 195-213 (2014); published online EpubFeb (10.1111/cmi.12214).

31. T. A. Mendum, H. Wu, A. M. Kierzek, G. R. Stewart, Lipid metabolism and Type VII secretion
systems dominate the genome scale virulence profile of Mycobacterium tuberculosis in human
dendritic cells. *BMC Genomics* **16**, 372 (2015); published online EpubMay 9 (10.1186/s12864-
015-1569-2

10.1186/s12864-015-1569-2 [pii]).
32. J. Sun, X. Wang, A. Lau, T. Y. Liao, C. Bucci, Z. Hmama, Mycobacterial nucleoside diphosphate
kinase blocks phagosome maturation in murine RAW 264.7 macrophages. *PloS one* **5**, e8769
(2010); published online EpubJan 19 (10.1371/journal.pone.0008769).
33. I. Vergne, R. A. Fratti, P. J. Hill, J. Chua, J. Belisle, V. Deretic, Mycobacterium tuberculosis
phagosome maturation arrest: mycobacterial phosphatidylinositol analog phosphatidylinositol
mannoside stimulates early endosomal fusion. *Mol Biol Cell* **15**, 751-760 (2004); published
online EpubFeb (10.1091/mbc.e03-05-0307

E03-05-0307 [pii]).
34. F. J. Martin, M. I. Gomez, D. M. Wetzel, G. Memmi, M. O'Seaghdha, G. Soong, C. Schindler, A.
Prince, Staphylococcus aureus activates type I IFN signaling in mice and humans through the Xr
repeated sequences of protein A. *The Journal of clinical investigation* **119**, 1931-1939 (2009);
published online EpubJul (
35. V. Castiglia, A. Piersigilli, F. Ebner, M. Janos, O. Goldmann, U. Dambock, A. Kroger, S. Weiss, S.
Knapp, A. M. Jamieson, C. Kirschning, U. Kalinke, B. Strobl, M. Muller, D. Stoiber, S. Lienenklaus,
P. Kovarik, Type I Interferon Signaling Prevents IL-1beta-Driven Lethal Systemic
Hyperinflammation during Invasive Bacterial Infection of Soft Tissue. *Cell host & microbe* **19**,
375-387 (2016); published online EpubMar 09 (10.1016/j.chom.2016.02.003).
36. T. Watanabe, N. Asano, S. Fichtner-Feigl, P. L. Gorelick, Y. Tsuji, Y. Matsumoto, T. Chiba, I. J. Fuss,
A. Kitani, W. Strober, NOD1 contributes to mouse host defense against Helicobacter pylori via
induction of type I IFN and activation of the ISGF3 signaling pathway. *The Journal of clinical
investigation* **120**, 1645-1662 (2010); published online EpubMay (10.1172/JCI39481).
37. K. Hansen, T. Prabakaran, A. Laustsen, S. E. Jorgensen, S. H. Rahbaek, S. B. Jensen, R. Nielsen, J.
H. Leber, T. Decker, K. A. Horan, M. R. Jakobsen, S. R. Paludan, Listeria monocytogenes induces
IFNbeta expression through an IFI16-, cGAS- and STING-dependent pathway. *The EMBO journal*
33, 1654-1666 (2014); published online EpubAug 01 (10.15252/embj.201488029).
38. Z. Abdullah, M. Schlee, S. Roth, M. A. Mraheil, W. Barchet, J. Bottcher, T. Hain, S. Geiger, Y.
Hayakawa, J. H. Fritz, F. Civril, K. P. Hopfner, C. Kurts, J. Ruland, G. Hartmann, T. Chakraborty, P.
A. Knolle, RIG-I detects infection with live Listeria by sensing secreted bacterial nucleic acids. *The
EMBO journal* **31**, 4153-4164 (2012); published online EpubNov 05 (10.1038/emboj.2012.274).
39. G. M. Boxx, G. Cheng, The Roles of Type I Interferon in Bacterial Infection. *Cell host & microbe*
19, 760-769 (2016); published online EpubJun 08 (10.1016/j.chom.2016.05.016).
40. M. Blanc, W. Y. Hsieh, K. A. Robertson, K. A. Kropp, T. Forster, G. Shui, P. Lacaze, S. Watterson, S.
J. Griffiths, N. J. Spann, A. Meljon, S. Talbot, K. Krishnan, D. F. Covey, M. R. Wenk, M. Craigon, Z.
Ruzsics, J. Haas, A. Angulo, W. J. Griffiths, C. K. Glass, Y. Wang, P. Ghazal, The transcription factor
STAT-1 couples macrophage synthesis of 25-hydroxycholesterol to the interferon antiviral
response. *Immunity* **38**, 106-118 (2013); published online EpubJan 24
(10.1016/j.immuni.2012.11.004

S1074-7613(12)00509-2 [pii]).
41. U. Diczfalusy, K. E. Olofsson, A. M. Carlsson, M. Gong, D. T. Golenbock, O. Rooyackers, U. Flaring,
H. Bjorkbacka, Marked upregulation of cholesterol 25-hydroxylase expression by
lipopolysaccharide. *J Lipid Res* **50**, 2258-2264 (2009); published online EpubNov
(10.1194/jlr.M900107-JLR200

M900107-JLR200 [pii]).

42.   C. Wilkins, M. Gale, Jr., Sterol-izing innate immunity. *Immunity* **38**, 3-5 (2013); published online
EpubJan 24 (10.1016/j.immuni.2013.01.002

S1074-7613(13)00006-X [pii]).
43.   M. Zimmermann, M. Kogadeeva, M. Gengenbacher, G. McEwen, H. J. Mollenkopf, N. Zamboni,
S. H. E. Kaufmann, U. Sauer, Integration of Metabolomics and Transcriptomics Reveals a
Complex Diet of Mycobacterium tuberculosis during Early Macrophage Infection. *mSystems* **2**,
(2017); published online EpubJul-Aug (e00057-17 [pii]

10.1128/mSystems.00057-17

mSystems00057-17 [pii]).
44.   X. Wang, T. Majumdar, P. Kessler, E. Ozhegov, Y. Zhang, S. Chattopadhyay, S. Barik, G. C. Sen,
STING Requires the Adaptor TRIF to Trigger Innate Immune Responses to Microbial Infection.
Cell host & microbe **20**, 329-341 (2016); published online EpubSep 14 (S1931-3128(16)30346-8
[pii]

10.1016/j.chom.2016.08.002).
45.   A. A. Herskovits, V. Auerbuch, D. A. Portnoy, Bacterial ligands generated in a phagosome are
targets of the cytosolic innate immune system. *PLoS pathogens* **3**, e51 (2007); published online
EpubMar (06-PLPA-RA-0531R2 [pii]

10.1371/journal.ppat.0030051).
46.   W. K. Ip, A. Sokolovska, G. M. Charriere, L. Boyer, S. Dejardin, M. P. Cappillino, L. M. Yantosca, K.
Takahashi, K. J. Moore, A. Lacy-Hulbert, L. M. Stuart, Phagocytosis and phagosome acidification
are required for pathogen processing and MyD88-dependent responses to Staphylococcus
aureus. *Journal of immunology* **184**, 7071-7081 (2010); published online EpubJun 15
(10.4049/jimmunol.1000110

jimmunol.1000110 [pii]).
47.   J. Gonzalo Asensio, C. Maia, N. L. Ferrer, N. Barilone, F. Laval, C. Y. Soto, N. Winter, M. Daffe, B.
Gicquel, C. Martin, M. Jackson, The virulence-associated two-component PhoP-PhoR system
controls the biosynthesis of polyketide-derived lipids in Mycobacterium tuberculosis. *The*
Journal of biological chemistry **281**, 1313-1316 (2006); published online EpubJan 20
(C500388200 [pii]

10.1074/jbc.C500388200).
48.   S. B. Walters, E. Dubnau, I. Kolesnikova, F. Laval, M. Daffe, I. Smith, The Mycobacterium
tuberculosis PhoPR two-component system regulates genes essential for virulence and complex
lipid biosynthesis. *Mol Microbiol* **60**, 312-330 (2006); published online EpubApr (MMI5102 [pii]

10.1111/j.1365-2958.2006.05102.x).
49.   P. Arumugam, D. Shankaran, A. Bothra, S. Gandotra, V. Rao, The MmpS6-MmpL6 Operon Is an
Oxidative Stress Response System Providing Selective Advantage to Mycobacterium tuberculosis
in Stress. *J Infect Dis* **219**, 459-469 (2019); published online EpubJan 9 (10.1093/infdis/jiy526

5092676 [pii]).
50.   A. Bothra, P. Arumugam, V. Panchal, D. Menon, S. Srivastava, D. Shankaran, A. Nandy, N.
Jaisinghani, A. Singh, R. S. Gokhale, S. Gandotra, V. Rao, Phospholipid homeostasis, membrane
tenacity and survival of Mtb in lipid rich conditions is determined by MmpL11 function. *Scientific*
reports **8**, 8317 (2018); published online EpubMay 29 (10.1038/s41598-018-26710-z

10.1038/s41598-018-26710-z [pii]).
51.   R. A. Slayden, C. E. Barry, 3rd, Analysis of the Lipids of Mycobacterium tuberculosis. *Methods*
Mol Med **54**, 229-245 (2001)10.1385/1-59259-147-7:229).

52.	V. Rao, N. Fujiwara, S. A. Porcelli, M. S. Glickman, Mycobacterium tuberculosis controls host
innate immune activation through cyclopropane modification of a glycolipid effector molecule. *J*
Exp Med **201**, 535-543 (2005); published online EpubFeb 21 (jem.20041668 [pii]
10.1084/jem.20041668).

Acknowledgements: The authors thank CSIR (VR-BSC0123), and DBT (GAP0096) funding

agency for supporting the study. The authors thank Mr. Manish Kumar and CSIR- BSC0403 for

the confocal microscopy facility. The Transmission Electron microscopy (TEM) and mass

spectrometry facilities are duly acknowledged. The institute BSL3 facility project (STS0016) is

duly acknowledged. The student fellowships from CSIR and DBT India are acknowledged. The

support provided by Academy of Scientific and Innovative research (AcSIR) is duly

acknowledged.

659
660	**Author Contributions**: DS, PA, AB, SG, VR were involved in conceptualizing and design of the

661	work, the work was performed by DS, PA and AB. VR, PA and DS were involved in the

662	manuscript preparation and proof reading.

Supplementary figures:

Fig S1: Relative luminescence in THP1 macrophages infected with H37Ra and H37Rv at 24h *p.i.* Values

represent average ratios $\pm$ SE of RLU in infected and control culture supernatants from triplicate wells of

two independent experiments [N=2].

Fig S2: Expression of genes involved in cholesterol biosynthesis in macrophages infected with N24

or N73 at 6h and 24h (shaded) estimated by qRTPCR. The relative expression values $\pm$ SE (with

respect to GAPDH) in triplicate assays of (N=2/3) is shown.

Fig.S3

Fig S3: Analysis of factors responsible for type I IFN induction in macrophages following Mtb infection:

A) ESAT6 expression in culture supernatant and lysates of Mtb N24 and N73. Expression was detected by immunoblotting with HYB76-8 monoclonal antibody.

B) Quantitation of mitochondrial ROS by mitosox staining following infection with mycobacteria or treatment of Antimycin A (+ control) at 6h and 24h.

Fig.S4

Fig S4: Mechanism of type I IFN induction in Mtb infected macrophages

A) Relative luminescence in RAW WT and KO macrophages infected with L3 and L1 Mtb at 24h post infection.

B) Dose dependent IRF induction by Mtb crude extracts at 6h.

Fig S5: Evaluation of a putative role for Mtb sulfolipids in IFN induction

  A) Relative luminescence in THP1 macrophages infected with Wt and MmpL8 mutant strains of Erdman at 24h p.i.