STELAR: A statistically consistent coalescent-based species tree estimation method by maximizing triplet consistency

Mazharul Islam[1], Kowshika Sarker[1], Trisha Das[1], Rezwana Reaz[2] and Md. Shamsuzzoha Bayzid[1,*]

[1]Department of Computer Science and Engineering
Bangladesh University of Engineering and Technology
Dhaka-1205, Bangladesh
[2]Department of Computer Science
The University of Texas at Austin
Texas 78712, USA
[*]Corresponding author

March 31, 2019

Abstract

Background: Species tree estimation is frequently based on phylogenomic approaches that use multiple genes from throughout the genome. However, estimating a species tree from a collection of gene trees can be complicated due to the presence of gene tree incongruence resulting from incomplete lineage sorting (ILS), which is modelled by the multi-species coalescent process. Maximum likelihood and Bayesian MCMC methods can potentially result in accurate trees, but they do not scale well to large datasets.

Results: We present STELAR (Species Tree Estimation by maximizing tripLet AgReement), a new fast and highly accurate statistically consistent coalescent-based method for estimating species trees from a collection of gene trees. We formalized the constrained triplet consensus (CTC) problem and showed that the solution to the CTC problem is a statistically consistent estimate of the species tree under the multi-species coalescent (MSC) model. STELAR is an efficient dynamic programming based solution to the CTC problem which is very fast and highly accurate. We evaluated the accuracy of STELAR in comparison with MP-EST and ASTRAL – two of the most popular and accurate coalescent-based methods. Experimental results suggest that STELAR matches the accuracy of ASTRAL and improves on MP-EST.

Conclusions: Theoretical and empirical results (on both simulated and real biological datasets) suggest that STELAR is a valuable technique for species tree estimation from gene tree distributions.

Availability: STELAR is freely available at https://islamazhar.github.io/STELAR/.

1 Introduction

Estimated species trees are useful in many biological analyses, but accurate estimation of species trees can be quite complicated. Species tree inference can potentially result in accurate evolutionary history using data from multiple loci. Therefore, due to the advent of modern sequencing technologies, it is increasingly common to infer trees by analyzing sequences from multiple loci. However, combining multi-locus data is difficult, especially in the presence of gene tree discordance. The traditional approach to species tree estimation is called concatenation (also known as combined analysis), which concatenates gene sequence alignments into a supergene matrix, and then estimates the species tree using a sequence based tree estimation technique (e.g., maximum parsimony, maximum likelihood etc.). Although it has been used in many biological analyses, concatenation can be problematic as it is agnostic to the topological differences among the gene trees, can be statistically inconsistent [1], and can return incorrect trees with high confidence [2, 3, 4, 5].

Recent modeling and computational advances have produced methods that explicitly take the gene tree discordance into account while combining multi-locus data to estimate species trees. Incomplete lineage sorting (ILS), also known as deep coalescence, is one of the most prevalent reasons for gene tree incongruence [6]. Rapid radiation is known to happen in many organisms (for example, birds) where ILS is likely to occur [7]. Due to this growing awareness that ILS can be present in many phylogenomic datasets, ILS-aware species tree estimation techniques have gained substantial attention from systematists. These types of method are usually referred to as "summary methods" as they summarize a set of gene trees to estimate the species trees.

Several species-tree estimation methods have been proven to be statistically consistent under the multi-species coalescent model, meaning that they will return the true species tree with high probability given a sufficiently large number of true gene trees sampled from the distribution defined by the species tree. Methods that are statistically consistent under the MSC include ASTRAL [8], MP-EST [9], *BEAST [10], NJst [11], BUCKy [12], GLASS [13], STEM [14], SNAPP [15], SVDquartets [16], STEAC [17] and ASTRID [18]. *BEAST, which is a Bayesian technique, can co-estimate both gene trees and species trees, and simulation studies suggest that *BEAST can be considered the best of the co-estimation methods with excellent accuracy, but it is computationally very intensive to run on even moderately-sized dataset [19, 20]. ASTRAL and MP-EST are two of the most accurate and widely used summary methods that are much faster than *BEAST. ASTRAL has been shown to be faster and more accurate than MP-EST and can handle large dataset containing hundreds of species which is not possible by MP-EST [8, 21].

ASTRAL finds a species tree that maximizes the number of consistent quartets between the species tree and the gene trees. MP-EST maximizes a pseudo-likelihood estimate utilizing the underlying triplet distribution of the gene trees. Quartet and triplet based methods are robust to the "anomaly zone" [22, 23] (a condition where the most probable gene tree topology may not be identical to the species tree topology) as there are no anomalous rooted three-taxon species trees [24] and no anomalous unrooted four-taxon species trees [25, 23].

We present STELAR (Species Tree Estimation by maximizing tripLet AgReement), a new colescent-based method which finds a species tree that agrees with the largest number

of triplets induced by the gene trees. STELAR is statistically consistent, fast (having a polynoimial running time), and highly accurate – enabling genome wide phylogenomic analyses.

2 Approach

Design of ASTRAL and other quartet-based methods [8, 26] and their proofs of being statistically consistent is based upon the fact that unrooted 4-taxon species trees do not contain anomaly zone [25, 27]. We use similar design in STELAR, utilizing the fact that rooted 3-taxon species trees do not contain anomaly zone [28].

2.1 Definitions and notations

Let T be a rooted tree, leaf-labelled by species set $\mathcal{X}$ with n taxa. Each internal node u in a tree T divides the set of taxa present in the subtree rooted at u into two distinct partitions. We call this a *subtree-bipartition* which was originally defined in [29]. We denote by T_u the subtree under node u of tree T. $L(T)$ denotes the leaves in T. r denotes an arbitrary set of three species $\{a, b, c\} \subset \mathcal{X}$, t_r is a rooted tree topology on r. We use $a|bc$ to show that b and c are siblings. Possible three topologies on $\{a, b, c\}$ are: $a|bc, b|ca, c|ab$. The triplet tree topology that appears more frequently than the two alternative topologies is called the *dominant* topology. $\mathcal{TR}(T)$ denotes the set of $\binom{n}{3}$ triplet topologies induced by the tree T. We denote by $T|r$ the triplet tree topology obtained by restricting T to the three species in r.

2.2 Problem definition

A natural optimization problem for estimating a species tree from a collection of gene trees under triplet consistency would be first decomposing the input gene trees into their induced triplets, and then estimating a species tree by amalgamating the dominant triplets. Building such a consensus tree from a set of triplets has been shown to be an NP-hard problem, and exact and heuristic algorithms have been proposed [30]. This approach considers only the dominant triplets which could be problematic when the frequencies of the alternate triplet topologies are very close to that of the dominant one, and when all the dominant triplets are not compatible with each other meaning that all of them do not agree with a single species tree. Moreover, this approach has to explicitly enumerate and consider $3 * \binom{n}{3}$ possible triplets, making it computationally expensive for larger values of n. An alternate approach (perhaps a better one) would be to consider the relative frequencies of all the triplets, and infer a species tree that agrees with the largest number of triplets induced by the gene trees. Methods that do not explicitly decompose the gene trees into induced triplets and quartets are preferred over the ones that take a set of quartets or triplets as input, since the latter types of methods demand high computational time and memory and thus are not scalable to large dataset [26, 8]. This is possibly one of the reasons why quartet amalgamation method like QFM [26], despite having excellent accuracy, is not as popular as ASTRAL. Taking these issues into consideration, we introduce a constrained version of the triplet consistency problem,

which we call the *Constrained Triplet Consensus* (CTC) problem, and formalize it as follows.

- **Input**: a set $\mathcal{G}$ of rooted gene trees on $\mathcal{X}$ and a set $\mathcal{SBP}$ of subtree-bipartitions.

- **Output**: a species tree ST on $\mathcal{X}$ that maximizes the triplet agreement with $\mathcal{G}$ and all its subtree-bipartitions are in $\mathcal{SBP}$.

Note that when $\mathcal{SBP}$ is set to all possible subtree-bipartitions on $\mathcal{X}$, the triplet consistency problem will find the globally optimal solution (exact solution). Otherwise, when $\mathcal{SBP}$ contains only the subtree-bipartitions in the input gene trees, the search space will include only those species trees where every subtree-bipartition is present in at least one of the gene trees in $\mathcal{G}$, and we call this the constrained version or heuristic version.

Theorem 2.1. *Given a set $\mathcal{G}$ of true gene trees, solution to the CTC problem (both exact and constrained version) is a statistically consistent estimator of the species tree topology under the MSC model.*

Proof. Let $\mathcal{G}$ be a set of sufficiently large number of true gene trees that has evolved within the model species tree T under the multi-species coalescent model. We know that a rooted 3-taxon species tree does not have any anomaly zone [28]. So, as we increase the number of gene trees, each triplet topology induced by the species tree will have more frequency in $\mathcal{G}$ than its alternatives, with probability that approaches 1. Let $w_{\mathcal{G}}(t_r)$ be the number of trees in $\mathcal{G}$ that agrees with t_r. Hence, for every triplet r and every possible tree T', $w_{\mathcal{G}}(T|r) \geq w_{\mathcal{G}}(T'|r)$ with probability that approaches 1 for a large enough number of gene trees. Therefore, if $\mathcal{TR}$ is the set of all possible triplets on $\mathcal{X}$, then

$$\sum_{r \epsilon \mathcal{TR}} w_{\mathcal{G}}(T|r) \geq \sum_{r \epsilon \mathcal{TR}} w_{\mathcal{G}}(T'|r).$$

By definition, the exact solution to the CTC problem will find a tree that will maximize the triplet agreement with $\mathcal{G}$. Therefore, for a sufficiently large number of true gene trees, the optimal tree under CTC will be identical to the true species tree T with probability approaching one.

We now show that the solution to the constrained version is also statistically consistent. If we increase the number of gene trees, with probability approaching 1, at least one of the gene tree topologies in $\mathcal{G}$ will be identical to the the true species tree topology. Therefore, with probability that tends to 1, the set $\mathcal{SBP}$ will contain all the subtree bipartitions from the true species tree, and therefore the solution to the CTC problem will be identical to the true species tree T.

$\square$

2.3   Algorithmic design of STELAR

We propose a dynamic programming (DP) solution to the CTC problem. The algorithmic design (and corresponding theoretical results) in STELAR is structurally similar to

ASTRAL. This sort of DP-based approach, which implicitly finds a maximum or minimum clique in a graph modelled from the input gene trees, was first used by Hallett and Lagergren and later was used in Phylonet [31, 32], DynaDup [29, 33] and ASTRAL. The key idea in STELAR is to find an equation for computing the number of triplets in the gene trees that agree with a given subtree in the species tree, which ultimately enables us to design a dynamic programming algorithm for estimating a species tree by maximizing the triplet agreement.

We call a triplet $tr = a|bc$ to be mapped to a subtree-bipartition at an internal node x in T $(SBP(T_x) = X_1|X_2)$ when $\{a, b, c\} \subseteq L(T_x)$, and the topology of tr is compatible (consistent) with T_u (see Fig. 1). Each triplet mapped to a subtree-bipartition $(X_1|X_2)$ will have either two leaves from X_1 and one leaf from X_2 or two leaves from X_2 and one leaf from X_1. Therefore, we have the following results.

Figure 1: **Mapping of a triplet to a subtree-bipartition.** Each internal node x in a rooted tree defines a subtree bipartition $(X_1|X_2)$. Each induced triplet in T maps to a subtree-bipartition in T. This figure shows the mapping of a triplet $a|bc$ to a subtree-bipartition $x = X_1|X_2$.

Lemma 2.2. *Total number of triplets mapped to a subtree-bipartition $x = (X_1|X_2)$ of a rooted binary tree T, where $|X_1| = n_1$ and $|X_2| = n_2$ is:*

$$\mathcal{NT}(n_1, n_2) = \binom{n_1}{2}\binom{n_2}{1} + \binom{n_2}{2}\binom{n_1}{1} = \frac{n_1 n_2 (n_1 + n_2 - 2)}{2} \tag{1}$$

Lemma 2.3. *For two subtree-bipartitions $x = (X_1|X_2)$ and $y = (Y_1|Y_2)$ where $(X_1 \cup X_2) \subseteq \mathcal{X}$ and $(Y_1 \cup Y_2) \subseteq \mathcal{X}$, number of triplets mapped to both x and y is given by:*

$$\mathcal{M}(x, y) = \mathcal{NT}(|X_1 \cap Y_1|, |X_2 \cap Y_2|) + \mathcal{NT}(|X_1 \cap Y_2|, |X_2 \cap Y_1|). \tag{2}$$

Proof. We can obtain two possible subtree bipartitions by taking pairwise intersection: $z_1 = (X_1 \cap Y_1 | X_2 \cap Y_2)$ and $z_2 = (X_1 \cap Y_2 | X_2 \cap Y_1)$. Note that each leaf in z_i $(i \in 1, 2)$ is also a leaf in both x and y, and so each triplet mapped to z_i is also mapped to x and y and the number of mapped triplets can be computed by Eqn. 1.

We now prove that Eqn. 2 counts a mapped triplet only once. Let $tr = ab|c$ is mapped to both x and y where $a, b \in X_1$, $c \in X_2$, $a, b \in Y_1$ and $c \in Y_2$. Eqn. 2 will count tr for $z_1 = (X_1 \cap Y_1 | X_2 \cap Y_2)$ only as z_2 will have no element from $\{a, b, c\}$. Thus the lemma follows. □

We now see how to compute the triplet consistency (TC) score of a species tree ST with respect to a set $\mathcal{G}$ of gene trees, which we denote by $TC_{\mathcal{G}}(ST)$. If $\mathcal{SBP}(T)$ is the set of subtree-bipartitions in T, for a subtree-bipartition x and a set of rooted binary trees $\mathcal{G}$, total number of triplets in $\mathcal{G}$ that are mapped to x is given by:

$$TC_{\mathcal{G}}(x) = \sum_{gt \in \mathcal{G}} \sum_{y \in \mathcal{SBP}(gt)} \mathcal{M}(x, y). \tag{3}$$

Thus, the TC score of an estimated species tree ST is obtained by,

$$TC_{\mathcal{G}}(ST) = \sum_{x \in \mathcal{SBP}(ST)} TC_{\mathcal{G}}(x) \tag{4}$$

Therefore, we can design a dynamic programming approach which starts with the whole species set $\mathcal{X}$ and recursively divides it into smaller subsets and picks the partition that maximizes the triplet consistency. At each level, we compute $V(A)$ which gives the score of an optimal subtree on leaf set $A \subseteq \mathcal{X}$. At the end level, we have computed $V(\mathcal{X})$ and sufficient information so that backtracking yields the corresponding subtree-bipartitions in the optimal species tree.

2.3.1 Dynamic Programming

Base case: $|A| = 1; V(A) = 0$
Recursive relation: $V(A) = \max_{(A'|A-A') \in \mathcal{X}} \{V(A') + V(A - A') + TC_{\mathcal{G}}(A'|A - A')\}$, where $A'|A - A' \in \mathcal{SBP}$.

2.3.2 Running time analysis

Lemma 2.4. *Given a set $\mathcal{G}$ of k gene trees on n taxa and a subtree-bipartition x, computing the triplet consistency score $TC_{\mathcal{G}}(x)$ takes $O(n^2 k)$ time.*

Proof. A gene tree with n taxa contains $O(n)$ subtree-bipartitions. Thus, for a set of k input gene trees, there will be $O(nk)$ subtree-bipartitions. To find the triplet consistency score $TC_{\mathcal{G}}(x)$ of a subtree-bipartition x using Equation 3, we need to score $\mathcal{M}(x, y)$ for all subtree-bipartions y present in the input gene trees. Each subtree-bipartion is represented by a bitset, hence calculating $\mathcal{M}(x, y)$ using Equation 2 takes $O(n)$ time. Since there are $O(nk)$ subtree-bipartions in $\mathcal{G}$, calculating $TC_{\mathcal{G}}(x)$ takes $O(n^2 k)$ time. □

Theorem 2.5. *For a given set $\mathcal{G}$ of k gene trees on n taxa and a set $\mathcal{SBP}$ of subtree-bipartitions, STELAR runs in $O(n^2 k |\mathcal{SBP}|^2)$ time.*

Proof. For a specific cluster A, $V(A)$ can be computed in $O(|\mathcal{SBP}| * n^2 k)$ time since at worst we need to look at every subtree-bipartition in $\mathcal{SBP}$, and computing $TC_{\mathcal{G}}(x)$

takes $O(n^2k)$ time. Note that this is a conservative analysis since the number of subtree-bipartitions that the DP algorithm needs to consider for computing $V(A)$ for a specific cluster A is, in reality, much less than $|\mathcal{SBP}|$ (especially for A where $|A| < |\mathcal{X}|$). There will be $O(|\mathcal{SBP}|)$ possible clusters for which we need to compute $V(A)$. Thus, the running time of STELAR is $O(n^2k|\mathcal{SBP}|^2)$.

$\square$

The running time of STELAR is mostly dependent on the number of subtree-bipartions considered by the DP algorithm. In the exact version, we consider all possible subtree-bipartions. Hence $|\mathcal{SBP}|$ and the running time of STELAR grows exponentially as we increase the number of taxa. In the heuristic version, we only consider $O(nk)$ subtree bipartions that are present in the input gene trees. Hence, from Thm. 2.5, heuristic version has a running time of $O(n^4k^3)$.

3   Experimental Studies

We compared STELAR with ASTRAL and MP-EST, two of the most widely used and accurate coalescent-based methods. ASTRAL has already been compared and shown to outperform many of the existing species tree estimation methods (e.g., BUCKy-pop, MRP, greedy consensus, CA-ML, etc.). So we did not include those methods in this study. We ran MP-EST with 10 random starting points and selected the species tree with the highest pseudo likelihood value. We used previously studied simulated and biological datasets to evaluate the performance of these methods. We compare the estimated species trees to the model species tree (for the simulated datasets) or to the scientific literature (for the biological datasets), to evaluate accuracy. We used False Negative (FN) rate to measure the tree error since all the trees estimated in this study are binary and so false negative and false positive rates are identical. We performed Wilcoxon signed-rank test (with $\alpha = 0.05$) to measure the statistical significance of the differences between two methods.

3.1   Datasets

We studied three collections of simulated datasets: one based on a biological dataset (37-taxon mammalian dataset) that was generated and subsequently analyzed in prior studies [8, 34, 21], and two other simulated datasets from [35, 34]. Table 1 presents a summary of these datasets, showing various model conditions with varying numbers of taxa (11 to 37), ILS levels (reflected in the average topological distance between true gene trees and true species tree) and gene sequence lengths. All datasets consist of gene sequence alignments generated under a multi-stage simulation process that begins with a species tree, simulates gene trees down the species tree under the multi-species coalescent model (and so can differ topologically from the species tree), and then simulates gene sequence alignments down the gene trees.

Mammalian dataset was simulated by taking the species tree estimated by MP-EST on the biological dataset studied in [36]. This species tree had branch lengths in coalescent units, which we used to produce a set of gene trees under the coalescent model. Thus, the mammalian simulation model tree has an ILS level based on a coalescent analysis of

Table 1: Properties of the simulated datasets. Level of ILS is presented in terms of the average topological distance between true gene trees and true species tree.

Dataset	ILS level	# genes	# sites	# Replicates	Ref.
11-taxon	85%	5 - 100	2000	20	[35]
15-taxon	82%	100 - 1000	100 - 1000	10	[34]
37-taxon	18%, 32%, 54%	25 - 800	250 -1000	20	[34]

the biological mammalian dataset, and other properties of the simulation that are set to reflect the biological sequences Song *et al.* studied. We explored the impact of varying amounts of ILS, varying numbers of genes ($25 \sim 800$), the impact of phylogenetic signal by varying the sequence length (250, 500, and 1000 base pair (bp)) for the markers. Three levels of ILS were simulated. The basic model species tree has branch lengths in coalescent units, and we produced other model species trees by multiplying or dividing all internal branch lengths in the model species tree by two. This rescaling varies the amount of ILS (shorter branches have more ILS), and also impacts the amount of gene tree estimation error in the estimated gene trees. The basic model condition with moderate amount of ILS is referred to as 1X and the model conditions with higher and lower amounts of ILS are denoted by 0.5X and 2X, respectively.

We used two other simulated datasets: 11-taxon dataset (generated by [35] and subsequently studied by [20, 37]), and 15-taxon datasets (generated and studied by [34]). For 11-taxon dataset, we analyzed only the model condition with high amount of ILS as the model condition with lower amount of ILS is very easy to analyze and even methods without any statistical guarantee produced highly accurate tree with just $25 \sim 50$ gene trees [20]. 15-taxon datasets vary in sequence lengths and has high amount of ILS.

We used two biological datasets: the 37-taxon mammalian datasets studied by Song *et al.* [36] with 424 genes, and the Amniota dataset from [38] containing 16 species and 248 genes.

4 Results

4.1 Results on mammalian simulated dataset

We analyzed the performance of ASTRAL, MP-EST and STELAR on various model conditions with varying amounts of ILS, numbers of genes and lengths of the sequences. Figure 2(a) shows the average FN rates of three methods on three model conditions with varying amounts of ILS. For all methods, FN rates increase as we increase the amount of ILS. STELAR and ASTRAL are better than MP-EST on all amounts of ILS and, in some cases, the differences are statistically significant ($P < 0.05$). STELAR and ASTRAL are comparable in terms of tree accuracy and no statistically significant difference was noticed.

Similar trends, in terms of relative performance, were observed when we varied the lengths of the sequences (Fig. 2(b)). As we increase the gene sequence lengths, the estimated gene trees become more accurate which results into lower FN rates of the trees estimated by STELAR, ASTRAL and MP-EST. ASTRAL and STELAR are better than MP-EST (in some cases the improvements are statistically significant). The lowest

Figure 2: **Comparison of ASTRAL, STELAR and MP-EST on 37-taxon simulated mammalian dataset.** We show the average FN rates with standard error bars over 20 replicates. (a) We fixed the sequence lenght to 500bp and number of genes to 200, and varied the amounts of ILS. 2X model condition contains the lowest amount of ILS while 0.5X refers to the model condition with the highest amount of ILS. (b) We varied the amount of gene tree estimation error by varying the sequence lengths from 250 to 1000 bp, while keeping the ILS level (moderate) and the number of genes (200) fixed. (c) We fixed the sequence length to 500bp and amount of ILS to 1X, and varied the numbers of genes (25, 50, 100, 200, 400, 800).

amount of error was obtained for true gene trees. STELAR matched the accuracy of ASTRAL.

Figure 2(c) shows the impact of the numbers of gene trees on the accuracy of the estimated species trees. All methods improved with increasing numbers of genes (this is expected as these methods are statistically significant and thus increasing the numbers of genes has resulted into improved topological accuracy). STELAR and ASTRAL achieved similar accuracy and performed significantly better than $MP_EST across all model conditions$.

4.2   Results on 11-taxon dataset

11-taxon dataset was simulated under a complex process to ensure substantial heterogeneity between genes and to deviate from the molecular clock [35]. We analyzed both the estimated and true gene trees. In both cases, we varied the number of genes from (5 $\sim$ 100). Figure 3 shows the results on 11-taxon dataset for both estimated and true gene trees. As expected, all methods improved with increased numbers of genes, and had their best accuracy on true gene trees. The accuracy of these three methods are almost similar with no statistically significant difference. On true gene trees, these methods recovered the true species trees across all the replicates with 50 or higher numbers of genes. Unlike mammalian dataset, MP-EST was slightly better than ASTRAL and STELAR on a couple of model conditions (albeit the differences are not statistically significant with P-values greater than 0.05).

Figure 3: **Average FN rates of ASTRAL, MP-EST and STELAR on the 11-taxon dataset with high amount of ILS.** We varied the numbers of genes from 5 to 100. We analyzed both the estimated and true gene trees. We show the average FN rates with standard error bars over 20 replicates.

4.3   Results on 15-taxon dataset

15-taxon dataset evolved under a caterpillar model species tree with very high level of ILS. We explored the performance on varying sequence lengths (100bp and 1000bp), and numbers of genes (100 and 1000). ASTRAL, STELAR and MP-EST improved as the the number of genes and sequence length increased, and had their best accuracy (FN rate = 0) on true gene trees (Fig. 4). On 100bp alignments, ASTRAL was better than STELAR and MP-EST and the difference was significant with 1000 genes ($P < 0.05$). On 1000bp alignments, no statistically significant difference was observed between STELAR and ASTRAL. MP-EST consistently had the highest error rates across all the model conditions on estimated gene trees. On true gene trees with 1000 genes, all the methods recovered the true species tree across all the replicates. With 100 genes, ASTRAL and STELAR reconstructed the true species tree on 9 replicates (out of 10), and MP-EST recovered the true species tree on 8 replicates.

4.4   Results on Biological datasets

Amniota dataset: We analyzed the Amniota dataset from Chiari *et al.* [38] containing 248 genes across 16 amniota taxa in order to resolve the position of turtles within the amniotes, especially relative to birds and crocodiles. We re-analyzed both the amino acid (AA) and DNA gene trees, obtained from the authors, using ASTRAL, MP-EST and STELAR. The sister relationship of crocodiles and birds (forming Archosaurs), and placement of the turtles as the sister clade to Archosaurs are considered reliable and were supported by previous studies [39, 40, 41].

For amino acid dataset, all the methods placed birds and crocodiles as sister groups (forming archosaurs) and turtles as sister to archosaurs clade. However, they differed in the placement of basal groups and resolution within the turtles. ASTRAL misplaced *Podarcis* as the basal group and thus did not form the Squamates. STELAR and MP-EST formed Squamates and rightly placed *Protopterus annectens* as the basal node [38]. We estimated quartet scores and triplet scores of these different tree topologies with respect to the corresponding gene trees (see Table 2). STELAR estimated tree achieved the highest triplet score, and interestingly it achieved the highest quartet score as well.

STELAR and MP-EST on the DNA gene trees produced the same tree which resolved bird/turtle/crocodile relationship as (birds,(turtles,crocodiles)) and thus did not form Archosaurs. However, ASTRAL on the DNA gene trees resolved the relationships as (birds,(turtles,crocodiles)). Therefore, maximizing triplet agreement and quartet agreement both recovered the (birds,(turtles,crocodiles)) relationships on AA gene trees, but differed on the DNA gene trees. MP-EST and STELAR rightly placed *Protopterus annectens* as the outgroup while ASTRAL placed *Ornithorhynchus anatinus* as the basal node. Moreover, ASTRAL did not recover the ((*Homo sapiens*, *Monodelphis domestica*), *Ornithorhynchus anatinus*)) clade which was supported by previous studies [38] and was also produced by STELAR and MP-EST. We note that previous study of this data using ASTRAL (with multi-locus bootsrapping (MLBS) [42] using 200 bootstrap replicates instead of the maximum likelihood gene trees) reported a tree that is different in terms of the placement of the basal nodes and resolving the relationships among *Homo sapiens*, *Monodelphis domestica* and *Ornithorhynchus anatinus* [8]. As expected, ASTRAL

Figure 4: **Comparison of ASTRAL, MP-EST and STELAR on the 15-taxon dataset**. We varied the numbers of estimated gene trees (100 and 1000) and sequence lengths (100bp and 1000bp). We also anlyzed the true gene trees. We show the average FN rates with standard errors over 10 replicates.

achieved the highest quartet agreement score and STELAR and MP-EST achieved the highest triplet agreement score (Table 3).

Figure 5: **Analyses of the Amniota dataset (both DNA and AA gene trees) using ASTRAL, MP-EST and STELAR.** The branches that are not consistent with existing biological beliefs are shown in red.

Mammalian dataset: We re-analyzed mammalian dataset from Song *et al.* containing 447 genes across 37 mammals [36] after removing 21 mislabeled genes (confirmed by the authors), and two other outlier genes. ASTRAL, MP-EST and STELAR recovered the same topology as shown in Fig. 6. The placement of tree shrews (*Tupaia belangeri*) is of great importance here as previous studies found that the phylogenetic placement of tree shrews is unstable as they can be classified as a sister group to Primates or to Glires [36, 43, 44, 45]. However, our analyses using ASTRAL, MP-EST and STELAR discovered a tree that has placed tree shrews as sister to Glires (see Fig. 6). This is consistent to the tree estimated by CA-ML (reported in [36]). However, previous studies with MP-EST, using multi-locus bootstrapping [42], on the same dataset recovered a tree which placed tree shrews as the sister to Primates [45, 8]. We investigated the quartet and triplet scores of these two alternative hypotheses and found that the tree that suggests (tree shrew, Gliers) clade has higher quartet and triplet scores (25526915 and 3042547, respectively) compared to the one that suggests (tree shrew, Primates) relationship which yields 25511503 and 3041747 quartet and triplet scores respectively.

Tool	Triplet score	Quartet score
ASTRAL	25132	83604
MP-EST	38385	83532
STELAR	38394	83604

Table 2: **Triplet and quartet scores of the species trees estimated by ASTRAL, MP-EST and STELAR on the amino acid (AA) gene trees in Amniota dataset.**

Tool	Triplet score	Quartet score
ASTRAL	31162	97890
MP-EST	42448	97744
STELAR	42448	97744

Table 3: **Triplet and quartet scores of the species trees estimated by ASTRAL, MP-EST and STELAR on the DNA gene trees in Amniota dataset.**

Therefore, our analyses strongly support the placement of tree shrews as sister to Glires.

4.4.1   Running time

We ran the experiments on a linux machine with 3.7 GiB RAM and 4 processors. Our experiments show that STELAR and ASTRAL are substantially faster than MP-EST. The running time of MP-EST is more sensitive to the number of species than the number of genes, and grows rapidly as we increase the number of taxa [21]. On 37-taxon simulated dataset with 200 genes (moderate ILS), both ASTRAL and STELAR took around 4 seconds whereas MP-EST (with 10 random starting points) took 3300 seconds (55 minutes). Running time of all these methods increased with increasing amounts of ILS. On 37-taxon, 0.5X dataset (high ILS) with 200 genes, ASTRAL took around 10 seconds, STELAR took 15 seconds and MP-EST took 4800 seconds (80 minutes). This is because the numbers of distinct quartets and triplets in the gene trees increase as we increase the amount of discordance among the gene trees. When we increased the numbers of genes from 200 to 800 in 1X model condition, the running time of ASTRAL increased from 4 seconds to 30 seconds, whereas STELAR took only 17 seconds. Therefore, STELAR was the fastest method on 37-taxon, 800-gene dataset. MP-EST took about 55 minutes and thus was not affected by the increased number of genes. On 11-taxon dataset, we ran the exact versions of ASTRAL and STELAR, and yet the running time is much less than MP-EST. Exact versions of ASTRAL and STELAR finished in less than a second on $5 \sim 50$ genes and less than two seconds on $50 \sim 100$ genes, whereas the running times of MP-EST range from 13 to 20 seconds. For 15-taxon dataset, MP-EST took $90 \sim 100$ seconds. ASTRAL and STELAR finished in less than 2 seconds on 100 gene dataset, and for 1000 genes they took about 50 and 80 seconds, respectively.

5   Discussion

In this study, we presented STELAR – a method for estimating species trees from a set of estimated gene trees which is statistically consistent under the multi-species coalescent

Figure 6: **Analyses of the mammalian dataset using ASTRAL, MP-EST and STELAR.** All these methods generated the same tree topology.

model. We formalized the constrained triplet consensus (CTC) problem and showed that the solution to the CTC problem is a statistically consistent estimator of the species tree under the MSC model. STELAR is a dynamic programming solution to the CTC problem. STELAR has an exact version (which provides the exact solution and takes exponential amount of time) as well as an efficient polynomial time heuristic version. We compared STELAR with ASTRAL and MP-EST (two of the most accurate and widely used coalescent-based species tree estimation methods) on both simulated and real biological dataset.

Our experiments on Amniota and mammalian biological dataset suggest that STELAR can produce reliable trees on real biological dataset. STELAR produced more reliable trees than ASTRAL on the Amniota dataset with AA gene trees, and achieved the highest triplet and quartet agreement scores. On the mammalian dataset, STELAR, MP-EST and ASTRAL returned the same tree topology. Extensive experimental studies on simulated dataset with varying model conditions suggest that STELAR improved on MP-EST in most of the cases and matched the accuracy of ASTRAL. Thus, within the scope of the experiments conducted in this study, maximum triplet agreement has been shown to be a better optimization criterion than the triplet frequency based maximum pseudo-likelihood used in MP-EST, and as good as the maximum quartet agreement criterion used in ASTRAL. We believe this study represents significant contributions, and STELAR would be considered as a useful tool in phylogenomics.

STELAR can consistently estimate the species tree topology and is fast enough to handle large numbers of genes and species, making it suitable for large scale phylogenomic analyses. Our algorithm also allows missing sequences for some genes. However, STELAR can be expanded in several directions. Future work will need to investigate how the estimation of branch lengths (in coalescent unit) can be incorporated in STELAR. STELAR, at this stage, cannot handle unrooted and non-binary gene trees. Future works on formalizing the CTC problem for unrooted and non-binary gene trees and developing appropriate algorithms will be of great importance.

References

[1] Sebastien Roch and Mike Steel. Likelihood-based tree reconstruction on a concatenation of aligned sequence data sets can be statistically inconsistent. *Theoretical population biology*, 100:56–62, 2015.

[2] L S Kubatko and J H Degnan. Inconsistency of phylogenetic estimates from concatenated data under coalescence. *Syst Biol*, 56:17, 2007.

[3] Scott V Edwards, Liang Liu, and Dennis K Pearl. High-resolution species trees without concatenation. *Proceedings of the National Academy of Sciences*, 104(14):5936–5941, 2007.

[4] A D Leaché and B Rannala. The accuracy of species tree estimation under simulation: a comparison of methods. *Syst Biol*, 60(2):126–137, 2011.

[5] Michael DeGiorgio and James H Degnan. Fast and consistent estimation of species trees using supermatrix rooted triples. *Molecular biology and evolution*, 27(3):552–569, 2009.

[6] W. P. Maddison. Gene trees in species trees. *Systematic Biology*, 46:523–536, 1997.

[7] Erich D Jarvis, Siavash Mirarab, Andre J Aberer, Bo Li, Peter Houde, Cai Li, Simon YW Ho, Brant C Faircloth, Benoit Nabholz, Jason T Howard, et al. Whole-genome analyses resolve early branches in the tree of life of modern birds. *Science*, 346(6215):1320–1331, 2014.

[8] Siavash Mirarab, Rezwana Reaz, Md S Bayzid, Théo Zimmermann, M Shel Swenson, and Tandy Warnow. ASTRAL: genome-scale coalescent-based species tree estimation. *Bioinformatics*, 30(17):i541–i548, 2014.

[9] Liang Liu, Lili Yu, and Scott V Edwards. A maximum pseudo-likelihood approach for estimating species trees under the coalescent model. *BMC Evolutinary Biology*, 10:302, 2010.

[10] J Heled and A J Drummond. Bayesian inference of species trees from multilocus data. *Mol Biol Evol*, 27:570–580, 2010.

[11] Liang Liu and Lili Yu. Estimating species trees from unrooted gene trees. *Systematic Biology*, 60(5):661–667, 2011.

[12] B Larget, S K Kotha, C N Dewey, and C Ané. BUCKy: Gene tree/species tree reconciliation with the Bayesian concordance analysis. *Bioinf*, 26(22):2910–2911, 2010.

[13] E Mossel and S Roch. Incomplete lineage sorting: consistent phylogeny estimation from multiple loci. *IEEE Comp Biol Bioinform*, 7(1):166–171, 2011.

[14] L S Kubatko, B C Carstens, and L L Knowles. Stem: Species tree estimation using maximum likelihood for gene trees under coalescence. *Bioinf*, 25:971–973, 2009.

[15] David Bryant, Remco Bouckaert, Joseph Felsenstein, Noah A Rosenberg, and Arindam RoyChoudhury. Inferring species trees directly from biallelic genetic markers: bypassing gene trees in a full coalescent analysis. *Molecular biology and evolution*, 29(8):1917–1932, 2012.

[16] Julia Chifman and Laura Kubatko. Quartet inference from snp data under the coalescent model. *Bioinformatics*, 30(23):3317–3324, 2014.

[17] Liang Liu, Lili Yu, Dennis K Pearl, and Scott V Edwards. Estimating species phylogenies using coalescence times among sequences. *Systematic biology*, 58(5):468–477, 2009.

[18] Pranjal Vachaspati and Tandy Warnow. Astrid: accurate species trees from internode distances. *BMC genomics*, 16(10):S3, 2015.

[19] Brian Tilston Smith, Michael G Harvey, Brant C Faircloth, Travis C Glenn, and Robb T Brumfield. Target capture and massively parallel sequencing of ultraconserved elements for comparative studies at shallow evolutionary time scales. *Systematic biology*, 63(1):83–95, 2013.

[20] Md Shamsuzzoha Bayzid and Tandy Warnow. Naive binning improves phylogenomic analyses. *Bioinformatics*, 29(18):2277–2284, 2013.

[21] Md Shamsuzzoha Bayzid, Tyler Hunt, and Tandy Warnow. Disk covering methods improve phylogenomic analyses. *BMC genomics*, 15(6):S7, 2014.

[22] J H Degnan and N A Rosenberg. Discordance of species trees with their most likely gene trees. *PLoS Genetics*, 2:762 – 768, 2006.

[23] James H Degnan. Anomalous unrooted gene trees. *Systematic biology*, 62(4):574–590, 2013.

[24] J H Degnan and N A Rosenberg. Gene tree discordance, phylogenetic inference and the multispecies coalescent. *Trends Ecology Evolution*, 26(6), 2009.

[25] E S Allman, J H Degnan, and J A Rhodes. Identifying the rooted species tree from the distribution of unrooted gene trees under the coalescent. *J Math Biol*, 62(6):833–862, 2011.

[26] Rezwana Reaz, Md Shamsuzzoha Bayzid, and M Sohel Rahman. Accurate phylogenetic tree reconstruction from quartets: A heuristic approach. *PLoS One*, 9(8):e104008, 2014.

[27] J. H. Degnan. Anomalous unrooted gene trees. *Systematic Biology*, 62(4):574–590, 2013.

[28] J H Degnan and N A Rosenberg. Gene tree discordance, phylogenetic inference and the multispecies coalescent. *Trends Ecology Evolution*, 26(6), 2009.

[29] M. S. Bayzid, S. Mirarab, and T. Warnow. Inferring optimal species trees under gene duplication and loss. In *Proc. of Pacific Symposium on Biocomputing (PSB)*, volume 18, pages 250–261, 2013.

[30] Bang Ye Wu. Constructing the maximum consensus tree from rooted triples. *Journal of Combinatorial Optimization*, 8(1):29–39, 2004.

[31] C. V. Than and L. Nakhleh. Species tree inference by minimizing deep coalescences. *PLoS Comp. Biol.*, 5(9), 2009.

[32] Yun Yu, Tandy Warnow, and Luay Nakhleh. Algorithms for mdc-based multi-locus phylogeny inference: beyond rooted binary gene trees on single alleles. *Journal of Computational Biology*, 18(11):1543–1559, 2011.

[33] M. S. Bayzid and T. Warnow. Gene tree parsimony for incomplete gene trees: addressing true biological loss. *Algorithms for Molecular Biology*, 13:1, 2018.

[34] Siavash Mirarab, Md Shamsuzzoha Bayzid, Bastien Boussau, and Tandy Warnow. Statistical binning enables an accurate coalescent-based estimation of the avian tree. *Science*, 346(6215):1250463, 2014.

[35] Y Chung and C Ané. Comparing two Bayesian methods for gene tree/species tree reconstruction: A simulation with incomplete lineage sorting and horizontal gene transfer. *Syst Biol*, 60(3):261–275, 2011.

[36] Sen Song, Liang Liu, Scott V Edwards, and Shaoyuan Wu. Resolving conflict in eutherian mammal phylogeny using phylogenomics and the multispecies coalescent model. *Proceedings of the National Academy of Sciences*, 109(37):14942–14947, 2012.

[37] Md Shamsuzzoha Bayzid and Tandy Warnow. Estimating optimal species trees from incomplete gene trees under deep coalescence. *Journal of Computational Biology*, 19(6):591–605, 2012.

[38] Ylenia Chiari, Vincent Cahais, Nicolas Galtier, and Frédéric Delsuc. Phylogenomic analyses support the position of turtles as the sister group of birds and crocodiles (archosauria). *Bmc Biology*, 10(1):65, 2012.

[39] Rafael Zardoya and Axel Meyer. Complete mitochondrial genome suggests diapsid affinities of turtles. *Proceedings of the National Academy of Sciences*, 95(24):14226–14231, 1998.

[40] Andrew F Hugall, Ralph Foster, and Michael SY Lee. Calibration choice, rate smoothing, and the pattern of tetrapod diversification according to the long nuclear gene rag-1. *Systematic biology*, 56(4):543–563, 2007.

[41] Naoyuki Iwabe, Yuichiro Hara, Yoshinori Kumazawa, Kaori Shibamoto, Yumi Saito, Takashi Miyata, and Kazutaka Katoh. Sister group relationship of turtles to the bird-crocodilian clade revealed by nuclear dna–coded proteins. *Molecular Biology and Evolution*, 22(4):810–813, 2004.

[42] Tae-Kun Seo. Calculating bootstrap probabilities of phylogeny using multilocus sequence data. *Molecular biology and evolution*, 25(5):960–971, 2008.

[43] B. Boussau, G. J. Szöllősi, L. Duret, M. Gouy, E. Tannier, and V. Daubin. Genome-scale coestimation of species and gene trees. *Genome Research*, 23(2):323–330, 2013.

[44] Vikas Kumar, Björn M Hallström, and Axel Janke. Coalescent-based genome analyses resolve the early branches of the euarchontoglires. *PLoS One*, 8(4):e60019, 2013.

[45] Siavash Mirarab, Md Shamsuzzoha Bayzid, and Tandy Warnow. Evaluating summary methods for multilocus species tree estimation in the presence of incomplete lineage sorting. *Systematic Biology*, 65(3):366–380, 2014.