1 **Longer is not always better: Optimizing barcode length for large-scale species**

2 **discovery and identification**

3 Running Title: mini-barcodes for species discovery

16 **Darren Yeo[1], Amrita Srivathsan[1], Rudolf Meier[1*]**

18 [1]*Department of Biological Sciences, National University of Singapore, 14 Science 8*

19 *Drive 4, Singapore 117543*

20 [*]*Corresponding author: meier@nus.edu.sg*

Abstract

New techniques for the species-level sorting of millions of specimens have to be developed in order to answer the question of how many species live on earth. These methods should be reliable, scalable, and cost-effective as well as largely insensitive to the low-quality genomic DNA commonly obtained from museum specimens. Mini-barcodes seem to satisfy these criteria, but it is unclear whether they are sufficiently informative for species-level sorting. This is here tested based on 20 datasets covering ca, 30,000 specimens of 5,500 species. All specimens were first sorted based on morphology before being barcoded with full-length *cox1* barcodes. Mini-barcodes of different lengths and positions were then obtained *in silico* from the full-length barcodes using nine published mini-barcode primers (length: 94 – 407-bp) and a sliding window approach (3 windows: 100-bp, 200-bp, 300-bp). Afterwards, we determined whether barcode length and/or position reduces congruence between morphospecies and molecular Operational Taxonomic Units (mOTUs) that were obtained using three different species delimitation techniques (ABGD, PTP, objective clustering). We also evaluate how useful the published mini-barcodes are for species identification with the "best close match" algorithm. We find that there is no significant difference in performance for both species delimitation and identification between full-length and mini-barcodes as long as they are of moderate length (>200-bp). Only very short mini-barcodes (<200-bp) perform poorly, especially when they are located near the 5' end of the Folmer region. Overall, congruence between morphospecies and mOTUs is ca. 80% for barcodes that are >200-bp. The congruent mOTUs contain ca. 75% of the specimens and we estimate that most of the conflict is caused by ca. 10% of the specimens that should be targeted for re-examination in order to resolve conflict efficiently. Overall, barcode length (>200-bp)

and species delimitation methods have minor effects on congruence. Our study

suggests that large-scale species discovery, identification and metabarcoding can

utilize mini-barcodes without substantial loss of information compared to full-length

barcodes. This is good news given that mini-barcodes can be obtained via cost-

effective tagged amplicon sequencing using short-read sequencing platforms

(Illumina: "NGS barcodes").

Keywords: DNA barcoding, mini-barcodes, species discovery, metabarcoding

Introduction

The question of how many species live on earth has intrigued biologists for a very long time, but we are nowhere close to having a robust answer. We do know that fewer than 2 million of the estimated 10-100 million multicellular species have been described (Roskov et al. 2018) and that many are currently being extirpated by the "sixth mass extinction" (Ceballos et al. 2015; Sánchez-Bayo & Wyckhuys 2019) with potentially catastrophic consequences for the environment (Cafaro 2015). Monitoring, halting, and perhaps even reversing this process is hampered by the "taxonomic impediment". This impediment is particularly severe for "invertebrates" that collectively contribute much of the animal biomass (e.g., arthropods, annelids, nematodes: Stork et al. 2015; Bar-On et al. 2018). Most biologists thus agree that there is a pressing need for accelerating species discovery and description. This is very likely to require the development of new molecular methods. We would argue that they should not only be accurate, but also (1) rapid, (2) cost-effective, and (3) and largely insensitive to DNA quality. These criteria are important because tackling the earth's biodiversity would likely require the processing of >500 million specimens even under the very conservative assumption that there are only 10 million species and a new species is discovered with every 50 specimens processed. Cost-effectiveness is similarly important because millions of species are found in countries with only basic research facilities. Fortunately, many species are already represented in museum holdings, but such specimens often yield degraded DNA (Cooper 1994). Therefore, methods that require DNA of high-quality and quantity are not likely to be suitable for large-scale species discovery in invertebrates.

Conceptually, species discovery and description can be broken up into three steps. The first is obtaining specimens, the second, species-level sorting, and the

third, species identification or description. Fortunately, centuries of collecting have

generated many of the specimens that are needed for large-scale species discovery.

Indeed, for many invertebrate groups it is likely that the museum collections contain

more specimens of undescribed than described species; i.e., this unsorted collection

material represents vast and still underutilized source for species discovery (Lister &

Climate Change Research Group 2011; Kemp 2015; Yeates et al. 2016). The

second step in species discovery/description is species-level sorting. This step is in

dire need for acceleration. Traditionally, it starts with the sorting of unsorted material

into major taxa (e.g., order-level in insects). This task can still be accomplished by

parataxonomists but may in the future be taken over by machines utilizing neural

networks (Valan et al. 2019). In contrast, the subsequent species-level sorting is

usually time-limiting because the specimens for many invertebrate taxa have to be

prepared and studied by highly-skilled specialists (e.g., dissected; slide-mounted)

before the material can be sorted into putative species. This means that the

traditional techniques are neither rapid nor cost-effective for many invertebrate

groups. This impediment is likely to be responsible for why certain taxa that are

known to be abundant and species-rich are particularly poorly studied (Bickel 1999).

An alternative way to sort specimens to species-level would be with DNA

sequences. This approach is particularly promising for metazoan species because

most multicellular animal species can be distinguished based on cytochrome c

oxidase subunit I (*cox1*) barcode sequences (Hebert et al. 2003). However, such

sorting requires that each specimen is barcoded. This creates cost- and scalability

problems when the barcodes are obtained with Sanger sequencing (see Taylor and

Harris 2012). Such sequencing is currently still the standard in barcoding studies

because the animal barcode was defined as a 658-bp long fragment of *cox1*

("Folmer region": Folmer et al. 1994), although sequences >500-bp with <1%

ambiguous bases are also considered BOLD-compliant (BOLDsystems.org). The

658-bp barcode was optimized for ABI capillary sequencers but it has become a

burden because it is not suitable for most new sequencing technologies, while

Sanger sequencing remains expensive and only scalable when expensive liquid-

handling robots are used. This approach is hence unlikely to become widely

available in those countries that harbour most of the species diversity. Due to these

constraints, very few studies have utilized DNA barcodes to sort entire samples into

putative species (but see Fagan-Jeffries et al. 2018). Instead, most studies use a

mixed approach where species-level sorting is carried out based on morphology

before a select few specimens per morphospecies are barcoded (e.g., Riedel et al.

2010). This two-step process requires considerable amounts of skilled manpower

and time.

Scalability and cost-effectiveness are hallmark features of the new short-read

high throughput sequencing technologies. In addition, these technologies are

particularly suitable for sequencing the kind of degraded DNA that is typical for

museum specimens. Indeed, anchored hybrid enrichment (AHE) has already been

optimized for the use with old museum specimens (Bi et al. 2013; Guschanski et al.

2013; Blaimer et al. 2016) and is likely to play a major role for the integration of rare

species into taxonomic and systematic projects. It will be difficult, however, to apply

AHE to millions of specimens because it requires time-consuming and expensive

molecular protocols (e.g., specimen-specific libraries). Fortunately, for most species

it is likely that species-level sorting does not require a very large number of markers.

We would thus argue that the initial species-level sorting can be achieved using

barcodes that can be obtained via "tagged amplicon sequencing" on next-generation

sequencing platforms ("NGS barcodes"; Wang et al. 2018; Yeo et al. 2018). Until

recently, obtaining full-length NGS barcodes via tagged amplicon sequencing was

difficult because the reads of most next-generation-sequencing platforms were too

short for sequencing the full-length barcode. This has now changed with the arrival

of third-generation platforms (ONT: MinION: Srivathsan et al. 2018; PacBio: Sequel:

Hebert et al. 2018). These platforms, however, come with drawbacks; *viz* elevated

sequencing error rates and higher cost. These problems are likely to be overcome in

the future (e.g., Yang et al. 2018), but such solutions would not solve the main

challenge posed by full-length barcodes; i.e., the problems with reliably obtaining

amplicons from museum specimens with degraded DNA (e.g., Hajibabaei et al.

2006). We thus submit that one should optimize the barcode length based on

empirical evidence; it should be only as long as needed for accurate pre-sorting of

specimens into putative species.

Barcodes that are shorter than the full-length barcode are called mini-

barcodes. They are obtained with primers that amplify shorter subsets of the original

barcode region and have several advantages. Firstly, such amplicons are easier to

obtain when the DNA in the sample is degraded (Hajibabaei & McKenna 2012).

Secondly, mini-barcodes can be sequenced at low cost using tagged amplicon

sequencing on short-read sequencing platforms (e.g., Illumina). Thirdly, mini-

barcode primers are available for a large number of species-rich metazoan clades

(Hajibabaei et al. 2006; Meusnier et al. 2008; Hebert et al. 2013; Little 2014) as well

as for specific taxa such as fruit flies, catfish and sharks (Fan et al. 2009;

Bhattacharjee & Ghosh 2014; Fields et al. 2015). It is thus not surprising that short

barcodes are already the barcodes of choice when the template DNA is degraded.

This is often the case for museum specimens (Zuccon et al. 2012; Hebert et al.

2013) or for environmental DNA which is usually analysed via metabarcoding (e.g.:

processed food: Armani et al. 2015; Shokralla et al. 2015; water, soil, fecal matter:

Epp et al. 2012; Srivathsan et al. 2015; Lim et al. 2016). Mini-barcodes were initially

obtained via Sanger sequencing, but they can now be sequenced much more

efficiently via tagged amplicon sequencing on short-read platforms ("NGS

barcoding": Wang et al. 2018: sequencing cost < 4 cents). Wang et al. (2018) could

thus implement a "reverse workflow" based on sequencing all specimens without any

species-level pre-sorting based on morphology. Four-thousand specimens of ants

were barcoded with a 313-bp mini-barcode. The specimens were then pre-sorted

into 89 molecular operational taxonomic units (mOTUs) that were largely congruent

with morphospecies (86 species). However, it remained unclear whether full-length

DNA barcodes would have further improved congruence, whether the results from

this study can be generalized, and which mini-barcode is optimal for large-scale pre-

sorting of specimens into putative species.

The answers to these questions are elusive, because mini-barcodes remain

insufficiently tested despite their ubiquitous use in metabarcoding. Arguably, existing

tests suffer from lack of scale (the largest study includes 6695 barcodes for 1587

species: Meusnier et al. 2008) and taxonomic scope (usually only 1-2 family-level

taxa: e.g. Hajibabaei et al. 2006; Yu & You 2010). Furthermore, the tests yielded

conflicting results. Hajibabaei et al. (2006) found high congruence with the full-length

barcode when species are delimited based on mini-barcodes and Meusnier et al.

(2008) find similar BLAST identification rates for mini-barcodes and full-length

barcodes in their *in silico* tests. However, Yu & You (2010) conceded that mini-

barcodes may have worse accuracy despite having close structural concordance

with the full-length barcode. In addition, Sultana et al. (2018) concluded that the

ability to identify species is compromised when the barcodes are too short (<150-bp),

but it remained unclear at which length and in which position mini-barcodes stop

performing poorly. Furthermore, published tests of mini-barcodes compare their

performance to results obtained with full-length barcodes. All conflict is then implicitly

considered evidence for the failure of mini-barcodes. However, results obtained with

longer barcodes should not automatically be assumed to be accurate given that the

Folmer region varies in nucleotide variability (Roe & Sperling 2007). Lastly, the

existing tests of mini-barcodes do not include a sufficiently large number of different

mini-barcodes in order to be able to detect positional and lengths effects across the

658-bp barcode region.

Here, we address the lack of scale by including 20 studies covering 5500

species represented by ca. 30,000 barcodes. We furthermore test a large number of

different mini-barcodes by applying a sliding window approach to generate mini-

barcodes of different lengths (100, 200, 300-bp window length, 60-bp intervals) and

compare the results to the performance of nine mini-barcodes with published primers

(mini-barcode length: 94 – 407-bp). The taxonomic scope of our study is broad and

includes a wide variety of metazoans ranging from earthworms to butterflies and

birds. Lastly, we do not assume that mOTUs based on full-length barcodes are

automatically more accurate than those obtained with mini-barcodes. Instead, we

use morphology as an external criterion for assessing whether mOTUs obtained with

different-length barcodes have different levels of congruence with morphospecies.

Note that this does not imply that morphology is more suitable for species

delimitation than molecular data. Instead, we test whether shortening barcodes

influences congruence with morphology; i.e. morphology is treated as a constant

while testing whether barcode length and/or position influences the number of

morphospecies that are recovered. Given that morphology is a generally accepted

type of data that can be used for species delimitation, mini-barcodes that significantly

lower congruence with morphospecies are unlikely to be useful for accurate species-

level sorting.

We also compare the performance of different species delimitation methods.

There has been substantial interest in developing algorithms for mOTU estimation,

leading to the emergence of various species delimitation algorithms over the past

decade (e.g., objective clustering: Meier et al. 2006; BPP: Yang & Rannala 2010;

jmOTU: Jones et al. 2011; ABGD; Puillandre et al. 2012; BINs: Ratnasingham &

Hebert 2013; PTP: Zhang et al. 2013; etc.). For the purposes of this study, we

selected three algorithms as representatives of distance and tree-based methods:

objective clustering, Automatic Barcode Gap Discovery (ABGD) and Poisson Tree

Process (PTP). Objective clustering utilizes an *a priori* distance threshold to group

sequences into clusters, ABGD groups sequences into clusters based on an initial

prior and recursively uses incremental priors to find stable partitions, while PTP

utilizes the branch lengths on the input phylogeny to delimit species units. Arguably,

barcode data may not be appropriate for the application of PTP because a single

marker is not likely to yield reliable phylogenetic trees (including branch lengths), but

PTP has been frequently applied to barcode data in the literature (e.g. Ermakov et al.

2015; Han et al. 2016; Hollatz et al. 2016) and is thus included here. There are

numerous additional techniques for species delimitation, but most require multiple

markers and/or are even more reliant on accurately reconstructed phylogenetic

trees. They may thus not easily scale to millions of specimens and are excluded

here.

Lastly, we examine the utility of mini-barcodes for species identification. This

is the original use of DNA barcodes (Hebert et al. 2003). The widespread use of the

metazoan *cox1* Folmer region barcode greatly contributed to the usefulness of DNA

barcodes for this purpose, with the value of barcodes increasing as the databases

grow. While not as relevant to large-scale species discovery, species identification

with barcodes is particularly valuable for targeted species detection and community

characterization in metabarcoding studies (Ficetola et al. 2008; Lim et al. 2016;

Morinière et al. 2019). Most of these studies tend to involve poor quality genetic

material (i.e., gut content, faecal matter, environmental DNA) and hence it would be

valuable to know whether short markers can reliably assign species identities to an

unknown sequence. We here test the performance of the barcodes in our test

datasets using "best close match", which considers those sequences correctly

identified when they match conspecific barcodes within a predefined threshold

(Meier et al. 2006).

Materials & Methods

Dataset selection

We surveyed the barcoding literature in order to identify publications that cited

the original barcode paper by Hebert et al. (2003) and met the following criteria: 1)

have pre-identified specimens where the barcoded specimens were pre-

sorted/identified based on morphology and 2) the dataset had at least 500

specimens with *cox1* barcodes >656-bp. We identified 20 most recent datasets

dating backwards from 2017 (Table S1); all had >500 barcoded specimens even

after removing those that were not sorted to species level (e.g., only identified to

genus or higher) or had short sequences <657-bp (the full-length barcode is

technically 658-bp long, but a 1-bp concession was made to prevent the loss of too

much data). The barcode sequences were downloaded from BOLDSystems or NCBI

GenBank and aligned with MAFFT v7 (Katoh & Standley 2013) with a gap opening

penalty of 5.0.

Using a custom python script, we generated three sets of mini-barcodes along

a "sliding window". They were of 100-, 200- and 300-bp lengths. The first iteration

begins with the first base pair of the 658-bp barcode and the shifting windows jump

60-bp at each iteration, generating ten 100-bp windows, eight 200-bp windows and

six 300-bp windows. Additionally, we identified nine mini-barcodes with published

primers within the *cox1* Folmer region (Fig. 1 & Table S2). These mini-barcodes

have been repeatedly used in the literature and were used for a broad range of taxa.

The primers for the various mini-barcodes were aligned to the homologous regions

of each dataset with MAFFT v7 --addfragments (Katoh & Standley 2013) in order to

identify the precise position of the mini-barcodes within the full-length barcode. The

mini-barcode subsets from each barcode were then identified after alignment to full-

length barcodes to mark the start and end-point of the amplicon. Note that most of

the published primers are in the 5' prime region of the full-length barcode.

Species delimitation

The mini-barcodes and the full-length barcodes were clustered into putative

species using three species delimitation algorithms: objective clustering (Meier et al.

2006), ABGD (Puillandre et al. 2012) and PTP (Zhang et al. 2013). For objective

clustering, the mOTUs were clustered at 2 – 4% uncorrected p-distance thresholds

(Srivathsan & Meier 2012) using a python script which reimplements the objective

clustering of Meier et al. 2006 and allows for batch processing. The p-distance

thresholds selected are the typical distance thresholds used for species delimitation

in the literature (Meier et al. 2006; Ratnasingham & Hebert 2013; Meier et al. 2016).

The same datasets were also clustered with ABGD (Puillandre et al. 2012) using the

default range of priors and with uncorrected p-distances, but the minimum slope

parameter (-X) was reduced in a stepwise manner (1.5, 1.0, 0.5, 0.1) if the algorithm

could not find a partition. We then considered the ABGD clusters at priors P=0.001,

P=0.01 and P=0.04. The priors (P) refer to the maximum intraspecific divergence

and function similarly to p-distance thresholds at the first iteration, before being

refined by recursive application of the ABGD algorithm. Lastly, in order to use PTP,

the datasets were used to generate maximum likelihood (ML) trees in RAxML v.8

(Stamatakis 2014) via rapid bootstrapping (-f a) and the GTRCAT model. The best

tree generated for each dataset was then used for species delimitation with PTP

(Zhang et al. 2013) under default parameters.

Performance assessment

We assess the performance of mini-barcodes by using morphospecies as an

external arbiter. Species-level congruence was quantified using match ratios

between molecular and morphological groups (Ahrens et al. 2016). The ratio is

defined as $\frac{2 \times N_{match}}{N_1 + N_2} \times 100$, where N_{match} is the number of clusters identical across

both mOTU delimitation methods/thresholds (N_1 & N_2). Incongruence between

morphospecies and mOTUs is usually caused by a few specimens that are assigned

to the "incorrect" mOTUs. Conflict at the specimen-level can thus be quantified as

the number of specimens that are in mOTUs that cause conflict with morphospecies.

In order to test whether barcode length is a significant predictor of congruence,

MANOVA tests were carried out in R (R Core Team 2017) with "match ratio"

(species-level congruence) as the response variable and "dataset" and "mini-

barcode" as categorical explanatory variables. We found that most of the variance in

our study was generated by the "dataset" variable ($P < 0.05$ in MANOVA tests).

Given that we were particularly interested in the effect of barcode length and

position, "dataset" was subsequently treated as a random effect "mini-barcode" as

the explanatory variable (categorical) in a linear mixed effects model (R package

lme4: Bates et al. 2015). The *emmeans* R package (Lenth 2018) was then used to

perform pairwise post-hoc Tukey tests between mini- and full-length barcodes so as

to assess whether either barcode was performing significantly better/worse. To

compare the differences in performance between objective clustering, ABGD and

PTP, ANOVA tests were performed in R. After which, pairwise Tukey tests were

used to determine which species delimitation method was responsible for significant

differences. Lastly, in order to explore the reasons for positional effects, the

proportion of conserved sites for each mini-barcode was obtained using MEGA6

(Tamura et al. 2013).

Match ratios indicate congruence at the species level, but it is also important

to determine how many specimens have been placed in congruent units. Species-

and specimen-level congruence are only identical when all mOTUs are represented

by the same number of specimens. However, specimen abundances are rarely equal

across species and hence match ratio is insufficient at characterizing congruence

between mOTUs and morphospecies. It is straightforward to determine the number

of congruent specimens as follows:

(1) Congruence Class I specimens: If $A = B$ then number of congruent specimens is

$\text{Nc}_1 = |A|$ OR $|B|$.

Incongruence is caused by morphospecies that are split, lumped, or split and

lumped in the mOTUs. This means that a single mis-sorted specimen placed into a

large-sized mOTU leads to all specimens in two mOTUs to be considered

"incongruent" according to the criterion outlined above. Yet, most specimens are

congruent and full congruence could be restored by re-examining the mis-sorted

specimen. It is therefore also desirable to determine the number of specimens that

require re-examination or, conversely, the number of specimens that would be

congruent if one were to remove a few incongruent specimens. This number of

specimens can be estimated by counting congruent specimens as follows:

(2) Congruence Class II specimens: Specimens that are in split or lumped mOTUs

relative to morphospecies. Here, the largest subset of congruently placed specimens

can be determined as follows. If $A_1 \cup A_2 \cup \ldots \cup A_x = B$: $\text{Nc}_2 = \max(|A_1|, |A_2| \ldots |A_x|)$

(3) Congruence Class III specimens: This covers specimens in sets of clusters that

are both split and lumped relative to morphospecies. Here, only those specimens are

considered potentially congruent that (1) are in one mOTU and one morphospecies

and (2) combined exceed the number of the other specimens in the set of clusters. In

detail, if $A_1 \cup A_2 \cup \ldots \cup A_x = B_1 \cup B_2 \cup \ldots \cup B_y$: $\text{Nc}_3 = \max(|A_1 \cap B_1|, |A_2 \cap$

$B_1| \ldots |A_x \cap B_y|)$ only if $\max(|A_1 \cap B_1|, |A_2 \cap B_1| \ldots |A_x \cap B_y|) > \frac{1}{2}(|A_1 \cup A_2 \ldots A_x|)$.

Note that these are approximate estimates because we did not re-cluster specimens

after the removal of the specimens that were causing incongruence during in the

original cluster sets.

Species identification

Using the full-length and mini-barcode sets derived from the published

primers, we assessed whether the barcodes were yielding conspecific matches

according to "best close match" implemented in SpeciesIdentifier program

(TaxonDNA: Meier et al. 2006). "Best close match" examines one barcode at a time.

It considers it unidentified and then determines whether the best-matching reference

sequence in the dataset is conspecific and within a user-defined distance threshold.

The distance threshold used for all assessments was 2%, which corresponds to the

best match ratio performance results for objective clustering in this study. Sequences

with conspecific matches within 2% are considered correctly identified, while

sequences that match allospecific sequences within 2% are considered incorrectly

identified. Sequences without matches within 2% are considered unidentified and

sequences that have both con- and allo-specific matches within the threshold are

considered to have an ambiguous identification. Mixed effects models were again

used to assess the performance of barcodes of different length. Similarly, pairwise

Tukey tests were used to test for statistical significance with the proportion of correct

matches as the response variable in order to determine whether the full-length

barcode performed significantly better than the mini-barcodes.

Results

For species delimitation with objective clustering, we found that the 2% p-

distance threshold yielded the highest congruence across the datasets. It was hence

used as the upper-bound estimator for species- and specimen-level congruence.

The corresponding results for the 3 and 4% p-distance clusters are reported in the

supplementary materials. For ABGD it was the P=0.001 prior that yielded the highest

average match ratio and hence the clusters generated by this prior were used in the

main analysis (see supplementary material for results under P=0.01 and P=0.04).

PTP does not require parameter choices post the input tree.

The MANOVA tests performed on all treatments (species delimitation method

and distance threshold/prior) indicated that the test variable "dataset" was

responsible for much of the observed variance in "match ratio". The choice of mini-

barcode or algorithm for generating mOTUs was of secondary importance (Table

S3). After accounting for "dataset", we find that only mini-barcodes <200-bp perform

significantly worse than full-length barcodes (Fig. 2); for all other mini-barcodes

(>200-bp) the congruence with morphospecies does not differ significantly and is

occasionally superior to what is observed for the full-length barcode. This is evident

from the large number of significant differences ($p < 0.05$ & $p < 0.001$) in pairwise

post-hoc Tukey tests applied to 100-bp mini- and 657-bp full-length barcodes. Only

short barcodes (<100-bp) have a mean performance that is worse (<0 match ratio

deviation) than the full-length barcode. Conversely, there is no significant difference

between the 200 and 300-bp mini-barcodes and the full-length barcode when

objective clustering or PTP are used to estimate mOTUs. Under ABGD, the mini-

barcodes outperform the full-length barcodes. For all mOTU delimitation methods,

the variance across datasets decline as the mini-barcode increases in length (Fig. 2).

The results obtained for *in silico* mini-barcodes are consistent with the performance

of mini-barcodes with published primers: the mini-barcodes of 94-bp, 130-bp, and

145-bp lengths tend to perform worse than the longer mini-barcodes (Fig. 3). The

results are also similar for specimen-level congruence (Table 1 & S8). However,

there are some exceptions including the performance improvements of short mini-

barcodes, for example, for European marine fish and Northwest Pacific molluscs

when grouped with objective clustering (Fig. 3).

When the performance of the three different clustering methods was

compared, significant differences ($p < 0.05$ in ANOVA test) were found only for the

100-bp mini-barcode set (Fig. 4). Here, pairwise post-hoc Tukey tests find that

objective clustering performs significantly better than the other delimitation methods

($p < 0.001$) while ABGD and PTP do not differ significantly ($p = 0.88$) but behave

erratically for short mini-barcodes (Fig. 2).

Mini-barcodes situated at the 5' end of the full-length barcode appear to

perform somewhat worse than those situated at the middle or at the 3' end (Fig. 2).

For example, the 100-bp mini-barcodes at the 5' end perform poorly for objective

clustering (mini-barcode midpoints at 50, 110 & 170-bp), ABGD (mini-barcode

midpoints at 110 & 170-bp) and PTP (mini-barcode midpoint at 110-bp). This effect

is, however, only statistically significant when the mini-barcodes are very short (100-

bp). This positional effect is present across all species delimitation techniques. Note

that the 5' end of the full-length barcode appears to contain a large proportion of

conserved sites, particularly around the 170-bp and 230-bp midpoint of the 100-bp

mini-barcode (Fig. 5). This positional effect averages out as the mini-barcodes

increase in length.

With regard to specimen-based congruence, we focused on the mini-

barcodes with published primers and here report the results for those that a barcode

length >200-bp. Approximately three quarters of all specimens are in the

"Congruence Class I" (Tables 1 & S8); i.e., their placement is congruent between

mOTUs and morphospecies (Average/Median: OC at 2%: 75/75%; ABGD P=0.001:

71/72%; PTP: 75/75%). The remaining specimens are placed in mOTUs that are

split, lumped, or split and lumped. The number of specimens that are responsible for

the splitting and lumping are classified as Congruence Class II and III specimens.

Overall, fewer than 10% of the specimens fall into these categories (Table 1: see

Class II specimens across species delimitation methods). These are the specimens

that should be studied first when addressing conflict between morphospecies and

mOTUs.

When comparing the suitability of full-length and mini-barcodes for

identification purposes, we find that full-length barcodes often have the highest

number of correct matches (Table 2), but the difference between mini-

barcodes >200-bp is not statistically significant and only very short mini-barcodes

had significantly lower proportions of correct matches when compared with the full-

length barcode: 94-bp: p = <0.0001, 130-bp: p = <0.0001, 145-bp: p = <0.0001, 164-

bp: p = 0.0243. The other mini-barcodes (189 – 407-bp) did not show statistically

significant differences (p = 0.0691 – 0.9996) to the full-length barcode in terms of

correct matches. However, for many datasets (e.g., Northwest Pacific Molluscs,

North American Birds, etc.), the proportion of ambiguous matches is higher for

shorter mini-barcodes (94 – 164-bp) compared to longer barcodes (189 – 657-bp).

Discussion

Accelerating species discovery and description is arguably one of the

foremost challenges in modern systematics. Specimens for many undescribed

species are already in world's natural history museums, but the samples need to be

sorted to species-level before they become available for species

identification/description and can be used for large-scale analyses of biodiversity

patterns. Pre-sorting specimens with DNA barcodes is a potentially promising

solution because it is scalable, can be applied to millions of specimens, and much of

the specimen handling can be automated. However, in order for this approach to be

suitable, a sufficiently large proportion of the pre-sorted units need to accurately

reflect species boundaries and the methods for obtaining the sequences need to be

suitable for processing specimens that yield degraded DNA.

The main source of variance in congruence: datasets

We here find that the average congruence between mOTUs and

morphospecies is 80% for all barcodes >200-bp (median: 83%), with the median

being higher (83%) because of outlier datasets with congruence <65% (OC at 2%;

ABGD P=0.001, PTP). These outlier datasets are also likely to be responsible for the

observation that much of the variance in congruence throughout our study is

explained by the variable "dataset". Despite the outliers, 72-75% (median) of the ca.

30,000 specimens are assigned to species that are supported by molecular and

morphological data. Overall, this is a very high proportion compared to species-level

congruence obtained with species-level sorting by parataxonomists (Krell 2004). The

specimen-based perspective on congruence between morphology and molecular

data is often underappreciated. However, specimen-level congruence is an important

criterion for evaluating the suitability of species-level sorting with barcodes. After all,

the basic units in a museum collection or an ecological survey are specimens and

not species. The correct placement of specimens into species is thus important for

systematists and biodiversity researchers alike given that the former would like to

see most of the specimens in a collection correctly placed and the latter often need

abundance and biomass information at species-level resolution. For our datasets,

congruence levels for specimens are slightly lower than for species.

The remaining ca. 25% of specimens are placed in mOTUs whose boundaries

do not agree with morphospecies. One may initially consider this an unacceptably

high proportion, but it is important to keep in mind that the misplacement of one

specimen (e.g., due to a contamination of a PCR) will render two mOTUs

incongruent; i.e., all specimens in these mOTUs will be considered incongruent and

contribute to the 25% of "incongruently" placed specimens. Arguably, one may

instead want to investigate how many specimens are causing the conflict. These are

the specimens that should be targeted for re-investigation in reconciliation studies.

The proportion across the 20 datasets in our study is fairly low and ranges from 10-

12% (median) depending on which mOTU delimitation technique is used.

Conflict between mOTUs and morphospecies can be caused by technical

error or biology. A typical technical factor would be accidental misplacement of

specimens due to lab contamination or error during morphospecies sorting. Indeed,

the literature is replete with cases where mOTUs that were initially in conflict with

morphospecies became congruent once the study of additional morphological

characters let to the revision of morphospecies boundaries (e.g., Smith et al. 2008;

Tan et al. 2010; Baldwin et al. 2011; Ang et al. 2017). But there are also numerous

biological reasons for why one should not expect perfect congruence between

mOTUs and species. Lineage sorting, fast speciation, large amounts of intraspecific

variability, and introgression are known to negatively affect the accuracy of DNA

barcodes (Will & Rubinoff 2004; Rubinoff et al. 2006; Meier 2008). It is thus

somewhat surprising that regardless of these issues, the final levels of congruence

between morphospecies and DNA sequences are often quite high in animals (Ball et

al. 2005; Cywinska et al. 2006; Renaud et al. 2012; Landi et al. 2014; Wang et al.

2018). This implies that the pre-sorting specimens to species-level units based on

barcodes is worth pursuing for many metazoan clades. High levels of congruence

are, however, not a universal observation across all of life. This approach to

specimen sorting is unlikely to be useful in groups with widespread barcode sharing

between species. This phenomenon occurs within Metazoa (e.g., Anthozoa: Huang

et al. 2008) and is likely to be the default outside of Metazoa (e.g., Chase & Fay

2009; Hollingsworth et al. 2011).

Barcode length and species delimitation methods

We here tested the widespread assumption that mOTUs based on full-length

barcodes are more reliable than those based on mini-barcodes (Burns et al. 2007;

Min & Hickey 2007). If this assumption was confirmed, then the use of mini-barcodes

would have to be discouraged despite higher amplification success rates and lower

cost. However, we find that the performance of *cox1* mini-barcodes with a

length >200-bp do not differ significantly from the performance of full-length

barcodes. Indeed, compared to the dataset effect, the choice of barcode length is

largely secondary. This conclusion is robust across 20 diverse datasets and holds for

different clustering algorithms.

We also find that the choice of species delimitation algorithm matters little for

mini-barcodes >200-bp (Fig. 4). This is fortunate as objective clustering and ABGD

algorithms are less computationally demanding than PTP, which necessitates the

reconstruction of a ML trees. However, there are some exceptions. Firstly, when the

mini-barcodes are extremely short (~100-bp), objective clustering tends to

outperform ABGD and PTP. PTP's poor performance for 100-bp mini-barcodes is not

surprising given that it relies on tree topologies which cannot be estimated with

confidence based on so little data. ABGD's poor performance is mostly observed for

certain priors (e.g., P=0.04: Fig. S5 & S6). Under these priors, ABGD tends to lump

most of the 100-bp barcodes into one or few large clusters. Prior-choice also affects

ABGD's performance for full-length barcodes. ABGD does not perform well with very

low priors (P = 0.001: Fig. 2 & 3 vs. P = 0.01; P = 0.04: Fig. S5). Overall, we

conclude that the selection of the best priors and/or clustering thresholds remains a

significant challenge for the study of largely unknown faunas that lack morphological

information as an *a posteriori* method for selecting priors/thresholds. Overall, we

recommend the use of multiple methods and thresholds in order to distinguish robust

from labile mOTUs that are heavily dependent on threshold- or prior-choice. The

latter should be targeted in reconciliation studies utilizing another type of data.

Barcode length and species identification

Similar to our tests on species delimitation, we find that mini-barcodes >200-

bp have no significant performance difference when compared with full-length

barcodes. However, it does appear that full-length barcodes have marginally – albeit

statistically not significantly – better performance than >200-bp mini-barcodes (Table

2: 0.00 – 4.15% difference, 0.37% median). This is consistent with findings by

Meusnier et al. (2008) who reported at least 95% identification success for 250-bp

barcodes while full-length barcodes performed the best (97% success). Overall, the

data suggest that mini-barcodes >200-bp are suitable for species identification when

matching sequences to barcode databases with identified barcodes. The poorer

performance of the shorter mini-barcodes (<200-bp) are largely due to an increase in

ambiguous matches rather than incorrect matches. These mini-barcodes are less

informative which could be due to short length or the placement of many short

barcodes in a more conserved part of the Folmer region as described below.

Positional effects

Overall, mini-barcodes at the 3' end of the Folmer region outperform mini-

barcodes at the 5' end. This is consistent across all three species delimitation

methods and was also mentioned by Shokralla et al. (2015) who concluded that

mini-barcodes at the 5' end have worse species resolution for fish species. This

positional effect is apparent when match ratios are compared across a "sliding

window" (Fig. 2). The lowest congruence with morphology is observed for 100-bp

mini-barcodes with midpoints at the 50, 110 and 170-bp marks. However, this

positional effect is only significant when the barcode lengths are very short (<200-

bp). Once the mini-barcodes are sufficiently long (>200-bp), there seems to be no

appreciable difference in performance, which is not surprising because sampling

more nucleotides helps with buffering against regional changes in nucleotide

variability. These changes in nucleotide variability may have functional reasons

related to the conformation of the Cox1 protein in the mitochondrion membrane. The

Folmer region of Cox1 contains six transmembrane α- helices that are connected by

five loops (Tsukihara et al. 1996; Pentinsaari et al. 2016). Pentinsaari et al. (2016)

compared 292 Cox1 sequences across 26 animal phyla and found high amino acid

variability in helix I and the loop connecting helix I and helix II (corresponding to

position 1-102 of *cox1*), as well as end of helix IV and loop connecting helix IV and V

(corresponding to positions ~448-498). These regions of high variability are distant

from the active sites and thus less likely to affect Cox1 function (Pentinsaari et al.

2016). This may lead to lower selection pressure and thus higher variability in these

areas which could impact the performance of mini-barcodes for species delimitation.

Accelerating biodiversity discovery and description

We had earlier argued that species discovery and description can be broken

up into three steps (1) obtaining specimens, (2) species-level sorting and (3) species

identification or description. We here only address species-level sorting. This means

that the impediments caused by slow rates of species identification and description

are apparently unresolved. However, this is only partially correct. Firstly, some

mOTUs delimited via barcodes can be identified via barcode databases. The

proportion of successful identification differs depending on how well a particular

fauna has been barcoded. This is illustrated by our recent work on dragon- and

damselflies (Odonata), ants (Formicidae), and non-biting midges (Chironomidae) of

Singapore (Wang et al. 2018, Yeo et al. 2018; Baloğlu et al. 2018). These studies

also document why it is important to distinguish between species- and specimen-

level identification because the results differ widely. For odonates, BLAST-searches

identified more than half of the 95 mOTUs and >75% of the specimens to species.

The corresponding numbers for ants and midges were ca. 20% and 10% at mOTU-

level, and 9% and 40% at the specimen-level. Secondly, mOTUs discovered via

barcodes can be readily compared across studies and borders (Ratnasingham, et al.

2013). In contrast, species newly discovered based on morphological evidence

usually remain unavailable to the scientific community until they are published. This

is a very significant difference because a large amount of downstream biodiversity

analysis can be carried out based on mOTUs instead of identified/described species.

This includes studying species richness and abundance over time which is a task

that is becoming increasingly important in the 21st century. This means that only

moderate harm is done if species identification or description are postponed.

Unfortunately, the analyses of biodiversity patterns will be impacted by

incorrectly delimited mOTUs and it is thus important to estimate this error. In our

study, we find that ca. 80% of the mOTUs are congruent with morphospecies. This is

prior to a reconciliation stage where the morphology of specimens with a conflicting

assignment is revisited in order to rule out that the morphological evidence was

misinterpreted and/or an insufficient number of characters was studied; i.e., we

would predict that the overall congruence levels after reconciliation will be higher.

Ideally, we would like to know which proportion of mOTUs that are in conflict with

morphospecies will eventually be rejected, but unfortunately we still know fairly little

about the congruence levels between morphology and barcodes after reconciliation.

This is because rigorous studies would have to be based on datasets with dense

taxon and geographic sampling where first morphological and DNA sequence

information is obtained for all specimens and then all cases of conflict are re-studied.

Unfortunately, there are very few datasets that satisfy these criteria. This is

presumably because the high cost of full-length barcodes has prevented biologists

from sequencing all specimens.

Conclusions

We here illustrate that mini-barcodes are as reliable for pre-sorting specimens

into putative species as full-length barcodes. Mini-barcodes are also suitable for

identifying unidentified barcodes to species based on barcode reference databases.

We would thus argue that mini-barcodes should be preferred over full-length

barcodes because (1) they can be obtained more readily for specimens that only

contain degraded DNA (Hajibabaei et al. 2006) and (2) are much more cost-

effective. In particular, we recommend the use of mini-barcodes >200-bp at the 3'

end of the Folmer region. It is encouraging that such mini-barcodes perform well

across a large range of metazoan taxa. These conclusions are based on three

species delimitation algorithms (objective clustering, ABGD and PTP) which, overall,

have no appreciable effect on the performance of mini-barcodes. If the DNA of the

specimens is so degraded that very short mini-barcodes have to be obtained, we

advise against the use of PTP and ABGD (especially with high priors) in order to

reduce the likelihood that morphospecies are lumped.

Funding

This work was supported by a Ministry of Education grant on biodiversity discovery

(R-154-000-A22-112).

Acknowledgements

We would like to acknowledge Athira Adom and Jonathan Ho for data processing

and Emily Hartop and Wendy Wang for commenting on the manuscript.

References

Ahrens D., Fujisawa T., Krammer H.J., Eberle J., Fabrizi S., & Vogler A.P. 2016.

Rarity and incomplete sampling in DNA-based species delimitation. *Systematic*

Biology. 65:478–494.

Ang Y., Rajaratnam G., Su K.F., & Meier R. 2017. Hidden in the urban parks of New

York City: *Themira lohmanus*, a new species of Sepsidae described based on

morphology, DNA sequences, mating behavior, and reproductive isolation

(Sepsidae, Diptera). *ZooKeys*. 698:95.

Armani A., Guardone L., Castigliego L., D'Amico P., Messina A., Malandra R.,

Gianfaldonia D. & Guidi A. 2015. DNA and Mini-DNA barcoding for the identification

of Porgies species (family Sparidae) of commercial interest on the international

market. *Food Control*. 50:589–596.

Baldwin C.C., Castillo C.I., & Weigt L.A. 2011. Seven new species within western

Atlantic *Starksia atlantica*, *S. lepicoelia*, and *S. sluiteri* (Teleostei, Labrisomidae),

with comments on congruence of DNA barcodes and species. *ZooKeys*. 79:21.

Ball S.L., Hebert P.D., Burian S.K., & Webb J.M. 2005. Biological identifications of

mayflies (Ephemeroptera) using DNA barcodes. *Journal of the North American*

Benthological Society. 24:508–524.

Baloğlu B., Clews E., & Meier R. 2018. NGS barcoding reveals high resistance of a

hyperdiverse chironomid (Diptera) swamp fauna against invasion from adjacent

freshwater reservoirs. *Frontiers in zoology*. 15:31.

Bar-On Y.M., Phillips R., & Milo R. 2018. The biomass distribution on

Earth. *Proceedings of the National Academy of Sciences*. 115:6506–6511.

Bates D., Mächler M., Bolker B., & Walker S. 2015. Fitting linear mixed-effects

models using *lme4*. *Journal of Statistical Software*. 67:1–48

Bhattacharjee M.J., & Ghosh S.K. 2014. Design of Mini-barcode for Catfishes for

assessment of archival biodiversity. *Molecular Ecology Resources*. 14:469–477.

Bi K., Linderoth T., Vanderpool D., Good J.M., Nielsen R., & Moritz C. 2013.

Unlocking the vault: next-generation museum population genomics. *Molecular*

ecology. 22:6018–6032.

Bickel D.J. 1999. What museum collections reveal about species accumulation,

richness, and rarity: an example from the Diptera. *The other 99%: the conservation*

and biodiversity of invertebrates. 174–181.

Blaimer B.B., Lloyd M.W., Guillory W.X. & Brady S.G. 2016. Sequence capture and

phylogenetic utility of genomic ultraconserved elements obtained from pinned insect

specimens. *PLoS One*. 11:e0161531.

Burns J.M., Janzen D.H., Hajibabaei M., Hallwachs W. & Hebert P.D. 2007. DNA

barcodes of closely related (but morphologically and ecologically distinct) species of

skipper butterflies (Hesperiidae) can differ by only one to three nucleotides. *Journal*

of the Lepidopterists Society. 61:138–153.

Cafaro P. 2015. Three ways to think about the sixth mass extinction. *Biological*

Conservation. 192:387–393.

Ceballos G., Ehrlich P.R., Barnosky A.D., García A., Pringle R.M. & Palmer T.M.

2015. Accelerated modern human–induced species losses: Entering the sixth mass

extinction. *Science Advances*. 1:e1400253.

Chase M.W. & Fay M.F. 2009. Barcoding of plants and fungi. *Science*. 325:682–683.

Cooper A. 1994. DNA from museum specimens. *Ancient DNA*. 149–165. Springer,

New York, NY.

Cywinska A., Hunter F.F. & Hebert P.D. 2006. Identifying Canadian mosquito

species through DNA barcodes. *Medical and Veterinary Entomology*. 20:413–424.

Epp L.S., Boessenkool S., Bellemain E.P., Haile J., Esposito A., Riaz T., Erséus C.,

Gusarov V.I., Edwards M.E., Johnsen A., Stenøien H.K., Hassel K., Kauserud H.,

Yoccoz N.G., Bråthen K.A., Willerslev E., Taberlet P., Coissac E. & Stenøien H.K.

2012. New environmental metabarcodes for analysing soil DNA: potential for

studying past and present ecosystems. *Molecular Ecology*. 21:1821–1833.

Ermakov O.A., Simonov E., Surin V.L., Titov S.V., Brandler O.V., Ivanova N.V. &

Borisenko A.V. 2015. Implications of hybridization, NUMTs, and overlooked diversity

for DNA barcoding of Eurasian ground squirrels. *PLoS One*. 10:e0117201.

Fagan-Jeffries E.P., Cooper S.J., Bertozzi T., Bradford T.M. & Austin A.D. 2018.

DNA barcoding of microgastrine parasitoid wasps (Hymenoptera: Braconidae) using

high-throughput methods more than doubles the number of species known for

Australia. *Molecular Ecology Resources*. 18:1132–1143.

Fan J.A., Gu H., Chen S., Mo B., Wen Y., He W., Liu W & Zeng X. 2009. Species

identification of 36 kinds of fruit flies based on minimalist-barcode. *Chinese Journal*

of Applied & Environmental Biology. 2:215–219.

Ficetola G.F., Miaud C., Pompanon F. & Taberlet P. 2008. Species detection using

environmental DNA from water samples. *Biology Letters*. 4:423–425.

Fields A.T., Abercrombie D.L., Eng R., Feldheim K. & Chapman D.D. 2015. A novel

mini-DNA barcoding assay to identify processed fins from internationally protected

shark species. *PloS One*. 10:e0114844.

Folmer O., Black M., Hoeh W., Lutz R. & Vrijenhoek R. 1994. DNA primers for

amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan

invertebrates. *Molecular Marine Biology and Biotechnology.* 3:294–299.

Guschanski K., Krause J., Sawyer S., Valente L.M., Bailey S., Finstermeier K., Sabin

R., Gilissen E., Sonet G., Nagy Z.T., Lenglet G., Mayer F. & Lenglet G. 2013. Next-

generation museomics disentangles one of the largest primate radiations.

Systematic Biology. 62:539–554.

Hajibabaei M., Smith M.A., Janzen D.H., Rodriguez J.J., Whitfield J.B. & Hebert P.D.

2006. A minimalist barcode can identify a specimen whose DNA is degraded.

Molecular Ecology Notes. 6:959–964.

Hajibabaei M. & McKenna C. 2012. DNA mini-barcodes. *DNA barcodes*. 339–353.

Humana Press, Totowa, NJ.

Han T., Lee W., Lee S., Park I.G. & Park H. 2016. Reassessment of species

diversity of the subfamily Denticollinae (Coleoptera: Elateridae) through DNA

Barcoding. *PloS One*. 11:e0148602.

Hebert P.D., Cywinska A., Ball S.L. & Dewaard J.R. 2003. Biological identifications

through DNA barcodes. *Proceedings of the Royal Society of London. Series B:*

Biological Sciences. 270:313–321.

Hebert P.D., Zakharov E.V., Prosser S.W., Sones J.E., McKeown J.T., Mantle B. &

La Salle J. 2013. A DNA 'Barcode Blitz': Rapid digitization and sequencing of a

natural history collection. *PLoS One*. 8:e68535.

Hebert P.D., Braukmann T.W., Prosser S.W., Ratnasingham S., Ivanova N.V.,

Janzen D.H., Hallwachs W., Naik S., Sones J.E. & Zakharov E.V. 2018. A Sequel to

Sanger: amplicon sequencing that scales. *BMC Genomics*. 19:219.

Hollatz C., Leite B.R., Lobo J., Froufe H., Egas C. & Costa F.O. 2016. Priming of a

DNA metabarcoding approach for species identification and inventory in marine

macrobenthic communities. *Genome*. 60:260–271.

Hollingsworth P.M., Graham S.W. & Little D.P. 2011. Choosing and using a plant

DNA barcode. *PloS One*. 6:e19254.

Huang D., Meier R., Todd P.A. & Chou L.M. 2008. Slow mitochondrial COI sequence

evolution at the base of the metazoan tree and its implications for DNA barcoding.

Journal of Molecular Evolution. 66:167–174.

Jones M., Ghoorah A. & Blaxter M. 2011. jmOTU and taxonerator: turning DNA

barcode sequences into annotated operational taxonomic units. *PLoS One.*

6:e19259.

Katoh K. & Standley D.M. 2013. MAFFT multiple sequence alignment software

version 7: improvements in performance and usability. *Molecular Biology and*

Evolution. 30:772–780.

Kemp C. 2015. Museums: The endangered dead. *Nature News*. 518:292.

Krell F.T. 2004. Parataxonomy vs. taxonomy in biodiversity studies–pitfalls and

applicability of 'morphospecies' sorting. *Biodiversity & Conservation*. 13:795–812.

Landi M., Dimech M., Arculeo M., Biondo G., Martins R., Carneiro M., Carvalho G.R.,

Brutto S.L. & Costa F.O. 2014. DNA barcoding for species assignment: the case of

Mediterranean marine fishes. *PLoS One*. 9:e106135.

Lenth R. 2018. *Emmeans*: Estimated marginal means, aka least-squares means. *R*

Package Version. 1.

Lim N.K.M., Tay Y.C., Srivathsan A., Tan J.W.T., Kwik J.T.B., Baloğlu B., Meier R &

Yeo D.C.J. 2016. Next-generation freshwater bioassessment: eDNA metabarcoding

with a conserved metazoan primer reveals species-rich and reservoir-specific

communities. *Royal Society Open Science*. 3:160635.

Lister A.M. & Climate Change Research Group. 2011. Natural history collections as

sources of long-term datasets. *Trends in Ecology & Evolution*. 26:153–154.

Little D.P. 2014. A DNA mini-barcode for land plants. *Molecular Ecology Resources*.

14:437–446.

Meier R., Shiyang K., Vaidya G. & Ng P.K. 2006. DNA barcoding and taxonomy in

Diptera: a tale of high intraspecific variability and low identification success.

Systematic Biology. 55:715–728.

Meier R. 2008. DNA sequences in taxonomy: opportunities and challenges. *The New*

Taxonomy (ed. Wheeler QD). 7:95–127. CRC Press, New York.

Meier R., Wong W., Srivathsan A. & Foo M. 2016. $1 DNA barcodes for

reconstructing complex phenomes and finding rare species in specimen-rich

samples. *Cladistics*. 32:100–110.

Meusnier I., Singer G.A., Landry J.F., Hickey D.A., Hebert P.D. & Hajibabaei M.

2008. A universal DNA mini-barcode for biodiversity analysis. *BMC Genomics*.

9:214.

Min X.J. & Hickey D.A. 2007. Assessing the effect of varying sequence length on

DNA barcoding of fungi. *Molecular Ecology Resources*. 7:365–373.

Morinière J., Balke M., Doczkal D., Geiger M.F., Hardulak L.A., Haszprunar G.,

Hausmann A., Hendrich L., Regalado L., Rulik B., Schmidt S., Wägele J.W. & Hebert

P.D.N. 2019. A DNA barcode library for 5,200 German flies and midges (Insecta:

Diptera) and its implications for metabarcoding-based biomonitoring. *Molecular*

Ecology Resources. doi: https://doi.org/10.1111/1755-0998.13022.

Pentinsaari M., Salmela H., Mutanen M. & Roslin T. 2016. Molecular evolution of a

widely-adopted taxonomic marker (COI) across the animal tree of life. *Scientific*

Reports. 6:35275.

Puillandre N., Lambert A., Brouillet S. & Achaz G. 2012. ABGD, Automatic Barcode

Gap Discovery for primary species delimitation. *Molecular Ecology*. 21:1864–1877.

R Core Team. 2017. R: A language and environment for statistical computing. *R*

Foundation for Statistical Computing. Vienna, Austria. URL http://www.R-project.org.

Ratnasingham S. & Hebert P.D. 2013. A DNA-based registry for all animal species:

the Barcode Index Number (BIN) system. *PloS One*. 8:e66213.

Renaud A.K., Savage J. & Adamowicz S.J. 2012. DNA barcoding of Northern

Nearctic Muscidae (Diptera) reveals high correspondence between morphological

and molecular species limits. *BMC Ecology*. 12:24.

Riedel A., Daawia D. & Balke M. 2010. Deep cox1 divergence and hyperdiversity of

Trigonopterus weevils in a New Guinea mountain range (Coleoptera,

Curculionidae). *Zoologica Scripta*. 39:63–74.

Roe A.D. & Sperling F.A. 2007. Patterns of evolution of mitochondrial cytochrome c

oxidase I and II DNA and implications for DNA barcoding. *Molecular Phylogenetics*

and Evolution. 44:325–345.

Roskov Y., Abucay L., Orrell T., Nicolson D., Bailly N., Kirk P.M., Bourgoin T.,

DeWalt R.E., Decock W., De Wever A., Nieukerken E. van, Zarucchi J., Penev L.

2017. Species 2000 & ITIS Catalogue of Life, 2017 Annual Checklist. Digital

resource at www.catalogueoflife.org/annual-checklist/2017. *Species 2000: Naturalis*.

Leiden, the Netherlands. ISSN 2405-884X.

Rubinoff D., Cameron S. & Will K. 2006. A genomic perspective on the shortcomings

of mitochondrial DNA for "barcoding" identification. *Journal of Heredity*. 97:581–594.

Sánchez-Bayo F. & Wyckhuys K.A. 2019. Worldwide decline of the entomofauna: A

review of its drivers. *Biological Conservation*. 232:8–27.

Shokralla S., Hellberg R.S., Handy S.M., King I. & Hajibabaei M. 2015. A DNA mini-

barcoding system for authentication of processed fish products. *Scientific Reports*.

5:15894.

Smith M.A., Rodriguez J.J., Whitfield J.B., Deans A.R., Janzen D.H., Hallwachs W. &

Hebert P.D. 2008. Extreme diversity of tropical parasitoid wasps exposed by iterative

integration of natural history, DNA barcoding, morphology, and

collections. *Proceedings of the National Academy of Sciences*. 105:12359–12364.

Srivathsan A. & Meier R. 2012. On the inappropriate use of Kimura-2-parameter

(K2P) divergences in the DNA-barcoding literature. *Cladistics*. 28:190–194.

Srivathsan A., Sha J., Vogler A.P. & Meier R. 2015. Comparing the effectiveness of

metagenomics and metabarcoding for diet analysis of a leaf-feeding monkey

(*Pygathrix nemaeus*). *Molecular Ecology Resources*. 15:250–261.

Srivathsan A., Baloğlu B., Wang W., Tan W.X., Bertrand D., Ng A.H.Q., Boey E.J.H.,

Koh J.J.Y., Nagarajan N. & Meier R. 2018. A Min ION™-based pipeline for fast and

cost-effective DNA barcoding. *Molecular Ecology Resources*. 0:1–15.

Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-

analysis of large phylogenies. *Bioinformatics*. 30:1312–1313.

Stork N.E., McBroom J., Gely C. & Hamilton A.J. 2015. New approaches narrow

global species estimates for beetles, insects, and terrestrial arthropods. *Proceedings*

of the National Academy of Sciences. 112:7519–7523.

Sultana S., Ali M.E., Hossain M.M., Naquiah N. & Zaidul I.S.M. 2018. Universal mini

COI barcode for the identification of fish species in processed products. *Food*

Research International. 105:19–28.

Tamura K., Stecher G., Peterson D., Filipski A. & Kumar S. 2013. MEGA6: molecular

evolutionary genetics analysis version 6.0. *Molecular Biology and Evolution*.

30:2725–2729.

Tan D.S., Ang Y., Lim G.S., Ismail M.R.B. & Meier R. 2010. From 'cryptic species' to

integrative taxonomy: an iterative process involving DNA sequences, morphology,

and behaviour leads to the resurrection of *Sepsis pyrrhosoma* (Sepsidae:

Diptera). *Zoologica Scripta*. 39:51–61.

Tamura K., Stecher G., Peterson D., Filipski A. & Kumar S. 2013. MEGA6: molecular

evolutionary genetics analysis version 6.0. *Molecular Biology and Evolution*.

30:2725–2729.

Taylor H.R. & Harris W.E. 2012. An emergent science on the brink of irrelevance: a

review of the past 8 years of DNA barcoding. *Molecular Ecology Resources.* 12:377–

388.

Tsukihara T., Aoyama H., Yamashita E., Tomizaki T., Yamaguchi H., Shinzawa-Itoh

K., Nakashima R., Yaono R. & Yoshikawa S. 1996. The whole structure of the 13-

subunit oxidized cytochrome c oxidase at 2.8 Å. *Science*. 272:1136–1144.

Valan M., Makonyi K., Maki A., Vondráček D. & Ronquist F. 2019. Automated

Taxonomic Identification of Insects with Expert-Level Accuracy Using Effective

Feature Transfer from Convolutional Networks. *Systematic Biology*, *syz014*,

https://doi.org/10.1093/sysbio/syz014.

Wang W.Y., Srivathsan A., Foo M., Yamane S.K. & Meier R. 2018. Sorting

specimen-rich invertebrate samples with cost-effective NGS barcodes: Validating a

reverse workflow for specimen processing. *Molecular Ecology Resources*. 18:490–

501.

Yang Z. & Rannala B. 2010. Bayesian species delimitation using multilocus

sequence data. *Proceedings of the National Academy of Sciences*. 107:9264–9269.

Yang C., Tan S., Meng G., Bourne D.G., O'brien P.A., Xu J., Liao S., Chen A., Chen X. & Liu S. 2018. Access COI barcode efficiently using high throughput Single End 400 bp sequencing. *BioRxiv*. 498618. doi: http://dx.doi.org/10.1101/498618.

Yeates D.K., Zwick A. & Mikheyev A.S. 2016. Museums are biobanks: unlocking the genetic potential of the three billion specimens in the world's biological collections. *Current Opinion in Insect Science*. 18:83–88.

Yeo D., Puniamoorthy J., Ngiam R.W.J. & Meier R. 2018. Towards holomorphology in entomology: rapid and cost-effective adult–larva matching using NGS barcodes. *Systematic Entomology*. 43:678–691.

Yu H.J. & You Z.H. 2010. Comparison of DNA truncated barcodes and full-barcodes for species identification. *Advanced Intelligent Computing Theories and Applications. With Aspects of Artificial Intelligence.* 108-114. Springer, Berlin, Heidelberg.

Zhang J., Kapli P., Pavlidis P. & Stamatakis A. 2013. A general species delimitation method with applications to phylogenetic placements. *Bioinformatics*. 29:2869–2876.

Zuccon D., Brisset J., Corbari L., Puillandre N., Utge J. & Samadi S. 2012. An optimised protocol for barcoding museum collections of decapod crustaceans: a case-study for a 10–40-years-old collection. *Invertebrate Systematics*. 26:592–600.

Figure Captions

Table 1. Proportion of specimens congruent between morphospecies and mOTU clusters under the three stringency classes. Values in brackets represent the estimated number of specimens causing conflict.

Table 2. Proportion of correct/incorrect/ambiguous/unidentified specimens in the "best close match" analysis.

Figure 1. Position of the mini-barcode with established primers in this study.

Figure 2. Performance of mini-barcodes along a sliding window (100, 200, 300-bp).

Mini-barcode position is indicated on the x-axis and congruence with morphology on

the y-axis. mOTUs were obtained with Objective Clustering (2%), ABGD P=0.001

prior), and PTP. Each line represents one data set while the boxplots summarise the

values across datasets. Significant deviations from the results obtained with full-

length barcodes are indicated with color-coded asterisks (* = $p < 0.05$; ** = $p <$

0.001; red = poorer and green = higher congruence with morphology).

Figure 3. Match ratios across three different species delimitation methods. Mini-

barcodes (columns) are sorted by primer length while the datasets (rows) are

grouped into 4 classes according to average match ratio. Colours are applied

separately to each class.

Figure 4. Comparison of species delimitation methods for full-length and mini-

barcodes generated by "sliding windows" (100-bp, 200-bp, 300-bp).

Figure 5. Proportion of conserved sites along the full-length barcode (sliding

windows of 100-bp, 200-bp, 300-bp).

Table 1. Proportion of specimens congruent between morphospecies and mOTU

clusters under the three stringency classes. Values in brackets represent the

estimated number of specimens causing conflict.

	Objective clustering, 2% p-distance						
Length	295	307	313	407	657	Average	Median
Midpoint	405	154	502	455	329		
Class I	74% (7475)	74% (7511)	75% (7275)	75% (7246)	76% (7118)	75% (7325)	75% (7372)
Class II	89% (2049)	89% (2214)	89% (2079)	89% (2030)	90% (1967)	89% (2068)	89% (2079)
Class III	90% (975)	90% (743)	90% (746)	90% (786)	91% (795)	90% (809)	90% (802)
	ABGD, P=0.001						
Length	295	307	313	407	657	Average	Median
Midpoint	405	154	502	455	329		
Class I	73% (7916)	74% (7685)	72% (8169)	71% (8583)	66% (9833)	71% (8437)	72% (8126)
Class II	89% (2178)	88% (2014)	89% (2138)	88% (2129)	87% (2604)	88% (2213)	88% (2057)
Class III	90% (806)	90% (895)	90% (849)	90% (915)	88% (881)	89% (869)	90% (862)
	PTP						
Length	295	307	313	407	657	Average	Median
Midpoint	405	154	502	455	329		
Class I	74% (7643)	74% (7744)	75% (7369)	76% (7130)	75% (7207)	75% (7419)	75% (7318)
Class II	89% (2362)	88% (2433)	89% (2227)	89% (2095)	89% (1965)	89% (2216)	89% (2179)
Class III	89% (732)	89% (904)	90% (702)	90% (721)	90% (897)	90% (791)	90% (733)

Table 2. Proportion of correct/incorrect/ambiguous/unidentified specimens in the "best close match" analysis.

Dataset	Published Primers									
	94	**130**	**145**	**164**	**189**	**295**	**307**	**313**	**407**	**657**
Great Barrier Reef Fish	85/1/0/15	83/1/3/13	83/1/2/14	85/0/0/14	86/1/0/13	86/0/0/14	85/1/0/14	86/0/0/14	86/0/0/14	86/0/0/14
South China Sea Fish	93/0/1/6	91/1/3/6	91/1/3/6	93/0/1/6	93/0/1/6	94/0/0/6	93/0/1/6	93/1/1/6	94/1/0/6	94/0/0/6
North Sea Molluscs	90/0/0/10	91/0/0/9	92/0/0/8	92/0/0/8	92/0/0/8	92/0/0/8	92/0/0/8	91/0/0/9	91/0/0/9	92/0/0/8
Canada Echinoderms	95/0/0/5	95/0/0/4	95/0/0/5	96/0/0/4	96/0/0/4	95/0/0/5	96/0/0/4	95/0/0/5	96/0/0/4	96/0/0/4
Ecuador Geometridae	84/0/0/15	85/1/0/13	85/0/0/14	85/0/0/14	85/0/0/14	86/0/0/14	86/0/0/14	86/0/0/14	86/0/0/14	86/0/0/14
Germany Aranea & Opiliones	88/0/5/7	89/0/5/5	89/0/5/6	90/1/4/5	90/0/4/5	92/0/2/6	91/1/3/5	92/0/2/5	92/1/2/5	93/1/2/5
European Marine Fish	92/0/7/1	91/0/7/1	92/0/6/2	96/0/3/1	96/0/3/1	97/0/1/1	97/0/1/1	96/0/2/1	97/0/1/1	98/0/1/1
South America Butterflies	89/0/4/7	91/1/2/6	92/0/1/6	92/0/1/6	93/0/0/6	93/0/0/6	94/0/0/6	94/0/0/6	94/0/0/6	94/0/0/6
Germany EPT	92/0/1/7	93/0/1/5	94/0/1/6	94/0/0/6	94/0/0/6	93/0/0/6	94/0/0/5	94/0/0/6	94/0/0/6	94/0/0/6
Northwest Pacific Molluscs	69/2/15/14	75/2/10/13	74/2/10/13	73/2/12/12	73/2/12/12	77/2/8/12	77/2/9/12	76/3/9/12	77/3/8/12	78/4/6/12
Amazon Moths	62/2/1/36	63/3/1/34	63/3/1/34	62/2/1/35	63/2/1/35	63/1/1/35	63/2/1/34	63/1/1/35	63/1/1/35	63/2/1/34
North America Birds	77/1/11/11	72/2/15/10	73/2/15/11	81/2/7/10	80/1/9/10	83/2/5/11	81/2/6/11	84/2/3/10	85/2/3/10	85/2/2/11
French Guiana Earthworms	96/0/0/4	97/0/0/3	97/0/0/3	96/0/0/4	96/0/0/4	97/0/0/3	97/0/0/3	97/0/0/3	97/0/0/3	96/0/0/4
Pakistan Lepidoptera	88/1/3/8	90/1/1/8	90/1/1/8	91/1/1/8	91/1/0/8	91/0/0/8	91/0/0/8	91/0/0/8	91/0/0/8	91/0/0/8
Tanytarsus	81/1/5/12	83/2/5/11	82/2/5/12	86/2/2/11	85/2/3/10	83/2/3/12	86/2/3/9	85/2/2/10	86/2/1/11	87/2/1/11
North Europe Tachinidae	64/2/14/19	65/3/16/17	65/3/15/17	66/3/13/18	65/3/15/17	67/3/11/18	68/3/12/17	68/3/10/18	69/3/10/18	71/4/8/17
Congo Fish	80/1/10/9	78/1/11/9	79/1/11/10	84/2/6/7	86/2/5/7	85/2/5/8	86/2/4/8	86/3/4/7	86/3/4/7	88/3/2/7
North America Pyraustinae	93/1/4/3	94/1/4/1	96/1/2/1	96/1/1/2	97/0/0/2	97/0/0/3	98/1/0/2	97/0/0/3	97/0/0/2	98/2/0/2
Ecuador Chrysomelidae	74/0/1/25	76/0/1/23	76/0/1/23	76/0/1/23	76/0/1/24	75/0/1/24	76/0/1/23	75/0/1/24	75/0/1/24	75/0/1/24
Iberia Butterflies	77/0/22/1	85/0/15/0	86/0/13/0	87/0/12/0	89/0/10/0	93/0/6/0	94/1/5/0	94/0/5/0	96/0/4/0	97/1/3/0

Figure 1. Position of the mini-barcode with established primers in this study.

Figure 2. Performance of mini-barcodes along a sliding window (100, 200, 300-bp).

Mini-barcode position is indicated on the x-axis and congruence with morphology on

the y-axis. mOTUs were obtained with Objective Clustering (2%), ABGD P=0.001

prior), and PTP. Each line represents one data set while the boxplots summarise the

values across datasets. Significant deviations from the results obtained with full-

947   length barcodes are indicated with color-coded asterisks (* = p < 0.05; ** = p <

948   0.001; red = poorer and green = higher congruence with morphology).

Objective clustering

Length	94	130	145	164	189	295	307	313	407	657	
Midpoint	257	65	73	243	213	405	154	502	455	329	Average
Great Barrier Reef Fish	94	91	92	95	95	97	94	97	97	97	95
South China Sea Fish	94	95	96	98	88	95	93	95	95	97	95
North Sea Molluscs	91	86	87	94	91	94	94	93	93	95	92
Canada Echinoderms	81	89	85	91	93	91	91	91	94	95	90
Ecuador Geometridae	78	70	73	85	98	99	99	99	99	99	90
Set Mean:			92	Set Range:			70	-	99		
Germany Aranea & Opiliones	83	89	88	90	84	84	85	84	85	85	86
European Marine Fish	87	88	89	90	81	83	85	83	83	84	85
South America Butterflies	72	74	73	77	90	92	93	92	93	93	85
Germany EPT	78	85	84	85	85	85	82	84	85	85	84
Northwest Pacific Molluscs	88	87	90	91	83	78	77	77	78	78	83
Set Mean:			84	Set Range:			72	-	93		
Amazon Moths	69	77	76	79	91	89	92	84	85	87	83
North America Birds	84	77	77	82	77	83	83	83	83	84	81
French Guiana Earthworms	77	85	84	90	78	74	75	78	77	84	80
Pakistan Lepidoptera	52	74	74	54	90	91	90	91	91	91	80
Tanytarsus	81	81	82	81	63	61	62	63	61	65	70
Set Mean:			79	Set Range:			52	-	92		
North Europe Tachinidae	47	51	49	57	77	80	82	82	80	81	68
Congo Fish	63	61	60	68	70	65	65	65	65	60	64
North America Pyraustinae	54	54	55	55	57	72	66	70	71	70	63
Ecuador Chrysomelidae	61	56	64	69	63	58	59	55	56	57	60
Iberia Butterflies	51	55	56	59	56	55	55	55	55	55	55
Set Mean:			62	Set Range:			47	-	82		

ABGD

Length	94	130	145	164	189	295	307	313	407	657	
Midpoint	257	65	73	243	213	405	154	502	455	329	Average
North Sea Molluscs	94	90	92	94	94	89	90	89	86	80	90
South China Sea Fish	92	78	83	89	93	94	92	93	91	90	89
Great Barrier Reef Fish	79	75	79	90	89	97	92	95	95	94	88
Canada Echinoderms	87	78	76	91	91	87	91	95	92	89	88
Pakistan Lepidoptera	61	67	75	84	89	95	91	95	96	94	85
Set Mean:			88	Set Range:			61	-	97		
Germany EPT	85	84	85	86	87	83	85	84	78	67	82
Ecuador Geometridae	48	58	57	96	70	99	99	99	99	97	82
South America Butterflies	60	70	77	84	84	89	91	91	91	81	82
European Marine Fish	80	72	72	85	86	83	84	80	78	69	79
North America Birds	61	67	73	74	76	81	83	83	84	87	77
Set Mean:			80	Set Range:			48	-	99		
Amazon Moths	72	43	42	88	75	87	90	90	91	89	77
North Europe Tachinidae	71	60	66	77	80	77	84	82	82	84	76
French Guiana Earthworms	78	89	93	81	81	73	80	70	59	55	76
Northwest Pacific Molluscs	76	75	75	77	77	76	77	76	76	63	75
Germany Aranea & Opiliones	76	18	79	85	82	71	74	73	65	52	67
Set Mean:			74	Set Range:			18	-	93		
Congo Fish	68	44	55	67	71	70	63	67	71	65	64
Tanytarsus	67	71	71	61	62	50	56	48	47	45	58
North America Pyraustinae	25	29	41	67	76	70	74	67	58	59	57
Ecuador Chrysomelidae	55	55	54	56	56	57	57	55	57	58	56
Iberia Butterflies	26	36	42	53	57	60	65	59	55	52	50
Set Mean:			57	Set Range:			25	-	76		

PTP

Length	94	130	145	164	189	295	307	313	407	657	
Midpoint	257	65	73	243	213	405	154	502	455	329	Average
Great Barrier Reef Fish	89	85	87	91	91	95	90	95	96	96	91
Pakistan Lepidoptera	95	83	82	87	90	95	95	95	95	94	91
Ecuador Geometridae	14	89	91	94	97	98	97	96	99	98	87
North Sea Molluscs	78	71	82	94	96	86	92	86	93	93	87
South China Sea Fish	86	75	77	87	88	92	89	89	92	92	87
Set Mean:			89	Set Range:			14	-	99		
Canada Echinoderms	75	71	78	83	90	89	90	87	91	89	84
Germany EPT	82	80	81	81	83	79	83	82	79	80	81
North Europe Tachinidae	77	70	73	82	80	84	83	83	82	85	80
Amazon Moths	32	88	38	89	88	91	89	91	90	90	79
Germany Aranea & Opiliones	72	73	77	77	79	82	80	77	81	79	78
Set Mean:			80	Set Range:			32	-	91		
North America Birds	72	67	66	77	78	81	82	81	85	85	77
French Guiana Earthworms	71	83	69	85	80	81	62	78	75	69	75
European Marine Fish	71	60	61	71	78	80	74	79	79	80	73
Northwest Pacific Molluscs	72	67	68	75	74	72	74	76	77	78	73
Congo Fish	57	44	50	67	69	68	64	68	71	69	63
Set Mean:			72	Set Range:			44	-	85		
Tanytarsus	51	58	64	61	58	68	63	64	66	63	62
North America Pyraustinae	45	58	54	56	49	58	71	80	76	58	61
Ecuador Chrysomelidae	55	57	55	56	55	55	54	55	55	55	55
Iberia Butterflies	32	47	48	47	48	59	58	61	58	54	51
South America Butterflies	0	0	0	0.5	0	89	90	93	91	91	45
Set Mean:			55	Set Range:			0	-	93		

Figure 3. Match ratios across three different species delimitation methods. Mini-barcodes (columns) are sorted by primer length while the datasets (rows) are grouped into 4 classes according to average match ratio. Colours are applied separately to each class.

Figure 4. Comparison of species delimitation methods for full-length and mini-

barcodes generated by "sliding windows" (100-bp, 200-bp, 300-bp).

Figure 5. Proportion of conserved sites along the full-length barcode (sliding

windows of 100-bp, 200-bp, 300-bp).