# Impaired speed encoding is associated with reduced grid cell periodicity in a mouse model of tauopathy

**Thomas Ridler [1], Jonathan Witton [2], Keith G. Phillips [3], Andrew D. Randall [1], Jonathan T. Brown* [1]**

1.  Institute of Biomedical and Clinical Sciences,
    University of Exeter Medical School,
    University of Exeter,
    Hatherly Laboratories,
    Prince of Wales Road,
    Exeter, EX4 4PS, UK

2.  School of Physiology, Pharmacology and Neuroscience,
    University of Bristol,
    University Walk,
    Bristol, BS8 1TD, UK

3.  Lilly UK
    Erl Wood Manor
    Windlesham,
    Surrey, GU20 6PH, UK

*Corresponding author: Dr Jonathan Brown, Institute of Biomedical and Clinical Sciences, University of Exeter Medical School, College of Medicine and Health, Hatherly Laboratories, Prince of Wales Road, Exeter, EX4 4PS, UK, Tel: +441392725504, Email: J.T.Brown@exeter.ac.uk.

**Abstract**

Dementia is associated with severe spatial memory deficits which arise from dysfunction in

hippocampal and parahippocampal circuits. For spatially-sensitive neurons, such as grid cells,

to faithfully represent the environment these circuits require precise encoding of direction and

velocity information. Here we have probed the firing rate coding properties of neurons in

medial entorhinal cortex (MEC) in a mouse model of tauopathy. We find that grid cell firing

patterns are largely absent in rTg4510 mice, while head direction tuning remains largely intact.

Conversely, neural representation of running speed information was significantly disturbed,

with smaller proportions of MEC cells having firing rates correlated with locomotion in rTg4510

mice. Additionally, the power of local field potential oscillations in the theta and gamma

frequency bands, which in wildtype mice are tightly linked to running speed, was invariant in

rTg4510 mice. These deficits in locomotor speed encoding likely severely impact path

integration systems in dementia.

**Introduction**

Accurate spatial navigation requires the integration of sensory information to generate neural representations of space. Various high-level representations of the external environment are expressed at a single cell level within the extended hippocampal formation (hippocampus proper, dentate gyrus, entorhinal cortex and subiculum) and connected brain areas, regions well known to be critical for spatial memory. For example, place cells in the hippocampus fire action potentials in specific spatial locations[1], whilst grid cells in the medial entorhinal cortex (MEC) fire in a highly organised, hexagonally distributed spatial pattern across an environment[2,3]. These directly spatially-sensitive neurons are collocated within the MEC with other functionally defined cell types, including head-direction[4–7], speed[8,9] and border cells[10,11]. Together these are thought to provide the crucial computational information required for effective path integration, the process of using idiothetic cues to continuously calculate update positional information[12].

Patients with Alzheimer's disease (AD) and other dementia-spectrum disorders exhibit profound disruption in spatial navigation and memory, even at very early stages of the diseas[13–17]. At a pathological level, misfolded tau deposition typically occurs first in the entorhinal cortex and hippocampus[18]. Taken together, these clinical signs strongly implicate pathology-induced circuit-level dysfunction in the hippocampal formation as a key early-stage functional deficit in AD. In this regard, there is substantial evidence from transgenic mouse models that dementia pathologies, such as β-amyloid deposition (Aβ) and hyperphosphorylation and misfolding of tau can disrupt the intrinsic properties[19–22] and synaptic microcircuits[23–25] of pyramidal cells in CA1 of hippocampus. Furthermore, there is growing evidence that place cells in the CA1 region of APP and tau overexpressing mice have reduced spatial-sensitivity[19,26,27], strongly suggesting a failure of some aspects of the upstream functional circuits involved in spatial cognition. Recent evidence suggests that the circuits required for generation of theta and gamma frequency oscillations in the dorsal entorhinal cortex are especially prone to dysfunction in a mouse model of tauopathy (rTg4510 mice)[28].

Furthermore, Fu *et al.*, (2017) showed that grid cell spatial periodicity is reduced in a mouse

in which tau overexpression is restricted to the entorhinal cortex. These data correlate with

human imaging studies which suggest deficits in grid cell-like activity in the entorhinal cortices

of people at genetic risk of developing AD[30].

Importantly, the effect of tau pathology on other functional components of the spatial

navigation system is not known. In this study we report for the first time that disruption to speed

encoding in the MEC of rTg4510 mice may underlie deficits in grid cell function. Using high-

density silicone probes and tetrode recording approaches in freely moving animals, we report

that in both the medial entorhinal cortex and the CA1 region of the hippocampus, encoding of

speed information at both the local field potential and cellular level is substantially impaired.

We propose a model whereby such deficiencies in speed encoding networks result in loss of

grid cell firing patterns in the MEC.

**Results**

*Tau pathology is associated with loss of oscillatory speed coding*

Neural coding of spatial information is likely to require the precise representation of locomotor

speed[12,31–33]. Velocity information can be represented in the brain via the dynamic regulation

of the power and frequency of neuronal network oscillations in the theta and gamma frequency

bands[34–38]. We hypothesised that the profound deficits in spatial memory that occur in

response to tau pathologies[19,29,39] arise from impaired representation of locomotor speed in

the hippocampal formation, ultimately leading to deficits in the encoding of spatial information.

To examine this hypothesis we implanted multi-site silicone-based recording probes in the

dorsal MEC of male wildtype (WT) and rTg4510 mice (6-7 months). Following a post-surgical

recovery period, mice were encouraged to explore a familiar linear track, baited by food

rewards at either end to encourage running between the two ends, whilst connected to a multi-

channel electrophysiological recording apparatus via a lightweight tether cable.

As expected, in WT mice, theta oscillation properties followed changes to locomotor activity

(fig 1A/B). Pooled data illustrate a clear relationship between running speed and theta

oscillation power (linear regression: $R^2 = 0.71$, p<0.001, n=7 mice) and frequency (linear

regression: $R^2 = 0.69$, p < 0.001, n = 7 mice) (fig 1Ci). In contrast, in rTg4510 mice, theta

oscillation amplitude was poorly correlated with locomotor activity and remained at consistent

levels throughout recording sessions (linear regression: $R^2 = 0.15$, p = 0.02, n = 8 mice, fig

1Ci). Importantly the overall Z-transformed correlation coefficient and slope relationships for

theta power-running speed relationships were significantly lower in rTg4510 mice when

compared to WT controls (2-way repeated measures ANOVA, see Table 1 and Figure 1D).

Although less pronounced than in WT animals, peak theta frequency was correlated with

running speed in rTg4510 mice (linear regression: $R^2 = 0.74$, p < 0.01, n = 8 mice, Figure 1Cii)

which, across the population was not significantly different from WT (Correlation (Z'); WT: 0.84

± 0.24, rTg4510: 0.40 ± 0.14, unpaired T-test, p = 0.13, n = 7/8 mice), with no difference in the

slope of the frequency-running speed relationship (Correlation (Hz/cms$^{-1}$'); WT: 0.045 ±

0.0077, rTg4510: 0.037 ± 0.011, unpaired T-test, p = 0.53, n = 7/8 mice) .

Similar running speed modulation has been observed in both in the fast (60-120 Hz) and slow

(30-50 Hz) gamma frequency bands[36,37]. In WT mice, both fast and slow gamma oscillation

power was positively correlated with running speed (linear regression; fast gamma: $R^2 = 0.90$,

p < 0.001, n = 7 mice; slow gamma: $R^2 = 0.17$ p = 0.013, n = 8 mice), although the slope of

this association was greater for fast gamma frequencies (slow gamma: 27.1 ± 8.4 mV$^2$/Hz.cm$^{-1}$

$^{1}$, fast gamma: 80.6 ± 19.2 mV$^2$/Hz.cm$^{-1}$; 2-way repeated measures ANOVA; Table 1 and

Figure 1D). rTg4510 mice did not show significant correlations for fast gamma frequency band

oscillations (linear regression; fast gamma: $R^2 = 0.03$, p = 0.67, n = 8 mice, Figure 1D), but did

for slow gamma frequencies ($R^2 = 0.43$, p < 0.01, n = 8 mice, Figure 1D). However, compared

to the WT population, this correlation was significantly lower only in the higher gamma

frequency range (2-way repeated measures ANOVA, Table 1 and Figure 1D, n = 7/8 mice).

*Loss of grid rhythmicity in rTg4510 mice*

Impairments to the theta oscillatory code for running speed may impair grid cell rhythmicity.

To establish whether this was the case, we used high density silicone probes to isolate a total

of 279 single-units in layer II/III of the dorsal MEC from 10 mice (WT:150 units from 5 mice,

rTg4510: 129 units from 5 mice). In this study we focused on the dorsal MEC, a subregion

which has previously been identified as being particularly vulnerable in this transgenic model[28].

By performing post-hoc electrolytic lesions on each recording shank to identify probe location,

we were able to estimate individual unit location by determining the largest average waveform

along the 200 µm probe for each isolated unit. Importantly, estimated recording location was

not different between genotypes (Median distance from dorsal entorhinal border (interquartile

range (IQR)): WT: 350(350) µm, rTg4510: 385(275) µm, U = 9660, p = 0.77, Mann-Whitney U

test; Figure S1A).

Considered across the entire cellular population, MEC single units recorded from rTg4510

mice exhibited a small decrease in mean firing rate (Median (IQR): WT: 2.97 (16.7) Hz,

rTg4510: 1.03 (3.0) Hz, U = 4472, p = 0.01, Mann-Whitney U, n = 150; 129 units, Figure S1E).

As in the CA1 region of the hippocampus[19], rTg4510 cells also showed a pronounced

decrease in theta modulation of firing patterns (Median (IQR) theta modulation index: WT:

9.17 (13.9), rTg4510: 2.29 (2.81), U = 2296, p < 0.001, Mann-Whitney U, n = 150;129 units,

Figure S1C). Individual units with a theta modulation index (TMI) greater than 5[40] were

assigned as 'theta modulated'. While the majority of WT MEC cells (65%) showed significant

theta modulation, only a small proportion (13%) passed threshold in rTg4510 mice ($\chi^2$ (1) =

64.1, p < 0.0001, Chi-Square test, Figure S1D). Extracellularly-recorded spike waveforms from

interneurons in the mEC have a significantly narrower spike widths when compared to

excitatory neurons[41]. The probability distributions of single units spike widths recorded from

the dorsal MEC in WT mice had a bimodal distribution (Figure S1F). Based on this distribution

and previously published work[41], we classed units with spike widths <0.35 ms as putative

interneurons and those >0.35 ms as putative excitatory cells. Using this approach we found

that 47/150 units (31%) were classed as putative interneurons in WT mice, where as in

rTg4510 mice a significantly smaller proportion of units were classed as putative interneurons

(8/129 units, 15%; $\chi^2$ (1) = 16.0, p < 0.001, Chi-Square test; Figure S1G).

Consistent with the literature we found that ~1/4 of cells in the WT MEC had grid-like spatial

firing patterns in a 0.8 m x 0.8 m square arena. We calculated a grid score for each cell, based

on the rotational symmetry of the 2D-autocorrelations and found that 36/150 cells (24%) had

grid scores higher than the 95[th] percentile of the distribution produced from shuffled spike

timestamps (threshold = 0.21; Figure 2B). In contrast, in rTg4510 mice there was an almost

complete breakdown of grid cell periodicity (3/129 units, 2.3%, threshold: 0.26), with animals

displaying irregular, non-uniform, firing fields across recording environments ($\chi^2$ (1) = 27.1, p

< 0.0001, Chi-Square test, Figure 2A/C). Furthermore, the distribution of grid scores of all cells

recorded from WT and rTg4510 mice was significantly different (Median (IQR) grid score: WT,

0.03 (0.3); rTg4510, 0 (0.12), n = 150/129 units; P < 0.001; Mann-Whitney U).

Conversely, when measuring the spatial information content (SI) of recorded cells[42], we found

that MEC neurons did not differ significantly between genotypes, either as a distribution (WT,

-0.05 (0.71), n = 150; rTg4510, 0 (0.24), n = 129 units; P > 0.05; Mann-Whitney U) or in the

proportion of cells that exceed threshold (WT: 10/150, rTg4510 6/129, $\chi^2$ (1) = 0.52, p = 0.47

Chi-Square test; data not shown).

*Tau pathology is associated with impaired speed coding in MEC single units*

Deficits in running speed-oscillation relationships at a local field potential level suggest

alterations in the neural representation of running speed in the MEC of rTg4510 mice. Recent

evidence suggests the existence of a separate population of cells in the MEC which express

a rate code for running speed ('speed cell')[8]. We calculated a speed score for all recorded

cells, whilst mice ran on an L-shaped track, by computing the Fisher-transformed correlation

coefficient (z) between instantaneous firing frequency and running speed (Figure 3A-B). Units

were considered 'speed-modulated' if they had a speed score outside the 5-95 centile range

of a shuffled distribution of data produced from 250 shuffles for each cell (Figure 3C). In WT

mice, 85/150 (57 %) cells had firing rates significantly modulated by running speed, whereas

in rTg4510 mice a significantly lower proportion of cells was speed modulated (17/129; 13%;

$\chi^2$ (1) = 56.55, p < 0.0001, Chi-Square test, Figure 3E). As an overall population, rTg4510

MEC units also displayed a significantly lower average speed score compared to WT mice

(Median (IQR): WT: z = 0.11(0.31), rTg4510: z = 0.009(0.01), U = 3603, p < 0.0001, Mann-

Whitney U, n = 150;129 units, Figure 3D), with cells recorded from rTg4510 mice, on average

displaying a running speed correlation close to zero.

Speed modulated MEC neurons can be broadly split into those that display linear and

saturating exponential relationships between running speed and firing rate[9]. To account for

this, speed scores were calculated for log transformed data and further classified as linear or

exponential by the best regression fit (Figure S2A). In WT animals speed-modulated cells

showed an approximately equal distribution between linear and saturating running speed

relationships (Figure S2B linear: 41/85 units, saturating: 44/85 units, which did not differ

significantly in the remaining rTg4510 MEC neurons (linear: 10/17 units, saturating: 7/17 units,

$\chi^2$ (1) = 0.6, p = 0.43 Chi-Square test, Figure S2B).

Previous studies have suggested that a small proportion of speed-modulated cells decreased

their firing frequency during locomotor activity[8,9]. On the linear track, in WT mice, this

population was observed to be consistent with previous reports (~13% of speed modulated

cells)[8,9]. In contrast, in rTg4510 mice the proportion of cells with negative speed relationships

was substantially and significantly higher (WT: 11/85 units, rTg4510: 7/17 units, $\chi^2$ (1) = 7.8,

p = 0.005, Chi-Square test, Figure S2C), meaning that speed modulated cells were split much

more evenly between positive and negative associations with firing rate. Given that many MEC

speed cells are fast spiking in nature[8], we also assessed the spiking properties the recorded

of speed-sensitive units. We found that in WT and rTg4510 mice, a similar proportion (41%

and 31%, respectively; $\chi^2$ (1) = 0.6, p = 0.4, Chi-Square test) of speed-modulated neurons

were fast-spiking (mean firing rate >10 Hz) (Figure S2E). The proportion of all cells that were

classified as fast-spiking was substantially lower in rTg4510 mice (13% vs 30% in WT; $\chi^2$ (1)

= 19.0, p <0.001, Chi-Square test; Figure S2E), although this likely reflects the overall

reduction in mean firing rate in these neurons (Figure S1E).

Speed modulated firing of single units has also been observed in the hippocampus[8,43,44], so

we next sought to determine whether deficits in speed tuning in rTg4510 mice were specific

to the MEC, or were also represented downstream in the hippocampus proper. For this

purpose, data were taken from previous single unit and local field potential recordings in the

hippocampal CA1 region of rTg4510 mice (Booth et al., 2016b) at a similar age point and

reanalysed to assess the contribution of locomotor activity to firing rate. In this region, CA1

theta band activity in the local field potential was also correlated with running speed in WT

mice (linear regression; theta power; WT: $R^2$ = 0.83, p < 0.001, n = 6 mice), but not in rTg4510

mice (linear regression: $R^2$ = -0.04, p = 0.6, n = 4 mice, Figure S3A). A significant proportion

of CA1 neurons had firing rates modulated by running speed in both WT and rTg4510 mice

(WT: 25/46 units, rTg4510: 27/52 units). Importantly however, as seen in MEC recordings, a

much greater proportion of CA1 cells were negatively modulated by locomotor activity in

rTg4510 mice than in WT controls (WT: 5/25 units, rTg4510:13/27 units; $\chi^2$ (1) = 4.5, p = 0.03,

Chi-Square test, Figure S3F).

*Normal head-direction tuning in rTg4510 mice*

Another critical component of the path integration system in the MEC are cells which

accurately encode directional heading information. Therefore, we next sought to establish

whether head direction cells[4,5] in the MEC were similarly disrupted in rTg4510 mice. We

calculated a head direction score (HD) by determining the mean vector length of circular firing

distributions and, as with other functional metrics, we compared observed HD scores with

shuffled distributions (Figure 4A&B). Importantly, a similar proportion of MEC cells in WT and

rTg4510 mice surpassed the 95th-centile threshold (WT: 19/150, rTg4510: 14/129, $\chi^2$ (1) =

0.79, p = 0.37, Chi-Square test, Figure 4C). Furthermore, as a population, the HD score was

actually slightly, but significantly, higher in rTg4510 cells compared to WT (Median (IQR)

vector length: WT: 0.074(0.091), rTg4510: 0.11(0.12), U = 6509, p = 0.0014, Mann-Whitney

U, n = 150;129 units, Figure 4B-E). These data suggest that head-direction tuning remains

intact in rTg4510 mice. Similarly, there was no change in the proportion of border cells

observed across genotypes (WT: 4/150, rTg4510: 6/129, $\chi^2$ (1) = 0.79, p =0.37, Chi-Square

test; data not shown), with relatively small numbers of cells passing threshold produced from

shuffled datasets. Neither was there a difference in the average border score between the two

populations (Median (IQR) vector length: WT: 0.097(0.67), rTg4510: -0.04(0.78), U = 6191, p

= 0.94, Mann-Whitney U, n = 150; 129 units; data not shown).

A proportion of cells within the MEC are known to demonstrate conjunctive functional

representation[5]. Using the tuning-curve approaches defined above, we found that ~60% of

WT cells passed the threshold for only a single functional class of neuron (i.e. grid-, speed- or

HD-cells). A further ~25% of WT cells passed the threshold for more than one functional class

and were therefore conjunctive cells, whilst the remaining ~20% of cells were unclassified

(Figure S4D). For example, in this study, only 3% of spatially sensitive cells were pure grid

cells, whilst the remaining grid cells also passed the threshold for speed- and HD-modulation

(Figure S4C). In contrast, due to the substantial reduction in numbers of speed and grid cells,

very few rTg4510 cells (~2%) expressed any conjunctive representation (Figure S4D).

| Correlation (Z) | | | | | |
|---|---|---|---|---|---|
| Source of Variation | Df | Sum-of-squares | Mean square | F | P |
| Frequency | 2 | 0.470 | 0.235 | 4.341 | 0.051 |
| Genotype | 1 | 0.602 | 0.602 | 5.927 | 0.038 |
| Interaction | 2 | 1.330 | 0.665 | 8.379 | 0.005 |
| Slope (mV2/Hz.cm$^{-1}$) | | | | | |
| Source of Variation | Df | Sum-of-squares | Mean square | F | P |
| Frequency | 2 | 0.004 | 0.002 | 8.973 | 0.004 |
| Genotype | 1 | 0.008 | 0.008 | 86.224 | >0.001 |
| Interaction | 2 | 0.005 | 0.003 | 16.623 | >0.001 |

**Table 1: Results of 2-Way repeated measure ANOVA for oscillation-running speed interaction.** Results of relevant *post-hoc* pairwise-comparisons displayed on Figure 1.

**Figure 1: Oscillation-running speed relationship is impaired in rTg4510 mice. A)** Local field potential from periods of slow (**i**) and fast (**ii**) running speed showing faster and larger theta oscillations during locomotor activity. **iii)** Power spectra of data shown in i and ii for slow (black) and fast (grey) running periods. **iv)** Example relationship between running speed and average theta oscillation power across recording session. **B)** Example plots showing animals running speed on linear track (black, left Y axis) showing high correlation with theta oscillation (grey, right Y axis) power (**i**) and frequency (**iii**) over several minutes of recording. Corresponding example from rTg4510 mouse showing with theta oscillation amplitude (**ii**) and frequency (**iv**) with decreased association with running speed. **C)** Running speed -theta oscillation relationships for power (**i**; normalised to 1-2 cm/s bin) and frequency (**ii**). Also shown are fast (**iii**) and slow (**iv**) gamma power-running speed relationships. **D)** Pooled data for each animal showing Z-transformed correlation coefficients (**i**) and slopes (**ii**) of running speed-oscillatory power relationships for different frequency bands (* p<0.05, ** p <0.01, *** p <0.001, ns = not significant, Bonferroni-corrected pairwise-multiple comparisons; for 2-way repeated measures ANOVA main effects and interactions see Table 1).

**Figure 2: Breakdown of grid cell periodicity in rTg4510 mice. A)** Example spatial firing patterns (left) and 2D autocorrelations (right) of cells from WT (**i**) and rTg4510 (**ii**) mice in a 0.8 x 0.8 m square arena, displayed with grid score (g), spatial information content (SI) and peak firing rate across recording environment. 5 cells with the highest grid score displayed for each genotype. **B)** Histograms of grid scores for all single units recorded from WT and rTg4510 mice. Observed data plotted as a solid line, shuffled data shown as grey bars. The 95[th] centile of shuffled distributions is plotted as a dashed line. **C)** Cumulative frequency plots for grid score; *inset* box plot illustrating average values for each genotype (dotted line: median, diamond: mean ± SEM, whiskers: 25[th]/75[th] centile), *** p<0.001, Mann-Whitney U test. **D)** Proportions of grid and spatial non-grid cells greater than threshold in WT and rTg4510 mice.

**Figure 3: Decreased speed modulation of MEC single units in rTg4510 mice.** Average waveforms of a single unit recorded from a 16 channel shank silicone probe, from a WT (**Ai**; black) and rTg4510 (**Bi**; blue) mouse. The running speed (grey) and cell firing rate (orange) (**ii**), correlation between running speed and firing rate for each time bin (40 ms) **(iii),** and average for each speed bin (1 cm/s) (**iv**) are shown for each of these example cells. Red line: linear fit for each. Scale bars: 0.3 ms, 25 μV. **C)** Distribution of speed scores for WT (**i**) and rTg4510 units (**ii**) with shuffled distribution of scores (grey bars); 5th/95th centile threshold: dotted lines. **D)** Average speed score for each recorded MEC unit. Box plots: dotted line: median, diamond: mean ± SEM, whiskers: 25th/75th centile), ** p<0.01, Mann-Whitney U test. **E)** Proportion of cells classified as speed modulated (>95th or <5th centile of shuffled distribution).

**Figure 4: rTg4510 mice retain MEC head-direction tuning. A)** Example head-direction tuning of 5 WT and 5 rTg450 units, displayed with head direction score (HD) and peak firing rate across head direction bins. **B)** Distribution of HD scores for WT (**i**) and rTg4510 mice (**ii**) with shuffled distribution of scores (grey), 95[th]-centile threshold: dotted line. **C)** Proportions of cells with head direction scores over threshold in WT and rTg4510 mice. **D)** Cumulative frequency distribution of HD scores (mean vector length). **E)** Mean vector length for all cells, showing a small but significant increase in head-direction tuning across the population in rTg410 mice (dotted line: median, diamond: mean ± SEM, whiskers: 25[th]/75[th] centile), * p<0.05 Mann-Whitney U test.

**Discussion**

Continuous integration of running speed information in the MEC has been proposed to be critically important for spatial navigation and path integration[5,8,31,45]. The current experiments clearly show that unlike WT animals, rTg4510 mice do not express adequate representations of locomotor activity in the MEC local field potential (LFP), since theta and gamma oscillations display blunted, or absent, relationships with running speed.

There has been some suggestion in the literature that dementia pathology affects the firing pattern of grid cells in the MEC both in mice[29] and grid-like neural representations in humans[30]. To date, however, no studies have examined changes to speed- or head direction-modulated firing of MEC single units in tauopathy. Our data suggest the hypothesis that reduced grid cell periodicity may be the result of the impaired integration of running speed information in the MEC.

*Vulnerability of grid cell firing*

These results highlight the importance of an intact path integration system in maintaining grid cell periodicity. Several studies have shown a breakdown of grid cell firing patterns after the inactivation of important spatial information streams. For example, inhibition of the medial septum, which controls theta rhythmicity, but also speed modulated inputs, produces a complete breakdown of grid cell periodicity[46,47]. Likewise, inactivation of the anterior thalamic nuclei (ATN) disrupts head direction tuning in the MEC, also impairing the grid cell signal[48]. Inactivation of reciprocal hippocampal inputs into the MEC is also sufficient to produce a breakdown in grid cell periodicity[49].

The almost complete breakdown of grid cell firing in rTg4510 mice contrasts with the effect on

spatial representation in the hippocampus of these animals. While several studies have shown

a reduction in the spatial information and stability of hippocampal place cells[19,26], firing fields

are still consistently present in these mice. Given the pattern of degeneration across the

hippocampal formation[39,50–52], poor place representation[19,26] may be the result of weakened

entorhinal inputs[53–55]. Since hippocampal place cells could be thought of as conjunctive

integrators of multiple spatial (and non-spatial) input streams, surviving inputs may facilitate

encoding of an impoverished spatial representation in these mice[19,26]. However, the loss of

grid/speed input prevents this from being anchored to an egocentric spatial reference frame,

resulting decreased spatial stability. Presumably, whatever representation is being encoded

by place cells is anchored to an allocentric reference frame. Decreasing the availability of

visual/contextual cues or implementing a pure path integration task would likely exacerbate

spatial learning/memory impairments. Reductions in grid and place field activity may therefore

mirror their appearance in neuronal development, where place cells appear before grid cells

but mature fully only after grid cell development[56,57]. The precise temporal nature of grid and

place cell impairment in this model is as yet unknown. However, taken together, these data

further suggest that hippocampal neurons can form place fields in the absence of effective

grid cell firing.

Grid cells are proposed to play a key role in path integration[31,32,58,59]. For example, mice lacking

GluA1-containing AMPA receptors have been shown to display reduced grid cell periodicity,

correlating with impairments on a path integration-based task[60]. Path integration has also been

suggested to be impaired in populations of dementia and MCI patients[13,15,17]. The direct

association between these two factors is still unclear; however, it is likely that grid cell deficits,

as described above, directly contribute to the deficits in memory in rTg4510 mice[19,50].

*Negative speed modulation of MEC neurons*

In WT animals, only a small population of MEC neurons display negative relationships with

running speed[8,9]. These data show that during recordings on the linear track, rT4510 mice

have a larger proportion of cells that decrease their firing rate with increases in running speed.

Interestingly, neurons in the medial septum also display a more even split between positive

and negative speed modulation[61], suggesting that, under normal physiological conditions, the

MEC does not simply recapitulate these firing patterns, but integrates this information

differently. The precise targets of negatively speed modulated inputs into MEC are unclear;

however it is possible that they are more likely to be conjunctive with other spatial information

systems, such as grid encoding or head direction. Indeed, negatively speed modulated cells

are more likely to be tuned to head direction[9], which is consistent with the observed increase

in mean vector length observed in rTg4510 neurons at a population level (Figure 4).

*Stability of head direction tuning*

A further feature of the inactivation studies described above[46–49] is the stability of head

direction tuning in the MEC. Blocking medial septum activity impairs grid cell firing, but not

head direction tuning[46,47]. Interestingly, the breakdown of grid periodicity after hippocampal

inactivation has also been shown to reveal head direction tuning in grid cells that would

previously not have been modulated by head direction[49]. It is possible that the small but

significant population increase in head direction tuning in rTg4510 mice may reflect this

unmasking of head direction inputs from cells that would have previously displayed grid

patterns. However, this is a hypothesis that would need to be addressed directly with long

term recordings of grid cells as tau pathology increased to critical levels to impair spatial firing

patterns. This approach is unlikely to be achieved with current technology. In any case, the

arrival of head direction information into the MEC is likely the result of an anatomically distinct

pathway[62] that appears to be relatively preserved in the face of  the tau pathology burden in

rTg4510 mice at this stage. This is most likely due to a greater dependence on subcortical

structures, such as the ATN, that integrate vestibular information[63–65].

*Impaired running speed representations in rTg4510 mice*

Theta oscillations show increases in both amplitude and frequency depending on running

speed (Figure 1). Spatially modulated MEC neurons are required to integrate large quantities

of multimodal sensory information from their environment. At fast running speeds, the time

window for this integration is smaller and it may therefore be necessary to increase the

sensitivity of such neurons during locomotion to accurately retrieve spatial associations from

memory. In rTg4510 mice, theta oscillation power appears to be independent of running speed

meaning that these animals are unable to integrate this information effectively.

In WT mice, the gamma oscillation power relationship with running speed shows a strong

predominance for fast gamma frequencies (Figure 1). This is consistent with the proposed

information flow across the hippocampal formation, in which the MEC provides the input

responsible for fast gamma frequencies in CA1 region of the hippocampus[66]. While the CA1

area has been shown to display two distinct peaks in the gamma band power spectra, MEC

LFP predominantly contains faster gamma frequencies only[66,67]. It is perhaps not surprising

then, that differences between genotypes (Figure 1) are only observed at these faster

frequencies. For this reason also, conclusions regarding the slow gamma oscillation frequency

should be made with caution. While slow gamma frequency has previously be shown to

display a negative speed relationship in the hippocampus, MEC correlations are generally

thought to be positive[36,68].

It is also worth noting that rTg4510 mice, like several other models of dementia, exhibit a

hyperactive phenotype[52,69–71], and hence altered patterns of locomotor activity. This may

reflect dysregulation of the septohippocampal pathway, activation of which has been shown

to stimulate movement or alternatively may reflect dysfunction in motor-related brain regions.

Nevertheless, whilst it should be noted that rTg4510 mice are experiencing their environment

at a generally higher state of arousal, at least in terms of locomotor activity, the key finding of

this study is that this locomotor activity is improperly represented in the hippocampal

formation.

*Conclusions*

Overall, these data show a clear breakdown in grid cell periodicity in rTg4510 mice compared

to WT controls. In addition, they suggest a role for the dysfunctional processing of locomotor

activity in this process, since the representation of running speed information in MEC single

units is severely disrupted in these mice, while head direction tuning remains constant, or

slightly increased. The changes to MEC single unit firing are likely to have profound

implications for the impairments in spatial memory observed in these mice and suggest

observable parameters to assess in dementia patient populations, for example through speed

modulated fMRI signals in virtual environments.

**Methods**

*Animals*

All procedures were carried out in accordance with the UK Animal (Scientific Procedures) Act 1986 and were approved by the Universities of Exeter and Bristol Animal Welfare and Ethical Review Body.

*Surgical implantation*

All surgical procedures were conducted using standard sterile and aseptic techniques. Animals were anaesthetized using isoflurane (4%) and fixed into a stereotaxic frame (ASI instruments). Anaesthesia was reduced and maintained at 1-2% during surgery. After careful cleaning of the skull surface, small screws (Antrin Miniature Specialities) were inserted into each bone plate in order to anchor the electrode array. Silver wire (World Precision Instruments) was soldered to a screw overlying the cerebellum to be used as a ground.

Probes were implanted at 0.2-0.3 mm anterior to the transverse sinus and 3-3.25 mm from midline. Linear probes were implanted and fixed 3 mm below the dura mater and angled at 10 degrees in the posterior to anterior direction in the sagittal plane in order to record consistently from layer II/III along the dorsal-ventral axis of the MEC. High density 16- (Neuronexus) or dual shank 32-channel (Cambridge Neurotech) silicone probes were implanted 0.3-0.5 mm below dura at an angle of 5 degrees, also in the posterior direction and subsequently moved slowly into the cortex using their attached microdrive (Cambridge NeuroTech). RelyX Unicem 2 dental cement with blue curing light (Henry Schein) were used to anchor the probe to the skull and anchor screws.

 *Data acquisition*

Animals were given at least 1 week of post-operative recovery before initial recording

sessions. Local field potential (LFP) signals were recorded using a Digital Lynx 10S recording

system (Neuralynx, Bozeman, MT, USA) tethered to a HS-18 or HS-36 unity gain headstage

and Cheetah 5 data acquisition software (Neuralynx). The headstage and tether were

counterbalanced using a moveable, weighted arm to allow for the maximum flexibility of

movement. Two light-emitting diodes (LEDs) on the headstage and an overhead video camera

(sample rate 25 Hz) were used to continuously track the animals' location using Cheetah's

built in video tracking software (VTS), allowing estimation of position and therefore running

speed. Once recorded, invalid tracking points, i.e. time-points where no light threshold was

reached, were excluded and the animal's position interpolated from the two nearest points.

Estimation of running speed was performed on binned position data, with erroneous bins,

above 50 cm/s, also removed. LFP data were recorded while animals explored either a linear

track (1.5 m) or square open field (0.8 m x 0.8 m).

*Analysis of local field potential (LFP) signal*

Data were continuously sampled at 2 kHz, band-pass filtered (1 – 500 Hz) and stored on a PC

for offline analysis. All LFP signals were analysed in MATLAB, using open-source toolboxes

or custom routines utilising built-in functions. Multi-tapered spectral analysis was performed

using the Chronux toolbox (available at http://chronux.org/).

Power and peak frequency of LFP frequency bands were compared to running speed. Spectral

analysis was conducted on 0.5 s bins of LFP data and compared to running speed calculated

from the same time frame. For running speed curves, locomotor activity was divided into 1

cm/s bins (between 1-30 cm/s) and oscillatory power and peak frequency averaged across all

relevant sections of data. Theta and gamma oscillation power was normalised to the power in

these frequency bands during non-movement, defined as speeds under 1 cm/s.

*Analysis of single unit data*

For single unit data, recordings were referenced to the ground electrode, continuously

sampled at 40 kHz, bandpass filtered between 1-30 kHz and saved unprocessed on a PC for

offline analysis. Each channel was referenced offline to a common-average of the opposite

16-channel shank (250 µm away) in order to eliminate signals common across the electrode

array such as noise and movement artefacts.  Extracellular spike activity was detected and

sorted using the klusta open source software package found at:

http://klusta.readthedocs.io/en/latest/[72].  Clusters were classified as either putative

interneurons or putative excitatory cells (pyramidal or stellate cells) by their spike half-width,

taken from the peak to the subsequent trough of the average extracellular waveform. While

the majority of cells recorded in the MEC are excitatory, a significant population can be

classified as inhibitory interneurons[41,73,74]. Using the average spike waveform, putative

interneurons were classified as displaying a spike-width less than 0.35 ms, based on the

extracellular properties of PV+ interneurons isolated optogenetically[41]. This approach was

taken alone, rather than in combination with average firing rare of neurons, since MEC

interneurons have been shown to vary widely in their spike frequency[41]. Neurons were

described by a theta modulation index (TMI), based on the fast Fourier transform (FFT) of

spike-train autocorrelations, using methods described previously[19,40,75]. Autocorrelations were

produced with ± 500 ms lags and 2 ms bin size. The peak at 0 lag was reduced to the next

maximal value and the entire function mean-normalised by subtracting the mean from all

values. The autocorrelation was tapered using a Hamming window to reduce spectral leakage

and FFT calculated. The power spectrum was calculated by dividing the square of the FFT by

the transform length ($2^{16}$, scaled to the length of the autocorrelation). TMI was defined as the

mean power within 1 Hz of each side of the peak in the theta frequency range (5-12 Hz) dived

by the mean power between 0 and 125 Hz. Cells were defined as 'theta modulated' if their

TMI was greater than 5.

*Analysis of speed modulated firing*

Speed modulation of single unit activity was calculated based on analysis described in Kropff

*et al.,* (2015). Running speed and firing rate of individual clusters were calculated for 40 ms

bins of data and smoothed across 500 ms using a Gaussian window function. Running speeds

from 2-30 cm/s and containing more than 0.5% of total recording duration were used for further

analysis. Speed modulation of cells was then defined by the correlation between all running

speed and firing rate bins and a speed score calculated using the Fisher-z transformation of

the correlation coefficient, r. Observed speed correlations were compared to a distribution of

randomly sampled correlations of shuffled data.  For shuffling, time stamps were forward-

shifted by a pseudorandom period between 20 s and the total trial length minus 20 s, with the

end of the trial wrapped to the beginning and reanalysed using the method above. Cells were

defined as 'speed modulated' if their speed score (z) was greater than the 95th percentile, or

less than the 5th percentile, of the global distribution of scores produced from at least 250

shuffled data sets for each unit (Figure 3).

*Analysis of head direction properties*

Head direction was determined by calculating the angle between two LEDs attached to the

animal's headstage. Time periods where neither, or only one, of the LEDs were observed

above threshold were discarded. Firing rate was calculated for 3º bins of head direction and

smoothed, using a Gaussian window over 14º.  A 'head direction score' was defined as the

resultant mean vector length, calculated from the smoothed firing rate maps. Observed mean

vector length was also compared to the 95th percentile of a distribution of shuffled data

produced as above.

 *Analysis of spatial firing properties*

Spike locations for each cell were obtained with a 2D histogram count, using the MATLAB

function *histcounts2*. Firing rate was calculated for 3 x 3 cm bins across recording

environments and smoothed using a 2D Gaussian function across 1.5 standard deviations.

'Gridness' was calculated using a 2D autocorrelation of smoothed firing rate maps[5]. Spatial

periodicity was determined by rotating autocorrelations in steps of 30°, between the central

peak and the 6 closest peaks, and correlating the rotated versions with the original. Grid score

was expressed as the difference between rotations at 30°, 90° and 150°, where if firing maps

show a hexagonal pattern give low correlations, and 60° and 120° where correlations will be

high.

The spatial information content (SI) of each cell was defined using the measure described by

Skaggs *et al.*, (1993) and expressed in terms of bits/spike. This approach measures the extent

to which a cell's firing rate can be used to predict the animal's location. By definition, this does

not assume spatial periodicity and has been used for quantifying place cell activity[19,27,55,76] as

well as spatially selective firing in the lateral EC[77].

*Hippocampal CA1 data*

Single unit and LFP data recorded from the CA1 pyramidal cell layer from previously published

datasets were additionally analysed for speed encoding (Booth *et al.*, (2016*b*)). These data

were collected using microdrives containing independently adjustable tetrodes, while animals

ran on a linear track. Running speed – firing rate relationships were determined using the

processing pathway described above.

*Electrode placement*

At the end of experiments, mice received a lethal overdose of sodium pentobarbital (Euthetal)

and electrolytic lesions were made at several electrode locations across the recording array.

Mice were then transcardially perfused with 4% v/v formaldehyde in 0.1 M phosphate buffered

saline (PBS). Brains were extracted from the skull and stored in 4% formaldehyde before being

cut in sagittal sections (50 μm) using a vibratome (Leica VT1000) and stained with cresyl

violet. The position of electrode sites was determined from digital pictures taken with a 2.5X

objective on a light microscope using QCapture pro 7 software (Qimaging). Probe electrode

location was expressed as distance from the most dorsal electrode site in layer II/III MEC.

**Acknowledgments**

This work was supported by an Alzheimer's Research UK Major Project Grant awarded to J.B. (ARUK-PG2017B-7). J.W. was an Alzheimer's Research UK Fellow (ARUK-RF2015-6). A.R. was a Royal Society Industry Fellow. K.P. is an employee of Eli Lilly and Company. Eli Lilly provided all the mice used in this study.

**Author Contributions**

Conceptualization, J.B., A.R., T.R; Methodology, K.P., T.R., J.W.; Investigation, T.R and J.W.; Writing – Original Draft, T.R. and JB.; Writing –Review & Editing, J.W., K.P., A.R., T.R. and J.B.; Funding Acquisition, J.B., J.W., A.R.; Resources, J.B., A.R., K.P.; Supervision, J.B., A.R.

**Declaration of Interests**

The authors declare no competing interests.

**References**

1. O'Keefe, J. & Dostrovsky, J. The hippocampus as a spatial map. Preliminary evidence from unit activity in the freely-moving rat. *Brain Res.* **34,** 171–175 (1971).

2. Fyhn, M., Molden, S., Witter, M. P., Moser, E. I. & Moser, M.-B. Spatial representation in the entorhinal cortex. *Science* **305,** 1258–64 (2004).

3. Hafting, T., Fyhn, M., Molden, S., Moser, M.-B. B. & Moser, E. I. Microstructure of a spatial map in the entorhinal cortex. *Nature* **436,** 801–6 (2005).

4. Giocomo, L. M. *et al.* Topography of head direction cells in medial entorhinal cortex. *Curr. Biol.* **24,** 252–62 (2014).

5. Sargolini, F. *et al.* Conjunctive representation of position, direction, and velocity in entorhinal cortex. *Science* **312,** 758–62 (2006).

6. Ranck, J. . Head-direction cells in the deep cell layers of dorsal presubiculum in freely moving rats. in *Society for Neuroscience Abstracts* **10,** (1984).

7. Valerio, S. & Taube, J. S. Path integration: how the head direction signal maintains and corrects spatial orientation. *Nat. Neurosci.* **15,** 1445–53 (2012).

8. Kropff, E., Carmichael, J. E., Moser, M.-B. & Moser, E. I. Speed cells in the medial entorhinal cortex. *Nature* **523,** 419–424 (2015).

9. Hinman, J. R., Brandon, M. P., Climer, J. R., Chapman, G. W. & Hasselmo, M. E. Multiple Running Speed Signals in Medial Entorhinal Cortex. *Neuron* **91,** 666–679 (2016).

10. Lever, C., Burton, S., Jeewajee, A., O'Keefe, J. & Burgess, N. Boundary vector cells in the subiculum of the hippocampal formation. *J. Neurosci.* **29,** 9771–7 (2009).

11. Solstad, T., Boccara, C. N., Kropff, E., Moser, M.-B. & Moser, E. I. Representation of geometric borders in the entorhinal cortex. *Science* **322,** 1865–8 (2008).

12. Barry, C. & Bush, D. From A to Z: a potential role for grid cells in spatial navigation. *Neural Syst. Circuits* **2,** 6 (2012).

13. Allison, S. L., Fagan, A. M., Morris, J. C. & Head, D. Spatial Navigation in Preclinical Alzheimer's Disease. *J. Alzheimer's Dis.* **52,** 77–90 (2016).

14. Lithfous, S., Dufour, A. & Després, O. Spatial navigation in normal aging and the prodromal stage of Alzheimer's disease: Insights from imaging and behavioral studies. *Ageing Res. Rev.* **12,** 201–213 (2013).

15. Hort, J. *et al.* Spatial navigation deficit in amnestic mild cognitive impairment. *Proc. Natl. Acad. Sci. U. S. A.* **104,** 4042–7 (2007).

16. Laczó, J. *et al.* Spatial navigation and APOE in amnestic mild cognitive impairment. *Neurodegener. Dis.* **8,** 169–77 (2011).

17. Mokrisova, I. *et al.* Real-space path integration is impaired in Alzheimer's disease and mild cognitive impairment. *Behav. Brain Res.* **307,** 150–158 (2016).

18. Pooler, A. M. *et al.* Propagation of tau pathology in Alzheimer's disease: identification of novel therapeutic targets. *Alzheimers. Res. Ther.* **5,** 49 (2013).

19. Booth, C. A. *et al.* Altered Intrinsic Pyramidal Neuron Properties and Pathway-Specific Synaptic Dysfunction Underlie Aberrant Hippocampal Network Function in a Mouse

Model of Tauopathy. *J. Neurosci.* **36,** 350–63 (2016).

20.   Tamagnini, F. *et al.* Hippocampal neurophysiology is modified by a disease-
associated C-terminal fragment of tau protein. *Neurobiol. Aging* **60,** 44–56 (2017).

21.   Brown, J. T., Chin, J., Leiser, S. C., Pangalos, M. N. & Randall, A. D. Altered intrinsic
neuronal excitability and reduced Na(+) currents in a mouse model of Alzheimer's
disease. *Neurobiol. Aging* **32,** 2109.e1-e2109.e14 (2011).

22.   Kerrigan, T. L., Brown, J. T. & Randall, A. D. Characterization of altered intrinsic
excitability in hippocampal CA1 pyramidal cells of the Aβ-overproducing PDAPP
mouse. *Neuropharmacology* **79,** 515–524 (2014).

23.   Hoover, B. R. *et al.* Tau Mislocalization to Dendritic Spines Mediates Synaptic
Dysfunction Independently of Neurodegeneration. *Neuron* **68,** 1067–1081 (2010).

24.   Brown, J. T., Richardson, J. C., Collingridge, G. L., Randall, A. D. & Davies, C. H.
Synaptic transmission and synchronous activity is disrupted in hippocampal slices
taken from aged TAS10 mice. *Hippocampus* **15,** 110–117 (2005).

25.   Fitzjohn, S. M. *et al.* Age-related impairment of synaptic transmission but normal long-
term potentiation in transgenic mice that overexpress the human APP695SWE mutant
form of amyloid precursor protein. *J. Neurosci.* **21,** 4691–8 (2001).

26.   Cheng, J. & Ji, D. Rigid firing sequences undermine spatial memory codes in a
neurodegenerative mouse model. *Elife* **2,** e00647 (2013).

27.   Cacucci, F., Yi, M., Wills, T. J., Chapman, P. & O'Keefe, J. Place cell firing correlates
with memory deficits and amyloid plaque burden in Tg2576 Alzheimer mouse model.
*Proc. Natl. Acad. Sci.* **105,** 7863–7868 (2008).

28.   Booth, C. A. *et al.* Electrical and Network Neuronal Properties Are Preferentially
Disrupted in Dorsal, But Not Ventral, Medial Entorhinal Cortex in a Mouse Model of
Tauopathy. *J. Neurosci.* **36,** 312–324 (2016).

29.   Fu, H. *et al.* Tau Pathology Induces Excitatory Neuron Loss, Grid Cell Dysfunction,
and Spatial Memory Deficits Reminiscent of Early Alzheimer's Disease. *Neuron* **93,**
533–541.e5 (2017).

30.   Kunz, L. *et al.* Reduced grid-cell–like representations in adults at genetic risk for
Alzheimer's disease. *Science (80-. ).* **350,** (2015).

31.   McNaughton, B. L., Battaglia, F. P., Jensen, O., Moser, E. I. & Moser, M.-B. B. Path
integration and the neural basis of the 'cognitive map'. *Nat. Rev. Neurosci.* **7,** 663–78
(2006).

32.   Burak, Y., Fiete, I. R., Trappenberg, T., Araujo, I. de & Wiggins, C. Accurate Path
Integration in Continuous Attractor Network Models of Grid Cells. *PLoS Comput. Biol.*
**5,** e1000291 (2009).

33.   Bush, D. & Burgess, N. A hybrid oscillatory interference/continuous attractor network
model of grid cell firing. *J. Neurosci.* **34,** 5065–79 (2014).

34.   Ahmed, O. J. & Mehta, M. R. Running Speed Alters the Frequency of Hippocampal
Gamma Oscillations. *J. Neurosci.* **32,** (2012).

35.   McFarland, W. L., Teitelbaum, H. & Hedges, E. K. Relationship between hippocampal
theta activity and running speed in the rat. *J. Comp. Physiol. Psychol.* **88,** 324–8
(1975).

36.   Zheng, C., Bieri, K. W., Trettel, S. G. & Colgin, L. L. The relationship between gamma

frequency and running speed differs for slow and fast gamma rhythms in freely
behaving rats. *Hippocampus* **25,** 924–38 (2015).

37. Chen, Z., Resnik, E., McFarland, J. M., Sakmann, B. & Mehta, M. R. Speed controls
the amplitude and timing of the hippocampal gamma rhythm. *PLoS One* **6,** e21408
(2011).

38. Sławińska, U. & Kasicki, S. The frequency of rat's hippocampal theta rhythm is related
to the speed of locomotion. *Brain Res.* **796,** 327–331 (1998).

39. Ramsden, M. *et al.* Age-dependent neurofibrillary tangle formation, neuron loss, and
memory impairment in a mouse model of human tauopathy (P301L). *J. Neurosci.* **25,**
10637–47 (2005).

40. Langston, R. F. *et al.* Development of the Spatial Representation System in the Rat.
*Science (80-. ).* **328,** (2010).

41. Buetfering, C., Allen, K. & Monyer, H. Parvalbumin interneurons provide grid cell–
driven recurrent inhibition in the medial entorhinal cortex. *Nat. Neurosci.* **17,** 710–8
(2014).

42. Skaggs, W. E., Mcnaughton, B. L., Gothard, K. M. & Markus, E. J. An Information-
Theoretic Approach to Deciphering the Hippocampal Code. *Neural Inf. Process. Syst.*
1030–1037 (1993).

43. McNaughton, B. L., Barnes, C. A. & O'Keefe, J. The contributions of position,
direction, and velocity to single unit activity in the hippocampus of freely-moving rats.
*Exp. Brain Res.* **52,** 41–49 (1983).

44. Lu, X. & Bilkey, D. K. The velocity-related firing property of hippocampal place cells is
dependent on self-movement. *Hippocampus* **20,** NA-NA (2009).

45. Burgess, N., Barry, C. & O'Keefe, J. An oscillatory interference model of grid cell
firing. *Hippocampus* **17,** 801–12 (2007).

46. Brandon, M. P. *et al.* Reduction of theta rhythm dissociates grid cell spatial periodicity
from directional tuning. *Science* **332,** 595–9 (2011).

47. Koenig, J., Linder, A. N., Leutgeb, J. K. & Leutgeb, S. The Spatial Periodicity of Grid
Cells Is Not Sustained During Reduced Theta Oscillations. *Science (80-. ).* **332,**
(2011).

48. Winter, S. S., Clark, B. J. & Taube, J. S. Disruption of the head direction cell network
impairs the parahippocampal grid cell signal. *Science (80-. ).* **347,** (2015).

49. Bonnevie, T. *et al.* Grid cells require excitatory drive from the hippocampus. *Nat.*
*Neurosci.* **16,** 309–317 (2013).

50. Santacruz, K. *et al.* Tau suppression in a neurodegenerative mouse model improves
memory function. *Science* **309,** 476–81 (2005).

51. Spires, T. L. *et al.* Region-specific dissociation of neuronal loss and neurofibrillary
pathology in a mouse model of tauopathy. *Am. J. Pathol.* **168,** 1598–607 (2006).

52. Blackmore, T. *et al.* Tracking progressive pathological and functional decline in the
rTg4510 mouse model of tauopathy. *Alzheimers. Res. Ther.* **9,** 77 (2017).

53. Hales, J. B. *et al.* Medial Entorhinal Cortex Lesions Only Partially Disrupt
Hippocampal Place Cells and Hippocampus-Dependent Place Memory. *Cell Rep.* **9,**
893–901 (2014).

54. Van Cauter, T., Poucet, B. & Save, E. Unstable CA1 place cell representation in rats
with entorhinal cortex lesions. *Eur. J. Neurosci.* **27,** 1933–46 (2008).

55. Brun, V. H. *et al.* Impaired Spatial Representation in CA1 after Lesion of Direct Input
from Entorhinal Cortex. *Neuron* **57,** 290–302 (2008).

56. Wills, T. J., Barry, C. & Cacucci, F. The abrupt development of adult-like grid cell firing
in the medial entorhinal cortex. *Front. Neural Circuits* **6,** 21 (2012).

57. Muessig, L., Hauser, J., Wills, T. J. & Cacucci, F. A Developmental Switch in Place
Cell Accuracy Coincides with Grid Cell Maturation. *Neuron* **86,** 1167–73 (2015).

58. Fuhs, M. C. & Touretzky, D. S. A Spin Glass Model of Path Integration in Rat Medial
Entorhinal Cortex. *J. Neurosci.* **26,** (2006).

59. Etienne, A. S. & Jeffery, K. J. Path integration in mammals. *Hippocampus* **14,** 180–
192 (2004).

60. Allen, K. *et al.* Impaired Path Integration and Grid Cell Spatial Periodicity in Mice
Lacking GluA1-Containing AMPA Receptors. *J. Neurosci.* **34,** (2014).

61. Justus, D. *et al.* Glutamatergic synaptic integration of locomotion speed via
septoentorhinal projections. *Nat. Neurosci.* **20,** 16–19 (2016).

62. Taube, J. S. The Head Direction Signal: Origins and Sensory-Motor Integration. *Annu.*
*Rev. Neurosci.* **30,** 181–207 (2007).

63. Stackman, R. W. & Taube, J. S. Firing properties of rat lateral mammillary single
units: head direction, head pitch, and angular head velocity. *J. Neurosci.* **18,** 9020–37
(1998).

64. Blair, H. T., Welday, A. C. & Zhang, K. Scale-invariant memory representations
emerge from moiré interference between grid fields that produce theta oscillations: a
computational model. *J. Neurosci.* **27,** 3211–29 (2007).

65. Sharp, P. E., Tinkelman, A. & Cho, J. Angular velocity and head direction signals
recorded from the dorsal tegmental nucleus of gudden in the rat: Implications for path
integration in the head direction cell circuit. *Behav. Neurosci.* **115,** 571–588 (2001).

66. Colgin, L. L. *et al.* Frequency of gamma oscillations routes flow of information in the
hippocampus. *Nature* **462,** 353–7 (2009).

67. Chrobak, J. J. & Buzsaki, G. Gamma Oscillations in the Entorhinal Cortex of the
Freely Behaving Rat. *J. Neurosci.* **18,** 388–398 (1998).

68. Kemere, C., Carr, M. F., Karlsson, M. P. & Frank, L. M. Rapid and continuous
modulation of hippocampal network state during exploration of new places. *PLoS One*
**8,** e73114 (2013).

69. Jul, P. *et al.* Hyperactivity with Agitative-Like Behavior in a Mouse Tauopathy Model.
*J. Alzheimer's Dis.* **49,** 783–795 (2015).

70. Cheng, D., Low, J. K., Logge, W., Garner, B. & Karl, T. Novel behavioural
characteristics of female APPSwe/PS1ΔE9 double transgenic mice. *Behav. Brain*
*Res.* **260,** 111–118 (2014).

71. Przybyla, M. *et al.* Disinhibition-like behavior in a P301S mutant tau transgenic mouse
model of frontotemporal dementia. *Neurosci. Lett.* **631,** 24–29 (2016).

72. Rossant, C. *et al.* Spike sorting for large, dense electrode arrays. *Nat. Neurosci.* **19,**
634–41 (2016).

73.　Beed, P. *et al.* Inhibitory gradient along the dorsoventral axis in the medial entorhinal
cortex. *Neuron* **79,** 1197–207 (2013).

74.　Miettinen, M., Koivisto, E., Riekkinen, P. & Miettinen, R. Coexistence of parvalbumin
and GABA in nonpyramidal neurons of the rat entorhinal cortex. *Brain Res.* **706,** 113–
122 (1996).

75.　Wills, T. J., Cacucci, F., Burgess, N. & O'Keefe, J. Development of the hippocampal
cognitive map in preweanling rats. *Science* **328,** 1573–6 (2010).

76.　Hussaini, S. A., Kempadoo, K. A., Thuault, S. J., Siegelbaum, S. A. & Kandel, E. R.
Increased size and stability of CA1 and CA3 place fields in HCN1 knockout mice.
*Neuron* **72,** 643–53 (2011).

77.　Deshmukh, S. S. & Knierim, J. J. Representation of non-spatial and spatial
information in the lateral entorhinal cortex. *Front. Behav. Neurosci.* **5,** 69 (2011).

 **Supplementary figures**

**Figure S1: Firing properties of mEC single units. A)** Example sections showing electrode lesions (red dot) of final probe location in mEC of WT and rTg4510 mice. **B)** Average waveforms from an example cell recorded from a 16 channel silicone probe shank for WT (**i**) and rTg4510 (**ii**) mouse, with firing autocorrelations. Scale bars: 0.4 ms, 50 µV. **C)** Cumulative frequency plot of theta modulation index for all recorded mEC single units, with average modulation for WT (black) and rTg4510 (blue) mice inset. **D)** Proportion of cells displaying theta modulation (threshold: TMI>5). **E)** Average firing frequency across entire recording session of mEC neurons for WT (black) and rTg4510 (blue) mice, average inset. **F)** Spike-width probability histogram for WT (black) and rTg4510 (blue) units. **G)** Proportion of cells classified as putative interneurons (spike-width <0.35 ms, dotted line in E) and putative excitatory. Box plots: dotted line: median, diamond: mean ± SEM, whiskers: 25[th]/75[th] centile), *** p <0.001 Mann-Whitney U test.

**Figure S2: Properties of MEC speed modulated cells. A**) Example negative (**i**), saturating **(ii)** and linear (**iii**) speed-modulated cells, with running speed (grey) and cell firing rate (orange), with average for each speed bin (1 cm/s) (right). Red line: linear/exponential fit for each. **B)** Proportion of speed modulated mEC cells that are best described by linear and saturating fits for WT and rTg4510 mice. **C**) Proportion of speed modulated units that show positive (>95[th] centile of shuffled distribution) and negative (<5[th] centile of shuffled distribution) speed modulation. **D)** Plot of speed score vs average firing rate, showing faster-spiking cells are more likely to have high speed modulation in WT, but not rTg4510 mice. **Ei)** Proportion of speed-modulated cells in WT and rTg4510 mice (replotted from Fig 3E). **ii)** Proportions of cells classified as fast-spiking (>10 Hz), with breakdown into speed and non-speed modulators for each genotype. **iii)** Proportion of fast spiking cells which display speed modulation, which shows decreased numbers in rTg4510 mice compared to WT. **iv)** Proportion of speed modulated cells which are classified as fast spiking shows only small decrease in rTg4510 mice.

**Figure S3: Speed modulation of single units in hippocampal CA1 pyramidal cell layer also shows similar increase in negatively speed modulated firing. A)** Normalised theta oscillation amplitude from WT and rTg4510 mice with increasing running speed, with average linear regression above (WT: $R^2 = 0.83$, p < 0.001, n = 6 mice; rTg4510: $R^2 = -0.04$, p = 0.6, n = 4 mice). **B)** Example speed-modulated cells recorded from the CA1 pyramidal cell layer. The cell on the left (i and ii) was positively speed-modulated, whilst the cell on the right (iii and iv) was negatively speed-modulated. **C)** histograms of speed scores for WT and rTg4510 mice. 5[th] and 95[th] centiles of shuffled data shown as dotted lines. **D)** box plot comparing speed scores between the two genotypes **E)** Proportion of cells passed criteria for speed modulated firing is approximately even between genotypes (WT: 25/46 units, rTg4510: 27/52 units). **F)** Increased proportion of negatively speed modulated units in CA1 in rTg4510 mice, compared to WT controls (WT: 5/25 units, rTg4510:13/27 units; $\chi^2$ (1) = 4.5, p = 0.03, Chi-Square test).

**Figure S4: Conjunctive representation of grid, head direction and running speed in WT and rTg4510 mice. A)** Example grid **(i)**, head direction **(ii)** and speed **(iii)** cells taken from the highest modulation scores from WT (top) and rTg4510 (bottom) mice, showing spatial firing patterns with 2D spatial autocorrelation, head direction tuning and running speed-firing rate relationship and corresponding score (SI: spatial information, HD: mean vector length, Z: speed score). **B)** Proportions of grid, head-direction and speed modulated cells recorded in mEC, showing reduced number of grid and speed, but not head-direction, cells passing threshold (95% centile of shuffled distribution). **C)** Conjunctive proportions of grid cells **(i)**, head direction cells **(ii)** and speed modulated cells **(iii)** recorded from WT and rTg4510 mice scaled to 100%. **D)** Breakdown of units from WT and rTg4510 mice that satisfied a single criteria (class), multiple criteria (conj) or no discernible firing pattern (other). Key, G: grid, S: speed, HD: head direction, G-S: grid-speed, G-HD: grid-head direction, G-S-HD: grid-speed-head direction, SnegHD: negative speed–head-direction, Sneg: negative speed, Class: classed as cell type, Conj: conjunctive representation (more than 1 classification), other: not classified as grid, speed or HD.