Sex-specific transcriptomic responses to changes in the nutritional environment

M. Florencia Camus[1], Matthew D.W. Piper[2] and Max Reuter[1]

[1] *Research Department of Genetics, Evolution and Environment, University College London, Gower Street, London, WC1E 6BT, United Kingdom*

[2] *School of Biological Sciences, Monash University, Clayton, Victoria, 3800, Australia*

Keywords: fitness, nutrition, reproduction, sexual dimorphism, nutritional plasticity

Abstract

Males and females typically pursue divergent reproductive strategies and accordingly require different dietary compositions to maximise their fitness. Here we move from identifying sex-specific optimal diets to understanding the molecular mechanisms that underlie male and female responses to dietary variation. We examine male and female gene expression on male-optimal (carbohydrate-rich) and female-optimal (protein-rich) diets. We find that the sexes share a large core of metabolic genes that are concordantly regulated in response to dietary composition. However, we also observe smaller sets of genes with divergent and opposing regulation, most notably in reproductive genes which are over-expressed on each sex's optimal diet. Our results suggest that nutrient sensing output emanating from a shared metabolic machinery are reversed in males and females, leading to opposing diet-dependent regulation of reproduction in males and females. Further analysis and experiments suggest that this reverse regulation occurs within the IIS/TOR network.

Introduction

Sex differences in life history, behaviour and physiology are pervasive in nature. These differences arise mainly from the divergent reproductive strategies between the sexes that are rooted in anisogamy [1]. Typically, males produce large numbers of small, cheap gametes and evolve traits that facilitate the acquisition of mates and the increase of fertilisation success. Females, on the other hand, produce fewer, energetically costlier gametes and tend to evolve traits that optimise rates of converting resources into offspring [2]. Given these fundamental differences between male and female reproductive investments, one of the key areas of divergence between the sexes concerns physiology, metabolism and responses to diet [3].

Studies in insect species [3-7] have shown that the two sexes require different diets to maximise fitness. Female fitness is typically maximised on a high concentration of protein, which fulfils the demands of producing and provisioning eggs. Males, in contrast, achieve optimal fitness with a diet consisting of more carbohydrate, which can fuel activities such as locating and attracting mates. Work on nutritional choices has shown that individuals tailor their diet in line with their physiological needs. In insects, females overall prefer diets with higher protein content, whereas males chose a more carbohydrate-rich diet [8, 9]. These choices are further adapted to reflect the individual's current condition and reproductive investment [9, 10]. For example, Camus et al. [11] found that the female preference for

protein in fruit flies was significantly higher in mated females (who require resources to

produce eggs) than virgins, while the preferences of males (who start producing sperm before

reaching sexual maturity) did not significantly differ between mated and virgin flies.

But individuals not only choose diets to suit their needs where possible, they also adapt

their physiology and reproductive investment in response to the quality and quantity of

nutrition available. This has been studied extensively using experiments that either alter the

macronutrients composition (carbohydrates vs. protein) of the diet while keeping the overall

caloric intent constant, or by manipulating the overall nutrient content of the food—dietary

restriction (DR). These studies have shown that a wide range of life history traits respond to

changes in both the composition of the food [7, 12, 13] and the quantity of nutrients supplied

[14-16]. For example, DR typically causes an extension of lifespan at the cost of reduced

reproduction [17], and a similar response can be triggered by a shift from protein to

carbohydrates in the diet [13].

Although most studies manipulating diet have concentrated on females only, those

including both sexes suggest that DR responses are broadly similar in males and females—

despite their large differences in optimal diet. In fruit flies, DR extends lifespan in both sexes

[18-20], even though the observed increase in longevity appears smaller in males than

females and the degree of DR that maximises lifespan can differ between the sexes [18].

Qualitatively similar results have been obtained when manipulating the macronutrient

composition of the diet. Studying field crickets, Maklakov et al. [5] found that shifting the

dietary balance away from protein and towards carbohydrates increased lifespan in both

sexes, even though the effect of nutrients on reproductive investment differed between the

sexes [5]. These quantitative sex differences in dietary lifespan effects can at least in part be

attributed to sex-biased responses in individual tissues. Thus, Regan and co-workers showed

that *D. melanogaster* males in which the gut had been genetically feminised had DR

responses more similar to those of females [15].

The contrast between large differences in optimal diet but similar responses to diet

manipulation raises the question of how males and females differ in their diet-dependent

regulation of metabolism and reproductive allocation. Due to the predominant focus on

female responses to nutrition, we currently know relatively little about the degree to which

regulation is shared or differs between the sexes [21], in particular at the molecular level.

Work in females has shown that nutrient-sensing pathways play a key role in the observed

DR phenotype [22-26]. Specifically, two evolutionarily conserved signaling pathways—

insulin/insulin-like growth factor 1 (IIS) and Target of Rapamycin (TOR)—are thought to

regulate longevity in a diet-dependent way [21, 27, 28]. Recent transcriptomic work in

female *D. melanogaster* has further shown that DR and rapamycin treatment (which inhibits

TORC1 activity) elicit similar changes in gene expression [29]. Both responses share a

significant number of overlapping genes, and are mediated by transcription factors in the

GATA family; in line with the involvement of these regulators in amino acid signaling and

lifespan modulation across eukaryotes [29].

While these data are starting to paint an increasingly detailed picture of nutrient-

dependent regulation in females, the lack of information on males severely limits our

understanding of how diet shapes metabolism and life history decisions. For example, it is

not clear to which degree the regulation identified in females reflects their specific dietary

requirements and physiology. Further, we cannot tell whether males and females differ in

their general metabolism and its nutrient-dependent regulation, or whether diet responses are

largely shared, and sex-specific effects limited to the regulation of reproductive investment.

Interestingly, perturbing the IIS/TOR network in virgin flies has been shown to elicit sex-

specific expression changes in males and females [30], but the link to nutrition and the effect

on reproductive investment remains unclear. Addressing these questions is important because

they have implications for the degree to which male and female physiology and its regulation

are uncoupled and able to independently evolve. Thus, a shared physiology and diet-

dependent regulation of metabolism across the two sexes would constrain the degree to which

each sex is able to independently optimise its life-history decisions in response to the current

nutritional environment.

Here, we are starting to address these fundamental questions by investigating male and

female diet responses in gene expression. We study this in the context of shifts of nutritional

composition (amino acid-to-carbohydrate ratio) between the male and female optima. This

manipulation is more subtle than classic dietary restriction, given we are changing the quality

of the diet whilst keeping caloric intake the same. This approach allows us to contrast, for

each sex, an optimal and a non-optimal condition, as well as, across sexes, a more amino

acid- and a more carbohydrate-rich diet. Furthermore, we can compare the female responses

to a smaller, more quantitative shift in diet composition to existing data on responses to DR.

We use nutritional geometry techniques to establish the male and female optimal diets in an

outbred *D. melanogaster* population and then examine the transcriptomic responses of both

sexes to the male-optimal diet (protein-to-carbohydrate ratio 1:4) and the female-optimal diet

(2:1). We then assess the degree to which expression changes from male- to female-optimal

diets are shared or divergent between the sexes, and how this relates to the function and

regulation of genes.

Our analysis reveals that most of the core metabolic gene network is shared between

the sexes, responding to diet changes in a sexually concordant manner. However, we also

find smaller sets of genes where male and female responses diverge, either by being restricted

to one sex or by males and females showing opposing diet-induced expression changes and

observe that sex-limited reproductive genes are generally up-regulated on each sex's optimal

diet. These results indicate that while males and females share a common, and concordantly

regulated metabolic machinery, the sexes diverge in how nutritional information is translated

into reproductive regulation. Further results allow us to link this divergent regulation to the

Tor pathway. First, we find that our genes with diet-dependent regulation overlap with genes

previously associated with responses to DR, rapamycin treatment and perturbation of the

IIS/TOR network and known targets of the TOR pathway. Second, we can show

experimentally that inhibiting TORC1 with rapamycin has a disproportionately negative

effect on reproductive fitness on each sex's optimal diet. These results are compatible with

the shared nutrient-sensing signal being inverted in males and females to produce

diametrically opposed Tor-dependent regulation of reproduction in the two sexes.

Materials and Methods

Fly Stocks and Maintenance

143 We used the *D. melanogaster* laboratory population LH_M for our experiments. This has been

144 sustained as a large outbred population for over 400 non-overlapping generations [31, 32],

145 maintained on a strict 14-day regime, with constant densities at larval (~175 larvae per vial)

146 and adult (56 vials of 16 male and 16 females) stages. All LH_M flies were reared at $25^{\circ}C$,

147 under a 12h:12h light:dark regime, on cornmeal-molasses-yeast-agar food medium.

Synthetic Diet

We used a modified liquid version of the synthetic diet described in Piper et al. [33], that is

prepared entirely from purified components to enable precise control over nutritional value

(see Table S1-S3). Previous studies have used diets based on natural components, typically

sugar as the carbon source and live or killed yeast as the protein source [34]. Such diets offer

only approximate control over their composition, because the yeast-based protein component

also contains carbohydrates and is required to provide other essential elements (vitamins,

minerals, cholesterol, etc.) that vary in relative abundance. As a consequence, phenotypic

responses to such diets cannot be straightforwardly interpreted in a carbohydrate-to-protein

framework as they are confounded by responses to other dietary components. Our use of a

holidic diet completely eliminates these problems without causing any apparent stress in the

flies [33].

Eight isocaloric artificial liquid diets were made that varied in the ratio of protein (P,

incorporated as individual amino acids) and carbohydrate (C, supplied as sucrose), while all

other nutritional components were provided in fixed concentrations. Nutrient ratios used were

[P:C] – 4:1, 2:1, 1:1, 1:2, 1:4, 1:8, 1:16 and 1:32, with the final concentration of each diet

(sum of sugar and amino acids) being 32.5g/L. These ratios span the P:C ratio of the

molasses medium on which the LH_M population is maintained. Based on the media recipe

used in our laboratory and the approximate protein and carbohydrate content of the

ingredients, we estimate our standard food to have a P:C ratio of about 1:8. The diets in our

experiments on the edges of our nutritional space, with the highest carbohydrate- or protein-

bias, can thus considered to be "extreme" in comparison to our standard laboratory media—

even taking into account the fact that ratios in synthetic and organic diets may not be directly

comparable, as nutrients in synthetic food appear to be more readily accessible [33].

For diet preference assay we used two diets; protein and carbohydrate. Each diet

contained all nutritional components (vitamins, minerals, lipids) at equal concentration, with

the protein diet containing amino acids and the carbohydrate diet containing sucrose.

Preliminary experiments established that flies would not eat purified amino acids with the

vitamin/mineral/lipid buffer, so we diluted our protein solution with 20% of a suspension of

dried yeast extract, made at the same protein concentration as the synthetic solution (16.25

g/L). Given that yeast extract also contains sugars, the final protein diet then included 4%

carbohydrate.

Experiment 1a: Identification of male and female optimal diets

<u>Experimental Setup and Diet Assay</u>

Flies from each sex were collected as virgins using CO_2 anaesthesia. Three virgin females

and three virgin males were randomly placed in individual vials containing culture medium

(molasses-yeast-agar) with no added live yeast. Twenty vials of hextets were collected for

each sex and diet treatment. Flies were left to mate for a period of 36 hours on molasses-

yeast-agar medium. Following this period, they were split by sex (now fly triplets), and

placed on 0.8% agar-water mixture. Agar-water vials provide water for the flies, but have no

nutritional value. Flies were kept in these vials overnight before being supplied with a 10µl

(females) or 5µl (males) microcapillary tube (ringcaps©, Hirschmann) containing one of the

eight allocated diets. These diets varied in their protein-to-carbohydrate ratios and captured

the following nutritional rails (P:C): 4:1, 2:1, 1:1, 1:2, 1:4, 1:8, 1:16, 1:32. Capillary tubes

were replaced daily, and food consumption for each fly trio was recorded for a total period of

four days. Consequently, our experiment design consisted of 2 sexes and 8 nutritional

environments, with 20 vials of fly triplets comprising each experimental unit (2 sexes × 8

diets × 20 vials = 320 vials, 960 flies). We chose to use capillary tubes of different sizes to

maximise the accuracy of our diet consumption measurements and minimise evaporation

errors. Larger capillary tubes increase evaporation rates; however, with a smaller capillary

tube we ran the risk that flies would consume all of the food leading to a subsequent slight

starvation response. For this reason, we found that a slightly larger capillary tube was ideal

for females because they ate more than males in a 24-hour period. Using this approach, we

found that flies never consumed all of the food from the capillary tubes. Flies were exposed

to diet treatments in a controlled temperature room (25°C), 12L:12D light cycle and high

relative humidity >80%. The rate of evaporation for all diet treatments was measured by

using five vials per diet that contained no flies, placed randomly in the constant temperature

chamber. The average evaporation per day was used to correct diet consumption for

evaporation. Following four days of feeding under these dietary regimes, flies were assayed

for fitness.

<u>Male Fitness Assay</u>

Male adult fitness was measured as the number of adult offspring produced in competitive

mating trials. Previous work in our laboratory has shown this to be a robust measure of

reproductive performance and, with lifetime adult production being largely determined by

mating success in our population [35].

We used an experimental approach similar to [36], whereby focal experimental males

competed with standard competitor males to mate with females. Following the experimental

feeding period described above, a focal trio of males was placed into a new vial (provided

with molasses-yeast-agar medium that did not contain live yeast, the main source of food for

both males and females [37, 38]), along with three virgin competitor males and six virgin

females. The competitor males and the females were of LH_M genetic background but

homozygous for the recessive bw^- eye-colour allele. Competitor flies were reared under the

same conditions as our experimental flies and were the same age as the experimental males.

The flies interacted, and female flies produced eggs for a period of 24 hours, after which the

adults were discarded from the vials. Eggs were left to develop for 12 days and the

subsequent adult offspring in each vial were counted and scored and assigned paternity to

either the focal experimental males (if the progeny had red eyes - wildtype) or the competitor

males (if the progeny had brown eyes).

<u>Female Fitness Assay</u>

Female adult fitness was measured as the number of eggs produced over a fixed period of

time. This performance proxy is expected to correlate closely with other fitness measures,

such as the total number of offspring [39, 40].

Following the feeding period, trios of mated females were placed in new agar vials and

presented with three males from the LH_M stock population. Flies were left to mate/oviposit

for 18 hours in vials containing *ad libitum* food corresponding to their diet treatment

provided via capillary tubes. All flies were removed after this 18-hour mating window.

Following removal of the flies, the total number of eggs laid were determined by taking

pictures of the agar surface and counting eggs using the software *QuantiFly* [41].

<u>Statistical Analyses</u>

First, we sought to investigate the effects of diet on sex-specific fitness. Separate models

were run for each sex, as the two datasets measured fitness in distinct ways. Female fitness

was measured as total number of eggs produced within a 18-hour timeframe following a

mating event. Given data followed a normal distribution, we used a linear model to analyse

the data. Number of eggs was the response variable, with mating status, and diet plus their

interaction as fixed factors. Male fitness was measured as the proportion of offspring sired

from the focal male. For this we modelled the response as a binomial vector comprising the

number of offspring sired by the focal male and the number sired by the competitor male and

diet composition as a categorical fixed effect. To examine whether the sexes varied in the

quantity they consumed of each diet, we used a linear model to investigate differences in

dietary consumption. We modelled total food consumption as a response variable with sex,

diet and their interaction as fixed effects. All models were performed using the *lm* function in

R version 3.3.2 [42].

To examine nutritional fitness landscapes, we combined fitness values with diet

consumption values for each sex. Before statistical analysis, we transformed the fitness data

as male and female fitness were measured in different units. For this, we standardised them

using Z-transformations for each sex across treatments. We used a multivariate response-

surface approach [43, 44] to estimate the linear and quadratic effects of protein and

carbohydrate intake on male and female fitness. The linear gradients for protein and

carbohydrate intake for each sex were estimated from a model containing only the linear

terms. The nonlinear gradients for nutrient intake were obtained from a model that contained

both linear and nonlinear terms. We used untransformed data to visualize nutritional

landscapes, using non-parametric thin-plate splines implemented with the *Fields* package in

R version 3.3.2 [42].

Experiment 1b: Dietary Preference Assay

Alongside the dietary setup used for measuring diet-dependant fitness, we tested what flies

preferred to eat, given the choice. For this, flies were supplied with two 5µl microcapillary

tubes (ringcaps©, Hirschmann); one containing the protein solution and the other the

carbohydrate solution. Capillary tubes were replaced daily, and food consumption for each

fly trio was recorded for a period of three days. As a control, the rate of evaporation for all

diet treatments was measured in six vials that contained the two solution-bearing capillary

tubes but no flies and placed randomly in the controlled temperature room. Their average

evaporation per day was used to correct diet consumption for evaporation.

<u>Statistical analysis</u>

To determine if male and female dietary choices differed between the sexes, we used a

multivariate analysis of variance (MANOVA). The main model had protein and carbohydrate

consumption as response variables, with sex as fixed effect. We performed subsequent

univariate analysis of variance (ANOVA) to determine which nutrient(s) contributed to the

overall multivariate effect. All analyses were performed using the *manova* function in R

version 3.3.2 [42]

Experiment 2: Transcriptional response

<u>Experimental Setup</u>

We followed the same experimental regime as previously stated, with the only exception of

using two diets instead of eight (Figure S1A). In brief, flies were collected in hextets; three

male and three female flies per vial, with 40 vials being setup. Following a period of 36 hours

where flies had the opportunity to mate, they were split by sex and placed onto agar medium

in triplets. Flies were allocated either a female-optimal diet (P:C=2:1) or a male-optimal diet

(P:C=1:4). Liquid food was provided using a 10ul capillary tube for females and a 5ul

capillary tube for males. Capillary tubes were replaced daily, and food consumption for each

fly trio was recorded for a total period of four days. Following this period, flies were flash-

frozen in their triplets.

We also set up 10 extra vials for each treatment alongside the RNA-Seq experiment

where we re-measured male and female fitness and preference. This was to verify the

repeatability of protocols for experiment 1 and 2.

Sample collection and RNA extraction

We generated 3 biological replicates for each of the experimental treatments (females on

female-optimal diet, females on male-optimal diet, males on female-optimal diet, males on

male-optimal diet), a total of 12 samples. For each replicate sample, we pooled 4 triplets (a

total of 12 flies) to ensure we collected sufficient amounts of RNA. Total RNA was extracted

using the Qiagen RNeasy Minikits (Qiagen BV, Venlo, The Netherlands). This kit includes

an on-column DNAse I digestion step. Quantity and quality of RNA was first inspected using

a Nanodrop 2000 spectrophotometer (Wilmington, USA), and later verified using an Agilent

Tapestation 2200 at the UCL Genomics facility.

Sequencing and read mapping

Library construction and sequencing were handled by the UCL Institute of Child Health

Genomics facility. cDNA libraries were constructed using the KAPA Hyper mRNA Library

prep kit. cDNA from all 12 libraries was mixed at equal concentrations and these multiplexed

samples were sequenced (43bp paired-end reads) on four flowcell lanes on an Illumina

Nextgen 500 instrument to an average of 18M reads per sample.

Having verified that there was no bias towards particular libraries across the sequencing

lanes using the Illumina Basespace online server, we merged reads from all four lanes.

Adaptors and low-quality base pairs were trimmed using trimmomatic v0.36 [45]. Trimmed

reads from each sample were independently mapped to the *Drosophila melanogaster* genome

release 6.19 using HISAT2 [46]. Mapped reads were manipulated using *samtools* [47].

Statistical analyses, identification of DE genes and enrichment analyses

Read counts for each annotated gene were performed using htseq-count [48], where reads are

counted at the exon level (using release 6.19 annotations obtained from the ENSEMBL

Biomart) and then summed across all exons within a single gene. Total read counts for each

gene for the 12 samples were then used for differential gene expression analysis using the

Bioconductor package edgeR [49] in R [42]. We first filtered read counts by expression and

removed lowly expressed genes. Read count data were normalised across libraries and

expression dispersion parameters calculated in edgeR using the entire dataset.

Subsequently, expression data was subsetted into three parts for separate analysis, i)

genes that were expressed in both sexes (transcripts detected in at least one replicate library

of each sex), ii) genes that were male-limited in expression (transcripts detected in at least

one replicate library from males, but none of the female libraries), and iii) genes that were

female-limited in expression (transcripts detected in at least one replicate library from

females, but none of the male libraries) (Figure S1B).

We tested for differential gene expression between our experimental groups using the

negative binomial models implemented in edgeR. For the shared gene dataset, we fitted a full

model where expression of each transcript is a function of sex, diet and their interaction. The

significance of each model term was tested using a specific contrast matrix. In order to obtain

estimates of expression fold changes between the two diets for each sex, we further fitted

models with diet as the sole model term separately to male and female data.

Gene ontogeny enrichment was performed using the Bioconductor package

clusterProfiler [50]. In order to assess whether genes that showed similar diet responses were

regulated by common transcription factors we used the Bioconductor package *RcisTarget*

[51], which tests for enrichment of cis-regulatory motifs upstream of a given gene sets. In all

analyses, we used a statistical significance threshold of 5% False Discovery Rate (FDR) [52].

For the smaller sex-specific gene sets, we ran enrichment analyses on the sets of genes with

significant diet responses, but also on the complete sets of sex-limited genes (irrespectively

of their responses to diet). This was to be able to identify (and remove) enriched binding

motifs that reflect general sex-specific regulation rather than diet responses.

We further compared our list of genes responding to diet (either via additive or

interactive effects) to previous work that has examined transcriptomic responses to dietary

restriction [53]. For this, we used the R package *GeneOverlap* [54] that implements a

contingency table test (Fisher's exact test) to identify greater than expected overlap between

gene lists. To compare our gene list to genes reported as significantly affected by

perturbation of IIS/TOR signalling by Graze *et al*. [30], we had to reanalyse their data using

our pipelines. This was required because their analysis was performed at the exon-level,

while we assessed transceription at the gene-level. We downloaded all raw data from the

SRA database (SRP137911). We aligned all reads to the same version of

the *Drosophila* nuclear genome we used for our analyses, and obtained gene-specific

expression patterns across all their samples. We then applied the same statistical framework

to these data that we had used for our own analysis, assessing the effect of sex, IIS

perturbation and their interaction to genes expressed in both sexes. Overlap between

classifications based on diet- and IIS/TOR perturbation-responses were assessed with X^2 tests

and only considered genes that showed male and female expression in both datasets

(N=8310).

Experiment 3: Fitness response to diet and rapamycin

Male and female flies were assayed for fitness in the same way as previously described for

Experiment 2. However, rather than just feeding either a protein-rich or a carbohydrate-rich

diet, we combined each of the two dietary treatments with one of four different

concentrations of the drug rapamycin (0μM, 5μM, 10μM, 50μM). Rapamycin is a drug that

very specifically inhibits TORC1, and hence TOR-signalling, with this function being highly

conserved from *S. cerevisiae* to humans [55]. Nutritional compositions and rapamycin levels

were combined in a full factorial design resulting in a total of eight different diets (two

nutritional compositions times four rapamycin levels, eight diets in total) for each sex. We

had approximately 20 vials for each experimental unit.

We performed a joint analysis on a dataset combining male and female fitness data.

Before statistical analyses, male and female fitness measures were transformed to obtain

normally distributed residual values. Female egg numbers were log-transformed, whereas

male competitive fertility data was arcsine-transformed. Moreover, to be able to compare

across sexes, male and female fitness measures were further centred and scaled (separately

for each sex) using Z-transformations. We fitted a linear fixed effects model to the

transformed fitness values with sex, diet and rapamycin concentrations (coded as a

categorical factor to accommodate possible non-linearity in the effect) and their interactions.

For the main analysis we categorised diet as optimal/non-optimal (where the nutritional

composition of the 'optimal' diet category is carbohydrate-rich for males and protein-rich for

females). This encoding makes it more straightforward to assess how rapamycin treatment

interacts with diet-quality in each sex. We also ran analysis where diet composition was

encoded as 'carbohydrate-rich' and 'protein-rich'.

Results

Dietary requirements and choice

We first examined the effects of diet composition on male and female fitness. We recovered previous results, finding that males and females differ significantly in their dietary requirements to maximise fitness (parametric bootstrap analysis: PB-stat = 78.002, $p < 0.001$). For females, the number of eggs produced differed significantly between diets (Analysis of Variance, $F_7 = 41.4703$, $p < 0.001$) and was maximised on the 2:1 (P:C) nutritional rail (Figures 1 and S3). Male competitive fertilisation success also differed between diets ($F_7 = 3.5927$, $p < 0.001$), but peaked at the 1:4 ratio (Figures 1 and S4). Dietary choices also differed between the sexes ($F_2 = 27.826$, $p < 0.001$). The choices of both sexes closely matched their previously established optimal composition, with females choosing to consume a more protein-rich diet than males (Figure 1). We also found that females, on average, tend to consume more liquid food than males but this relationship depends on the diet (sex×diet: $F_7=5.66$, $p < 0.001$, Figure S2).

Transcriptional responses to diet

We measured gene expression in males and females maintained on food of either the female-optimal (2:1) or male-optimal (1:4) protein-to-carbohydrate ratio. We separately analysed transcriptomic responses in genes that were expressed in both males and females (hereafter 'shared genes', N = 8888) and those that showed sex-limited expression ($N_{male-limited} = 1879$ and $N_{female-limited} = 165$, see File S1 for full gene lists). For each shared gene, we tested for the effect of sex, diet and the sex-by-diet interaction on expression level. As expected, we found evidence for sex-differences in expression for a large number of genes (a total of 8318 genes with significant sex effect). In addition, we found large-scale transcriptomic responses to diet (806 genes with significant diet effect). Despite the large differences between male and female dietary requirements and food choices, the largest part of the transcriptional responses to diet is shared between the sexes (significant diet effect but no interaction, category 'D' in Table 1, 639 genes). Here, males and females show parallel shifts in expression (although in most cases from a sexually dimorphic baseline expression) when reared on high-carbohydrate vs. high-protein food, and fold-changes between the two diets are strongly positively correlated between males and females (Figure 2; $r = 0.76$, $p < 0.001$).

In addition to these sexually concordant responses, however, we also find a significant number of genes where the sexes show different responses to diet shifts (significant sex-by-diet interaction). For some of these genes, male and female expression change in opposing directions (category 'D×S' in Table 1, 51 genes). Thus, genes that are more highly expressed

on a protein-rich diet in one sex are more lowly expressed on that diet in the other sex,

resulting in negatively correlated fold-changes in the two sexes (Figure 2; r = -0.75, p <

0.001). For another, larger group of genes (category 'D+D×S', 116 genes), both sexes tend to

show expression shifts in the same direction (significant diet effect) but differ in the

magnitude of their responses (significant interaction term). These genes typically show a

large expression response in one sex, but only a small or no response in the other sex, with

overall a lower correlation of fold changes across sexes (r = 0.53, p < 0.001). For the most

part, the dominant expression change occurs in females, but there is a small number of genes

where only male expression responds to diet (Figure 2).

We next analysed diet responses in genes with sex-limited expression. Similar to shared

genes, we also observed significant expression changes in response to diet. Thus, 56 out of

165 female-limited genes showed significant expression change between carbohydrate- and

protein-rich diets. The majority of these (50 genes) had higher expression on the protein-rich

diet preferred by females, while only a small number (6 genes) had higher expression on the

less beneficial carbohydrate-rich diet (Figure 3). In males, 30 out of the 1879 genes with

male-limited expression showed significant diet responses. All of these had higher expression

in the males' preferred carbohydrate-rich diet, compared to the less beneficial protein-rich

media (Figure 3). Taken together, these results show that both sexes respond to their

nutritional environment by upregulating sex-limited genes on their respective optimal diets.

Functional enrichment of dietary responses

We used several approaches to investigate the functions of the genes showing diet responses.

First, we performed Gene Ontology (GO) enrichment analyses for the shared genes of the

three categories (D, D×S, D+D×S) defined above. We found distinct and significant

enrichment in each class, with a predominance of GO terms relating to neuronal and

metabolic biological processes (Figure 4). Second, we took a more targeted approach and

analysed male and female expression changes for genes with specific GO annotations. With

this analysis we aimed to assess how metabolic genes responded to diet manipulation,

compared to the rest of the genome. For this, we fist created a "baseline" of gene expression

by extracting a list of genes that fall under the parent term "Biological Process"

(GO:0008150). From that list, we then removed the genes in the offspring category

"Metabolic Process" to create a set of genes performing biological functions unrelated to

metabolism. We then compared this baseline to genes that fell within the following GO

categories: "Metabolic Process" (GO: 0008152), "Glycolysis" (GO:0006096) and "TCA

cycle" (GO:0006099). The latter two were chosen as core processes in carbohydrate and

protein metabolism. For the sets of genes in each of these categories that showed shared

expression across the sexes, we found positive correlations between male and female fold

changes between the two diet treatments ($R_{MP} = 0.35$, $R_{GLY} = 0.74$, $R_{TCA} = 0.6$, Figure 5A).

These correlations were significantly more positive than the (also slightly positive)

correlation observed in the non-metabolic baseline gene set, despite the fact that correlations

for the small Glycolysis and TCA gene sets have wide confidence intervals (Figure 5B). This

indicates that, even though there is a general shared response to diet between males and

females, male and female responses are more similar in genes involved in core metabolic

processes than the rest of the genome.

For the sex-limited differentially expressed genes, we unsurprisingly found an

enrichment of GO terms involved in reproduction (Figure 6). In females, differentially

expressed genes were enriched for functions associated with egg production (chorion-

containing eggshell formation), but also hormonal control (ecdysone biosynthetic pathway

and hormone synthetic pathway). Male differentially expressed genes were enriched for

sperm function (sperm competition). Since responses in both sexes consisted predominantly

of up-regulation of genes under their respective optimal diets, these results show that for both

males and females, the expression of reproductive genes is increased in the condition that

maximises the fitness of that sex.

Regulation of dietary responses

In order to infer the regulators that drive the observed expression responses to diet, we

searched for enrichment of transcription factor binding motifs upstream of the genes in the

three categories. Our analyses revealed significant enrichment of regulatory motifs in each

group (see File S2 for a full list). Genes that showed only significant diet responses

(concordant response between the sexes, D), presented an overrepresentation of binding

motifs for the transcription factors *CrebB* and *lola*. Genes that showed opposing changes in

males and female (D×S) were enriched for motifs for *vri* and GATA transcription factors

(*grn*, *pnr*, *srp*, *GATAd*, *GATAe*). Finally, genes that showed diet responses largely restricted

to one sex (D+D×S) were enriched primarily for GATA motifs, irrespectively of whether the

response occurred predominantly in females or predominantly in males. Female-specific

genes were mostly enriched for the transcription factors *Blimp-1*, *slbo* and *Dfd*, whereas

male-specific genes were enriched for regulation by *pan* and *Sox*.

494
Overlap with previously described diet and nutrient-signalling responses

We used comparisons to previously published transcriptomic datasets to relate the responses

to shifts in diet quality observed here to those triggered by dietary restriction and

perturbations of nutrient signalling. First, we compared genes in our three categories of diet-

dependent regulation overlapped significantly with sets of genes that change expression in

response to dietary restriction and rapamycin in females, analysed separately for brain,

thorax, gut, and fat body [29]. We found significant overlap in the majority of comparisons

made (Table 3A and B). Non-significant results were only obtained for some comparisons

involving the list of genes in the D×S category, where males and females show opposing

responses to diet. While this might reflect biological reality, it has to be noted that the

numbers of genes—and hence statistical power to detect overlap—are smallest in the D×S

category. Overall, the results of these comparisons demonstrate that transcriptional responses

to the more subtle changes in dietary composition that we apply here generally mirror those

that have previously been observed under dietary restriction.

We then compared our gene categories with a dataset from heads of virgin males and

females in which *IIS/TOR* signalling had been perturbed by expressing a dominant-negative

allele of the insulin receptor InR^{DN} [30]. Reanalysing this dataset (see Methods) we obtained

a list of genes that were altered by an IIS/TOR perturbation across both sexes (N = 5200

genes) similar to the results obtained in the original paper. However, subjecting the data to an

analysis analogous to that we performed on our own, we further found that IIS/TOR

perturbation causes expression changes similar to those observed for our diet treatments.

Thus, a large number of genes show concordant responses to altered insulin signalling in

males (significant InR effect) and females, while a second set shows opposing responses

(InR-by-sex interaction, InR×S) and a third shows largely sex-specific responses

(InR+InR×S) (Figure S5, TableS3). Furthermore, we detect parallelism in the effects of diet

manipulation and InR perturbation on several levels. At the most basic level, the genes that

are significantly affected by IIS/TOR perturbation overlap significantly with the genes that

are significantly affected by diet quality (489 genes observed, 351 expected, 39% excess,

Fisher's exact test, p < 0.001). Second, genes that show a significant diet effect ('D') are more

likely to also show a significant effect of InR perturbation ('InR') (436 genes with both terms

significant, 37% excess, Fisher's exact test, p < 0.001) and genes with a significant diet-by-

sex interaction are more likely to also show a significant InR-by-sex interaction (51 genes,

108% excess, Fisher's exact test, p < 0.001). Third, a full comparison based on a contingency

table containing all possible combinations of classes also showed a significant

correspondence (Chi-squared test, $X_9^2 = 248.53$, p < 0.001), with excess overlap in most

combinations of classes as well in genes that are classified in neither analyses (Table S5).

And finally, fold changes in male and female gene expression in response to IIS/TOR

perturbation correlate positively with those in response to diet manipulations (see Figure S6,

File S3), despite the fact that the two datasets analyse different tissues (head vs. whole body).

These results indicate that manipulating diet quality and manipulating IIS/TOR signalling

produces parallel and overlapping expression responses.

We also investigated the overlap between our diet-responsive genes and genes that have

been identified as TORC1-regulated due to their dependence on REPTOR and REPTOR-BP

[56]. While based on expression in S2 cells only, this to our knowledge is the best

characterised set of TOR-responsive genes. In line with the similarity between expression

responses to diet and IIS/TOR-manipulation described above, we find significant overlap

between our gene categories and genes with REPTOR- or REPTOR-BP–dependent

expression, specifically in our category that responds to diet ('D', 28 genes) and our sex-

biased category ('D+D×S', 9 genes, Table 3C, File S3).

Effect of rapamycin treatment on diet-specific fitness

The overlap with previously described responses raises the potential for the IIS/TOR

network, and specifically TORC1, mediating the diet-dependent phenotypes that we observe

here. This appears plausible for the modulation of female fecundity in response to diet, where

the role of TORC1 is well established, but has not been assessed in males. We therefore

directly tested the phenotypic effect of varying doses of rapamycin and its interaction with

diet, on our proxies for male and female fitness. Our experiment showed that, across the two

sexes, rapamycin leads to a reduction in reproductive output (rapamycin effect: p < 0.001,

Figures 7 and S7, Table S4). More importantly, however, we also found a significant

interaction between diet and rapamycin treatment that was shared across males and females,

where rapamycin lead to a larger reduction in reproductive output on each sex's optimal diet

(sex×rapamycin: p = 0.001). Finally, our experiment revealed possible quantitative

differences between the sexes in the effect of rapamycin on reproduction

(sex×rapamycin×diet: p = 0.068); while the effect of the treatment in females correlated

roughly with the dose administered, males showed a threshold response where all rapamycin

levels in the optimal diet resulted in a reduction in reproductive output to the level observed

on the non-optimal diet.

Discussion

Our study examined the transcriptomic response of male and female *D. melanogaster* to

variation in dietary composition, being exposed to either a male-optimal (protein-to-

carbohydrate ratio 1:4) or a female-optimal (2:1) diet. Our results provide interesting insights

into nutritional effects on male and female fitness in relation to sex- and diet-dependent

expression levels, function and regulation. We show that both sexes share a large metabolic

core transcriptome that is regulated in a sexually concordant way. Nevertheless, smaller parts

of the transcriptome are sex-specifically regulated to diet, including sex-limited reproductive

genes. Together with the observed effects of rapamycin in the two sexes, this suggests that

male and female reproduction is inversely regulated in response to diet composition.

A shared metabolic core transcriptome

Our analyses demonstrated the existence of a core metabolic transcriptome that shows

sexually concordant regulation in response to diet. Overall, expression fold changes from

carbohydrate- to protein-rich food among metabolic genes are positively correlated between

the sexes, and significantly more so than for the transcriptomic background. This indicates

that gene expression in males and females responds generally similarly to changes in dietary

composition. In line with this interpretation, the large majority of genes with diet-dependent

expression show significant changes only in response to diet, independently of sex (639 out

of 806 genes, 79%). Functionally, genes in this core metabolic transcriptome are enriched for

carboxylic acid metabolism and neurological biological processes. Carboxylic acid

metabolism is an integral part of both protein and carbohydrate processing—for instance, part

of the components of amino acids are carboxylic acid sidechains. The prominence of

neurological biological processes, on the other hand, supports the notion of a neural gut-brain

connection that is conserved evolutionarily [57] and shared between the sexes. Specifically,

the sensory mechanisms in the gastrointestinal tract convey information about the nutritional

status to regulate satiety (and thereby feeding behaviour), metabolism, and digestion [58] in a

way that is similar between males and females.

We were also able to infer key regulators of sexually concordant, diet-dependent gene

expression, using motif enrichment tools. Upstream regions of genes with sexually

concordant diet responses were enriched for motifs of two main transcription factors *CreB*

and *lola* transcription factors. *CrebB* is involved in diurnal rhythms and memory formation

[59, 60], but also in energy homeostasis and starvation resistance, mediated by insulin

signalling [61]. The *lola* transcription factor, on the other hand, is mainly involved in axon

guidance in *Drosophila* [62, 63]. But interestingly, some protein isoforms have also been

associated with octopamine synthesis pathways which are essential for nutrient sensing [64].

Sex-specific diet responses in gene regulation

Besides the large, shared core metabolic transcriptome, we also identified smaller groups of

genes with sex-specific expression responses to diet. A first group showed opposing diet

responses in males and female (D×S, 51 out of 806 genes, 6.3%). These genes are enriched

for transport functions and synapse assembly/organisation. One of our candidate antagonistic

genes is *fit* (*female-specific independent of transformer*). Known to be sexually dimorphic in

expression, *fit* has been found to be rapidly upregulated in male heads during the process of

male courtship and mating, along with another antagonistic candidate *Odorant binding*

protein 99b, *Obp99B* [65, 66]. Interestingly, *fit* has also been implicated in protein satiety in a

sex-specific manner [67]. Following the ingestion of protein-rich food, *fit* expression

increases in both sexes (although more so in females than males), but only supresses protein

appetite in females [67]. Both *fit* and *Obp99B* were found to be significantly altered in a sex-

specific way when flies were starved, further cementing their role in nutrient response [68].

Together with previous work, our results therefore cement the tight link between nutritional

sensory mechanisms and reproduction, however this response is sex-specific.

Another group of genes showed mostly responses in one sex (D+D×S, 116 genes,

14.4%). Most of the genes observed in this category show expression changes in females

(with little change in male expression levels) and are mainly involved in carbohydrate

metabolism and female receptivity. One notable gene in this category is the transcription

factor *doublesex,* which plays a key role in sexual differentiation and the regulation of sex-

specific behavioural traits [69]. Expression levels of this gene are higher in females that are

fed a high-protein diet (unless the difference in *dsx* mRNA levels is due to growth in a

sexually dimorphic, and hence *dsx*-expressing, tissue type). Of interest among the few genes

with male-limited diet response (Figure 2) is *Adenosylhomosysteinase* (*Ahcy*), which we find

males to express at lower levels on the carbohydrate-rich (optimal) diet. *Ahcy* is involved in

methionine metabolism and has been linked to male lifespan regulation. *Ahcy* knock-outs

were shorter lived, while knock-outs for two putative *Ahcy*-repressors extended male life- and

health-span [70]. These effects are in line with the under-expression we observe on high
carbohydrate, under the assumption that greater investment in current reproduction is
associated with decreased lifespan (which may not generally hold in the context of nutrient
manipulation [3]).
Both the genes with opposing (D×S) and those with sex-limited diet-dependent
regulation (D+D×S) show significant enrichment for GATA transcription factors. This class
of transcription factors has been previously implicated in female nutritional and reproductive
control. For example, the ovary-specific *dGATAb* binds upstream of both yolk protein genes
Yp1 and *Yp2* [71]. GATA-related motifs have also previously been shown to be enriched in
genes showing differential expression in response to DR and rapamycin treatment in female
flies [29]. The shared regulation is further supported by the fact that the diet-responsive genes
we identify here also overlap significantly with those previously inferred to respond to DR-
and rapamycin-treatment. These results suggest that changing the *quality* of the diet elicits a
similar response as changing the *quantity* via protein dilution. This may not be surprising, if
DR is considered a response mainly to the quantity of protein ingested [8, 72], and fits with
previous work that found the ratio of macronutrients—not caloric intake—to be the main
determinant of healthy ageing in mice [13]. However, the overlap highlights that DR-
phenotypes are not an all-or-nothing response but instead are part of a continuum of life
history adjustments in response to how suitable the dietary environment is for current
reproduction.
647
Diet-specific regulation of male and female reproduction
We also found diet responses in reproductive genes that are exclusively expressed in either
males or females. Regulation largely reflects diet-dependent reproductive investment, with
most genes being more highly expressed on a sex's optimal diet with lower expression on the
suboptimal diet. In females, a significant number of these genes are involved in egg
production and thus linked to diet-dependent reproductive investment [2]. Also among the
genes is *insulin-like peptide-7* (*dILP-7*), one of a family of peptides known to having the
functional as hormones and neuropeptides [73] involved in nutrient foraging control [74].
More specifically, *dILP-7* is expressed in neurons that play an active role in female fertility.
These neurons have been linked with the egg-laying decision process [75, 76] and *dILP-7* is
among a number of genes show sexually dimorphic expression in these neuronal cells [77].
Interestingly, IIS/TOR perturbation also results in sex-specific changes in dILP peptides
(dILP2, 3, 5 and 6) in the head [30] (where dILP7 is not expressed [78]).

Mirroring expression responses in females, we also find higher expression of
reproductive genes on the optimal diet in males. This is surprising—based on the view that
male fitness is limited by the acquisition of mates and the supposedly low investment
required for sperm production [2], one could expect that males do not modulate their
reproductive investment in response to the nutritional environment but remain primed to
maximally exploit any mating opportunity. Assuming that expression of these genes reflects
reproductive investment, the fact that they do respond to the nutritional environment suggests
that male reproductive strategies are maybe more subtle, and their investment more costly,
than previously appreciated. This is plausible, as work on other insects has shown that the
production of high quality sperm is costly [79] (but courtship activity does not appear to carry
a significant cost, at least in fruit flies [80]).

Superficially, it may seem obvious that male and female reproductive genes are
upregulated on each sex's respective optimal diet. In the presence of a largely shared and
concordantly regulated metabolic machinery, however, this pattern implies that the output of
nutrient sensing pathways is used in different, and potentially inversed ways in males and
females. While our analyses do not allow us to identify the exact point of reversal within the
regulatory hierarchy, our data provide some interesting insights. First, it is noteworthy that
GATA transcription factors are inferred to be regulating genes that show a wide range of
expression patterns, being enriched among genes with opposing expression changes in males
and females (the D×S set), as well as those that show largely sex-limited responses (D+D×S).
This could imply that the main role played by these factors is to convey information about the
metabolic and nutritional state of the animal (similar to homeotic genes in development),
which is then incorporated combinatorially with additional factors to give rise to the sex- and
diet-specific expression patterns that we observe.

Second, several lines of bioinformatic evidence suggest that the expression changes that
we describe here are at least in part regulated by IIS/TOR signalling. Thus, the genes that we
find to respond to diet manipulation significantly overlap with genes affected by
manipulation of IIS/TOR signalling as described by Graze *et al.* [30], a dataset that our
reanalysis reveals to show a similar structure of genes with sexually concordant, sexually
opposing and sex-biased expression changes. This pattern and the overlap with our data is all
the more noteworthy as Graze *et al.* assessed the effect of IIS/TOR perturbation in virgin
flies, where males and females have more similar dietary requirements, and hence
presumably more similar physiological states, than in mated flies [11]. In addition to showing
parallels with IIS/TOR-dependent expression, our diet-dependent genes also significantly

overlap with the arguably best-defined set of TORC1-dependent genes currently available

[56]. These results suggest that diet-dependent expression responses, and their sex-specific

differences, are mediated by IIS and the TOR pathway.

This conclusion is corroborated by the results of our experiment combining diet

manipulation with rapamycin treatments, which are consistent with TORC1-dependent

upregulation of reproduction on optimal diets in both sexes. Here we find that while

rapamycin generally lowers reproductive output, this effect is more pronounced on the

respective optimal diet of each sex. This is expected in females, where a large body of work

implicates the IIS/TOR network in life-history shifts between reproduction and longevity

[81]. Accordingly, a nutritionally favourable environment should lead to increased TORC1

activity and elevated reproductive output. What our data show, however, is that a parallel

effect of increased reproduction on the optimal diet is detectable in males, even though the

composition of that diet is the one that is unfavourable in females, leading to low TORC1 and

reduced reproduction. Across the sexes, TORC1 activity would thus not reflect a specific

dietary composition but a measure of nutritional optimality and regulate reproductive

investment accordingly.

We note that, while tantalising, these inferences will require further careful validation.

Due to the focus on females, diet-dependent regulation of male reproduction has been little

explored. Knock-down of *Tor* and *raptor* in males has been found to result in an

accumulation of germline stem cells, combined with deficient differentiation [82]. Future

work will need to assess the effect of these changes on male reproductive output and, more

importantly, whether and how the signal of the nutrient sensing mechanisms that feed into the

Tor pathway are modulated in a sex-specific way. Independently of how the regulatory

reversal is achieved mechanistically, our data also suggest that the relationship between the

composition of the diet consumed and reproductive output does not merely reflect the passive

effect of metabolic conversion rates from nutritional components to gametes and energy but

is at least in part the result if an active regulation of immediate reproductive investment. This

has important implications for our interpretation of variation in diet-specific reproductive

success, which has been documented in the population studied here [83]. Thus, variation

between genotypes in the dietary composition that maximises, for example, male

reproductive fitness is therefore probably at least partly caused by genetic variation in how

nutrients are sensed or how this sensory output is used to regulate reproductive investment.

Studying this variation in more detail will provide a fruitful avenue to better understand the

regulatory mechanisms involved, as well as the selective forces that shape variation in its

components.

Acknowledgements. We would like to thank Nazif Alic and Adam Dobson for their help

with planning the experiments, analysis, and for insightful comments on previous versions of

this manuscript. In addition, we thank Kevin Fowler, Filip Ruzicka and Michael Jardine for

useful discussions and Rebecca Finlay for help with laboratory work. This study was funded

by a Marie Sklodowska-Curie Research Fellowship [#708362] to MFC.

Data, Code and Materials. Data from this manuscript will be uploaded to Dryad upon

acceptance of this manuscript.

Authors' contributions. MFC, MWDP and MR conceived the study and wrote the

manuscript. MFC performed all experimentation. MFC and MR analysed the data.

Competing interests. The authors declare no competing interests.

References

1.   Chapman, T. (2006). Evolutionary conflicts of interest between males and females.
Current Biology *16*, R744-R754.

2.   Trivers, R., and Campbell, B. (1972). Parental investment and sexual selection. In
Sexual selection and the descent of man 1871-1971. (Chicago (IL) Aldine-Atherton), pp.
136-179.

3.   Jensen, K., McClure, C., Priest, N.K., and Hunt, J. (2015). Sex-specific effects of
protein and carbohydrate intake on reproduction but not lifespan in Drosophila melanogaster.
Aging Cell *14*, 605-615.

4.   Reddiex, A.J., Gosden, T.P., Bonduriansky, R., and Chenoweth, S.F. (2013). Sex-
specific fitness consequences of nutrient intake and the evolvability of diet preferences.
American Naturalist *182*, 91-102.

5.   Maklakov, A.A., Simpson, S.J., Zajitschek, F., Hall, M.D., Dessmann, J., Clissold, F.,
Raubenheimer, D., Bonduriansky, R., and Brooks, R.C. (2008). Sex-specific fitness effects of
nutrient intake on reproduction and lifespan. Current Biology *18*, 1062-1066.

6.   Maklakov, A.A., Hall, M.D., Simpson, S.J., Dessmann, J., Clissold, F.J., Zajitschek,
F., Lailvaux, S.P., Raubenheimer, D., Bonduriansky, R., and Brooks, R.C. (2009). Sex
differences in nutrient-dependent reproductive ageing. Aging Cell *8*, 324-330.

7.   Simpson, S.J., and Raubenheimer, D. (2011). The nature of nutrition: a unifying
framework. Aust J Zool *59*, 350-368.

8.   Lee, K.P., Simpson, S.J., Clissold, F.J., Brooks, R., Ballard, J.W.O., Taylor, P.W.,
Soran, N., and Raubenheimer, D. (2008). Lifespan and reproduction in *Drosophila*: New
insights from nutritional geometry. Proceedings of the National Academy of Sciences of the
United States of America *105*, 2498-2503.

9.   Corrales-Carvajal, V.M., Faisal, A.A., and Ribeiro, C. (2016). Internal states drive
nutrient homeostasis by modulating exploration-exploitation trade-off. eLife *5*, e19920.

10.   Ribeiro, C., and Dickson, B.J. (2010). Sex peptide receptor and neuronal TOR/S6K
signaling modulate nutrient balancing in *Drosophila*. Current Biology *20*, 1000-1005.

11.   Camus, M.F., Huang, C.C., Reuter, M., and Fowler, K. (2018). Dietary choices are
influenced by genotype, mating status, and sex in *Drosophila melanogaster*. Ecology and
Evolution *8*, 5385-5393.

12.   Moatt, J.P., Fyfe, M.A., Heap, E., Mitchell, L.J.M., Moon, F., and Walling, C.A.
(2019). Reconciling nutritional geometry with classical dietary restriction: Effects of nutrient
intake, not calories, on survival and reproduction. Aging Cell *18*, e12868.

13.   Solon-Biet, Samantha M., McMahon, Aisling C., Ballard, J.William O., Ruohonen,
K., Wu, Lindsay E., Cogger, Victoria C., Warren, A., Huang, X., Pichaud, N., Melvin,
Richard G., et al. (2014). The ratio of macronutrients, not caloric intake, dictates
cardiometabolic health, aging, and longevity in *ad libitum*-fed mice. Cell Metab *19*, 418-430.

14.   Piper, M.D., Skorupa, D., and Partridge, L. (2005). Diet, metabolism and lifespan in
Drosophila. Experimental Gerontology *40*, 857-862.

15.   Regan, J.C., Khericha, M., Dobson, A.J., Bolukbasi, E., Rattanavirotkul, N., and
Partridge, L. (2016). Sex difference in pathology of the ageing gut mediates the greater
response of female lifespan to dietary restriction. elife *5*, e10956.

16.   Piper, M.D.W., and Partridge, L. (2018). Drosophila as a model for ageing. Bba-Mol
Basis Dis *1864*, 2707-2717.

17.   Partridge, L., Gems, D., and Withers, D.J. (2005). Sex and death: What is the
connection? Cell *120*, 461-472.

18.   Magwere, T., Chapman, T., and Partridge, L. (2004). Sex differences in the effect of
dietary restriction on life span and mortality rates in female and male *Drosophila*
melanogaster. Journal of Gerontology: Series A *59*, 3-9.

19.   Zajitschek, F., Jin, T., Colchero, F., and Maklakov, A.A. (2014). Aging differently:
diet- and sex-dependent late-life mortality patterns in *Drosophila melanogaster*. Journal of
Gerontology: Series A *69*, 666-674.

20.   Zajitschek, F., Zajitschek, S.R., Friberg, U., and Maklakov, A.A. (2013). Interactive
effects of sex, social environment, dietary restriction, and methionine on survival and
reproduction in fruit flies. Age *35*, 1193-1204.

21.   Hoedjes, K.M., Rodrigues, M.A., and Flatt, T. (2017). Amino acid modulation of
lifespan and reproduction in *Drosophila*. Current Opinion in Insect Science *23*, 118-122.

22.   Clancy, D.J., Gems, D., Hafen, E., Leevers, S.J., and Partridge, L. (2002). Dietary
restriction in long-lived dwarf flies. Science *296*, 319.

23.   Slack, C., Giannakou, M.E., Foley, A., Goss, M., and Partridge, L. (2011). dFOXO-
independent effects of reduced insulin-like signaling in *Drosophila*. Aging Cell *10*, 735-748.

24.   Zandveld, J., van den Heuvel, J., Zwaan, B.J., and Piper, M.D.W. (2017). Both
overlapping and independent mechanisms determine how diet and insulin-ligand knockouts
extend lifespan of Drosophila melanogaster. Npj Aging and Mechanisms of Disease *3*.

25.   Emran, S., Yang, M.Y., He, X.L., Zandveld, J., and Piper, M.D.W. (2014). Target of
rapamycin signalling mediates the lifespan-extending effects of dietary restriction by
essential amino acid alteration. Aging-Us *6*, 390-398.

26.   Bjedov, I., Toivonen, J.M., Kerr, F., Slack, C., Jacobson, J., Foley, A., and Partridge,
L. (2010). Mechanisms of Life Span Extension by Rapamycin in the Fruit Fly Drosophila
melanogaster. Cell Metab *11*, 35-46.

27.   Alic, N., and Partridge, L. (2011). Death and dessert: nutrient signalling pathways and
ageing. Curr Opin Cell Biol *23*, 738-743.

28.   Gallinetti, J., Harputlugil, E., and Mitchell, J.R. (2013). Amino acid sensing in
dietary-restriction-mediated longevity: roles of signal-transducing kinases GCN2 and TOR.
Biochemical Journal *449*, 1-10.

29.   Dobson, A.J., He, X., Blanc, E., Bolukbasi, E., Feseha, Y., Yang, M., and Piper,
M.D.W. (2018). Tissue-specific transcriptome profiling of *Drosophila* reveals roles for
GATA transcription factors in longevity by dietary restriction. Aging and Mechanisms of
Disease *4*, 5.

30.   Graze, R.M., Tzeng, R.Y., Howard, T.S., and Arbeitman, M.N. (2018). Perturbation
of IIS/TOR signaling alters the landscape of sex-differential gene expression in *Drosophila*.
BMC Genomics *19*, 893.

31.   Chippindale, A.K., Gibson, J.R., and Rice, W.R. (2001). Negative genetic correlation
for adult fitness between sexes reveals ontogenetic conflict in *Drosophila*. Proceedings of the
National Academy of Sciences of the United States of America *98*, 1671-1675.

32.   Rice, W.R. (1996). Sexually antagonistic male adaptation triggered by experimental
arrest of female evolution. Nature *381*, 232-234.

33.   Piper, M.D.W., Blanc, E., Leitao-Goncalves, R., Yang, M.Y., He, X.L., Linford, N.J.,
Hoddinott, M.P., Hopfen, C., Soultoukis, G.A., Niemeyer, C., et al. (2014). A holidic
medium for *Drosophila melanogaster*. Nat Methods *11*, 100-105.

34.   Piper, M.D.W., and Partridge, L. (2007). Dietary restriction in *Drosophila*: Delayed
aging or experimental artefact? PLoS Genetics *3*, e57.

35.   Pischedda, A., and Rice, W.R. (2012). Partitioning sexual selection into its mating
success and fertilization success components. Proceedings of the National Academy of
Sciences of the United States of America *109*, 2049-2053.

36.   Collet, J.M., Fuentes, S., Hesketh, J., Hill, M.S., Innocenti, P., Morrow, E.H., Fowler,
K., and Reuter, M. (2016). Rapid evolution of the intersexual genetic correlation for fitness in
Drosophila melanogaster. Evolution *70*, 781-795.

37.   Sang, J.H. (1978). The nutritional requirements of *Drosophila melanogaster*. In The
Genetics and Biology of *Drosophila*. (Academic Press, London), pp. 159-192.

38.   Colinet, H., and Renault, D. (2014). Dietary live yeast alters metabolic profiles,
protein biosynthesis and thermal stress tolerance of *Drosophila melanogaster*. Comparative
Biochemistry and Physiology Part A Molecular & Integrative Physiology *170*, 6-14.

39.   Tanaka, T., and Yamazaki, T. (1990). Fitness and its components in *Drosophila*
melanogaster. Japanese Journal of Genetics *65*, 417-426.

40.   Hoffmann, A.A., and Harshman, L.G. (1985). Male effects on fecundity in
Drosophila melanogaster. Evolution *39*, 638-644.

41.   Waithe, D., Rennert, P., Brostow, G., and Piper, M.D.W. (2015). QuantiFly: Robust
trainable software for automated *Drosophila* egg counting. PLoS ONE *10*, e0127659.

42.   Team, R.C. (2016). R: A Language and Environment for Statistical Computing.

43.   Lande, R., and Arnold, S.J. (1983). The measurement of selection on correlated
characters. Evolution *37*, 1210-1226.

44.   Chenoweth, S.F., and Blows, M.W. (2005). Contrasting mutual sexual selection on
homologous signal traits in *Drosophila serrata*. American Naturalist *165*, 281-289.

45.   Bolger, A.M., Lohse, M., and Usadel, B. (2014). Trimmomatic: a flexible trimmer for
Illumina sequence data. Bioinformatics *30*, 2114-2120.

46.   Kim, D., Langmead, B., and Salzberg, S.L. (2015). HISAT: a fast spliced aligner with
low memory requirements. Nat Methods *12*, 357-360.

47.   Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G.,
Abecasis, G., Durbin, R., and Genome Project Data Processing, S. (2009). The Sequence
Alignment/Map format and SAMtools. Bioinformatics *25*, 2078-2079.

48.   Anders, S., Pyl, P.T., and Huber, W. (2015). HTSeq - a Python framework to work
with high-throughput sequencing data. Bioinformatics *31*, 166-169.

49.   Robinson, M.D., McCarthy, D.J., and Smyth, G.K. (2010). edgeR: a Bioconductor
package for differential expression analysis of digital gene expression data. Bioinformatics
26, 139-140.

50.   Yu, G., Wang, L.-G., Han, Y., and He, Q.-Y. (2012). clusterProfiler: an R package for
comparing biological themes among gene clusters. OMICS: A Journal of Integrative Biology
16, 284-287.

51.   Aibar, S., Gonzalez-Blas, C.B., Moerman, T., Huynh-Thu, V.A., Imrichova, H.,
Hulselmans, G., Rambow, F., Marine, J.C., Geurts, P., Aerts, J., et al. (2017). SCENIC:
single-cell regulatory network inference and clustering. Nat Methods *14*, 1083-1086.

52.   Benjamini, Y., and Hochberg, Y. (1995). Controlling the false discovery rate: A
practical and powerful approach to multiple testing. Journal of the Royal Statistical Society.
Series B (Methodological) *57*, 289-300.

53.   Ding, F., Gil, M.P., Franklin, M., Ferreira, J., Tatar, M., Helfand, S.L., and Neretti, N.
(2014). Transcriptional response to dietary restriction in *Drosophila melanogaster*. J Insect
Physiol *69*, 101-106.

54.   Shen, L., and Sinai, M. (2018). GeneOverlap: Test and visualize gene overlaps. R
package version 1.18.0.

55.   Crespo, J.L., and Hall, M.N. (2002). Elucidating TOR signaling and rapamycin
action: lessons from Saccharomyces cerevisiae. Microbiol Mol Biol Rev *66*, 579-591, table
of contents.

56.   Tiebe, M., Lutz, M., De La Garza, A., Buechling, T., Boutros, M., and Teleman, A.A.
(2015). REPTOR and REPTOR-BP Regulate Organismal Metabolism and Transcription
Downstream of TORC1. Dev Cell *33*, 272-284.

57. Kaelberer, M.M., and Bohorquez, D.V. (2018). The now and then of gut-brain
signaling. Brain Res *1693*, 192-196.

58. Kaelberer, M.M., Buchanan, K.L., Klein, M.E., Barth, B.B., Montoya, M.M., Shen,
X., and Bohorquez, D.V. (2018). A gut-brain neural circuit for nutrient sensory transduction.
Science *361*.

59. Bittinger, M.A., McWhinnie, E., Meltzer, J., Iourgenko, V., Latario, B., Liu, X.,
Chen, C.H., Song, C., Garza, D., and Labow, M. (2004). Activation of cAMP response
element-mediated gene expression by regulated nuclear transport of TORC proteins. Curr
Biol *14*, 2156-2161.

60. Kogan, J.H., Frankland, P.W., Blendy, J.A., Coblentz, J., Marowitz, Z., Schutz, G.,
and Silva, A.J. (1997). Spaced training induces normal long-term memory in CREB mutant
mice. Current Biology *7*, 1-11.

61. Wang, B., Goode, J., Best, J., Meltzer, J., Schilman, P.E., Chen, J., Garza, D.,
Thomas, J.B., and Montminy, M. (2008). The insulin-regulated CREB coactivator TORC
promotes stress resistance in *Drosophila*. Cell Metab *7*, 434-444.

62. Horiuchi, T., Giniger, E., and Aigaki, T. (2003). Alternative trans-splicing of constant
and variable exons of a *Drosophila* axon guidance gene, lola. Genes & Development *17*,
2496-2501.

63. Goeke, S., Greene, E.A., Grant, P.K., Gates, M.A., Crowner, D., Aigaki, T., and
Giniger, E. (2003). Alternative splicing of lola generates 19 transcription factors controlling
axon guidance in *Drosophila*. Nature Neuroscience *6*, 917-924.

64. Dinges, N., Morin, V., Kreim, N., Southall, T.D., and Roignant, J.Y. (2017).
Comprehensive characterization of the complex lola locus reveals a novel role in the
octopaminergic pathway via tyramine beta-hydroxylase regulation. Cell Reports *21*, 2911-
2925.

65. Ellis, L.L., and Carney, G.E. (2010). Mating alters gene expression patterns in
Drosophila melanogaster male heads. BMC Genomics *11*, 558.

66. Carney, G.E. (2007). A rapid genome-wide response to *Drosophila melanogaster*
social interactions. BMC Genomics *8*, 288.

67. Sun, J.H., Liu, C., Bai, X.B., Li, X.T., Li, J.Y., Zhang, Z.P., Zhang, Y.P., Guo, J., and
Li, Y. (2017). Drosophila FIT is a protein-specific satiety hormone essential for feeding
control. Nat Commun *8*.

68. Fujikawa, K., Takahashi, A., Nishimura, A., Itoh, M., Takano-Shimizu, T., and Ozaki,
M. (2009). Characteristics of genes up-regulated and down-regulated after 24 h starvation in
the head of Drosophila. Gene *446*, 11-17.

69.   Shirangi, T.R., Taylor, B.J., and McKeown, M. (2006). A double-switch system
regulates male courtship behavior in male and female Drosophila melanogaster. Nature
Genetics *38*, 1435-1439.

70.   Parkhitko, A.A., Binari, R., Zhang, N., Asara, J.M., Demontis, F., and Perrimon, N.
(2016). Tissue-specific down-regulation of S-adenosyl-homocysteine via suppression of
dAhcyL1/dAhcyL2 extends health span and life span in *Drosophila*. Genes & Development
30, 1409-1422.

71.   Lossky, M., and Wensink, P.C. (1995). Regulation of *Drosophila* yolk protein genes
by an ovary-specific GATA factor. Molecular Cell Biology *15*, 6943-6952.

72.   Grandison, R.C., Piper, M.D.W., and Partridge, L. (2009). Amino-acid imbalance
explains extension of lifespan by dietary restriction in *Drosophila*. Nature *462*, 1061-U1121.

73.   Sisodia, S., and Singh, B.N. (2012). Experimental evidence for nutrition regulated
stress resistance in *Drosophila ananassae*. PLoS One *7*, e46131.

74.   Shim, J., Gururaja-Rao, S., and Banerjee, U. (2013). Nutritional regulation of stem
and progenitor cells in *Drosophila*. Development *140*, 4647-4656.

75.   Yang, C.H., Belawat, P., Hafen, E., Jan, L.Y., and Jan, Y.N. (2008). *Drosophila* egg-
laying site selection as a system to study simple decision-making processes. Science *319*,
1679-1683.

76.   Lihoreau, M., Poissonnier, L.A., Isabel, G., and Dussutour, A. (2016). *Drosophila*
females trade off good nutrition with high-quality oviposition sites when choosing foods. J
Exp Biol *219*, 2514-2524.

77.   Castellanos, M.C., Tang, J.C., and Allan, D.W. (2013). Female-biased dimorphism
underlies a female-specific role for post-embryonic Ilp7 neurons in *Drosophila* fertility.
Development *140*, 3915-3926.

78.   Nassel, D.R., and Vanden Broeck, J. (2016). Insulin/IGF signaling in Drosophila and
other insects: factors that regulate production, release and post-release action of the insulin-
like peptides. Cell Mol Life Sci *73*, 271-290.

79.   Bunning, H., Rapkin, J., Belcher, L., Archer, C.R., Jensen, K., and Hunt, J. (2015).
Protein and carbohydrate intake influence sperm number and fertility in male cockroaches,
but not sperm viability. Proceedings of the Royal Society B-Biological Sciences *282*.

80.   Flintham, E.O., Yoshida, T., Smith, S., Pavlou, H.J., Goodwin, S.F., Carazo, P., and
Wigby, S. (2018). Interactions between the sexual identity of the nervous system and the
social environment mediate lifespan in Drosophila melanogaster. Proceedings of the Royal
Society B-Biological Sciences *285*.

81.   Wullschleger, S., Loewith, R., and Hall, M.N. (2006). TOR signaling in growth and
metabolism. Cell *124*, 471-484.

82.   Liu, Y., Ge, Q., Chan, B., Liu, H., Singh, S.R., Manley, J., Lee, J., Weideman, A.M.,
Hou, G., and Hou, S.X. (2016). Whole-animal genome-wide RNAi screen identifies networks
regulating male germline stem cells in Drosophila. Nat Commun *7*, 12149.

83.   Camus, M.F., Fowler, K., Piper, M.W.D., and Reuter, M. (2017). Sex- and genotype-
effects on nutrient-dependent fitness landscapes in *Drosophila melanogaster*. Proceedings of
the Royal Society B-Biological Sciences *284*, 20172237.
970
971

Tables and Figures

Table 1: Shared transcriptomic response – Number of genes that are influenced by sex (S),

diet (D), and their interaction (D×S). From this method, we were able to cluster genes into 3

main categories. Categories highlighted in orange encompass genes that show an additive

effect to diet (D), whereas clusters in blue show interactive effects (D×S). Green rows show

both additive and interactive effects (D+D×S)

significance (FDR<0.05)			n. genes
S	D	D×S	
-	-	-	545
-	-	Y	3
-	Y	-	18
-	Y	Y	4
Y	-	-	7537
Y	-	Y	48
Y	Y	-	621
Y	Y	Y	112

984   **Table 2: Sex-specific transcriptomic response** – Number of genes that are differentially

985   expressed when moving from a carbohydrate-rich environment to a protein-rich environment

986   in females and males (FDR < 0.1).

Sex	Contrast	UP	ns	DOWN	Total
Female	Carb → Protein	50	109	6	56
Male	Carb → Protein	0	1845	34	34

Table 3: Gene overlap between our three categories (D, D×S, D+D×S) and results from three
previously published papers. The first study (**A+B**) examines female transcriptomic response
to dietary restriction and rapamycin across six different tissues [29]. The second study (**C**)
characterises genes that respond to TORC1 inhibition via the transcription factors REPTOR
and REPTOR-BP [56]. In italics we show the total number of genes in that category, with
bold counts showing the significant (P < 0.05) overlaps between two categories. Overlap is
assessed with Fisher's exact tests, p-values are provided below the counts.
996
997

A. Dietary Restriction

	Brain *(167)*	Thorax *(193)*	Gut *(25)*	Fatbody *(358)*
D *(639)*	**27** $p < 0.001$	**51** $p < 0.001$	**14** $p < 0.001$	**58** $p < 0.001$
D×S *(51)*	**5** $p = 0.0026$	**5** $p = 0.0048$	0 $p = 1$	**7** $p = 0.0041$
D+D×S *(116)*	**10** $p < 0.001$	**19** $p < 0.001$	**3** $p = 0.004$	**20** $p < 0.001$

B. Rapamycin

	Brain *(58)*	Thorax *(38)*	Gut *(76)*	Fatbody *(222)*
D *(639)*	**14** $p < 0.001$	**9** $p = 0.0012$	**17** $p < 0.001$	**57** $p < 0.001$
D×S *(51)*	**5** $p < 0.001$	**2** $p = 0.02$	2 $p = 0.07$	3 $p = 0.13$
D+D×S *(116)*	**6** $p < 0.001$	**7** $p < 0.001$	**4** $p = 0.017$	**16** $p < 0.001$

C. TORC1

	REPTOR/REPTOR-BP *(212)*
D *(639)*	**28** $p = 0.019$
D×S *(51)*	1 $p = 0.78$
D+D×S *(116)*	**9** $p = 0.0068$

998
999

1000

1001

1002

1003 **Figure 1:** Nutritional landscapes for female (left) and male (right) fitness in the LH_M

1004 population. Small grey dots represent the dietary coordinates of individual fitness measures.

1005 Dietary choices for each sex are also plotted (white dot). The red arrow denotes the female

1006 optimal nutritional rail (P:C = 2:1), whereas the blue arrow is the male optimal nutritional rail

1007 (P:C = 1:4). For each nutritional rail we samples 120 flies of each sex.

1008

1009   **Figure 2**: Male and female expression responses (log$_2$-fold change) for genes classified as

1010   showing only a diet effect (Diet), only a diet-by-sex interaction (Diet×Sex) or both (Diet +

1011   Diet×Sex). Expression changes are measured from the carbohydrate- to protein-rich diet.

1012   Colours represent genes with significant differential expression (at 5% FDR) only in females

1013   (red), only in males (blue), in both sexes (yellow) or in neither sex (grey).

1014

1015   **Figure 3:** Volcano plot of the sex-specific gene sets. Yellow data points denote genes that

1016   were identified as differentially expressed at a 5% FDR cut-off.

1017

1018 **Figure 4:** GO enrichment for the shared transcriptomic response. Enrichment for "biological

1019 process" was performed for all categories, and p-values were adjusted for FDR<0.05

1020 ('p.adjust').

1021

1022

1023

Figure 5: (A) Male and female $\log_2$ fold changes in gene expression going from

carbohydrate to protein diets for selected GO terms. Red data points are genes that are found

within the chosen GO terms (Metabolic process, TCA cycle and Glycolysis), whereas grey

datapoints all other genes annotated with the first-level parent term "biological process". This

background set provides a transcriptome-wide expression baseline between the sexes. Lines

represent the regression of female on male $\log_2$ fold change for the target term (red) and the

background set (blue). **(B)** Bootstrapped correlation coefficients selected GO categories (red)

and the baseline (grey).

1032
1033 **Figure 6:** GO enrichment for sex-specific genes. Enrichment for differentially expressed
1034 genes was performed using "biological process" and p-values were adjusted for FDR
1035 ('p.adjust').

Figure 7: Male and female fitness measures across the two diets and for rapamycin
treatments. Sample size for each experimental treatment is 60 flies.