Data Descriptor

A DNA barcode reference library of the French Polynesian shore fishes

Erwan Delrieu-Trottin[1,2,3,4], Jeffrey T. Williams[5], Diane Pitassy[5], Amy Driskell[6], Nicolas Hubert[1], Jérémie

Viviani[3,7,8], Thomas H. Cribb[9], Benoit Espiau[3,4], René Galzin[3,4], Michel Kulbicki[10], Thierry Lison de Loma[3,4],

Christopher Meyer[11], Johann Mourier[3,4,12], Gérard Mou-Tham[10], Valeriano Parravicini[3,4], Patrick

Plantard[3,4], Pierre Sasal[3,4], Gilles Siu[3,4], Nathalie Tolou[3,4], Michel Veuille[4,13], Lee Weigt[6] & Serge Planes[3,4]

1. Institut de Recherche pour le Développement, UMR 226 ISEM (UM2-CNRS-IRD-EPHE), Université de

Montpellier, Place Eugène Bataillon, CC 065, F-34095 Montpellier cedex 05, France

2. Museum für Naturkunde, Leibniz-Institut für Evolutions-und Biodiversitätsforschung an der Humboldt-

Universität zu Berlin, Invalidenstrasse 43, Berlin 10115, Germany

3. PSL Research University, EPHE-UPVD-CNRS, USR 3278 CRIOBE, Université de Perpignan, 58 Avenue

Paul Alduy, 66860, Perpignan, France.

4. Laboratoire d'Excellence «CORAIL», Papetoai, Moorea, French Polynesia , France.

5. Division of Fishes, Department of Vertebrate Zoology, National Museum of Natural History,

Smithsonian Institution, 4210 Silver Hill Road, Suitland, MD 20746, USA

6. Laboratories of Analytical Biology, National Museum of Natural History, Smithsonian Institution,

Washington, D.C., 20013, United States of America

7. Département de Biologie, École Normale Supérieure de Lyon, Université de Lyon, UCB Lyon1, 46 Allée

d'Italie, Lyon, FRANCE

8. Team Evolution of Vertebrate Dentition, Institute of Functional Genomics of Lyon, ENS de Lyon, CNRS

UMR 5242, Université de UCB Lyon1, 46 allée d'Italie, Lyon, FRANCE

9. School of Biological Sciences, The University of Queensland, Brisbane, Australia 4072

10. Institut de Recherche pour le Développement – UR 227 CoReUs, LABEX "CORAIL", UPVD, 66000

Perpignan, France

11. Department of Invertebrate Zoology, National Museum of Natural History, National Museum of

Natural History, Smithsonian Institution, Washington, D.C., 20560-0163, United States of America

12. UMR 248 MARBEC (IRD, Ifremer, Univ. Montpellier, CNRS), Station Ifremer de Sète, Av Jean Monnet,

CS 30171 34203 Sète cedex, France

13. Institut Systématique, Évolution, Biodiversité (ISYEB), UMR 7205, CNRS, MNHN, UPMC, EPHE. Ecole

Pratique des Hautes Etudes, Paris Sciences Lettres (PSL), 57 rue Cuvier, CP39, F-75005, Paris, France.

Correspondence and requests for materials should be addressed to E.D.T. (email:

erwan.delrieu.trottin@gmail.com) or S.P. (email: planes@univ-perp.fr) or J.T.W. (williamsjt@si.edu)

Abstract

The emergence of DNA barcoding and metabarcoding opened new ways to study biological diversity, however, the completion of DNA barcode libraries is fundamental for such approaches to succeed. This dataset is a DNA barcode reference library (fragment of Cytochrome Oxydase I gene) for 2,190 specimens representing at least 540 species of shore fishes collected over 10 years at 154 sites across the four volcanic archipelagos of French Polynesia; the Austral, Gambier, Marquesas and Society Islands, a 5,000,000 km^2 area. At present, 65% of the known shore fish species of these archipelagoes possess a DNA barcode associated with preserved, photographed, tissue sampled and cataloged specimens, and extensive collection locality data. This dataset represents one of the most comprehensive DNA barcoding efforts for a vertebrate fauna to date. Considering the challenges associated with the conservation of coral reef fishes and the difficulties of accurately identifying species using morphological characters, this publicly available library is expected to be helpful for both authorities and academics in various fields.

Background & Summary

DNA barcoding aims to identify individuals to the species level by using a short and standardized portion of a gene as a species tag[1]. This standardized procedure has revolutionized how biodiversity can be surveyed as the identification of a species then becomes independent of the level of taxonomic expertise of the collector[2], the life stage of the species[3,4] or the state of conservation of the specimen[5,6]. Due to its large spectrum of potential applications, DNA barcoding has been employed in a large array of scientific fields such as taxonomy[7], biogeography, biodiversity inventories[8] and ecology[9]; but see Hubert and Hanner for a review[10]. In the genomic era, this approach has been successfully applied to the simultaneous identification of multiple samples (*i.e.* the metabarcoding approach), extending its applications to surveys of whole ecological communities[11], but also monitoring species diet[12,13], identifying the presence of specific species in a region[14], or studying changes in the community through time by sampling environmental DNA[15,16].

By design, DNA barcoding has proved to be fast and accurate, but its accuracy is highly dependent on the completeness of DNA barcode reference libraries. These libraries turn surveys of Operational Taxonomic Units (OTUs) into species surveys through the assignment of species names to OTUs[17,18], hence giving meaning to data for ecologists, evolutionary biologists and stakeholders. Taxonomists increasingly provide DNA barcodes of new species they are describing; but thousands of species of shore fishes still lack this diagnostic molecular marker.

In the South Pacific, an early initiative led by the CRIOBE Laboratory was successfully carried out for French Polynesian coral reef fishes at the scale of one island, Moorea (Society Island)[19]. The fish fauna of Moorea's waters is one of the best known of the region given the historical operation of research laboratories and long term surveys[20,21]. The Moorea project revealed a high level of cryptic diversity in Moorea's fishes[19] and motivated the CRIOBE Laboratory to extend this biodiversity survey of shore fishes to the remaining islands of French Polynesia. French Polynesia (FP) is a 5,000,000 km^2 region located between 7° and 27° South Latitude that constitutes a priority area for conducting a barcoding survey. This region is species rich due to its position at the junction of several biogeographic areas with varying levels of endemism. For example, the Marquesas Islands (northeastern FP) rank as the third highest region of endemism for coral reef fishes in the Indo-Pacific (13.7%[22]). The Austral Islands (southwestern FP) and Gambier Islands (southeastern FP) host numerous southern subtropical endemic species[23–25]. Finally, the

Society Islands (western FP) possess the highest species richness (877 species) and the highest number of
widespread species in French Polynesia[26].
Here, we present the result of a large-scale effort to DNA barcode the shore fishes in French
Polynesia. Conducted between 2008 and 2014, a total of 154 sites were inventoried across these four
archipelagoes. Islands of varying ages and topographies were visited ranging from low-lying atolls to high
islands surrounded by a barrier reef, or solely fringing reefs. Furthermore, inventories were conducted
across different habitats at each island (*i.e.* sand bank, coral reefs, rubble, rocky, etc.). In total, 2,190
specimens were identified, preserved, photographed, tissue sampled, DNA barcoded and cataloged with
extensive metadata to build a library representing at least 540 species, 232 genera and 61 families of
fishes (Fig. 1). Merged with previous sampling efforts at Moorea, a total of 3,131 specimens now possess
a DNA barcode representing at least 645 nominal species for a coverage of approximately 65% of the
known shore fish species diversity of these four archipelagoes. These biodiversity surveys have already
resulted in the publication of updated species checklists[22,26] and in the description of 17 new species[27–34].
This comprehensive library for French Polynesia shore fishes will certainly benefit a wide community of
users with different interests, ranging from basic to applied science, and including fisheries management,
functional ecology, taxonomy and conservation. Furthermore, many newly detected taxa for science are
revealed here, along with complete collection data and DNA barcodes, which should facilitate their
formal description as new species. While shedding new light on the species diversity of the Pacific region,
this publicly available library is expected to fuel the development of DNA barcode libraries in the Pacific
Ocean and to provide more accurate results for the growing number of studies using DNA
metabarcoding in the Indo-West Pacific.
101

Methods

Sampling strategy

We explored a diversity of habitats across the four corners of French Polynesia with shallow and deep
SCUBA dives (down to 50–55 m) for a total of 154 sampled sites (Fig. 2, Table 1). A total of 2,190
specimens, representing at least 540 species, 232 genera and 61 families (Fig. 3a) have been collected
across four archipelagos representing the four corners of French Polynesia (FP), through six scientific
expeditions: Marquesas Islands (1) in 2008 at Mohotani and (2) in 2011 at every island of the archipelago
aboard the M.V. Braveheart (Clark Bank, Motu One, Hatutaa, Eiao, Motu Iti, Nuku-Hiva, Ua-Huka, Ua-
Pou, Fatu-Huku, Hiva-Oa, Tahuata, Fatu-Hiva; 52 sites), (3) in 2010 at Gambier Islands aboard the M.V.
Claymore (Mangareva, Taravai, Akamaru, and all along the barrier reef; 53 sites), (4) at Austral Islands in
2013 aboard the Golden Shadow (Raivavae, Tubuai, Rurutu, Rimatara, Maria Islands; 25 sites), (5) at
westernmost atolls of the Society Islands in 2014 aboard the M.V. Braveheart (Manuae and Maupiha'a;
20 sites). A sixth scientific expedition took place on Moorea's deep reefs in 2008 (Society Islands) as a
small scale scientific expedition that included the exploration and sampling of some of the deep reefs of
Moorea (53 to 56 m depth; 4 sites) (Fig. 2).
117

Specimen collection

Specimens were captured using rotenone (powdered root of the Derris plant) and spear guns while
SCUBA diving. These complementary sampling methods[35] allowed us to sample both the cryptic and
small fish fauna as well as the larger specimens of species not susceptible to rotenone collecting. Four

individuals per species were collected on average. Fishes were sorted and identified onboard to the
species level using identification keys and taxonomic references[23,36] and representative specimens of all
species collected were photographed in a fish photo tank to capture fresh color patterns, labeled and
tissue sampled for genetic analyses (fin clip or muscle biopsies preserved in 96% ethanol). The
photographed/sampled voucher specimens were preserved in 10% formalin (3.7% formaldehyde
solution) and later transferred into 75% ethanol for permanent archival storage. Preserved voucher
specimens and tissues were deposited and cataloged into the fish collection at the Museum Support
Center, National Museum of Natural History, Smithsonian Institution, Suitland, Maryland, USA.
Nomenclature follows Randall[23] and we followed recent taxonomic changes using the California
Academy of Sciences Online Eschmeyer's Catalog of Fishes[37].
132
DNA barcode sequencing
We extracted whole genomic DNA using QIAxtractor (QIAGEN, Crawley) and Autogen AutoGenPrep 965
according to manufacturer's protocols. A 655bp fragment of the cytochrome oxidase I gene (COI) was
amplified using Fish COI primers FISHCOILBC (TCAACYAATCAYAAAGATATYGGCAC) and FISHCOIHBC
(ACTTCYGGGTGRCCRAARAATCA) and Polymerase Chain Reaction (PCR) and Sanger sequencing protocols
as in Weigt et al.[38]. PCR products were Sanger sequenced bidirectionally and run on an ABI3730XL in the
Laboratories of Analytical Biology (National Museum of Natural History, Smithsonian Institution).
Sequences were edited using Sequencher 5.4 (Gene Codes) and aligned with Clustal W as implemented
in Barcode Of Life Datasystem (BOLD, http://www.boldsystems.org). Alignments were unambiguous with
no indels or frameshift mutations. A total of 2,190 DNA barcodes have been generated.
143
Specimen identification
All morphological identifications were revised as needed after the specimens were deposited in the
archival specimen collection to confirm initial identifications made in the field. Specimens of specific
groups like Antennaridae, Bythitidae, Chlopsidae or Muraenidae were revised by additional taxonomist
specialists (David Smith, John McCosker, Leslie W. Knapp, Werner Schwarzhans). After the morphological
identification, we used the Taxon-ID Tree tool and Barcode Index Numbers (BIN) discordance tools as
implemented in the Sequence Analysis module of BOLD to check every identification using the DNA
barcodes generated. The Taxon-ID tool consists of the construction of a neighbor-joining (NJ) tree using
K2P (Kimura 2 Parameter) distances by BOLD to provide a graphic representation of the species
divergence[39]. The BIN discordance tool uses the Refined Single Linkage algorithm (RESL[40]) to provide a
total number of OTUs.
155

Data Records

157 This library is composed of three main components: (1) voucher specimens archived in the national fish
158 collection at the Smithsonian Institution (Washington, DC), which were photographed in the field, (2)
159 complete collection data associated with each voucher specimen, and (3) DNA barcodes (Fig. 1).
160
161 All photographs, voucher collection numbers, DNA barcodes and collection data are publicly available in
162 BOLD[41] in the Container INDOF "Fish of French Polynesia" or by scientific expedition ("AUSTR", "GAMBA",

163 "MARQ", "MOH", "MOOP" and "SCILL") and in Figshare[42]. DNA barcodes have also been made available
164 in GenBank[43] and this database is accessible through the CRIOBE portal (http://fishbardb.criobe.pf).
165
The library fulfills the BARCODE data standard[44,45] which requires: 1) Species name, 2) Voucher data, 3)
Collection data, 4) Identifier of the specimen, 5) COI sequence of at least 500 bp, 6) PCR primers used to
generate the amplicon, 7) Trace files. In BOLD, each record in a project represents a voucher specimen
with its photographs, voucher collection numbers, associated sequences and extensive collection data
related to (1) the Voucher: Sample ID, Field ID, Museum ID, Institution Storing; (2) the Taxonomy:
Phylum, Class, Order, Family, Subfamily, Genus, species, Identifier, Identifier E-mail, Taxonomy Notes; (3)
Specimen Details: Sex, Reproduction, Life Stage, FAO Zone, Notes such as sizes of the specimens,
Voucher Status, and (4) Collection Data: Collectors, Collection Date, Continent, Country/Ocean*,
State/Province, Region, Sector, Exact Site, GPS Coordinates, Elevation, Depth, Depth Precision, GPS
Source, and Collection Notes[42].
176

Technical Validation
To test the robustness of our library, we first computed the distribution of the interspecific and
intraspecific variability for all the described species (Fig. 3b, c, d). We found that there is little to no
overlap in the distribution of divergence within and between species for the vast majority of the species
identified morphologically (mean intra-specific divergence 0.66, min: 0.00, max: 21.56; mean inter-
specific divergence 12.28, min: 0.00, max: 24.01). The RESL algorithm identified more BINs (617) than
nominal species identified morphologically (540). The morphological reexamination of specimens in light
of these results suggest that 65 taxa could be new species for science awaiting a formal description
(Online-only Table 1) as they are morphologically distinguishable from other species and possess unique
BIN numbers. Taxonomic paraphyly (*i.e.* potentially cryptic species) has been found for 18 additional
species (Table 2) as they are divided in 37 different BINs, while no morphological character has been
found so far to distinguish them. Finally, mixed genealogies between sister-species were observed for 17
species (Table 3), mostly between some of the Marquesan endemics and their closest relatives that are
not currently observed in the Marquesas Islands. Considering the maternal inheritance of the
mitochondrial genes and the very shallow genealogies involved (maximum K2P genetic distances lower
than 2%), both incomplete lineage sorting and past introgressive hybridization might be responsible of
the mixing of species genealogies in those 17 cases. In summary, 94% of the BINs match species
identified using morphological characters, meaning that it was possible to successfully identify a species
using DNA barcodes in 94% of the cases.
196

Usage Notes
This Barcode release dataset is freely available to use in barcoding or metabarcoding surveys for
specimen identification. Several approaches can be considered:
(1) directly downloading the sequences in fasta format, and working offline by merging this dataset with
an ongoing barcoding project;
(2) working online, through the BOLD website (registration is free), and merging the Container INDOF
"Fish of French Polynesia" or parts of the scientific expeditions (Table 1) with an ongoing BOLD project;
(3) through online identification tools, as data are indexed in both BOLD and Genbank databases. This

library will be considered when any queries of molecular identification will be made through the identification engine of BOLD (http://www.boldsystems.org/index.php/IDS-OpenIdEngine) or the standard nucleotide Basic Local Alignment Search Tool (BLAST, https://blast.ncbi.nlm.nih.gov/). In the same manner, this dataset should also be indexed in the MIDORI database[46,47]. Composed of both endemic and widespread species, this library is expected to benefit a large community from academics to authorities who use molecular data to monitor and survey biodiversity.

Acknowledgements

This work was financially supported by the French National Agency for Marine Protected Area in France ('Pakaihi I Te Moana' expedition), the ANR IMODEL and Contrat de Projet Etat-Territoire in French Polynesia and the French Ministry for Environment, Sustainable Development and Transport (MEDDTL) ('CORALSPOT' expeditions), the Living Oceans Foundation ('Australs' expedition) and the LabEx CORAIL and the GOPS, ('Scilly' expedition) and the Gordon and Betty Moore Foundation (Mo'orea Biocode Project). Additional funding was provided by the IFRECOR in French Polynesia and the TOTAL Foundation. We are grateful to T. Frogier, P. Mery and the Centre Plongée Marquises (Xavier (Pipapo) and Marie Curvat), for their field assistance in the Gambier, the Marquesas, the Australs and Scilly Islands along with the crew of the *Claymore II*, *Braveheart* and the *Golden Shadow*. We thank the Ministère de l'Environnement de Polynésie, the Délégation à la Recherche Polynésie, the Mairie of Nuku-Hiva, and the people of the Marquesas Islands for their kind and generous support of the project as we traveled throughout the islands. We thank Jerry Finan, Erika Wilbur, Shirleen Smith, Kris Murphy, David Smith and Sandra Raredon of the Division of Fishes (National Museum of Natural History) for assistance in preparations for the trip and processing specimens and Jeffrey Hunt and Kenneth Macdonald III and Meaghan Parker Forney of the Laboratories of Analytical Biology (Smithsonian Institution) for assistance in molecular analysis of samples. Finally, we thank the staff of the CRIOBE and particularly Yannick Chancerelle for logistical support in French Polynesia. Specimens were collected under the permit "Permanent agreement, Délégation à la Recherche, French Polynesia".

Author contributions

E.D.T. drafted the first manuscript, J.T.W., D.P., A.D., and N.H. provided extensive edits and comments. E.D.T, J.T.W., T.C., R.G., M.K., T.L.M., J.M., G.M.-T., V.P., P.P., P.S., G.S., N.T., M.V. and S.P. collected fish specimens. E.D.T., J.T.W., D.P., A.D., N.H., J.V., B.E., C. M., L.W. and S.P. produced DNA barcodes and cleaned the database. R.G. and S.P. financed the scientific expeditions. C.M. and S.P. financed the sequencing. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Figure Legends

Figure 1. Overview of data generation. From collection of specimen to the validation of data generation.

Figure 2. Sampling localities across French Polynesia. The number of sampling sites and the number of specimens collected are displayed for each archipelago. Several sampling localities may be represented by a single dot due to the geographic scale of French Polynesia. Map data: Google, DigitalGlobe.

Figure 3. **Species diversity and distribution of genetic distance across the DNA barcode library.** (a) Species diversity by family for the four archipelagoes sampled; (b) Distribution of maximum intraspecific distances (K2P, percent); (c) Distribution of nearest neighbor distances (K2P, percent); (d) Relationship between maximum intraspecific and nearest neighbor distances. Points above the diagonal line indicate species with a barcode gap.

Tables Legends

Table 1. Overview of the dataset. Number of specimens and species collected for each scientific expedition. Sampling effort expressed in number of sampling days and number of sites.

Table 2. Potential cryptic species. Species with number of specimens collected displaying taxonomic paraphyly most likely representing undescribed cryptic species. Sample ID includes sampling location (AUST: Austral Islands, GAMB: Gambier Islands, MARQ and MOH: Marquesas Islands, SCIL and MOOP: Society Islands).

Table 3. Species displaying either incomplete lineage sorting or shallow inter-species divergence. Mean and Maximum intra-Species distances (Mean Intra-Sp and Max Intra-Sp), and Kimura 2 Parameter distances from the Nearest Neighbour (NN).

Online-only Table 1. Potential new species detected in the dataset. Specimens which were identified only to the genus level and which represent potentially new species waiting to be described. Number of specimens included in each Barcode Index Number (BIN).

268 **Tables**

269 **Table 1.**

BOLD project	Geographical location	No. of specimen collected	No. of species collected	Sampling effort (No. of sampling days / No. of sites)
AUSTR	Austral Islands	560	263	12 / 25
GAMBA	Gambier Islands	705	290	18 / 53
MARQ	Marquesas Islands	386	182	18 / 41
MOH	Marquesas Islands	190	107	5 / 11
MOOP	Society Islands	42	27	4 / 4
SCIL	Society Islands	309	213	8 / 20

 Table 2.

BINs	Taxa	No. of specimens
BOLD:AAF8427	*Apogon crassiceps*	2
BOLD:ABW7007	*Apogon crassiceps*	4
BOLD:ACE7901	*Apogon crassiceps*	1
BOLD:ACX1964	*Apogon doryssa*	1
BOLD:ABW8494	*Apogon doryssa*	2
BOLD:AAF5636	*Aporops bilinearis*	1
BOLD:AAF5637	*Aporops bilinearis*	4
BOLD:AAD2580	*Centropyge flavissima*	2
BOLD:AAD9019	*Centropyge flavissima*	6
BOLD:ACD1956	*Fusigobius duospilus*	5
BOLD:AAD1050	*Fusigobius duospilus*	1
BOLD:AAA6306	*Gnatholepis cauerensis*	9
BOLD:AAC6155	*Gnatholepis cauerensis*	5
BOLD:ACC5235	*Gymnothorax melatremus*	3
BOLD:AAC8364	*Gymnothorax melatremus*	5
BOLD:AAF0704	*Leiuranus semicinctus*	3
BOLD:AAL6561	*Leiuranus semicinctus*	2
BOLD:ACD1820	*Myrophis microchir*	1
BOLD:AAE0976	*Myrophis microchir*	2
BOLD:AAB3862	*Parupeneus multifasciatus*	6
BOLD:ACD1989	*Parupeneus multifasciatus*	3
BOLD:ACD1988	*Priolepis triops*	3
BOLD:AAX7961	*Priolepis triops*	1
BOLD:AAB4082	*Pristiapogon kallopterus*	1
BOLD:ABZ7996	*Pristiapogon kallopterus*	7
BOLD:ACC5180	*Pseudocheilinus octotaenia*	10
BOLD:AAD3038	*Pseudocheilinus octotaenia*	9
BOLD:AAB4821	*Pterocaesio tile*	4
BOLD:ACK9118	*Pterocaesio tile*	1
BOLD:ACP9778	*Scolecenchelys gymnota*	1
BOLD:AAJ8783	*Scolecenchelys gymnota*	2
BOLD:AAC7090	*Stegastes fasciolatus*	11
BOLD:ABZ0285	*Stegastes fasciolatus*	2
BOLD:ACC5053	*Uropterygius kamar*	1
BOLD:ACC5109	*Uropterygius kamar*	1
BOLD:ACD1642	*Uropterygius macrocephalus*	1
BOLD:AAU1965	*Uropterygius macrocephalus*	2

271 **Table 3.**

Family	Species	Mean Intra-Sp	Max Intra-Sp	Nearest Neighbour	Nearest Species	Distance to NN
Acanthuridae	*Acanthurus reversus*	0.08	0.15	AUSTR453-13	*Acanthurus olivaceus*	0
Holocentridae	*Myripristis earlei*	0.28	0.62	SCILL065-15	*Myripristis berndti*	0
Monacanthidae	*Pervagor marginalis*	0.36	0.62	SCILL083-15	*Pervagor aspricaudus*	0
Tetraodontidae	*Canthigaster criobe*	0	0	MOH030-16	*Canthigaster janthinoptera*	0
Mullidae	*Mulloidichthys mimicus*	0.52	0.52	AUSTR089-13	*Mulloidichthys vanicolensis*	0.17
Pomacentridae	*Chromis abrupta*	0	0	SCILL209-15	*Chromis margaritifer*	0.31
Labridae	*Coris marquesensis*	0	0	SCILL040-15	*Coris gaimard*	0.46
Apogonidae	*Ostorhinchus relativus*	N/A	0	SCILL142-15	*Ostorhinchus angustatus*	0.93
Tetraodontidae	*Canthigaster rapaensis*	0.21	0.31	MARQ456-12	*Canthigaster marquesensis*	1.1
Pomacentridae	*Abudefduf conformis*	0.15	0.15	GAMBA844-12	*Abudefduf sexfasciatus*	1.24
Monacanthidae	*Cantherhines nukuhiva*	0.15	0.31	GAMBA711-12	*Cantherhines sandwichiensis*	1.4
Pomacentridae	*Plectroglyphidodon sagmarius*	0.08	0.15	AUSTR222-13	*Plectroglyphidodon imparipennis*	1.56
Holocentridae	*Sargocentron caudimaculatum*	0.68	1.1	SCILL104-15	*Sargocentron tiere*	1.57
Acanthuridae	*Zebrasoma rostratum*	0	0	AUSTR376-13	*Zebrasoma scopas*	1.72
Apogonidae	*Apogon marquesensis*	0.23	0.31	GAMBA657-12	*Apogon susanae*	1.88
Chaetodontidae	*Chaetodon flavirostris*	0.08	0.15	SCILL269-15	*Chaetodon lunula*	1.88
Chaetodontidae	*Chaetodon lunula*	0.1	0.15	GAMBA555-12	*Chaetodon flavirostris*	1.88

Online-only Table 1. Potential new species detected in the dataset. Specimens which were identified only to the genus level and which represent potentially new species waiting to be described. Number of specimens included in each Barcode Index Number (BIN).

Sample ID	Family	Genus	Species	No. of specimens
AUST-419	Antennariidae	*Antennatus*	sp1	1
AUST-570, GAM-355	Antennariidae	*Antennatus*	sp2	2
MARQ-022, MOH-087	Antennariidae	*Antennatus*	sp3	2
SCIL-293	Antennariidae	*Antennatus*	sp4	1
MARQ-105, MARQ-106, MARQ-107	Apogonidae	*Fowleria*	sp	3
MARQ-380, MARQ-381, MOH-068	Apogonidae	*Gymnapogon*	sp	3
AUST-142, AUST-143	Apogonidae	*Pseudamiops*	sp	2
AUST-470, AUST-471	Apogonidae	*Siphamia*	sp	2
MARQ-177, MARQ-208, MOH-040	Blenniidae	*Blenniella*	sp	3
AUST-242, GAM-791, GAM-792, MARQ-139, MARQ-140	Blenniidae	*Cirripectes*	*sp*	5
GAM-278, GAM-279, SCIL-017, SCIL-021, SCIL-058, SCIL-288, SCIL-322	Blenniidae	*Enchelyurus*	*sp*	7
AUST-407, AUST-408, AUST-409	Blenniidae	*Entomacrodus*	sp	3
MARQ-184, MARQ-187, MARQ-378, MARQ-379	Blenniidae	*Rhabdoblennius*	sp	4
MOOP-028	Chlopsidae	*Kaupichthys*	sp	1
AUST-600	Congridae	*Ariosoma*	sp1	1
MARQ-318	Congridae	*Ariosoma*	sp2	1
MARQ-314, MARQ-315, MARQ-316	Congridae	*Gnathophis*	sp	3
MARQ-397, MOH-062	Creediidae	*Chalixodytes*	sp	1
AUST-427, AUST-538, AUST-539	Creediidae	*Crystallodytes*	*sp*	3
AUST-413	Gobiesocidae	*NA*	sp	1
AUST-159, AUST-305	Gobiesocidae	*Pherallodus*	sp1	2
AUST-532, AUST-533, AUST-534	Gobiesocidae	*Propherallodus*	sp	3
AUST-082	Gobiidae	*Bryaninops*	sp	1
GAM-374, GAM-375	Gobiidae	*Cabillus*	sp	2
MARQ-430, MOH-129, MOH-211	Gobiidae	*Callogobius*	sp	3
AUST-303, GAM-379	Gobiidae	*Eviota*	sp1	1
GAM-697	Gobiidae	*Eviota*	sp2	1
SCIL-038	Gobiidae	*Eviota*	sp3	1
AUST-346, AUST-347	Gobiidae	*Gobiodon*	sp	2
MARQ-097, MARQ-098	Gobiidae	*Gobiodon*	sp	2
SCIL-240	Gobiidae	*Gobiodon*	sp	1
AUST-566, AUST-567, AUST-568, GAM-364	Gobiidae	*Paragobiodon*	sp	4
MARQ-363, MARQ-364, MARQ-365	Gobiidae	*Pleurosicya*	sp1	3
MOOP-007	Gobiidae	*Pleurosicya*	sp2	1
SCIL-237	Gobiidae	*Priolepis*	sp	1

AUST-032	Gobiidae	*Silhouettea*	sp	1
MOOP-047	Gobiidae	*Sueviota*	sp	1
AUST-446	Gobiidae	*Trimma*	sp1	1
MARQ-435, MARQ-436	Gobiidae	*Trimma*	sp2	2
GAM-32	Gobiidae	*Trimmatom*	sp1	1
GAM-33	Gobiidae	*Trimmatom*	sp2	1
MOOP-002	Gobiidae	*Trimmatom*	sp3	1
AUST-423, AUST-424, AUST-425, AUST-544	Isonidae	*Iso*	sp1	4
GAM-521, GAM-522	Isonidae	*Iso*	sp2	2
MOH-079	Lethrinidae	*Gymnocranius*	sp	1
AUST-383, GAM-176, GAM-177, GAM-178, MOH-200, SCIL-299	Moringuidae	*Moringua*	sp1	6
SCIL-300	Moringuidae	*Moringua*	sp2	1
MARQ-505	Muraenidae	*Gymnothorax*	sp1	1
SCIL-335	Muraenidae	*Gymnothorax*	sp2	1
AUST-573, GAM-709	Ophidiidae	*Brotula*	sp	2
AUST-297, AUST-298, AUST-299, AUST-300, GAM-761	Pempheridae	*Pempheris*	sp1	5
MARQ-063, MARQ-165, MARQ-166, MARQ-167, MARQ-276, MARQ-382, MOH-178, MOH-179	Pempheridae	*Pempheris*	sp2	8
AUST-308, AUST-309, AUST-310, AUST-311	Pomacentridae	*Stegastes*	sp	4
MOOP-018, MOOP-019, MOOP-034, MOOP-035	Pseudochromidae	*Lubbockichthys*	sp	4
GAM-599, GAM-600	Scorpaenidae	*Scorpaenodes*	sp1	2
MOH-137, MOH-151	Scorpaenidae	*Scorpaenodes*	sp2	2
GAM-56, GAM-569, GAM-574, GAM-58	Scorpaenidae	*Sebastapistes*	sp1	4
MARQ-328, SCIL-114	Scorpaenidae	*Sebastapistes*	sp2	1
MARQ-046, MOH-027	Syngnathidae	*Doryrhamphus*	sp	1
MARQ-321	Synodontidae	*Synodus*	sp	1
AUST-011, AUST-048	Tripterygiidae	*Enneapterygius*	sp1	1
AUST-360, AUST-057, AUST-058, AUST-059	Tripterygiidae	*Enneapterygius*	sp2	4
GAM-68, GAM-69, GAM-123, GAM-124, GAM-125	Tripterygiidae	*Enneapterygius*	sp3	3
GAM-002	Tripterygiidae	*Enneapterygius*	sp4	1
SCIL-112, SCIL-133, SCIL-156	Tripterygiidae	*Enneapterygius*	sp5	3

References

1. Hebert, P. D. N. & Gregory, T. R. The promise of DNA barcoding for taxonomy. *Syst. Biol.* **54,** 852–859 (2005).

2. Garnett, S. T. & Christidis, L. Taxonomy anarchy hampers conservation. *Nat. News* **546,** 25 (2017).

3. Hubert, N., Delrieu-Trottin, E., Irisson, J. O., Meyer, C. & Planes, S. Identifying coral reef fish larvae through DNA barcoding: A test case with the families Acanthuridae and Holocentridae. *Mol. Phylogenet. Evol.* **55,** 1195–1203 (2010).

4. Ko, H.-L. *et al.* Evaluating the accuracy of morphological identification of larval fishes by applying DNA barcoding. *PLoS One* **8,** e53451 (2013).

5. Wong, E. H.-K. & Hanner, R. H. DNA barcoding detects market substitution in North American seafood. *Food Res. Int.* **41,** 828–837 (2008).

6. Holmes, B. H., Steinke, D. & Ward, R. D. Identification of shark and ray fins using DNA barcoding. *Fish. Res.* **95,** 280–288 (2009).

7. Riedel, A., Sagata, K., Suhardjono, Y. R., Tänzler, R. & Balke, M. Integrative taxonomy on the fast track-towards more sustainability in biodiversity research. *Front. Zool.* **10,** 15 (2013).

8. Monaghan, M. T. *et al.* Accelerated species inventory on Madagascar using coalescent-based models of species delineation. *Syst. Biol.* **58,** 298–311 (2009).

9. Tänzler, R., Sagata, K., Surbakti, S., Balke, M. & Riedel, A. DNA barcoding for community ecology-how to tackle a hyperdiverse, mostly undescribed Melanesian fauna. *PLoS One* **7,** e28832 (2012).

10. Hubert, N. & Hanner, R. DNA barcoding, species delineation and taxonomy: a historical perspective. *DNA barcodes* **3,** 44–58 (2015).

11. Leray, M. & Knowlton, N. DNA barcoding and metabarcoding of standardized samples reveal patterns of marine benthic diversity. *Proc. Natl. Acad. Sci.* **112,** 2076–2081 (2015).

12. Leray, M., Boehm, J. T., Mills, S. C. & Meyer, C. P. Moorea BIOCODE barcode library as a tool for understanding predator-prey interactions: insights into the diet of common predatory coral reef fishes. *Coral Reefs* **31,** 383–388 (2012).

13. Leray, M., Meyer, C. P. & Mills, S. C. Metabarcoding dietary analysis of coral dwelling predatory fish demonstrates the minor contribution of coral mutualists to their highly partitioned, generalist diet. *PeerJ* **3,** e1047 (2015).

14. Bakker, J. *et al.* Environmental DNA reveals tropical shark diversity in contrasting levels of anthropogenic impact. *Sci. Rep.* **7,** 16886 (2017).

15. Stat, M. *et al.* Ecosystem biomonitoring with eDNA: metabarcoding across the tree of life in a tropical marine environment. *Sci. Rep.* **7,** 12240 (2017).

16. DiBattista, J. D. *et al.* Assessing the utility of eDNA as a tool to survey reef-fish communities in the Red Sea. *Coral Reefs* **36,** 1245–1252 (2017).

17. Hebert, P. D. N. *et al.* A DNA 'Barcode Blitz': Rapid digitization and sequencing of a natural history collection. *PLoS One* **8,** e68535 (2013).

18. Schmidt, S., Schmid-Egger, C., Morinière, J., Haszprunar, G. & Hebert, P. D. N. DNA barcoding largely supports 250 years of classical taxonomy: identifications for Central European bees (Hymenoptera, Apoidea *partim*). *Mol. Ecol. Resour.* **15,** 985–1000 (2015).

19. Hubert, N. *et al.* Cryptic diversity in indo-pacific coral-reef fishes revealed by DNA-barcoding provides new support to the centre-of-overlap hypothesis. *PLoS One* **7,** (2012).

314   20.   Lamy, T., Legendre, P., Chancerelle, Y., Siu, G. & Claudet, J. Understanding the spatio-temporal
315   response of coral reef fish communities to natural disturbances: insights from beta-diversity
316   decomposition. *PLoS One* **10,** e0138696 (2015).
317   21.   Galzin, R. *et al.* Long term monitoring of coral and fish assemblages (1983 - 2014) in Tiahura reefs,
318   Moorea, French Polynesia. *Cybium* **40,** 31–41 (2016).
319   22.   Delrieu-Trottin, E. *et al.* Shore fishes of the Marquesas Islands, an updated checklist with new
320   records and new percentage of endemic species. *Check List* **11,** (2015).
321   23.   Randall, J. E. *Reef and Shore Fishes of the South Pacific : New Caledonia to Tahiti and the Pitcairn
322   Islands*. **1,** (University of Hawai'i Press Honolulu, 2005)
323   24.   Randall, J. E. & Cea, A. *Shore Fishes of Easter Island*. (University of Hawai'i Press, 2011)
324   25.   Delrieu-Trottin, E. *et al.* Evidence of cryptic species in the blenniid *Cirripectes alboapicalis* species
325   complex, with zoogeographic implications for the South Pacific. *Zookeys* **810,** 127–138 (2018).
326   26.   Siu, G. *et al.* Shore fishes of French polynesia. *Cybium* **41,** 1–34 (2017).
327   27.   Williams, J. T., Delrieu-Trottin, E. & Planes, S. A new species of Indo-Pacific fish, *Canthigaster
328   criobe*, with comments on other Canthigaster (Tetraodontiformes: Tetraodontidae) at the
329   Gambier Archipelago. *Zootaxa* **3523,** 80–88 (2012).
330   28.   Tornabene, L., Ahmadia, G. N. & Williams, J. T. Four new species of dwarfgobies (Teleostei:
331   Gobiidae: *Eviota*) from the Austral, Gambier, Marquesas and Society Archipelagos, French
332   Polynesia. *Syst. Biodivers.* **11,** 363–380 (2013).
333   29.   Williams, J. T., Delrieu-Trottin, E. & Planes, S. Two new fish species of the subfamily Anthiinae
334   (Perciformes, Serranidae) from the Marquesas. *Zootaxa* **3647,** 167–180 (2013).
335   30.   Delrieu-Trottin, E., Williams, J. T. & Planes, S. *Macropharyngodon pakoko*, a new species of
336   wrasse (Teleostei: Labridae) endemic to the Marquesas Islands, French polynesia. *Zootaxa* **3857,**
337   433–443 (2014).
338   31.   McCosker, J. E. & Hibino, Y. A review of the finless snake eels of the genus *Apterichtus*
339   (Anguilliformes: Ophichthidae), with the description of five new species. *Zootaxa* **3941,** 49–78
340   (2015).
341   32.   Polanco, F. A., Acero, P. A. & Betancur-R., R. No longer a circumtropical species: revision of the
342   lizardfishes in the *Trachinocephalus myops* species complex, with description of a new species
343   from the Marquesas Islands. *J. Fish Biol.* **89,** 1302–1323 (2016).
344   33.   Viviani, J., Williams, J. T. & Planes, S. Two new pygmygobies (Percomorpha : Gobiidae : *Trimma*)
345   from French Polynesia. *J. Ocean Sci. Found.* **23,** 1–11 (2016).
346   34.   Williams, J. T. & Viviani, J. *Pseudogramma polyacantha* complex (Serranidae, tribe Grammistini):
347   DNA barcoding results lead to the discovery of three cryptic species, including two new species
348   from French Polynesia. *Zootaxa* **4111,** 246–260 (2016).
349   35.   Williams, J. T. *et al.* Checklist of the shorefishes of Wallis Islands (Wallis and Futuna French
350   Territories , South-Central Pacific). *Cybium* **30,** 247–260 (2006).
351   36.   Bacchet, P., Zysman, T. & Lefevre, Y. *Guide des poissons de Tahiti et ses iles.* (Au vent des îles,
352   2006)
353   37.   Eschmeyer, W. N., Fricke, R. & van der Laan, R. Eschmeyer's Catalog of Fishes electronic version.
354   *http://researcharchive.calacademy.org/research/Ichthyology/catalog/fishcatmain.asp* (2018).
38.   Weigt, L. A., Driskell, A. C., Baldwin, C. C. & Ormos, A. in *DNA barcodes* 109–126 (Springer, 2012)

39.   Ratnasingham, S. & Hebert, P. D. N. BOLD: The Barcode of Life Data System
(www.barcodinglife.org). *Mol. Ecol. Notes* **7,** 355–364 (2007).
40.   Ratnasingham, S. & Hebert, P. D. N. A DNA-based registry for all animal species: the Barcode
Index Number (BIN) system. *PLoS One* **8,** e66213 (2013).
41.   *BOLD DS-INDOF* https://doi.org/10.5883/DS-INDOF (2019).
42.   Delrieu-Trottin et al. Figshare https://doi.org/10.6084/m9.figshare.7923905 (2019).
43.   NCBI Nucleotide https://www.ncbi.nlm.nih.gov/nucleotide/ Accession numbers KC567661 -
KC567663, KC684990, KC684991, KU905709-KU905727, KY570698, KY570703 - KY570705,
KY570708, KY683549, MH707846 - MH707881, MK566774 - MK567153, MK656969 - MK658713
(2019)
44.   Hubert, N. *et al.* Identifying Canadian freshwater fishes through DNA barcodes. *PLoS One* **3,**
(2008).
45.   Hubert, N. & Hanner, R. DNA Barcoding, species delineation and taxonomy: a historical
perspective. *DNA Barcodes* **3,** 44–58 (2015).
46.   Machida, R. J., Leray, M., Ho, S. L. & Knowlton, N. Metazoan mitochondrial gene sequence
reference datasets for taxonomic assignment of environmental samples. *Sci. Data* **4,** 1–7 (2017).
47.   Leray, M., Ho, S. L., Lin, I. J. & Machida, R. J. MIDORI server: a webserver for taxonomic
assignment of unknown metazoan mitochondrial-encoded sequences using a curated database.
Bioinformatics **34,** 3753–3754 (2018).
375

N

Motu one

Clark Bank

Motu Iti

Marquesas Islands
(52 sites - 574 specimens)

Hiva Oa

Fatu Hiva

Society Islands
(24 sites - 351 specimens)

Manuae

Maupiha'a

Moorea

Austral Islands
(25 sites - 560 specimens)

Maria

Rimatara

Rurutu

100 km

Tubuai

Raivavae

Taravai

Mangareva

Gambier Islands
(53 sites - 705 specimens)