The biochemical activities of the Saccharomyces cerevisiae Pif1 helicase are regulated by its

N-terminal domain

David G. Nickens[†], Christopher W. Sausen[†], and Matthew L. Bochman*

Affiliation: Molecular and Cellular Biochemistry Department, Indiana University, Bloomington, IN

47405 USA

[†]**These authors contributed equally to this work.**

*Corresponding author:

Matthew L. Bochman

212 S. Hawthorne Dr.

Simon Hall MSB1, room 405B

Bloomington, IN 47405

bochman@indiana.edu

Abstract

PIF1 family helicases represent a highly conserved class of enzymes involved in multiple aspects of

genome maintenance. Many PIF1 helicase are multi-domain proteins, but the functions of their

non-helicase domains are poorly understood. Here, we characterized how the N-terminal domain

(NTD) of the *Saccharomyces cerevisiae* Pif1 helicase affects its functions both *in vivo* and *in vitro*.

Removal of the Pif1 NTD alleviated the toxicity associated with Pif1 over-expression in yeast.

Biochemically, the N-terminally truncated Pif1 (Pif1ΔN) retained *in vitro* DNA binding, DNA

unwinding, and telomerase regulation activities, but these activities differed markedly from those

displayed by full-length recombinant Pif1. However, Pif1ΔN was still able to synergize with the

Hrq1 helicase to inhibit telomerase activity *in vitro*, similar to full-length Pif1. These data impact our

understanding of PIF1 helicase evolution and the roles of these enzymes in the maintenance of

genome integrity.

Key words: DNA helicase, *Saccharomyces cerevisiae*, Pif1, telomerase, telomere

1. Introduction

DNA helicases are enzymes that couple DNA binding and ATP hydrolysis to unwind

double-stranded DNA (dsDNA) into its component single strands [1]. This activity is vital to many

processes involved in the maintenance of genome integrity, including DNA replication,

recombination, and repair, transcription, and telomere maintenance [2]. As such, the genomes of all

organisms encode many helicases to fulfill these roles, from approximately 10 in typical prokaryotes

to 100 in eukaryotes [3]. These enzymes can be grouped into superfamilies based on the primary

sequence, structural, and functional homology of their helicase domains [4].

However, many DNA helicases are multi-domain enzymes with additional activities conferred

by functional modules other than their helicase motors. For instance, some helicases in the RecQ

family also contain an exonuclease domain (*e.g.*, WRN [5]), a DNA strand exchange domain (*e.g.*,

RECQL4 [6]), and/or domains involved in DNA and protein interactions (RQC and HRDC domains,

respectively) [7]. These accessory domains may be used in tandem with helicase activity to ensure

genome integrity, or they may be functionally separable. The latter is the case with RECQL4, which
has an N-terminal domain (NTD) necessary for the initiation of DNA replication with no
dependence on helicase activity [8].
Similar to RecQ helicases, PIF1 family helicases are also often multi-domain proteins, especially
those found in eukaryotes [9, 10]. They tend to have a central helicase domain with NTDs and
C-terminal domains (CTDs) that significantly vary in length and sequence conservation, even
among closely related homologs. Further, with very few exceptions, these accessory domains have
no identified or predicted function. Among the two *Saccharomyces cerevisiae* PIF1 family helicases,
Rrm3 and Pif1, it is known that the NTD of Rrm3 is essential for its functions *in vivo* because
truncation of the Rrm3 NTD phenocopies the null allele [11]. However, the Rrm3 NTD can be fused
to the *S. cerevisiae* Pif1 helicase domain or the helicase domain of bacterial Pif1 proteins, and these
chimeras can rescue the synthetic lethality of cells lacking Rrm3 and, for instance, the gene encoding
the Srs2 helicase [12]. Thus, in this experimental set up, the PIF1 helicase domains are generic and
interchangeable motor modules, while the Rrm3 NTD is vital for genome integrity.
We recently characterized the PIF1 helicase from the thermophilic bacterium *Thermotoga elfii*
(TePif1) and investigated the function of its C-terminal WYL domain [12]. We found that the WYL
domain contains a single-stranded DNA (ssDNA) binding motif that regulates and couples the
ATPase and helicase activities of the enzyme. Here, we sought to extend this line of investigation to
the NTD of the *S. cerevisiae* Pif1 helicase, which currently has an unknown function. Using both *in*
vivo and *in vitro* approaches, we found that the Pif1 NTD is involved in the toxicity of Pif1
over-expression and impacts the ability of Pif1 to regulate telomerase activity. These data and
similar investigations will further illuminate the conserved and divergent functions of PIF1 family
helicases across species, leading to a better understanding of the roles of these important enzymes in
genome integrity.
67
2. Materials and Methods
2.1. Yeast strains, media, and other reagents
S. cerevisiae strain JBBY26, a derivative of BCY123 (*MATa can1 ade2 trp1 ura3-52 his3 leu2-3, 112*
pep4::HIS3 prb1::LEU2 bar1::HISG, lys2::pGAL1/10-GAL4) [13], harbors a dual overexpression plasmid
for *TLC1* and *EST2*. The other yeast strains used in this work are derived from W303 (*MATa*
leu2-3,112 trp1-1 can1-100 ura3-1 ade2-1 his3-11,15), which was a gift from Peter Walter (UCSF).
Escherichia coli strain Rosetta 2 (DE3) pLysS (Novagen) was used for the overexpression of
SUMO-tagged Pif1, Pif1ΔN, and SUMO protease (see Table 1 for expression plasmid data). For
propagation and Est2/*TLC1* overproduction, JBBY26 cultures were grown in SC-Ura drop-out
media. Rosetta cells were maintained on LB medium supplemented with 50 μg/mL kanamycin and
34 μg/mL chloramphenicol. Liquid cultures were grown in 2× YT medium (1.6% w/v tryptone, 1%
w/v yeast extract, 0.5% NaCl, pH 7.0) for protein overproduction and supplemented with the same
antibiotics. Radiolabeled [α-^{32}P]TTP and [γ-^{32}P]ATP were purchased from PerkinElmer Life Sciences.
All dNTPs were purchased from New England Biolabs (Ipswich, MA). Oligonucleotides were
purchased from IDT (Coralville, IA), and the Tel15, Tel30, and Tel50 primers used for quantitative
telomerase assays were PAGE-purified. The sequences of all of the oligonucleotides used are listed
in Table 2. Chemical reagents were purchased from Sigma or DOT Scientific (Burton, MI). All
recombinant proteins were purified as described in [14].

Table 1. Plasmids used in this study.

Name	Description
pESC-URA	Multi-copy vector enabling epitope tagging of genes cloned under the control of the bidirectional *GAL1,10* promoter
pMB526	Pif1 cloned into pESC-URA, enabling galactose induction and epitope-tagged with FLAG
pMB540	Pif1ΔN cloned into pESC-URA, enabling galactose induction and epitope-tagged with FLAG
pSUMO-Pif1	Nuclear isoform of Pif1 cloned into the pSUMO vector
pSUMO-Pif1ΔN	Pif1ΔN cloned into the pSUMO vector

89 **Table 2. Oligonucleotides used in this study.**

Name	Sequence (5′-3′)	Purpose
MB1571	*rGrGrUrGrUrGrGrUrGrUrGrGrGrUrGrUrG rGrUrGrUrGrGrGrUrCrArCrArGrUrGrArGr UrGrUrArUrCrGrCrArArG	RNA-DNA hybrid fork substrate
MB1572	GAACGCTATGTGAGTGACACTGGGTCAC CACACCCACACCACACC	Fork substrates
MB1596	GGTGTGGTGTGGGTGTGGTGTGGGTCAC AGTGAGTGTATCGCAAG	DNA-DNA fork substrate
Tel15	TGTGGTGTGTGTGGG	EMSAs and *in vitro* telomerase assays
Tel30	CGCCATGCTGATCCGTGTGGTGTGTGTGG G	EMSAs
Tel50	GTGTGGGTGTGGTGTGGGTGTGGTGTGG GTGTGTGGTGTGGTGTGTGTGG	*In vitro* telomerase assays

* "r" denotes an RNA base; EMSA = electrophoretic mobility shift assay

2.2. Recombinant proteins

Plasmid pSUMO-Pif1 (Table 1) was used for the over-expression of SUMO-tagged Pif1 and was

a gift from Kevin Raney. This plasmid was also used as the template to create pSUMO-Pif1ΔN to

over-express an N-terminal truncation of Pif1 lacking the first 233 amino acids of the helicase

(Pif1ΔN) in *E. coli*. The over-expression and purification of these proteins proceeded as previously

described for full-length Pif1 [14]. Recombinant Hrq1 was generated using baculovirus-infected

insect cells as previously described [6, 15]. All recombinant helicase preparations were tested for

ATPase activity and the absence of contaminating nuclease activity before use in biochemical assays.

2.3. Telomerase Assays

In vitro telomerase assays were performed as described in [14] using telomerase-enriched
extracts from *S. cerevisiae* prepared by DEAE fractionation of clarified lysates [16, 17]. Each
telomerase preparation was titrated to standardize activity levels before use in experiments.
Reaction products were separated on 16% 19:1 acrylamide:bis-acrylamide gels containing 6 M urea.
The gels were run at 2500 V for 120 min, dried, and imaged and quantified using a Typhoon 9500
scanner with ImageQuant software. Total activity was measured by quantifying the densitometry
for each telomerase extension product band on a gel using ImageQuant. The sum of the measured
densitometry values in a lane is reported as the total activity. Bands were corrected for the number
of dT residues (*i.e.*, the amount of α-^{32}P-dTTP incorporation) and normalized to a loading control to
generate corrected pixel values.

2.4. Helicase Assays

Fork substrates for helicase assays were constructed by incubating two partially
complementary oligonucleotides (both at 1 μM) overnight at 37°C for annealing. In each case, the
helicase loading strand was end-labeled with [γ-^{32}P]ATP by T4 PNK (NEB) using the manufacturer's
instructions. Labeled DNA was separated from unincorporated ^{32}P using G-50 micro-columns (GE
Healthcare) and added to unlabeled partially complementary oligonucleotides in annealing buffer
(20 mM Tris–HCl, pH 8.0, 4% glycerol, 0.1 mM EDTA, 40 μg/ml BSA, and 10 mM DTT) [15]. The
DNA-DNA fork and its RNA-DNA fork analog were prepared by heating partially complementary
oligonucleotides (MB1572/MB1596 and MB1571/MB1572, respectively) (Table 2) to 75°C, followed by
slow cooling for ~2 h to room temperature. RNase inhibitors (NEB) were used during the
preparation of RNA-containing substrates and during helicase assays with RNA/DNA hybrid fork
substrates. All reagents were prepared with DEPC-treated water. Helicase reactions were performed
at 30°C for 30 min in 1x binding buffer (25 mM HEPES (pH 8.0), 5% glycerol (w/v), 50 mM NaOAc,
150 μM NaCl, 7.5 mM MgCl$_2$, and 0.01% Tween-20 (w/v)) supplemented with 5 mM ATP. Reactions
were stopped by the addition of 5x dye-free load buffer (50 mM Tris, pH 8.0, and 25% w/v glycerol)
and placed on ice. Labeled fork substrates were added to a final concentration of 0.2 nM. Helicase
reaction products were separated on native 12% 19:1 acrylamide:bis-acrylamide gels supplemented
with 10 mM MgOAc and 5% glycerol. The Tris-Borate-EDTA running buffer (45 mM Tris-borate and
1 mM EDTA, pH 8.0) was supplemented with 2.5 mM MgOAc. Gels were run at 100 V for 30-45 min,
dried, and imaged and quantified using a Typhoon 9500 scanner with ImageQuant software.

2.5. Electrophoretic Mobility Shift Assays (EMSAs)

Substrates for EMSAs were prepared by end-labeling oligonucleotides. The Tel15 and Tel30
substrates were used in ssDNA binding reactions, and forked dsDNA substrates were prepared as
described above for helicase assays. All "Tel" oligonucleotides contained the *S. cerevisiae* telomere
repeat sequence TG$_{1-3}$. Oligonucleotides were labeled with T4 polynucleotide kinase and [γ-^{32}P]ATP
under standard conditions. Labeled oligonucleotides were separated from unincorporated label
using G50 micro-columns. Binding reactions were performed in 1x binding buffer. Radiolabeled
substrates were boiled, placed on ice, and added to binding reactions to a final concentration 0.2 nM.
When present, ATP was added to binding reactions to a final concentration of 4 mM. Binding
reactions were incubated at 30°C for 30 min and mixed with 5x dye-free loading buffer (50 mM Tris
(pH 8.0) and 25% glycerol (w/v)). The reactions were separated on native 4% 37.5:1
acrylamide:bis-acrylamide gels in 1x Tris-glycine running buffer (25 mM Tris (pH 8.0) and 185 mM
glycine, pH 8.8). Gels were run at 100 V for 30-45 min, dried, and imaged and quantified using a

Typhoon 9500 scanner with ImageQuant software. All data were plotted and, where appropriate, fit
with curves using GraphPad software.

2.6. Statistical analysis

All data were analyzed and graphed using GraphPad Prism 6 software. The reported values
are averages of ≥ 3 independent experiments, and the error bars are the standard deviation. p values
were calculated as described in the figure legends, and we defined statistical significance as $p < 0.01$.

3. Results

3.1. Removal of the Pif1 NTD relieves the toxicity of Pif1 over-expression

It has previously been reported that the over-expression of Pif1 is toxic to *S. cerevisiae* [18].
This was also the case in our strain background (Pif1 glucose *vs.* Pif1 galactose, $p < 2.2 \times 10^{-6}$; Fig. 1A),
and we wanted to test whether the Pif1 NTD is involved in this effect. To investigate this question,
we over-expressed a Pif1 construct lacking the N-terminal 233 amino acids of the helicase (Pif1ΔN).
Although cell growth did not completely recover to levels observed in the absence of Pif1
over-expression (Pif1 glucose *vs.* Pif1ΔN galactose, $p = 0.0002$), we still found that the toxicity of
over-expression was significantly relieved in the absence of the Pif1 NTD (Pif1 galactose *vs.* Pif1ΔN
galactose, $p < 1.2 \times 10^{-6}$; Fig. 1A). Pif1's large NTD is of unknown function, both *in vivo* and *in vitro*. It
is predicted to be natively disordered, with the exception of five putative short alpha helices (Fig.
1B,C). Further, Pif1 is a multi-functional helicase [9], and truncation of its NTD may impact one or
more of its *in vivo* roles. Because the Pif1 helicase has been firmly established as a regulator of
telomerase activity and a suite of assays already exist to probe this function, we next decided to
determine how Pif1ΔN behaved in this role.

3.2. In vitro, Pif1ΔN is a more potent telomerase inhibitor than full-length Pif1

We recently found that Pif1 impacts *in vitro* telomerase activity in a biphasic manner: low
concentrations of Pif1 increase telomerase activity, but high Pif1 concentrations decrease telomerase
activity [14]. Using a 15-nt telomerase repeat sequence (TG_{1-3}) oligonucleotide substrate (Tel15), an *S.*
cerevisiae extract enriched for telomerase activity, and recombinant Pif1, we demonstrated a similar
phenomenon here in Figures 2A and B. The addition of 25-125 nM Pif1 significantly ($p < 0.01$)
increased both Type I processivity (*i.e.*, direct extension of Tel15 by 1-7 nt to generate products
T16-T22) and Type II processivity (*i.e.*, further elongation after Type I processivity to generate
products T23-T34) [19] by telomerase (Fig. 2B). However, with increasing Pif1 concentration from
25-125 nM, this stimulation of telomerase activity decreased in a dose-dependent manner for Type I
processivity products. At 250 nM Pif1, total telomerase activity was further decreased to levels
indistinguishable from reactions lacking Pif1 (Fig. 2A, B), and additional increases in Pif1
concentration up to 360 nM significantly decrease telomerase activity relative to reactions lacking
Pif1 (data not shown and [14]).

We next performed similar *in vitro* telomerase reactions containing Pif1ΔN. Compared to
full-length Pif1, the effect of Pif1ΔN on telomerase activity was quite different (Fig. 2). In this case,
Pif1ΔN only stimulated telomerase activity at the lowest helicase concentration tested (25 nM), and
this stimulation was only significant ($p < 0.0001$) for Type I processivity (Fig. 2C, D). At higher
concentrations of Pif1ΔN, the helicase significantly inhibited both Type I and especially Type II
processivity by ≥ 50% relative to reactions lacking added helicase. Thus, Pif1ΔN was a more effective
inhibitor of *in vitro* telomerase activity than full-length Pif1.

Figure 1. **The disordered N-terminus of Pif1 is involved in Pif1 over-expression toxicity. A)** Removal of the Pif1 NTD relieves the toxicity of Pif1 over-expression in *S. cerevisiae*. Galactose-inducible expression vectors for Pif1 and Pif1ΔN, as well as empty vector (EV) were transformed into wild-type cells, and the growth of these strains was monitored at an optical density of 660 nm (OD_{660}) for 48 h. The mean OD_{660} during the entire time course of the cells containing the Pif1 or Pif1ΔN expression vectors was normalized to that of cells containing the empty vector to compare growth in the absence of induction (glu) and upon chronic over-expression (gal). The graphed values represent the average of ≥ 3 independent experiments, and the error bars correspond to the standard deviation. *** $p < 0.0001$. **B)** The NTD of Pif1 is predicted to be natively disordered. The primary sequence of *S. cerevisiae* Pif1 was analyzed using the DISOPRED3 disorder prediction program [20], and the predicted disorder is plotted for each amino acid. The confidence score represents the likelihood that the amino acid resides in a disordered region. **C)** Domain schematics of full-length Pif1 and Pif1ΔN. The first ATP hydrolysis motif of the Pif1 helicase domain begins at residue 241 [9], but sequence alignments, domain prediction, and disorder prediction indicate that the helicase domain itself includes residues 234-240 as well. The predicted secondary structure within the Pif1 NTD is shown below, with cylinders denoting ⍺-helices and the lines connecting them denoting random coil.

Figure 2. Pif1ΔN is a potent *in vitro* telomerase inhibitor. A) *In vitro* telomerase extension of the Tel15 substrate in the absence or presence of the indicated concentration of recombinant Pif1. The Type I (T16-T22) and Type II (T23-T34) extension products are noted on the left. LC, loading control. B) Quantification of the telomerase activity shown in A; mean values and standard deviation are shown. Total activity refers to the total amount of signal in the lanes. C) *In vitro* telomerase extension of the Tel15 substrate in the absence or presence of the indicated concentration of recombinant Pif1ΔN. D) Quantification of the telomerase activity shown in C. Significant differences were determined by multiple *t* tests using the Holm-Sidak method, with $\alpha = 5\%$ and without assuming a consistent S.D. ($n = 3$). *, $p < 0.01$; **, $p < 0.001$; and ***, $p < 0.0001$; all comparisons were made against the reactions lacking added helicase. For a graphical description of the *in vitro* telomerase activity, including type I and type II processivity and the roles of Pif1 in this process, please see [14].

3.3. Chronic over-expression of Pif1ΔN does not lead to telomere length crisis

In the absence of telomerase activity, telomere length decreases during every cell generation until the telomeres reach a critically short point known as crisis [21]. When crisis occurs, most cells senesce. Another way to shorten telomeres is to over-express Pif1, which decreases telomere length in a dose-dependent manner [18, 22]. Our *in vitro* data above indicate that Pif1ΔN is a more potent inhibitor of telomerase activity than the full-length Pif1 helicase, and thus, the expression of Pif1ΔN *in vivo* may lead to rapid shortening of telomeres and crisis even in the presence of functional telomerase. To investigate this hypothesis, we chronically over-expressed Pif1 and Pif1ΔN by galactose induction from a multi-copy vector in wild-type yeast.

In the absence of induction, cultures of cells carrying either the Pif1- or Pif1ΔN-encoding
vector grew just as well as cultures harboring empty vector over three successive restreaks (~75
generations [23]) (Fig. 3). In the presence of galactose, cultures of Pif1 over-expressing cells grew
more slowly than cells containing the Pif1ΔN or empty vector, as expected based on our results in
Figure 1A. However, upon serial restreaking of the cells over-expressing Pif1, they ceased to grow
by streak three, suggesting that cells had undergone telomere length crisis between 50 and 75
generations (Fig. 3). In contrast, cultures of cells over-expressing Pif1ΔN displayed no gross growth
defect relative to cells containing the empty vector. Thus, as opposed to the *in vitro* data (Fig. 2),
these data indicate that Pif1ΔN is a less potent inhibitor of telomerase activity than full-length Pif1.

3.4. Truncation of the Pif1 N-terminus affects DNA binding and unwinding by the helicase

To better understand the biochemical function of the Pif1 NTD, we compared DNA binding
and unwinding by recombinant Pif1 and Pif1ΔN. Using the Tel15 substrate from the *in vitro*
telomerase assays in EMSAs, we found that the affinity of Pif1ΔN for this ssDNA (k_d = 390 ± 58 pM)
was nearly 300-fold greater than full-length Pif1 (k_d = 112 ± 12.7 nM) (Fig. 4A). We previously found
that Pif1 bound to the longer Tel30 substrate with higher affinity than Tel15 (k_d = 33.5 *vs.* 113 nM,
respectively [14]), so we also assessed binding of Pif1ΔN to Tel30. In contrast to the full-length
helicase, Pif1ΔN displayed similarly tight binding to Tel30 ssDNA (k_d = 410 ± 31 pM) compared to
Tel15 (Fig. S1A), indicating that the NTD was responsible for the > 3-fold increase in k_d for Tel30 *vs.*
Tel15 exhibited by Pif1 [14].

Next, we investigated DNA unwinding by Pif1 and Pif1ΔN. Both helicases displayed similar
levels of activity on a forked substrate with a short dsDNA region and ssDNA telomeric repeat
sequence arms (Pif1 $k_{1/2}$ = 2.2 nM *vs.* Pif1ΔN $k_{1/2}$ = 1.1 nM) (Fig. 4B, C). It is known that RNA-DNA
hybrids stimulate Pif1 helicase activity when the enzyme tracks along the DNA strand of the
substrate due to an increase in processivity [24]. To determine if Pif1ΔN can still be stimulated by
RNA-DNA hybrids, we repeated the unwinding reactions using a hybrid fork. Here, we were able to
recapitulate the stimulation in activity displayed by full-length Pif1 (Fig. 4B), but this effect was
largely abrogated with Pif1ΔN (Fig. 4C). Indeed, where Pif1 DNA unwinding was stimulated >
7-fold by the RNA-DNA hybrid substrate, Pif1ΔN activity was only stimulated 2.6-fold.

To determine what may have caused this decrease in activity on RNA-DNA hybrids, we
performed helicase time course assays. However, we found that both Pif1 and Pif1ΔN unwind the
hybrid fork with similar kinetics ($t_{1/2}$ ≈ 18 min) (Fig. S1B). Thus, the decreased activity of Pif1ΔN
relative to Pif1 was not due to a slower rate of unwinding. Subsequently, we measured the affinity of
Pif1 and Pif1ΔN for the RNA-DNA hybrid fork. As with the Tel15 ssDNA (Fig. 4A), Pif1ΔN bound
more tightly to the hybrid fork than Pif1 (Fig. 4D) but only by ~5-fold. In this case however, both Pif1
and Pif1ΔN bound the hybrid fork substrate with greater affinity than binding to Tel15 ssDNA (Pif1
k_d = 0.67 ± 0.085 nM *vs.* 112 ± 12.7 nM, respectively; Pif1ΔN k_d = 0.135 ± 0.013 nM *vs.* 0.390 ± 0.058 nM,
respectively). Therefore, the > 160-fold increase in binding affinity for the RNA-DNA hybrid fork
relative to ssDNA by Pif1 underpinned the stimulation in hybrid fork unwinding relative to the
DNA-DNA fork (Fig. 4B), likely due to increased processivity [24]. Pif1ΔN displayed a much more
modest < 3-fold increase in k_d for the RNA-DNA hybrid fork relative to ssDNA and, consequently,
was less stimulated by the hybrid fork in helicase assays.

Figure 3. **Chronic over-expression of Pif1 leads to cellular senescence.** Wild-type yeast cells were transformed with a multi-copy empty vector (Vector) or vector encoding Pif1 or Pif1ΔN under the control of the inducible *GAL1/10* promoter. When grown on media containing glucose to inhibit over-expression, all strains produced colonies with similar growth kinetics over many successive restreaks. When grown on media containing galactose to induce over-expression of Pif1 or Pif1ΔN, the Pif1 over-expressing cells grew very slowly and failed to grow during the third restreak, indicating that these cells had senesced. Cells over-expressing Pif1ΔN, however, displayed no overt growth defect.

Figure 4. The NTD of Pif1 affects DNA binding and helicase activities. A) Pif1ΔN binds the Tel15

ssDNA substrate much more tightly than full-length Pif1. The plotted values represent the results of

EMSAs using radiolabeled Tel15 and the indicated concentrations of Pif1 and Pif1ΔN protein. **B)** Pif1

helicase activity is stimulated by RNA-DNA hybrids. Unwinding of a radiolabeled DNA-DNA fork
substrate and an analogous RNA-DNA fork were monitored with native gels at the indicated
concentrations of Pif1. **C**) Pif1ΔN displays minor helicase activity stimulation in the presence of the
RNA-DNA hybrid fork. **D**) Both Pif1 and Pif1ΔN have higher affinity for the RNA-DNA hybrid fork
substrate the than Tel15 ssDNA. All data points represent the average of ≥ 3 independent
experiments, and the error bars correspond to the standard deviation. The plotted data were fit with
hyperbolic curves using GraphPad Prism software.

3.5. Is the Pif1 NTD necessary to interact with Hrq1?
We previously demonstrated that Pif1 and the RecQ family helicase Hrq1 synergistically
regulate telomerase activity both *in vivo* [25] and *in vitro* [14], and this may be due to a direct physical
interaction between the helicases. As disordered regions in proteins are often sites of protein-protein
interaction [26], we wondered if truncation of the disordered NTD of Pif1 would affect the
synergism with Hrq1 in *in vitro* telomerase assays. Using a 50-nt ssDNA substrate (Tel50) to enable
binding of both helicases and titrating equimolar amounts of Hrq1 and Pif1ΔN into telomerase
activity reactions, we found that the Hrq1 + Pif1ΔN combination was a potent inhibitor of telomerase
activity at all concentrations tested (Fig. 5A, B). This is in contrast to Hrq1 alone, which mildly
stimulates telomerase activity at high concentration [14], and Pif1ΔN alone, which stimulates
telomerase activity at low concentrations and inhibits it at greater concentrations (Fig. 3C, D).
Therefore, Hrq1 and Pif1 can still interact to regulate telomerase activity in the absence of the Pif1
NTD.

309
310
Figure 5. Hrq1 and Pif1ΔN are still able to interact to regulate telomerase activity. A) *In vitro*
telomerase extension of the Tel50 substrate in the absence or presence of the indicated concentration
of both recombinant Hrq1 and Pif1ΔN (*e.g.*, 45 nM indicates that 45 nM of each helicase was added).
The direct extensions (Type I; T51-T58) and nuclease-cleaved extension products (T26-T32) [14] are
noted on the left. The positions of the Tel30 and Tel50 markers are shown on the right. **B)**
Quantification of the telomerase activity shown in A; mean values and standard deviation are

shown. Total activity refers to the total amount of signal in the lanes. Significant differences were
determined by multiple *t* tests using the Holm-Sidak method, with $\alpha = 5\%$ and without assuming a
consistent S.D. ($n = 3$). *, $p < 0.01$ and **, $p < 0.001$; all comparisons were made against the reactions
lacking added helicase.
321

4. Discussion

To begin to determine the roles of the non-helicase domains of Pif1, we compared the activity of
full-length Pif1 and a Pif1 truncation lacking the entirety of the natively disordered NTD (Pif1ΔN).
Both *in vivo* and *in vitro*, Pif1 and Pif1ΔN displayed obvious differences in assays for cell growth,
telomerase inhibition, and biochemical activities, indicating that the Pif1 NTD is critical for the
proper functioning of the helicase. The implications of the results presented above are discussed
below.

4.1. The Pif1 NTD is involved in regulating telomerase activity

It has been shown that Pif1 induces replication stress at telomeres, with excess Pif1 activity
yielding detrimental increases in telomeric ssDNA [18]. The results in Figure 1A suggest that
removing the Pif1 NTD alleviates the majority of the cellular stress associated with Pif1
over-expression, indicating that Pif1ΔN may have less of an impact on telomere biology than
full-length Pif1, despite increased helicase activity. However, *in vitro* telomerase assays
demonstrated that Pif1ΔN is a more active inhibitor of telomerase activity (Fig. 2C, D) than Pif1 (Fig.
2A, B). What explains this apparent contradiction?

We have found that the standard *in vitro* telomerase inhibition reaction classically used to
characterize Pif1 activity [17] is actually a poor reflection of telomerase regulation *in vivo* [14, 25].
Indeed, with the minimal system (*i.e.*, a telomeric repeat sequence ssDNA substrate, telomerase
activity-enriched cellular extract, and recombinant helicase), one fails to accurately demonstrate the
activities of both the Hrq1 and Pif1 helicases. Based on *in vivo* data [25, 27], these helicases are known
telomerase inhibitors, but using them individually in reactions indicates that Hrq1 stimulates
telomerase activity, while Pif1 both stimulates and inhibits telomere lengthening based on helicase
concentration [14]. When combined, however, the helicases function synergistically *in vitro* to mimic
the telomerase inhibition demonstrated *in vivo*. In other words, the *in vitro* telomerase activity assay
is too biochemically reductive without supplementing the telomerase-enriched extract with
additional recombinant proteins to reflect physiological conditions. That appears to be the case here
with Pif1ΔN. When recombinant Hrq1 was added to telomerase activity assays along with Pif1ΔN,
the combination of these helicases yielded strong telomerase activity inhibition (Fig. 5A, B).

Alternatively, the differences between the *in vitro* and *in vivo* data could be due to lack of
post-translational modifications (PTMs) in our recombinant Pif1 and Pif1ΔN preparations. These
enzymes were prepared by over-expression in *E. coli* and thus lack eukaryotic-type PTMs. Indeed, *in*
vivo, it is known that the activity of Pif1 is affected by PTMs [28-30]. For instance, phosphorylation of
the C-terminus of Pif1 by is necessary for the helicase to inhibit telomerase at DNA double-strand
breaks [28]. To address this issue, future work should investigate the spectrum of PTMs found on
Pif1 *vs.* Pif1ΔN using mass spectrometry.

4.2. Is the Pif1 NTD a site of protein-protein interaction?

Because natively disordered regions of proteins are often sites of protein binding [26], we
initially suspected that the Pif1 NTD may be responsible for the putative physical interaction with

the Hrq1 helicase that affects telomere length homeostasis [14]. However, as stated above,
recombinant Hrq1 and Pif1ΔN together still inhibited *in vitro* telomerase activity to a greater extent
than either enzyme individually (Fig. 2C, D; Fig. 5A, B; and [14]). This could indicate that Pif1 and
Hrq1 do not need to physically interact to regulate telomerase activity but merely to bind to the same
DNA substrate. Alternatively, the Hrq1 interaction motif on Pif1 may be located outside of the NTD,
and thus, the Pif1ΔN truncated protein is still able to bind to Hrq1. Further, the concentrations of
recombinant protein in the telomerase assays may be artificially high enough to force an interaction
between Hrq1 and Pif1ΔN that would otherwise be decreased or abolished in the absence of the Pif1
NTD.
We attempted to distinguish between these possibilities by assaying for an Hrq1-Pif1
interaction using recombinant Hrq1, Pif1, and Pif1ΔN and the bifunctional protein crosslinking
compound DSSO. Unfortunately, we failed to identify Hrq1-Pif1 crosslinks by mass spectrometry
(data not shown) and therefore had no comparison for potential Hrq1-Pif1ΔN interactions or lack
thereof. Crosslinking compounds vary in the chemistry of their reactive groups and linker length
that the crosslinks can span [31], so a survey of additional crosslinkers will be conducted in an
attempt to find *in vitro* conditions suitable for Hrq1-Pif1 crosslinking. Such reactions will also be
performed in the presence of a ssDNA substrate and/or telomerase-enriched extract if a ternary or
quaternary complex is necessary to stabilize the putative Hrq1-Pif1 interaction. Finally, we also plan
to examine *in vivo* telomere length in wild-type and *hrq1Δ* strains expressing Pif1ΔN at
Pif1-endogenous and over-expression levels to directly gauge its effects on telomerase activity.
4.3. Is the Pif1 NTD a site of nucleic acid interaction?
We previously demonstrated that the non-helicase CTD (a predicted WYL domain [32]) of the
bacterial TePif1 helicase is an accessory ssDNA binding domain [12]. This WYL domain impacts
both ATP hydrolysis and DNA unwinding, coupling the two to regulate the biochemistry of the
helicase. Here, the Pif1 NTD was similarly found to affect DNA binding and helicase activities.
However, Pif1ΔN bound both ssDNA (Fig. 4A) and forked DNA (Fig. 4D) substrates with higher
affinity than full-length Pif1. This suggests that the NTD inhibits ssDNA binding by the helicase.
This inhibition could be due to native disorder in the NTD that requires a binding partner to attain
the active Pif1 cellular conformation and proper ssDNA binding affinity.
It should also be noted, however, that Pif1 can interact with RNA in the context of
RNA-DNA hybrids, and that this interaction stimulates helicase activity by increasing Pif1
processivity [24]. Although Pif1ΔN DNA unwinding was stimulated by an RNA-DNA hybrid fork
relative to activity on an analogous DNA-DNA fork (Fig. 4C), the extent of this stimulation was
diminished relative to full-length Pif1 (Fig. 4B). Based on all of these data, we therefore hypothesize
that the Pif1 NTD is involved in single-strand nucleic acid interaction, both ssDNA and ssRNA, and
that these interactions also affect similar interactions of the helicase domain with single-stranded
nucleic acids. Ideally, one could observe these interactions by comparing atomic-resolution
structures of apo- and substrate-bound Pif1. To date, however, no high-resolution X-ray crystal or
cryo-electron microscopy structures exists for a multi-domain PIF1 helicase bound to a nucleic acid
substrate.
4.4. PIF1 perspectives
This work represents a first step toward the dissection of the functions of the non-helicase
domains of *S. cerevisiae* Pif1. We found that the Pif1 NTD affects telomerase regulation and the basic

biochemistry of the helicase. However, much work remains to be done, including structure-function
analyses, surveying the effects of the Pif1ΔN truncation on known genetic interactions of *PIF1* (*e.g.*,
the synthetic lethality of *pif1Δ pol2-16* [33]), and determining the effects of Pif1ΔN on the myriad of
other processes that Pif1 is involved in *in vivo* (*e.g.*, Okazaki fragment processing [34] and
break-induced repair [29]). Similar sets of assays should also be performed with Pif1ΔC constructs
lacking the CTD, which is also predicted to be natively disordered (Fig. 1B), as well as truncations of
both the NTD and CTD to yield the helicase domain in isolation.

Additionally, these types of experiments will be necessary across the PIF1 helicase family. For
instance, the second PIF1 helicase in *S. cerevisiae*, Rrm3, also has largely uncharacterized N- and
CTDs, but Pif1 and Rrm3 perform very different functions *in vivo* [9]. Thus, the functions of the Pif1
NTD and/or CTD on Pif1 biology will likely not mirror those of the Rrm3 NTD and CTD on its
activities. Further, these investigations will yield information about the evolutionary conservation of
the functions of non-helicase domains in the PIF1 family. As many of these uncharacterized domains
are predicted to be natively disordered but have divergent sequences [9, 10], it will be interesting to
determine if conservation of the disorder is important for function, allowing drift in the primary
sequence to occur at a higher rate than in the globular helicase domain. Finally, the human PIF1
helicase (hPIF1) is disease-linked when mutated [35], and understanding the roles of its non-helicase
domains will shed further light on why hPIF1 mutations are related to carcinogenesis and perhaps
suggest treatment strategies.

Supplementary Materials
Supplemental Figure S1. Pif1ΔN binds ssDNA tightly and unwinds DNA with similar kinetics to
full-length Pif1.

Acknowledgments
We thank members of the Bochman lab for thoughtful comments on this work, as well as Kevin
Raney and Hengyao Niu for providing experimental reagents. This work was supported by funds
from the College of Arts and Sciences, Indiana University, a Collaboration in Translational Research
Pilot Grant from the Indiana Clinical and Translational Sciences Institute, and American Cancer
Society Grant RSG-16-180-01-DMC.

Author Contributions
Conceptualization, C.W.S., D.G.N., and M.L.B.; Methodology, C.W.S., D.G.N., and M.L.B.;
Validation, C.W.S., D.G.N., and M.L.B.; Formal Analysis, C.W.S., D.G.N., and M.L.B.; Investigation,
C.W.S., D.G.N., and M.L.B.; Resources, C.W.S., D.G.N., and M.L.B.; Data Curation, M.L.B.; Writing –
Original Draft Preparation, C.W.S., D.G.N., and M.L.B.; Writing – Review & Editing, C.W.S., D.G.N.,
and M.L.B.; Visualization, M.L.B.; Supervision, M.L.B.; Project Administration, M.L.B.; Funding
Acquisition, M.L.B.

Conflicts of Interest
The authors declare no conflict of interest.

References

1.   Abdelhaleem M: **Helicases: an overview.** *Methods Mol Biol* 2010, **587**:1-12.
2.   Bochman ML: **Roles of DNA helicases in the maintenance of genome integrity.** *Mol Cell*
Oncol 2014, **1**:e963429.
3.   Eki T: **Genome-Wide Survey and Comparative Study of Helicase Superfamily Members in**
Sequences Genomes. In *Advances in Genetics Research. Volume* 3. Edited by Urbano KV:
Nova Science Publishers; 2010: 168-203
4.   Berger JM: **SnapShot: nucleic acid helicases and translocases.** *Cell* 2008, **134**:888-888
e881.
5.   Huang S, Li B, Gray MD, Oshima J, Mian IS, Campisi J: **The premature ageing syndrome**
protein, WRN, is a 3'-->5' exonuclease. *Nat Genet* 1998, **20**:114-116.
6.   Rogers CM, Wang JC, Noguchi H, Imasaki T, Takagi Y, Bochman ML: **Yeast Hrq1 shares**
structural and functional homology with the disease-linked human RecQ4 helicase.
Nucleic Acids Res 2017, **45**:5217-5230.
7.   Bernstein KA, Gangloff S, Rothstein R: **The RecQ DNA helicases in DNA repair.** *Annu Rev*
Genet 2010, **44**:393-417.
8.   Liu Y: **Rothmund-Thomson syndrome helicase, RECQ4: On the crossroad between DNA**
replication and repair. *DNA Repair (Amst)* 2010, **9**:325-330.
9.   Bochman ML, Sabouri N, Zakian VA: **Unwinding the functions of the Pif1 family helicases.**
DNA Repair (Amst) 2010, **9**:237-249.
10.   Bochman ML, Judge CP, Zakian VA: **The Pif1 family in prokaryotes: what are our helicases**
doing in your bacteria? *Molecular biology of the cell* 2011, **22**:1955-1959.
11.   Bessler JB, Zakian VA: **The amino terminus of the** *Saccharomyces cerevisiae* **DNA helicase**
Rrm3p modulates protein function altering replication and checkpoint activity. *Genetics*
2004, **168**:1205-1218.
12.   Andis NM, Sausen CW, Alladin A, Bochman ML: **The WYL Domain of the PIF1 Helicase**
from the Thermophilic Bacterium Thermotoga elfii is an Accessory Single-Stranded DNA
Binding Module. *Biochemistry* 2018, **57**:1108-1118.
13.   Zhao S, Douglas NW, Heine MJ, Williams GM, Winther-Larsen HC, Meaden PG: **The STL1**
gene of Saccharomyces cerevisiae is predicted to encode a sugar transporter-like
protein. *Gene* 1994, **146**:215-219.
14.   Nickens DG, Rogers CM, Bochman ML: **The Saccharomyces cerevisiae Hrq1 and Pif1 DNA**
helicases synergistically modulate telomerase activity in vitro. *J Biol Chem* 2018,
293:14481-14496.
15.   Rogers CM, Bochman ML: **Saccharomyces cerevisiae Hrq1 helicase activity is affected by**
the sequence but not the length of single-stranded DNA. *Biochem Biophys Res Commun*
2017, **486**:1116-1121.
16.   Boule J, Vega L, Zakian V: **The Yeast Pif1p helicase removes telomerase from DNA.** *Nature*
2005, **438**:57-61.
17.   Boule JB, Zakian VA: **Characterization of the Helicase Activity and Anti-Telomerase**
Properties of Yeast Pif1p *in vitro*. In *Helicases Methods and Protocols, Methods in*
Molecular Biology Series. Volume 587: Humana Press, ed; 2009: 359-376
18.   Chang M, Luke B, Kraft C, Li Z, Peter M, Lingner J, Rothstein R: **Telomerase is essential to**
alleviate pif1-induced replication stress at telomeres. *Genetics* 2009, **183**:779-791.

19.   Bosoy D, Lue NF: **Yeast telomerase is capable of limited repeat addition processivity.**
490   *Nucleic Acids Res* 2004, **32**:93-101.
491   20.   Jones DT, Cozzetto D: **DISOPRED3: precise disordered region predictions with annotated**
492   **protein-binding activity.** *Bioinformatics* 2015, **31**:857-863.
493   21.   Wellinger RJ, Zakian VA: **Everything you ever wanted to know about Saccharomyces**
494   **cerevisiae telomeres: beginning to end.** *Genetics* 2012, **191**:1073-1105.
495   22.   Zhou J-Q, Monson EM, Teng S-C, Schulz VP, Zakian VA: **The Pif1p helicase, a catalytic**
496   **inhibitor of telomerase lengthening of yeast telomeres.** *Science* 2000, **289**:771-774.
497   23.   Joseph SB, Hall DW: **Spontaneous mutations in diploid Saccharomyces cerevisiae: more**
498   **beneficial than expected.** *Genetics* 2004, **168**:1817-1825.
499   24.   Chib S, Byrd AK, Raney KD: **Yeast Helicase Pif1 Unwinds RNA:DNA Hybrids with Higher**
500   **Processivity than DNA:DNA Duplexes.** *J Biol Chem* 2016, **291**:5889-5901.
501   25.   Bochman ML, Paeschke K, Chan A, Zakian VA: **Hrq1, a Homolog of the Human RecQ4**
502   **Helicase, Acts Catalytically and Structurally to Promote Genome Integrity.** *Cell Rep* 2014,
503   **6**:346-356.
504   26.   Fong JH, Shoemaker BA, Garbuzynskiy SO, Lobanov MY, Galzitskaya OV, Panchenko AR:
505   **Intrinsic disorder in protein interactions: insights from a comprehensive structural**
506   **analysis.** *PLoS Comput Biol* 2009, **5**:e1000316.
507   27.   Schulz VP, Zakian VA: **The *Saccharomyces PIF1* DNA helicase inhibits telomere elongation**
508   **and *de novo* telomere formation.** *Cell* 1994, **76**:145-155.
509   28.   Makovets S, Blackburn EH: **DNA damage signalling prevents deleterious telomere**
510   **addition at DNA breaks.** *Nat Cell Biol* 2009, **11**:1383-1386.
511   29.   Vasianovich Y, Harrington LA, Makovets S: **Break-induced replication requires DNA**
512   **damage-induced phosphorylation of Pif1 and leads to telomere lengthening.** *PLoS Genet*
513   2014, **10**:e1004679.
514   30.   Rossi SE, Ajazi A, Carotenuto W, Foiani M, Giannattasio M: **Rad53-Mediated Regulation of**
515   **Rrm3 and Pif1 DNA Helicases Contributes to Prevention of Aberrant Fork Transitions**
516   **under Replication Stress.** *Cell Rep* 2015, **13**:80-92.
517   31.   Arora B, Tandon R, Attri P, Bhatia R: **Chemical Crosslinking: Role in Protein and Peptide**
518   **Science.** *Curr Protein Pept Sci* 2017, **18**:946-955.
519   32.   Makarova KS, Anantharaman V, Grishin NV, Koonin EV, Aravind L: **CARF and WYL domains:**
520   **ligand-binding regulators of prokaryotic defense systems.** *Front Genet* 2014, **5**:102.
521   33.   Garbacz MA, Lujan SA, Burkholder AB, Cox PB, Wu Q, Zhou ZX, Haber JE, Kunkel TA:
522   **Evidence that DNA polymerase delta contributes to initiating leading strand DNA**
523   **replication in Saccharomyces cerevisiae.** *Nat Commun* 2018, **9**:858.
524   34.   Rossi ML, Pike JE, Wang W, Burgers PM, Campbell JL, Bambara RA: **Pif1 helicase directs**
525   **eukaryotic Okazaki fragments toward the two-nuclease cleavage pathway for primer**
526   **removal.** *J Biol Chem* 2008, **283**:27483-27493.
527   35.   Chisholm KM, Aubert SD, Freese KP, Zakian VA, King MC, Welcsh PL: **A Genomewide**
528   **Screen for Suppressors of Alu-Mediated Rearrangements Reveals a Role for PIF1.** *PloS*
529   *one* 2012, **7**:e30748.