1

5 **_SSD1_ suppresses phenotypes induced by the lack of Elongator-dependent**

6 **tRNA modifications**

11 Fu Xu, Anders S. Byström*, Marcus J.O. Johansson*

15 Department of Molecular Biology, Umeå University, 901 87 Umeå, Sweden

* Corresponding author

E-mail: marcus.johansson@umu.se or anders.bystrom@umu.se

Abstract

The Elongator complex promotes formation of 5-methoxycarbonylmethyl (mcm^5) and

5-carbamoylmethyl (ncm^5) side-chains on uridines at the wobble position of cytosolic

eukaryotic tRNAs. In all eukaryotic organisms tested to date, the inactivation of

Elongator not only leads to the lack of mcm^5/ncm^5 groups in tRNAs, but also a wide

variety of phenotypes. Although the phenotypes are most likely caused by a

translational defect induced by reduced functionality of the hypomodified tRNAs, the

mechanism(s) underlying individual phenotypes are poorly understood. In this study,

we show that the genetic background modulates the phenotypes induced by the lack

of mcm^5/ncm^5 groups in *Saccharomyces cerevisiae.* We show that the stress-

induced growth defects of Elongator mutants are stronger in the W303 than in the

closely related S288C genetic background and that the phenotypic differences are

caused by the known polymorphism at the locus for the mRNA binding protein Ssd1.

Moreover, the mutant *ssd1* allele found in W303 cells is required for the reported

histone H3 acetylation and telomeric gene silencing defects of Elongator mutants.

The difference at the *SSD1* locus also partially explains why the simultaneous lack of

mcm^5 and 2-thio groups at wobble uridines is lethal in the W303 but not in the

S288C background. Collectively, our results demonstrate that the *SSD1* locus

modulates phenotypes induced by the lack of Elongator-dependent tRNA

modifications.

Author Summary

Modified nucleosides in the anticodon region of tRNAs are important for the

efficiency and fidelity of translation. The Elongator complex promotes formation of

several related modified uridine residues at the wobble position of eukaryotic tRNAs.

In yeast, plants, worms, mice and humans, mutations in genes for Elongator

subunits lead to a wide variety of different phenotypes. Here, we show that the

genetic background modulates the phenotypic consequences of the inactivation of

budding yeast Elongator. This background effect is largely a consequence of a

polymorphism at the *SSD1* locus, encoding a RNA binding protein that influences

translation, stability and/or localization of mRNAs. We show that several phenotypes

reported for yeast Elongator mutants are either significantly stronger or only

detectable in strains harboring a mutant *ssd1* allele. Thus, *SSD1* is a suppressor of

the phenotypes induced by the hypomodification of tRNAs.

Introduction

A general feature of tRNA molecules is that a subset of their nucleosides harbors post-transcriptional modifications. Modified nucleosides are frequently found in the anticodon region of tRNAs, especially at position 34 (the wobble nucleoside) and 37. Modifications at these positions typically influence the decoding properties of tRNAs by improving or restricting anticodon-codon interactions [1, 2]. Uridines present at the wobble position in eukaryotic cytoplasmic tRNAs often harbor a 5-methoxycarbonylmethyl (mcm^5) or 5-carbamoylmethyl (ncm^5) side-chain and sometimes also a 2-thio (s^2) or 2'-O-methyl group [3, 4]. The first step in the synthesis of the mcm^5 and ncm^5 side-chains requires the Elongator complex, which is composed of six Elp proteins (Elp1-Elp6) [5-9]. Elongator is thought to catalyze the addition of a carboxymethyl (cm) group to the 5-position of the uridine which is then converted to mcm^5 by the Trm9/Trm112 complex or to ncm^5 by a yet unidentified mechanism [5, 8-12].

In the budding yeast *Saccharomyces cerevisiae*, the inactivation of any of the six *ELP* genes (*ELP1-ELP6*) not only leads to the lack of the mcm^5/ncm^5 groups but also slower growth rate and numerous additional phenotypes [5, 13]. These phenotypes include increased sensitivity to elevated temperatures and various chemical stresses as well as defects in transcription, exocytosis, telomeric gene silencing, and protein homeostasis [14-18]. Even though Elongator mutants lack mcm^5/ncm^5 groups in 11 tRNA species [5, 19], the pleiotropic phenotypes are suppressed by increased expression of various combinations of the hypomodified forms of the three tRNA species that normally harbor a $mcm^5s^2U_{34}$ residue, $tRNA_{UUU}^{Lys}$, $tRNA_{UUG}^{Gln}$ and $tRNA_{UUC}^{Glu}$ [18, 20, 21]. These findings suggest the pleiotropic phenotypes of Elongator mutants are caused by a reduced functionality of the

hypomodified $tRNA_{UUU}^{Lys}$, $tRNA_{UUG}^{Gln}$ and $tRNA_{UUC}^{Glu}$ in translation [20, 21]. The

importance of the modified wobble residue in these tRNAs was supported by the

finding that strains lacking the s^2 group show the same but slightly weaker

phenotypes that are also suppressed by increased expression of the three tRNAs

[20, 21]. Moreover, ribosome profiling experiments have shown that the inactivation

of Elongator causes an accumulation of ribosomes with AAA, CAA or GAA codons in

the ribosomal A-site [18, 22, 23]. However, the pausing at the codons appears to be

relatively small [18, 22] and the mechanism(s) underlying the pleiotropic phenotypes

of Elongator mutants are poorly understood.

In yeast, the cell wall stress that arises during normal growth or through

environmental challenges is sensed and responded to by the cell wall integrity (CWI)

pathway [24, 25]. The CWI pathway is induced by several different types of stresses,

including growth at elevated temperatures, hypo-osmotic shock, and exposure to

various cell wall stressing agents [25]. A family of cell surface sensors (Wsc1-Wsc3,

Mid2 and Mtl1) detects the cell wall stress and recruits the guanine nucleotide

exchange factors Rom1 and Rom2 which activate the small GTPase Rho1. Rho1-

GTP binds and activates several effectors, including the kinase Pkc1. Pkc1 activates

a downstream MAP kinase cascade comprised of the MAPKKK Bck1, the two

redundant MAPKK Mkk1 and Mkk2, and the MAPK Mpk1 (Slt2). The phosphorylated

Mpk1 then activates factors that promote transcription of genes important for cell wall

biosynthesis and remodeling.

In addition to the CWI pathway, several other factors and pathways are known

to influence the cell wall remodeling that occurs upon stress, e.g. the mRNA-binding

protein Ssd1. Ssd1 has been reported to bind and influence the translation, stability

and/or localization of a subset of cellular mRNAs of which many encode proteins

important for cell wall biosynthesis and remodeling, [26-31]. The wild-type *SSD1*

gene was originally identified as a suppressor of the lethality induced by a deletion of

the *SIT4* gene, which encodes a phosphatase involved in a wide range of cellular

processes [32]. The study also led to the finding that some wild-type *S. cerevisiae*

laboratory strains harbor a mutation at the *SSD1* locus that is synthetic lethal with

the *sit4Δ* allele [32]. The *SSD1* locus has since been genetically implicated in many

cellular processes, including cell wall integrity, various signal transduction pathways,

cell morphogenesis, cellular aging, virulence, and transcription by RNA polymerase I,

II and III [33-38]. Although the mechanisms by which Ssd1 influences these

processes are poorly understood, they possibly involve both direct and indirect

effects of Ssd1´s influence on messenger ribonucleoprotein (mRNP) complexes [28,

29, 39]. With respect to the transcripts that encode cell wall remodeling factors, Ssd1

seems to act as a translational repressor and this function is controlled by the protein

kinase Cbk1, which is a component in the RAM (Regulation of Ace2 and cellular

morphogenesis) network [28]. In addition to relieving the translational repression, the

phosphorylation of Ssd1 appears to promote polarized localization of some Ssd1-

associated mRNAs [28, 31].

In this study, we show that increased activation of the CWI signaling pathway

counteracts the temperature sensitive (Ts) growth defect of *elp3Δ* mutants in the

W303 but not in the S288C genetic background. Further, the stress-induced growth

phenotypes caused by the tRNA modification defect are generally stronger in W303-

than in S288C-derived strains. We show that the phenotypic differences are due to

the allelic variation at the *SSD1* locus, i.e. the phenotypes are aggravated by the

nonsense *ssd1-d2* allele found in the W303 background. We also show that the

phenotypes linking the tRNA modification defect to histone acetylation and telomeric

gene silencing are caused by a synergistic interaction with the *ssd1-d2* allele. The

difference at the *SSD1* locus also provide a partial explanation to the finding [18, 40,

41] that cells lacking both the mcm^5 and s^2 group are viable in the S288C but not in

the W303 background.

Results

The Ts phenotype of W303-derived *elp3Δ* cells is partially suppressed by increased expression of factors in the CWI signaling pathway

The observation that Elongator mutants are sensitive to cell wall stressing agents, e.g. calcofluor white and caffeine, implies a defect in cell wall integrity [14]. This notion is further supported by the finding that the Ts growth defect of Elongator-deficient cells is partially suppressed by osmotic support (1 M sorbitol) [42]. As the caffeine sensitivity and Ts growth defect are suppressed by increased levels of the hypomodified $tRNA_{UUU}^{Lys}$ and $tRNA_{UUG}^{Gln}$ [20], the phenotypes are likely caused by reduced functionality of these tRNAs in translation. To further define the wall integrity defect in Elongator mutants, we determined, in the W303 genetic background, if the Ts phenotype of *elp3Δ* cells is suppressed by increased expression of factors in the CWI signaling pathway. The analyses revealed that the introduction of a high-copy *MID2*, *WSC2*, *ROM1,* or *PKC1* plasmid into the *elp3Δ* strain partially suppressed the growth defect at 37°C (Fig 1A**)**. No suppression of the phenotype was observed when the cells carried a high-copy *RHO1, BCK1* or *MPK1* plasmid (Fig 1A**)**. As the overexpression of neither the upstream GTPase (Rho1) nor the downstream kinases (Bck1 and Mpk1) suppressed the Ts phenotype, we considered the possibility that the levels of these factors may be too high when expressed from a high-copy plasmid. Accordingly, low-copy *RHO1, BCK1* and *MPK1* plasmids suppress the Ts phenotype of *elp3Δ* cells to a level similar to that observed for the high-copy *MID2, WSC2, ROM1,* and *PKC1* plasmids (Fig 1A). The level of suppression is, however, smaller than that observed for increased *tK(UUU)* and *tQ(UUG)* dosage, encoding $tRNA_{UUU}^{Lys}$ and $tRNA_{UUG}^{Gln}$, respectively (Fig 1A).

Since the Ts phenotype of *elp3Δ* cells is counteracted by elevated $tRNA_{UUU}^{Lys}$

and $tRNA_{UUG}^{Gln}$ levels [20], it was possible that the activation of the CWI pathway leads

to an increase in their relative abundance. To investigate this possibility, we used

northern blotting to analyze the effect of increased *PKC1* dosage on the levels of

$tRNA_{UUU}^{Lys}$ and $tRNA_{UUG}^{Gln}$ in *elp3Δ* cells grown at either 30°C or 37°C. The blots were

also probed for $tRNA_i^{Met}$, which served as the loading control. In contrast to the ≈2.5-

fold increase in $tRNA_{UUU}^{Lys}$ and $tRNA_{UUG}^{Gln}$ levels induced by increased *tK(UUU)* and

tQ(UUG) dosage (Fig 1B and S1 Table), the abundance of the tRNAs were largely

unaffected by increased *PKC1* dosage at both 30°C and 37°C (Fig 1B and S1 Table).

Collectively, our results suggest that the Ts phenotype of *elp3Δ* cells is, at least in

the W303 genetic background, partially suppressed by increased activation of the

CWI signaling pathway.

The allelic variant at the *SSD1* locus modulates the growth phenotypes of

***elp3Δ* cells**

Phenotypes caused by a mutation in an individual gene can be modulated by the

genetic background of the cell. In fact, the Ts phenotype induced by an *elp3Δ* allele

is more pronounced in strains derived from W303 than in those from S288C (Fig 2A).

As the inactivation of Elongator causes a lack of wobble mcm^5/ncm^5 groups in both

strain backgrounds [5, 43], the Ts phenotype is likely modulated by genetic variation

between W303 and S288C. The difference in phenotype (Fig 2A) prompted us to

investigate if increased *MID2*, *WSC2*, *ROM1, RHO1, PKC1, BCK1* or *MPK1* dosage

counteracts the Ts phenotype of *elp3Δ* cells in the S288C background. The analyses

showed that none of the plasmids counteracted the phenotype (S1A Fig). Moreover,

the growth defect of *elp3Δ* cells at 37°C is counteracted by osmotic support (1 M

sorbitol) in the W303, but not in the S288C background (S1B Fig). Importantly, the

Ts phenotype of *elp3Δ* cells is suppressed by increased *tK(UUU)* and *tQ(UUG)*

dosage in both genetic backgrounds (Fig 1A and S1A Fig), showing that the

underlying cause is the hypomodified $tRNA_{UUU}^{Lys}$ and $tRNA_{UUG}^{Gln}$. We conclude that the

genetic background influences the phenotypes linking the tRNA modification defect

to cell wall integrity.

Although W303 is closely related to S288C, comparisons of the genomes

identified polymorphisms in ≈800 genes that lead to variations in the amino acid

sequence [44, 45]. To identify the cause of the phenotypic differences between the

elp3Δ strains, we examined polymorphisms that have been shown to be

physiologically relevant. The polymorphism at the *SSD1* locus was a good candidate

as *SSD1* has been genetically implicated in many cellular processes, including the

maintenance of cellular integrity [32, 33, 46-50]. Ssd1 is a RNA binding protein that

associates with a subset of mRNAs of which many encode proteins important for cell

wall biosynthesis and remodeling [26-28]. The *SSD1* allele in the S288C background

encodes the full-length Ssd1 protein (1250 amino acids) whereas the allele in W303,

designated *ssd1-d2,* contains a nonsense mutation that introduces a premature stop

codon at the 698[th] codon of the open reading frame [32, 35]. To investigate if the

allele at the *SSD1* locus contributes to the phenotypic differences between the *elp3Δ*

mutants, we analyzed congenic *ssd1-d2, SSD1, ssd1-d2 elp3Δ*, and *SSD1 elp3Δ*

strains in both strain backgrounds. HPLC analyses of the nucleoside composition of

total tRNA from these strains showed that the levels of ncm^5U, mcm^5U, and

mcm^5s^2U are comparable in the *ssd1-d2* and *SSD1* strains and not detectable in the

ssd1-d2 elp3Δ and *SSD1 elp3Δ* mutants (S2 Table). The HPLC analyses also

showed that the allele at the *SSD1* locus does not appear to influence the

abundance of other modified nucleosides present in tRNAs (S2 Fig). By analyzing

the growth of the strains, we found that the *ssd1-d2* allele augments the Ts

phenotype of *elp3Δ* cells in both backgrounds (Fig 2B and S3 Table). The *ssd1-d2*

allele also appears to cause a slightly reduced growth rate of strains with a wild-type

ELP3 gene at the elevated temperature (S3 Table). Even though the *elp3Δ* mutants

grow slower than the *ELP3* strains at 30°C, the effect of the *ssd1-d2* allele is less

pronounced at that temperature (S3 Table). In both genetic backgrounds, the

increased activation of the CWI signaling pathway, through increased *PKC1* dosage,

suppresses the Ts phenotype of the *ssd1-d2 elp3Δ*, but not the *SSD1 elp3Δ* strains

(Fig 2C). Although the *elp3Δ* strains show increased sensitivity to caffeine,

irrespective of the allele at the *SSD1* locus, the *ssd1-d2 elp3Δ* cells are in both

backgrounds more caffeine-sensitive than the *SSD1 elp3Δ* cells (Fig 2D). This

observation is consistent with the previous finding that the *ssd1-d2* allele enhances

the growth inhibitory effects of caffeine [32].

The inactivation of Elongator not only leads to increased sensitivity to caffeine,

but also to other stress-inducing agents [14, 17, 18]. To investigate if the *ssd1-d2*

allele influences these phenotypes, we analyzed the growth of the *ssd1-d2, SSD1,*

ssd1-d2 elp3Δ, and *SSD1 elp3Δ* strains on medium containing rapamycin,

hydroxyurea or diamide. The analyses revealed that the *ssd1-d2* allele, irrespective

of background, also increases the sensitivity of *elp3Δ* cells to these compounds (Fig

2D). Northern blot analyses revealed that the relative abundance of tRNA$_{UUU}^{Lys}$ and

tRNA$_{UUG}^{Gln}$ is largely unaffected by allelic variant at the *SSD1* locus (S3 Fig and S4

Table), indicating that the *ssd1-d2* allele influences the phenotypes by a different

mechanism. Collectively, these results show that the allele at the *SSD1* locus

influences stress-induced growth defects of *elp3Δ* cells.

The *ssd1-d2* allele is required for the histone acetylation and telomeric gene silencing defects of *elp3Δ* mutants

In the W303 background, *elp3Δ* mutants show reduced acetylation of histone H3 [15]. Although the phenotype was originally thought to reflect a function of Elongator in RNA polymerase II transcription [15], the reduced acetylation of lysine-14 (K14) in histone H3 was subsequently shown to be an indirect consequence of the tRNA modification defect [20]. To investigate if the *ssd1-d2* allele contributes to the phenotype, we analyzed, in the W303 background, the histone H3 K14 acetylation levels in the *ssd1-d2, ssd1-d2 elp3Δ, SSD1,* and *SSD1 elp3Δ* strains. As previously shown [15, 20], the level of K14 acetylation is lower in the *ssd1-d2 elp3Δ* mutant than in the *ssd1-d2* strain (Fig 3A). However, the *SSD1* and *SSD1 elp3Δ* strains show comparable levels of K14 acetylated histone H3, indicating that it is the combination of the *elp3Δ* and *ssd1-d2* alleles that induces the histone H3 acetylation defect.

W303-derived Elongator mutants have also been reported to show delayed transcriptional activation of the *GAL1* and *GAL10* genes upon a shift from raffinose- to galactose-containing medium [20, 51]. To determine if the *ssd1-d2* allele influences this phenotype, we analyzed the induction of the *GAL1* mRNA in the W303-derived *ssd1-d2, ssd1-d2 elp3Δ, SSD1,* and *SSD1 elp3Δ* strains. Unexpectedly, we observed no obvious delay in the induction of *GAL1* transcripts in the *elp3Δ* strains regardless of the nature of the allele at the *SSD1* locus (S4 Fig). As the phenotype is thought to reflect a reduced ability of Elongator mutants to adapt to new growth conditions [51], it is possible that differences in media or the handling of

the cultures can explain why *elp3Δ* cells show rapid *GAL1* induction in our

experiments.

Another phenotype reported for Elongator mutants in the W303 background is

a defect in telomeric gene silencing [17, 21]. The telomere silencing defect of

Elongator mutants was inferred from experiments where the expression of a *URA3*

gene integrated close to the left telomere of chromosome VII was assayed [17, 21].

The defect in telomeric gene silencing leads to increased expression of the *URA3*

gene, which is scored as reduced growth on plates containing 5-fluoroorotic acid (5-

FOA) [17, 21]. Even though the integration of an *SSD1* allele into *ssd1-d2* cells does

not influence the level of telomeric gene silencing [36], the inactivation of *SSD1* does

increase the expression of a reporter gene at the silent mating type locus HMR [52].

The latter finding implies that the *ssd1-d2* allele may influence the assembly of silent

chromatin and consequently contribute to the silencing defect in Elongator mutants.

Accordingly, the introduction of a low-copy *SSD1* plasmid into the *ssd1-d2 elp3Δ*

TELVIIL::URA3 strain [21] complemented the 5-FOA sensitivity to a level similar to

that observed with a plasmid carrying the wild-type *ELP3* gene (Fig 3B). Thus, the

telomeric gene silencing defect in Elongator mutants is caused by a synergistic

interaction between the *ssd1-d2* and *elp3Δ* alleles.

The *ssd1-d2* allele augments phenotypes induced by the simultaneous lack of

mcm^5 and s^2 groups

In the formation of mcm^{5}s^{2}U$_{34}$, Elongator promotes synthesis of the mcm^5 side-chain

whereas the thiolation of the 2-position is catalyzed by the Ncs2/Ncs6 complex [40,

43, 53-55]. The simultaneous lack of mcm^5 and s^2 groups was originally reported to

be lethal [40]. However, those experiments were performed in the W303 background

and more recent studies have shown that strains lacking both groups are viable in

the S288C background [18, 41]. To investigate if the allele at the *SSD1* locus

accounts for the difference in viability, we constructed *ssd1-d2 elp3Δ ncs2Δ* and

SSD1 elp3Δ ncs2Δ strains in both backgrounds all carrying a wild-type *ELP3* gene

on a low-copy *URA3* plasmid. Analyses of the strains revealed that the *elp3Δ ncs2Δ*

double mutant is viable in the W303 background if it encompasses a *SSD1* allele

(Fig 4A). In the S288C background, the *ssd1-d2 elp3Δ ncs2Δ* strain is viable but it

grows slower than the *SSD1 elp3Δ ncs2Δ* strain (Fig 4A; S5 Fig shows the same

plates after 2 days of incubation). These observations not only show that allele at the

SSD1 locus influences the viability of *elp3Δ ncs2Δ* cells, but they also indicate that

the growth phenotypes are modulated by additional genetic factors.

The lack of the mcm^5 and/or s^2 groups has, in the S288C background, been

shown to correlate with an increased accumulation of protein aggregates [18]. This

effect was most pronounced in a strain lacking both groups and the phenotype was

suggested to be a consequence of co-translational misfolding due to slower

decoding of AAA and CAA codons by the hypomodified tRNA$_{UUU}^{Lys}$ and tRNA$_{UUG}^{Gln}$ [18].

Moreover, the increased load of aggregates during normal growth was proposed to

account for observation that the double mutant is impaired in clearing diamide-

induced protein aggregates [18]. As strains deleted for *SSD1* show a defect in the

disaggregation of heat-shock-induced protein aggregates [56], it seemed possible

that the *ssd1-d2* allele would augment the protein homeostasis defect in *elp3Δ*

ncs2Δ cells. To investigate this possibility, we isolated aggregates [18, 57] from the

ssd1-d2, SSD1, ssd1-d2 elp3Δ ncs2Δ, and *SSD1 elp3Δ ncs2Δ* strains. Analyses of

the insoluble fractions revealed that the levels of aggregated proteins are

comparable in the *ssd1-d2* and *SSD1* strains (Fig 4B). However*, the ssd1-d2 elp3Δ*

ncs2Δ mutant shows increased accumulation of aggregates compared to the *SSD1*

elp3Δ ncs2Δ strain (Fig 4B). Thus, the allele at the *SSD1* locus modulates the

protein homeostasis defect induced by the simultaneous lack of the wobble mcm^5

and s^2 groups.

Discussion

The phenotypic penetrance of a mutation is often impacted by the genetic background, a phenomenon frequently observed in monogenic diseases [58, 59]. In this study, we investigate the effect of genetic background on the phenotypes of *S. cerevisiae* mutants defective in the formation of modified wobble uridines in tRNA. We show that the phenotypes of Elongator mutants are augmented by the *ssd1-d2* allele found in some wild-type laboratory strains. Moreover, the histone H3 acetylation and telomeric gene silencing defects reported for Elongator mutants are only observed in cells harboring the *ssd1-d2* allele. Thus, the *ssd1-d2* allele sensitizes yeast cells to the effects induced by the lack mcm^5/ncm^5 groups in U_{34}-containing tRNAs.

Although the pleiotropic phenotypes of Elongator mutants are largely caused by the reduced functionality of the hypomodified $tRNA_{UUU}^{Lys}$, $tRNA_{UUG}^{Gln}$ and $tRNA_{UUC}^{Glu}$, the basis for individual phenotypes is poorly understood. Several not necessarily mutually exclusive models have been proposed to explain how the lack of the mcm^5/ncm^5 groups can lead to a particular phenotype. One model postulates that phenotypes can be induced by inefficient translation of mRNAs enriched for AAA, CAA and/or GAA codons and the consequent effects on the abundance of the encoded factors [21, 60-62]. In this model, the inefficient decoding of the mRNAs leads to reduced protein output without affecting transcript abundance. The mechanism by which the slower decoding of the codons leads to reduced protein levels is unclear, but it may involve the inhibition of translation initiation by the queuing of ribosomes. Alternative models suggest that the phenotypes can be caused by defects in protein homeostasis and/or by indirect effects on transcription [18, 22]. The inactivation of *SSD1* not only influences translation and stability of the

transcripts normally bound by Ssd1, but it also leads to altered abundance of many

transcripts that do not appear to be Ssd1-associated [28, 39]. Thus, the effects of the

ssd1-d2 allele on the phenotypes of *elp3Δ* cells could be due to either direct or

indirect effects on gene expression. Moreover, the ribosome profiling experiments of

Elongator mutants have been performed in the S288C background [18, 22, 23] and it

remains possible that the lack of Ssd1 influences the pausing at the AAA, CAA, and

GAA codons.

While the difference at the *SSD1* locus partially explains the nonviability of

elp3Δ ncs2Δ cells in the W303 background, the W303-derived *SSD1 elp3Δ ncs2Δ*

strain grows slower than the corresponding strain in the S288C background (S5 Fig).

Moreover, the *ssd1-d2 elp3Δ ncs2Δ* strain is viable, although with a growth defect, in

the S288C background. These findings indicate that the growth phenotypes of *elp3Δ*

ncs2Δ cells strains are modulated by additional genetic factors. Consistent with the

finding that *ssd1Δ* cells show a defect in Hsp104-mediated protein disaggregation

[56], the *ssd1-d2 elp3Δ ncs2Δ* strain shows increased accumulation of protein

aggregates compared to the *SSD1 elp3Δ ncs2Δ* strain in the S288C background. It

is, however, unclear if this increase is the cause or the consequence of the reduced

growth of the *ssd1-d2 elp3Δ ncs2Δ* strain.

Materials and Methods

Yeast strains, plasmids, media and genetic procedures

Strains and plasmids used in this study are listed in S5 Table and S6 Table. Yeast

media were prepared as described [63, 64]. The medium was where appropriate

supplemented with 2.5 ng/ml rapamycin (R0395, Sigma-Aldrich), 7 mM caffeine

(C0750, Sigma-Aldrich), 100 mM hydroxyurea (H8627, Sigma-Aldrich), 0.25 ng/ml

diamide (D3648, Sigma-Aldrich), or 1 mg/ml 5-fluoroorotic acid (R0812, Thermo

Fisher).

To generate *ssd1-d2* derivatives of BY4741 and BY4742 (S288C

background)[65], we first replaced the sequence between position 2907 and 3315 of

the *SSD1* ORF with a *URA3* gene PCR-amplified from pRS316 [66]. The

oligonucleotides used for strain constructions are described in S7 Table. The

generated strains were transformed with an *ssd1-d2* DNA fragment PCR-amplified

from W303-1A [67]. Following selection on 5-fluoroorotic acid (5-FOA)-containing

plates and subsequent single cell streaks, individual clones were screened for the

integration of *ssd1-d2* allele by PCR and DNA sequencing. The generated strains

(UMY4432 and UMY4433) were allowed to mate producing the homozygous *ssd1-*

d2/ssd1-d2 strain (UMY4434).

Strains deleted for *ELP3* or *NCS2* were constructed by transforming the

appropriate diploid (UMY3387, UMY2836 or UMY4434) with an *elp3::KanMX4* or

ncs2::KanMX4 DNA fragment PCR-amplified from UMY3269 (*elp3::KanMX4*) or

UMY3442 (*ncs2::KanMX4)* with appropriate homologies. Following PCR confirmation

of the deletion, the generated heterozygous diploids were allowed to sporulate and

the UMY4456 (*elp3Δ SSD1*, W303), MJY1019 (*ncs2Δ SSD1*, W303)*,* UMY4439

(*elp3Δ ssd1-d2*, S288C), UMY4442 (*ncs2Δ ssd1-d2*, S288C), MJY1036 (*elp3Δ*

SSD1, S288C), and MJY1021 (*ncs2Δ SSD1,* S288C) strains were obtained from

tetrads. The *elp3Δ ncs2Δ SSD1* mutants (MJY1058 and UMY4467) were obtained

from crosses between the relevant strains. The diploids used to generate the *elp3Δ*

ncs2Δ ssd1-d2 strains (MJY1159 and UMY4454) were transformed with pRS316-

ELP3 [40] before sporulation. MJY1159 was able to lose the plasmid generating

strain UMY4449.

To construct plasmids carrying individual genes for factors in the CWI

signaling pathway, we PCR-amplified the gene of interest using oligonucleotides that

introduce appropriate restriction sites (S7 Table). The DNA fragment was then

cloned into the corresponding sites of pRS425 [68] or pRS315 [66].

RNA methods

The abundance of individual tRNA species was determined in total RNA isolated

from exponentially growing cultures at an optical density at 600 nm (OD_{600}) of ≈0.5

[64]. Samples containing 10 μg of total RNA were separated on 8M urea-containing

8% polyacrylamide gels followed by electroblotting to Zeta-probe membranes (Bio-

Rad). The blots were sequentially probed for $tRNA_{UUU}^{Lys}$, $tRNA_{UUG}^{Gln}$ and $tRNA_{i}^{Met}$ using

^{32}P-labeled oligonucleotides (S7 Table). Signals were detected and analyzed by

phosphorimaging using a Typhoon FLA 9500 biomolecular imager and Quantity One

software.

To analyze the induction of *GAL1* transcripts, cells were grown at 30°C in 50

ml synthetic complete (SC) medium containing 2% raffinose (SC/Raf) to OD_{600}≈0.45.

The culture was harvested by centrifugation at 1,500 x g for 5 min at room

temperature, and the cell pellet resuspended in 15 ml pre-warmed (30°C) SC/Raf

medium. Following reincubation in the shaking water bath for 10 min, transcription of

GAL1 was induced by the addition of 1.5 ml pre-warmed 20% galactose. Aliquots

were harvested [64] at various time points after the addition of galactose and the cell

pellets frozen on dry ice. The procedures for determining mRNA levels have been

described [64].

To analyze the nucleoside composition of total tRNA, the tRNA was isolated

from exponentially growing cultures at $OD_{600} \approx 0.8$ [21]. The tRNA was digested to

nucleosides using nuclease P1 (Sigma-Aldrich, N8630) and bacterial alkaline

phosphatase (Sigma-Aldrich, P4252) and the hydrolysate analyzed by HPLC [69, 70].

The compositions of the elution buffers were as described [70] with the difference

that methanol concentration in buffer A was changed to 5% (v/v).

Histone preparation and immunoblot analyses

Histones were isolated from cells grown in SC medium at 30°C to $OD_{600} \approx 0.8$. Cells

representing 100 OD_{600} units were harvested, washed once with water, resuspended

in 30 ml of buffer A (0.1 mM Tris-HCl at pH 9.4, 10 mM DTT), and incubated on a

rotator at 30°C for 15 min. Cells were collected, washed with 30 ml buffer B (1 M

Sorbitol, 20 mM HEPES at pH7.4) and resuspended in 25 ml of buffer B containing

600 U yeast lytic enzyme. After 1 hour incubation on a rotator at 30°C, the sample

was mixed with 25 ml of ice-cold buffer C (1 M Sorbitol, 20 mM PIPES at pH 6.8, 1

mM $MgCl_2$) followed by centrifugation at 1,500 x g for 5 min. The pellet was

resuspended in 40 ml nuclei isolation buffer [71] and the suspension incubated with

gentle mixing at 4°C for 30 min. Cell debris were removed by centrifugation at 1,500

x g for 5 min. The supernatant was homogenized with 5 strokes in a Dounce

homogenizer followed by centrifugation at 20,000 x g for 10 min. Histones in the

pelleted nuclei were extracted by re-suspension in 5 ml of cold 0.2 M H_2SO_4 and

overnight incubation on a rotator at 4°C. After centrifugation at 10,000 x g for 10 min,

proteins in the supernatant were precipitated by adding 0.5 volumes of 100%

trichloroacetic and 30 min of incubation on ice. Following centrifugation, the pellet

was washed twice with acetone and then dissolved in 200 µl 10 mM Tris-HCl at pH

8.0. Fractions (10 µl) were resolved by 15% SDS-PAGE and transferred to

Immobilon-P (Millipore) membranes. The blots were incubated with rabbit anti-acetyl-

histone H3 (Lys14) antibodies (1:1,000 dilution, Millipore, 07-353) and then with

horseradish peroxidase-linked donkey anti-rabbit IgG (NA934, GE Healthcare). Blots

were stripped and reprobed with rabbit anti-histone H3 antibodes (1:5,000 dilution,

Millipore, 07-690). Proteins were detected using ECL Western blotting detection

reagents (GE Healthcare, RPN2209) and Amersham Hyperfilm ECL (GE Healthcare,

28906836).

Analysis of protein aggregates

Protein aggregates were analyzed in exponentially growing cultures in SC medium at

30°C. Cells representing 50 OD_{600} units were harvested at $OD_{600}{\approx}0.5$ and protein

aggregates were isolated [18, 57] from samples containing 5 mg of total protein. 1/10

of the aggregates and 5µg of total protein were resolved on a 4-12% NuPAGE Bis-

Tris gel (Thermo Fisher, NP0321BOX) followed by staining with the Colloidal Blue

Staining Kit (Thermo Fisher, LC6025).

Acknowledgments

We thank members of M.J.'s and A.B.'s laboratories for valuable discussions.

References

1. Björk GR, Hagervall TG. Transfer RNA Modification: Presence, Synthesis, and Function. EcoSal Plus. 2014;6. PMID: 26442937

2. Agris PF, Narendran A, Sarachan K, Vare VYP, Eruysal E. The Importance of Being Modified: The Role of RNA Modifications in Translational Fidelity. Enzymes. 2017;41: 1-50. PMID: 28601219

3. Machnicka MA, Olchowik A, Grosjean H, Bujnicki JM. Distribution and frequencies of post-transcriptional modifications in tRNAs. RNA Biol. 2014;11: 1619-29. PMID: 25611331

4. Phizicky EM, Hopper AK. tRNA biology charges to the front. Genes Dev. 2010;24: 1832-60. PMID: 20810645

5. Huang B, Johansson MJO, Byström AS. An early step in wobble uridine tRNA modification requires the Elongator complex. RNA. 2005;11: 424-36. PMID: 15769872

6. Winkler GS, Petrakis TG, Ethelberg S, Tokunaga M, Erdjument-Bromage H, Tempst P, et al. RNA polymerase II elongator holoenzyme is composed of two discrete subcomplexes. J Biol Chem. 2001;276: 32743-9. PMID: 11435442

7. Dauden MI, Jaciuk M, Muller CW, Glatt S. Structural asymmetry in the eukaryotic Elongator complex. FEBS Lett. 2018;592: 502-15. PMID: 28960290

8. Kolaj-Robin O, Seraphin B. Structures and Activities of the Elongator Complex and Its Cofactors. Enzymes. 2017;41: 117-49. PMID: 28601220

9. Johansson MJO, Xu F, Byström AS. Elongator-a tRNA modifying complex that promotes efficient translational decoding. Biochim Biophys Acta Gene Regul Mech. 2018;1861: 401-8. PMID: 29170010

10. Kalhor HR, Clarke S. Novel methyltransferase for modified uridine residues at

the wobble position of tRNA. Mol Cell Biol. 2003;23: 9283-92. PMID: 14645538

11. Mazauric MH, Dirick L, Purushothaman SK, Björk GR, Lapeyre B. Trm112p is a

15-kDa zinc finger protein essential for the activity of two tRNA and one protein

methyltransferases in yeast. J Biol Chem. 2010;285: 18505-15. PMID: 20400505

12. Chen C, Huang B, Anderson JT, Byström AS. Unexpected accumulation of

ncm(5)U and ncm(5)S(2) (U) in a trm9 mutant suggests an additional step in the

synthesis of mcm(5)U and mcm(5)S(2)U. PLoS One. 2011;6: e20783. PMID:

21687733

13. Karlsborn T, Tukenmez H, Mahmud AK, Xu F, Xu H, Byström AS. Elongator, a

conserved complex required for wobble uridine modifications in eukaryotes. RNA

Biol. 2014;11: 1519-28. PMID: 25607684

14. Frohloff F, Fichtner L, Jablonowski D, Breunig KD, Schaffrath R. *Saccharomyces*

cerevisiae Elongator mutations confer resistance to the *Kluyveromyces lactis*

zymocin. EMBO J. 2001;20: 1993-2003. PMID: 11296232

15. Winkler GS, Kristjuhan A, Erdjument-Bromage H, Tempst P, Svejstrup JQ.

Elongator is a histone H3 and H4 acetyltransferase important for normal histone

acetylation levels in vivo. Proc Natl Acad Sci U S A. 2002;99: 3517-22. PMID:

11904415

16. Rahl PB, Chen CZ, Collins RN. Elp1p, the yeast homolog of the FD disease

syndrome protein, negatively regulates exocytosis independently of

transcriptional elongation. Mol Cell. 2005;17: 841-53. PMID: 15780940

17. Li Q, Fazly AM, Zhou H, Huang S, Zhang Z, Stillman B. The elongator complex

interacts with PCNA and modulates transcriptional silencing and sensitivity to

DNA damage agents. PLoS Genet. 2009;5: e1000684. PMID: 19834596

18. Nedialkova DD, Leidel SA. Optimization of Codon Translation Rates via tRNA

Modifications Maintains Proteome Integrity. Cell. 2015;161: 1606-18. PMID:

26052047

19. Johansson MJO, Esberg A, Huang B, Björk GR, Byström AS. Eukaryotic wobble

uridine modifications promote a functionally redundant decoding system. Mol

Cell Biol. 2008;28: 3301-12. PMID: 18332122

20. Esberg A, Huang B, Johansson MJO, Byström AS. Elevated levels of two tRNA

species bypass the requirement for elongator complex in transcription and

exocytosis. Mol Cell. 2006;24: 139-48. PMID: 17018299

21. Chen C, Huang B, Eliasson M, Ryden P, Byström AS. Elongator complex

influences telomeric gene silencing and DNA damage response by its role in

wobble uridine tRNA modification. PLoS Genet. 2011;7: e1002258. PMID:

21912530

22. Zinshteyn B, Gilbert WV. Loss of a conserved tRNA anticodon modification

perturbs cellular signaling. PLoS Genet. 2013;9: e1003675. PMID: 23935536

23. Chou HJ, Donnard E, Gustafsson HT, Garber M, Rando OJ. Transcriptome-wide

Analysis of Roles for tRNA Modifications in Translational Regulation. Mol Cell.

2017;68: 978-92 e4. PMID: 29198561

24. Levin DE. Cell wall integrity signaling in *Saccharomyces cerevisiae*. Microbiol

Mol Biol Rev. 2005;69: 262-91. PMID: 15944456

25. Levin DE. Regulation of cell wall biogenesis in *Saccharomyces cerevisiae*: the

cell wall integrity signaling pathway. Genetics. 2011;189: 1145-75. PMID:

22174182

26. Uesono Y, Toh-e A, Kikuchi Y. Ssd1p of *Saccharomyces cerevisiae* associates

with RNA. J Biol Chem. 1997;272: 16103-9. PMID: 9195905

27. Hogan DJ, Riordan DP, Gerber AP, Herschlag D, Brown PO. Diverse RNA-

binding proteins interact with functionally related sets of RNAs, suggesting an

extensive regulatory system. PLoS Biol. 2008;6: e255. PMID: 18959479

28. Jansen JM, Wanless AG, Seidel CW, Weiss EL. Cbk1 regulation of the RNA-

binding protein Ssd1 integrates cell fate with translational control. Curr Biol.

2009;19: 2114-20. PMID: 19962308

29. Ohyama Y, Kasahara K, Kokubo T. *Saccharomyces cerevisiae* Ssd1p promotes

CLN2 expression by binding to the 5'-untranslated region of CLN2 mRNA.

Genes Cells. 2010;15: 1169-88. PMID: 20977549

30. Wanless AG, Lin Y, Weiss EL. Cell morphogenesis proteins are translationally

controlled through UTRs by the Ndr/LATS target Ssd1. PLoS One. 2014;9:

e85212. PMID: 24465507

31. Kurischko C, Kim HK, Kuravi VK, Pratzka J, Luca FC. The yeast Cbk1 kinase

regulates mRNA localization via the mRNA-binding protein Ssd1. J Cell Biol.

2011;192: 583-98. PMID: 21339329

32. Sutton A, Immanuel D, Arndt KT. The SIT4 protein phosphatase functions in late

G1 for progression into S phase. Mol Cell Biol. 1991;11: 2133-48. PMID:

1848673

33. Kaeberlein M, Guarente L. *Saccharomyces cerevisiae MPT5* and *SSD1* function

in parallel pathways to promote cell wall integrity. Genetics. 2002;160: 83-95.

PMID: 11805047

34. Wilson RB, Brenner AA, White TB, Engler MJ, Gaughran JP, Tatchell K. The

Saccharomyces cerevisiae SRK1 gene, a suppressor of bcy1 and ins1, may be

involved in protein phosphatase function. Mol Cell Biol. 1991;11: 3369-73. PMID:

1645449

35. Jorgensen P, Nelson B, Robinson MD, Chen Y, Andrews B, Tyers M, et al. High-

resolution genetic mapping with ordered arrays of *Saccharomyces cerevisiae*

deletion mutants. Genetics. 2002;162: 1091-9. PMID: 12454058

36. Kaeberlein M, Andalis AA, Liszt GB, Fink GR, Guarente L. *Saccharomyces*

cerevisiae SSD1-V confers longevity by a Sir2p-independent mechanism.

Genetics. 2004;166: 1661-72. PMID: 15126388

37. Stettler S, Chiannilkulchai N, Hermann-Le Denmat S, Lalo D, Lacroute F,

Sentenac A, et al. A general suppressor of RNA polymerase I, II and III

mutations in *Saccharomyces cerevisiae*. Mol Gen Genet. 1993;239: 169-76.

PMID: 8510644

38. Wheeler RT, Kupiec M, Magnelli P, Abeijon C, Fink GR. A *Saccharomyces*

cerevisiae mutant with increased virulence. Proc Natl Acad Sci U S A. 2003;100:

2766-70. PMID: 12589024

39. Li L, Lu Y, Qin LX, Bar-Joseph Z, Werner-Washburne M, Breeden LL. Budding

yeast *SSD1-V* regulates transcript levels of many longevity genes and extends

chronological life span in purified quiescent cells. Mol Biol Cell. 2009;20: 3851-

64. PMID: 19570907

40. Björk GR, Huang B, Persson OP, Byström AS. A conserved modified wobble

nucleoside (mcm5s2U) in lysyl-tRNA is required for viability in yeast. RNA.

2007;13: 1245-55. PMID: 17592039

41. Klassen R, Grunewald P, Thuring KL, Eichler C, Helm M, Schaffrath R. Loss of

anticodon wobble uridine modifications affects tRNA(Lys) function and protein

levels in *Saccharomyces cerevisiae*. PLoS One. 2015;10: e0119261. PMID:

25747122

42. Jablonowski D, Frohloff F, Fichtner L, Stark MJ, Schaffrath R. *Kluyveromyces*

lactis zymocin mode of action is linked to RNA polymerase II function via

Elongator. Mol Microbiol. 2001;42: 1095-105. PMID: 11737649

43. Huang B, Lu J, Byström AS. A genome-wide screen identifies genes required for

formation of the wobble nucleoside 5-methoxycarbonylmethyl-2-thiouridine in

Saccharomyces cerevisiae. RNA. 2008;14: 2183-94. PMID: 18755837

44. Ralser M, Kuhl H, Ralser M, Werber M, Lehrach H, Breitenbach M, et al. The

Saccharomyces cerevisiae W303-K6001 cross-platform genome sequence:

insights into ancestry and physiology of a laboratory mutt. Open Biol. 2012;2:

120093. PMID: 22977733

45. Matheson K, Parsons L, Gammie A. Whole-Genome Sequence and Variant

Analysis of W303, a Widely-Used Strain of *Saccharomyces cerevisiae*. G3

(Bethesda). 2017;7: 2219-26. PMID: 28584079

46. Costigan C, Gehrung S, Snyder M. A synthetic lethal screen identifies SLK1, a

novel protein kinase homolog implicated in yeast cell morphogenesis and cell

growth. Mol Cell Biol. 1992;12: 1162-78. PMID: 1545797

47. Lee KS, Irie K, Gotoh Y, Watanabe Y, Araki H, Nishida E, et al. A yeast mitogen-

activated protein kinase homolog (Mpk1p) mediates signalling by protein kinase

C. Mol Cell Biol. 1993;13: 3067-75. PMID: 8386319

48. Mazzoni C, Zarzov P, Rambourg A, Mann C. The Slt2 (Mpk1) map kinase

homolog is involved in polarized cell growth in *Saccharomyces cerevisiae*. J Cell

Biol. 1993;123: 1821-33. PMID: 8276900

49. Martin H, Castellanos MC, Cenamor R, Sanchez M, Molina M, Nombela C.

Molecular and functional characterization of a mutant allele of the mitogen-

activated protein-kinase gene *SLT2*(*MPK1*) rescued from yeast autolytic mutants. Curr Genet. 1996;29: 516-22. PMID: 8662190

50. Ibeas JI, Yun DJ, Damsz B, Narasimhan ML, Uesono Y, Ribas JC, et al. Resistance to the plant PR-5 protein osmotin in the model fungus *Saccharomyces cerevisiae* is mediated by the regulatory effects of SSD1 on cell wall composition. Plant J. 2001;25: 271-80. PMID: 11208019

51. Otero G, Fellows J, Li Y, de Bizemont T, Dirac AM, Gustafsson CM, et al. Elongator, a multisubunit component of a novel RNA polymerase II holoenzyme for transcriptional elongation. Mol Cell. 1999;3: 109-18. PMID: 10024884

52. Raisner RM, Madhani HD. Genomewide screen for negative regulators of sirtuin activity in *Saccharomyces cerevisiae* reveals 40 loci and links to metabolism. Genetics. 2008;179: 1933-44. PMID: 18689887

53. Noma A, Sakaguchi Y, Suzuki T. Mechanistic characterization of the sulfur-relay system for eukaryotic 2-thiouridine biogenesis at tRNA wobble positions. Nucleic Acids Res. 2009;37: 1335-52. PMID: 19151091

54. Leidel S, Pedrioli PG, Bucher T, Brost R, Costanzo M, Schmidt A, et al. Ubiquitin-related modifier Urm1 acts as a sulphur carrier in thiolation of eukaryotic transfer RNA. Nature. 2009;458: 228-32. PMID: 19145231

55. Nakai Y, Nakai M, Hayashi H. Thio-modification of yeast cytosolic tRNA requires a ubiquitin-related system that resembles bacterial sulfur transfer systems. J Biol Chem. 2008;283: 27469-76. PMID: 18664566

56. Mir SS, Fiedler D, Cashikar AG. Ssd1 is required for thermotolerance and Hsp104-mediated protein disaggregation in *Saccharomyces cerevisiae*. Mol Cell Biol. 2009;29: 187-200. PMID: 18936161

57. Koplin A, Preissler S, Ilina Y, Koch M, Scior A, Erhardt M, et al. A dual function

for chaperones SSB-RAC and the NAC nascent polypeptide-associated complex

on ribosomes. J Cell Biol. 2010;189: 57-68. PMID: 20368618

58. Kammenga JE. The background puzzle: how identical mutations in the same

gene lead to different disease symptoms. FEBS J. 2017;284: 3362-73. PMID:

28390082

59. Hou J, van Leeuwen J, Andrews BJ, Boone C. Genetic Network Complexity

Shapes Background-Dependent Phenotypic Expression. Trends Genet. 2018;34:

578-86. PMID: 29903533

60. Rezgui VA, Tyagi K, Ranjan N, Konevega AL, Mittelstaet J, Rodnina MV, et al.

tRNA tKUUU, tQUUG, and tEUUC wobble position modifications fine-tune

protein translation by promoting ribosome A-site binding. Proc Natl Acad Sci U S

A. 2013;110: 12289-94. PMID: 23836657

61. Bauer F, Matsuyama A, Candiracci J, Dieu M, Scheliga J, Wolf DA, et al.

Translational control of cell division by Elongator. Cell Rep. 2012;1: 424-33.

PMID: 22768388

62. Fernandez-Vazquez J, Vargas-Perez I, Sanso M, Buhne K, Carmona M, Paulo E,

et al. Modification of tRNA(Lys) UUU by elongator is essential for efficient

translation of stress mRNAs. PLoS Genet. 2013;9: e1003647. PMID: 23874237

63. Amberg DC, Burke DJ, Strathern JN. Methods in Yeast Genetics. Cold Spring

Harbor, N Y: Cold Spring Harbor Laboratory Press; 2005.

64. Johansson MJO. Determining if an mRNA is a Substrate of Nonsense-Mediated

mRNA Decay in *Saccharomyces cerevisiae*. Methods Mol Biol. 2017;1507: 169-

77. PMID: 27832540

65. Brachmann CB, Davies A, Cost GJ, Caputo E, Li J, Hieter P, et al. Designer

deletion strains derived from *Saccharomyces cerevisiae* S288C: a useful set of

strains and plasmids for PCR-mediated gene disruption and other applications.

Yeast. 1998;14: 115-32. PMID: 9483801

66. Sikorski RS, Hieter P. A system of shuttle vectors and yeast host strains

designed for efficient manipulation of DNA in Saccharomyces cerevisiae.

Genetics. 1989;122: 19-27. PMID: 2659436

67. Fiorentini P, Huang KN, Tishkoff DX, Kolodner RD, Symington LS. Exonuclease

I of *Saccharomyces cerevisiae* functions in mitotic recombination in vivo and in

vitro. Mol Cell Biol. 1997;17: 2764-73. PMID: 9111347

68. Christianson TW, Sikorski RS, Dante M, Shero JH, Hieter P. Multifunctional

yeast high-copy-number shuttle vectors. Gene. 1992;110: 119-22. PMID:

1544568

69. Gehrke CW, Kuo KC. Ribonucleoside Analysis by Reversed-Phase High

Performance Liquid Chromatography. In: Gehrke CW, Kuo KC, editors. Journal

of Chromatography Library. Chromatography and Modification of Nucleosides

Analytical Methods for Major and Modified Nucleosides: HPLC, GC, MS, NMR,

UV and FT-IR. Amsterdam: Elsevier; 1990. pp. A3-A71.

70. Xu F, Zhou Y, Byström AS, Johansson MJO. Identification of factors that

promote biogenesis of tRNACGA(Ser). RNA Biol. 2018;15: 1286-94. PMID:

30269676

71. Edmondson DG, Smith MM, Roth SY. Repression domain of the yeast global

repressor Tup1 interacts directly with histones H3 and H4. Genes Dev. 1996;10:

1247-59. PMID: 8675011

72. Lu J, Huang B, Esberg A, Johansson MJO, Byström AS. The *Kluyveromyces*

lactis gamma-toxin targets tRNA anticodons. RNA. 2005;11: 1648-54. PMID:

16244131

 Supporting information

 S1 Fig. The S288C-derived *elp3Δ* strain does not appear to have cell wall integrity

defect. (A) Growth of the *elp3Δ* (MJY1036) strain carrying the indicated high-copy

(h.c.) or low-copy (l.c.) *LEU2* plasmids. Cells were grown over-night at 30°C in liquid

SC-leu medium, serially diluted, spotted on SC-leu plates, and incubated at 30°C or

37°C for 3 days. (B) The wild-type (W303-1A and BY4741) and *elp3Δ* (UMY3269

and MJY1036) strains were grown over-night at 30°C in liquid SC medium, serially

diluted, spotted on SC plates and SC plates supplemented with 1M sorbitol. The

plates were incubated for 3 days at 30°C or 37°C.

S2 Fig. HPLC analyses of the nucleoside composition of total tRNA from various

ssd1-d2 and *SSD1* strains. The peaks representing ncm^5U, mcm^5U, mcm^5s^2U,

pseudouridine (Ψ), cytidine (C), uridine (U), guanosine (G), and adenosine (A) are

indicated. The asterisk indicates a peak that is a contamination from the bacterial

alkaline phosphatase.

S3 Fig. Effects of *ssd1-d2* and *SSD1* alleles on the abundance of $tRNA_{UUU}^{Lys}$ and

$tRNA_{UUG}^{Gln}$. Northern analysis of total RNA isolated from the *ssd1-d2* (W303-1A and

UMY4432), *ssd1-d2 elp3Δ* (UMY3269 and UMY4439), *SSD1* (UMY3385 and

BY4741), and *SSD1 elp3Δ* (UMY4456 and MJY1036) strains grown in SC medium at

30°C. The blot was probed for $tRNA_{UUU}^{Lys}$, $tRNA_{UUG}^{Gln}$, and $tRNA_i^{Met}$ using radiolabeled

oligonucleotides.

S4 Fig. The transcriptional activation of *GAL1* is not impaired in *elp3Δ* mutants.

Northern analysis of total RNA isolated from the *ssd1-d2* (W303-1A), *ssd1-d2 elp3Δ*

(UMY3269), *SSD1* (UMY3385) and *SSD1 elp3Δ* (UMY4456) strains. Cells were

grown in SC medium containing 2% raffinose followed by induction of *GAL1*

transcription by the addition of 0.1 volumes 20% galactose. Time points after the

addition of galactose are indicated above the lanes. The blot was probed for *GAL1*

transcripts using a randomly labelled DNA fragment. 18S rRNA was detected using a

oligonucleotide probe. The blot is a representative of two independent experiments.

S5 Fig. Influence of the *ssd1-d2* allele on the growth of *elp3Δ ncs2Δ* cells. The figure

shows a shorter incubation (2 days) of the plates in Fig. 4A.

S1 Table. Steady-state tRNA levels in *elp3Δ* cells carrying the indicated plasmids.

S2 Table. Relative amounts of ncm^5U, mcm^5U and mcm^5s^2U in total tRNA isolated

from various strains.

S3 Table. Generation times of indicated strains grown at 30°C or 37°C.

S4 Table. Steady-state tRNA levels in the indicated strains.

S5 Table. Yeast strains used in this study.

S6 Table. Plasmids used in this study.

S7 Table. Oligonucleotides used in this study.

Figures

Fig 1. Increased expression of factors in the CWI signaling pathway counteracts the Ts phenotype of W303-derived *elp3Δ* cells.

(A) Growth of the *elp3Δ* (UMY3269) strain carrying the indicated high-copy (h.c.) or low-copy (l.c.) *LEU2* plasmids. The high copy plasmid carrying the *tK(UUU)* and *tQ(UUG)* genes [72] is abbreviated as h.c. *tK+tQ*. Cells were grown over-night at 30°C in liquid synthetic complete medium lacking leucine (SC-leu), serially diluted, spotted on SC-leu plates, and incubated at 30°C or 37°C for 3 days.

(B) Northern analysis of total RNA isolated from *elp3Δ* (UMY3269) cells carrying the indicated plasmids. The cells were grown in SC-leu medium at 30°C or 37°C. The blot was probed for $tRNA_{UUU}^{Lys}$, $tRNA_{UUG}^{Gln}$ and $tRNA_i^{Met}$ using radiolabeled oligonucleotides. $tRNA_i^{Met}$ serves as the loading control.

Fig 2. The growth phenotypes *elp3Δ* cells are modulated by the allele at the *SSD1* locus.

(A) Growth of *elp3Δ* mutants in the W303 and S288C genetic backgrounds. The wild-type (W303-1A and BY4741) and *elp3Δ* strains (UMY3269 and MJY1036) were streaked on SC plates and incubated at 30°C or 37°C for 3 days.

(B) Effects of the *ssd1-d2*/*SSD1* alleles on the growth of *elp3Δ* strains in the W303 and S288C genetic backgrounds. The *ssd1-d2* (W303-1A and UMY4432), *SSD1* (UMY3385 and BY4741), *ssd1-d2 elp3Δ* (UMY3269 and UMY4439) and *SSD1 elp3Δ* (UMY4456 and MJY1036) strains were grown over-night at 30°C in liquid SC medium, serially diluted, spotted on SC plates, and incubated at 30°C or 37°C for 3 days.

(C) Effects of increased *PKC1* dosage on the growth of *elp3Δ ssd1-d2* and *elp3Δ SSD1* strains. The relevant strains (UMY3269,UMY4456, UMY4439, and MJY1036) carrying the indicated plasmids were grown over-night at 30°C in liquid SC-leu medium, serially diluted, spotted on SC-leu plates, and incubated for 3 days at 30°C or 37°C.

(D) Influence of the *ssd1-d2* allele on growth phenotypes induced by various stress-inducing agents. The strains from B were grown over-night at 30°C in liquid SC medium, serially diluted, spotted on SC plates and SC plates supplemented with caffeine, rapamycin, hydroxyurea or diamide. The plates were incubated for 3 days at 30°C.

Fig 3. The *ssd1-d2* allele is required for the histone acetylation and telomeric gene silencing defects of *elp3Δ* cells.

(A) Western analysis of histones isolated from the *ssd1-d2* (W303-1A), *ssd1-d2 elp3Δ* (UMY3269), *SSD1* (UMY3385) and *SSD1 elp3Δ* (UMY4456) strains grown in SC medium at 30°C. Polyclonal anti-acetyl-histone H3 and anti-histone H3 antibodies were used to detect the indicated proteins. The blot is a representative of two independent experiments.

 (B) Influence of the *ssd1-d2* allele on telomeric gene silencing in *elp3Δ* cells. The s*sd1-d2 TELVIIL::URA3* (UMY2584) and *ssd1-d2 elp3Δ TELVIIL::URA3* (UMY3790) strains carrying the indicated plasmids were grown over-night at 30°C in liquid SC-leu medium, serially diluted, spotted on SC-leu and SC-leu+5-FOA plates, and incubated for 3 days at 30°C.

Fig 4. The growth and protein homeostasis defects of *elp3Δ ncs2Δ* cells are augmented by the *ssd1-d2* allele.

(A) Influence of the *ssd1-d2* allele on the growth of *elp3Δ ncs2Δ* cells. The s*sd1-d2 elp3Δ ncs2Δ* (UMY4454 and MJY1159) and *SSD1 elp3Δ ncs2Δ* (UMY4467 and MJY1058) strains carrying the l.c. *URA3* plasmid pRS316-*ELP3* were grown over-night at 30°C in SC medium, serially diluted, spotted on SC and SC+5-FOA plates, and incubated for 3 days at 30°C.

(B) Effects of the *ssd1-d2* allele on protein aggregation in *elp3Δ ncs2Δ* cells. Total protein and protein aggregates was analyzed from the *ssd1-d2* (UMY4432), *ssd1-d2 elp3Δ ncs2Δ* (UMY4449), *SSD1* (BY4741) and *SSD1 elp3Δ ncs2Δ* (MJY1058) strains grown in SC medium at 30°C. The gel is a representative of two independent experiments.

Figure 1

Figure 2

Figure 3

A

l.c. *ELP3*

	SC	SC+5-FOA	
ssd1-d2 elp3Δ ncs2Δ			W303
SSD1 elp3Δ ncs2Δ			
ssd1-d2 elp3Δ ncs2Δ			S288C
SSD1 elp3Δ ncs2Δ			

B

Figure 4