Real-time integration of cell death mechanisms and proliferation kinetics at the

single-cell and population-level using high-throughput live-cell imaging

Jesse D. Gelles[1,2,3], Jarvier N. Mohammed[1,2,3], Luis C. Santos[1,2], Diana Legarda[4],

Adrian T. Ting[4], and Jerry E. Chipuk[1,2,3,5,6],*

[1] Department of Oncological Sciences,

[2] The Tisch Cancer Institute,

[3] The Graduate School of Biomedical Sciences,

[4] Precision Immunology Institute,

[5] Department of Dermatology,

[6] The Diabetes, Obesity, and Metabolism Institute,

Icahn School of Medicine at Mount Sinai, One Gustave L. Levy Place, Box 1130, New

York, New York 10029 USA

*Corresponding author: jerry.chipuk@mssm.edu

Summary

Quantifying cytostatic and cytotoxic outcomes are integral components of characterizing

perturbagens used as research tools and/or in drug discovery pipelines. Furthermore,

data-rich acquisition coupled with robust methods for analysis is required to properly

assess the function and impact of these perturbagens. Here, we present a detailed and

versatile method for Single-cell and Population-level Analyses using Real-time Kinetic

Labeling (SPARKL). SPARKL integrates high-content live-cell imaging with automated

detection and analysis of fluorescent reporters of cell death. We outline several examples

of zero-handling, non-disruptive protocols for detailing cell death mechanisms and

proliferation profiles. Additionally, we suggest several methods for mathematically

analyzing these data to best utilize the collected kinetic data. Compared to traditional

methods of detection and analysis, SPARKL is more sensitive, accurate, and high-

throughput while substantially eliminating sample processing and providing richer data.

Keywords

Apoptosis, Automated image analysis, Cell death, High-throughput, Live-cell imaging,

Real-time assay

Acknowledgements

This work was supported by: NIH grants CA157740 (J.E.C.), CA206005 (J.E.C.), AI52417 (A.T.T.), and F31AA024681 (J.D.G.), the JJR Foundation, the William A. Spivak Fund, the Fridolin Charitable Trust, an American Cancer Society Research Scholar Award, a Leukemia & Lymphoma Society Career Development Award, and an Irma T. Hirschl/Monique WeillCaulier Trust Research Award. This work was also supported in part by two research grants (5FY1174 and 1FY13416) from the March of Dimes Foundation, and the Developmental Research Pilot Project Program within the Department of Oncological Sciences at the Icahn School of Medicine at Mount Sinai, and the Tisch Cancer Institute Cancer Center Support Grant (P30 CA196521). *Mlkl*[-/-] mice were a gift to Adrian Ting from Warren Alexander at the University of Melbourne via Douglas Green at St. Jude Children's Research Hospital.

Conflict of Interest

The authors declare no conflict of interest.

Abbreviations

AEF: adult ear fibroblast; BAK: BCL-2 homologous antagonist killer; BAX: BCL-2-associated X protein; BCL-2: B-cell lymphoma 2; BCL-xL: B-cell lymphoma-extra large; Brp: Birinapant; CFSE: Carboxyfluorescein succinimidyl ester; CHX: cycloheximide; cIAP1/2: cellular inhibitor of apoptosis-1/2; CYLD: cylindromatosis; DMEM: Dulbecco's modified Eagle's medium; DR: death receptor; FBS: fetal bovine serum; FITC: fluorescein isothiocyanate; GST: glutathione S-transferase; MEF: mouse embryonic fibroblast; NHS

Esters: N-hydroxysuccinimide esters; PARP: poly ADP ribose polymerase; PBS:

phosphate-buffered saline; PI: propidium iodide; PS: phosphatidylserine; RIPK1/3:

receptor-interacting serine/threonine-protein kinase-1/3; ROI: region of interest; SMAC:

second mitochondria-derived activator of caspases; STS: staurosporine; SV40: simian

vacuolating virus 40; TNFα: tumor necrosis factor α; TRAIL: TNF-related apoptosis-

inducing ligand; UPR: unfolded protein response; zVAD-fmk: carbobenzoxy-valyl-alanyl-

aspartyl-[O-methyl]-fluoromethylketone.

Author Contributions

Conceptualization: J.D.G.; Funding Acquisition: J.D.G., A.T.T., and J.E.C.; Investigation:

J.D.G. and J.N.M.; Methodology: J.D.G.; Project Administration: J.D.G.; Resources: D.L.;

Software: J.D.G. and L.C.S.; Supervision: A.T.T. and J.E.C.; Validation: J.N.M.,

Visualization: J.D.G.; Writing: J.D.G., J.N.M., and J.E.C.

Introduction

Programmed cell death pathways are conserved signaling mechanisms which developed early in the evolution of metazoan (Oberst et al., 2008). One aspect shared between many programmed cell death pathways is a variable lag phase between exposure to a perturbagen and the commitment to a cell death program. This lag phase is the consequence of intersecting intracellular pro-death and pro-survival signal transduction and provides a cell with an opportunity to resolve the stress signal and repair accumulated damage (Biton and Ashkenazi, 2011). If these damages are not resolved, the pro-death signaling contributions will overwhelm the pro-survival reserve and trigger biological events committing the cell to death. Importantly, in apoptosis this lag phase also contains an orchestrated and systematic dissolution of organelles and cellular components conducive to efficient clearance with minimal perturbation to neighboring cells. This process is exemplified in apoptosis by the BCL-2 family of proteins, consisting of pro-apoptotic effector proteins (e.g., BAX, BAK) and anti-apoptotic proteins (e.g., BCL-2, BCL-xL), which ultimately serve to regulate the permeabilization of the outer mitochondrial membrane and subsequent activation of the caspase cascade (Wei et al., 2001, Chipuk et al., 2010). However, the kinetics and perpetuation of cell death signaling is highly variable between perturbagens, cell types, death pathways, and between sister cells within a population (Spencer et al., 2009; Gaudet et al., 2012). Elucidating the underlying biology which causes this variability remains a principle focus within the fields of cell death, cell biology, disease etiology, and drug discovery (Kepp et al., 2011). To this end, development of technologies to properly observe and analyze cell death is crucial to progress these fields.

Current standard methods used to observe and quantify cell death remain

outdated, suffer from limited throughput, and generate limited datasets for interpretation.

The detection and quantification of dead or dying cells is most commonly accomplished

by flow cytometry, which requires non-trivial cell numbers, extensive sample handling,

sample exposure to significant mechanical and chemical stress, and creates considerable

delays between sample harvesting and analyses (Koopman et al., 1994). For example,

experiments must be terminated in order to be analyzed, and therefore only provide static

endpoint data, requiring considerable effort to optimize the experimental design.

Commonly used reagents involve cell-impermeable "viability" dyes (such as

propidium iodide, DRAQ7, SYTOX, and YOYO3), which label cells following loss of

plasma membrane integrity and permeabilization. Reliance on this feature for

quantification does not distinguish between distinct pathways and labels cells at the tail-

end of the dismantling process, thereby failing to capture the time in which cells undergo

key biological processes (Vanden Berghe et al., 2010; Dillon et al., 2014). Additionally,

labeling with viability dyes is not stoichiometric and often results in pseudo-binary labeling

profiles following the first instance of membrane instability. Enzymatically-cleaved

fluorescently-conjugated probes (e.g., DEVD-containing caspase-target peptides) are

another common strategy despite their cost, difficulty of use, poor signal, and differential

and non-specific activation (Yu et al., 2001; McStay et al., 2014). Alternative

methodologies use metabolic activity or biochemical measures as surrogate readouts for

cell viability, but interpretations from this data are obfuscated by the underlying biology of

the perturbagens and ultimately do not directly rely on cell death machinery (Chan et al.,

2013).

Design

Here, we integrate and advance multiple previously described methods for

observing and quantifying cell proliferation and cell death using live-cell high-content

imagers and refer to this workflow as Single-cell and Population-level Analyses using

Real-time Kinetic Labeling (SPARKL). We utilize fluorescently-labeled Annexin V due to

stability, signal strength and longevity, stoichiometric binding, and relevance to cell death

machinery. Additionally, we test and verify several fluorescent labeling reagents

compatible with long-term incubation and time-course studies as well as providing

examples of how they can be used to investigate novel aspect of cell death kinetics which

were not feasible with analogous workflows. Our non-toxic label-and-go methods

outperform previous methodologies while maintaining high-sensitivity, high-accuracy, and

a high-throughput zero-handling rapid protocol. Furthermore, we expand on the versatility

of our method by providing several example adaptations and mathematical analyses to

explore the depth of the collected kinetic data. Collectively, we will demonstrate how data

gathered and analyzed using SPARKL can be used to investigate fundamental aspects

of cell death biology by applying these workflows to models of death receptor-mediated

apoptosis, the mitochondrial pathway of apoptosis, ferroptosis, and necroptosis.

Results and Discussion

Automated live-cell imaging and detection of fluorescent labeling provides rich datasets for analyzing, characterizing, and identifying cell death in response to perturbagens.

We developed SPARKL chiefly to capture kinetic cell death data since traditional methods utilizing flow cytometry are limited to endpoint data. Additionally, we aimed to design a workflow compatible with high-throughput studies while requiring minimal handling and retaining high sensitivity and specificity. Here, we capitalize on the advent of high-content in-incubator live-cell imagers to capture cellular phenotypes, measure fluorescent signals, and analyze data in real-time through regular scanning schedules. This technology eliminates the requirement for disruptive handling (e.g., trypsinization, pipet-induced shear forces, centrifugation, incubation in labeling buffers, etc.) and is non-invasive to cells in culture. The repeated observation of samples and coupled automated real-time analysis also collects richer datasets in a workflow that is less labor- and time-intensive than analogous workflows using flow cytometry (Figure 1A).

One significant benefit of this method is that cells are incubated in normal growth media containing fluorescent probes in order to eliminate stepwise processing to label samples. Therefore, detection reagents must be non-disruptive, highly specific, culture-stable, non-labile, and readily detectable when incubated with cells for prolonged periods of time. Most fluorescent probes utilized in flow cytometry assays depend on the brief incubation period and single-scan detection for their utility and are not compatible in kinetic assays due to inherent cyto-static/-toxic effects with prolonged incubation and rapid photobleaching with successive detection. Therefore, we sought to find reagents which are compatible with kinetic assays by demonstrating no negative effects on cells in

culture while exhibiting potent and stable fluorescent signal. Annexin V labels apoptotic

cells by detecting phosphatidylserine (PS) on the outer leaflet of the plasma membrane,

which is exposed during caspase-dependent cell death (Devaux, 1991). We began these

studies by testing FITC-labeled recombinant Annexin V and cell-impermeable viability

dye YOYO3 for compatibility within our method (Logue et al., 2009; Gelles and Chipuk,

2016). MEFs were cultured in growth media containing Annexin V and YOYO3, co-treated

with tumor necrosis factor α (TNFα) and cycloheximide (CHX) to promote apoptosis, and

subjected to SPARKL using a high-content in-incubator live-cell imager. The kinetic data

captured the lag phase followed by the robust labeling of MEFs undergoing death (Figure

1B). Here, cells were cultured in media containing sufficient calcium for robust Annexin V

labeling, but alternative media formulations may require calcium supplementation (Meers

and Mealy, 1993). Importantly, cells not treated with TNFα and CHX exhibited no toxicity

as a consequence of either fluorescent probe at the highest concentrations (Figure 1B,

"UT" line). As expected, the stoichiometric method of Annexin V-labeling was observed

as the concentration-dependent decrease in number of positive events. Said another way,

low concentrations of Annexin V will bind to cells exposing PS but may not generate

sufficient signal (as defined by the intensity, size, and segmentation of the fluorescent

event) according to the user-defined processing definition (see Materials and Methods).

However, both Annexin V and YOYO3 were capable of strong and sustained labeling in

live-cell imagers at concentrations approximately 10-fold below what is needed in flow

cytometric assays (Gelles and Chipuk, 2016). To investigate if growth media containing

colorimetric indicators might affect detection of fluorescent signals, we compared signal

detection of MEFs instigated to die when incubated in DMEM lacking or containing phenol

red. Events of cells labeled with either Annexin V-FITC or YOYO3 exhibited diminished

average fluorescent intensity when culture media contained phenol red (Figure 1C).

Therefore, all future applications of SPARKL utilized phenol red-free culture media.

The foundation of this method is the collection of high-content images of cells in

culture at regular intervals which provide phenotypic characterization of cell populations,

confluency data, and detection of fluorescent pixels. To demonstrate the diversity and

specificity of data collected with SPARKL, $Bax^{+/+}Bak^{+/+}$ and $Bax^{-/-}Bak^{-/-}$ (WT and DKO,

respectively) MEFs were cultured in media containing Annexin V-FITC and YOYO3,

treated with perturbagens, and subjected to the SPARKL workflow. Images from select

timepoints were manually reviewed to observe cellular morphology and detection of

fluorescent labeling (Figure 1D). Green and red pixels from each collected image were

quantified using a trained algorithm and reported as the number of Annexin V- and

YOYO3-positive events observed at each timepoint (Figure 1E). To assess the selectivity

our fluorescent probes, MEFs were instigated to die through the extrinsic pathway of

apoptosis (via co-treatment of TNFα+CHX), the intrinsic pathway of apoptosis (via

topoisomerase inhibitor, VP16), or non-apoptotic pathway of ferroptosis (via cysteine-

glutamate antiport inhibitor, erastin) (Peltzer et al., 2016; Dixon et al., 2012). Both Annexin

V and YOYO3 demonstrated accuracy and selectivity by labeling cells in a time-

dependent manner. Interestingly, VP16-treated MEFs exhibited the longest lag phase

before onset of death and demonstrated Annexin V-positivity more rapidly than YOYO3-

positivity, which reflects the biology of PS exposure prior to loss of plasma membrane

integrity during apoptosis. To directly compare Annexin V-positivity between WT and DKO

MEFs, data collected at specific timepoints were extracted and visualized side-by-side

(Figure 1F). It is worth noting that WT and DKO MEFs proliferate at different rates and

therefore result in vastly different populations over the course of a short experiment

(exampled in Figure 1D). As such, the y-axes for the two cell lines are not congruent but

are graphed appropriately based on internal controls (see Materials and Methods); we

explore methods for appropriate data normalization in a later section. An important aspect

of this methodology is that all data are collected via cell imaging which means all data

generated via SPARKL can be backtracked to the original visualization of the cells in

culture. As an example, collection and visualization of these data from DKO MEFs treated

with VP16 reveals that the morphological phenotypes observed in these cells does not

coincide with probe labeling and therefore is not indicative of cell death (Figures 1D-F).

In this way, we have demonstrated the depth of data collected using SPARKL and how it

can be visualized for thorough interpretation of experimental results.

Investigations of apoptotic cell death commonly employ Annexin V-binding assays

due to the exposure of PS on the outer leaflet of the plasma membrane following caspase-

mediated cleavage of flippases (Segawa et al., 2014). Exposure of PS is indicative of the

underlying apoptotic signaling, and precedes loss of plasma membrane integrity caused

by breakdown of cellular homeostatic processes. Therefore, apoptotic studies

occasionally utilize a dual-labeling methodology of Annexin V with a cell-impermeable

viability dye (such as propidium iodide or DRAQ7) in order to characterize cells in early-

or late-apoptosis by dichotomizing single-positive and dual-positive events (Jiang et al.,

2016). However, cells undergoing non-apoptotic cell death often do not exhibit this

sequential labeling because loss of plasma membrane integrity occurs without the

preceding exposure of PS. As a consequence, cells label with the viability dye and

Annexin V simultaneously, which is indistinguishable from cells in late-apoptosis when

analyzed by flow cytometry (Figure 2A). Therefore, we investigated whether the kinetic

data of SPARKL could distinguish between apoptotic and non-apoptotic cell death

following treatment of cellular perturbagens.

In order to assess the capability of SPARKL to dichotomize kinetic labeling

patterns of cells undergoing apoptotic or non-apoptotic cell death, we cultured WT MEFs

in growth media containing Annexin V-FITC, YOYO3, and perturbagens engaging either

the extrinsic pathway of apoptosis (TNFα+CHX), intrinsic pathway of apoptosis (CHX or

VP16), or non-apoptotic ferroptosis (erastin) (Figure 2B). Cell populations instigated to

die by TNFα+CHX co-treatment or VP16 exhibited Annexin V-positivity prior to YOYO3-

positivity, indicative of apoptotic cell death. By contrast, cells instigated to die by high

concentrations of CHX alone or erastin demonstrated similar kinetics of labeling with

Annexin V and YOYO3. Additionally, we co-treated cells with ABT-737 (an inhibitor to

multiple anti-apoptotic BCL-2 proteins) and/or zVAD-fmk (a pan-caspase inhibitor) to

further characterize mechanisms of cell death in response to each perturbagen using the

SPARKL assay (Oltersdorf et al., 2005; McStay et al., 2008). Cells dying via apoptosis

demonstrated accelerated or attenuated rates of Annexin V-labeling in response to co-

treatment with ABT-737 or zVAD-fmk, respectively; cells dying via ferroptosis

demonstrated no change in kinetics of Annexin V labeling. By comparison, kinetics of

YOYO3 labeling exhibited less resolution and only modest changes in cells co-treated

with ABT-737 or zVAD-fmk. Death receptor (DR)-mediated apoptosis can occur either

through a mitochondrial independent or dependent pathway, termed Type I and Type II

respectively, and the latter signals through members of the BCL-2 family of proteins (Jost

et al., 2009). Kinetic data collected by the SPARKL workflow clearly revealed that WT

MEFs treated with TNFα+CHX signal primarily via the Type II pathway as evidenced by

the synergic labeling of Annexin V in cells co-treated with ABT-737. To further interrogate

the mechanism of cell death engaged by each perturbagen, we replicated this treatment

strategy in DKO MEFs (Figure 2C). DKO MEFs treated with VP16 demonstrated minimal

labeling due to the requirement of BAX and BAK for mitochondrial outer membrane

permeabilization in the intrinsic pathway of apoptosis. By contrast, DKO MEFs treated

with erastin exhibited similar kinetics of death compared to WT MEFs. DKO MEFs treated

with TNFα+CHX died with similar kinetics as their WT counterparts consistent with death

receptor signaling which is independent of BAX and BAK. Furthermore, DKO MEFs

exhibited similar kinetics for both Annexin V and YOYO3. Interestingly, DKO MEFs

treated with CHX alone exhibited death similar to analogously treated WT MEFs but did

not synergize with ABT-737, suggesting that CHX treatment engaged multiple pathways

of cell death which were selectively amplified with particular co-treatments or in different

model systems.

Until this point, we have generalized programmed cell death pathways as either

apoptotic or non-apoptotic, the latter of which has been modeled exclusively by

ferroptosis thus far. However, death receptor signaling can initiate non-apoptotic cell

death programs, such as necroptosis, which are instigated when caspases are inhibited

(Vercammen et al., 1998; Kawahara et al., 1998). Interestingly, while WT MEFs treated

with TNFα+CHX demonstrated attenuated cell death when co-treated with zVAD-fmk,

suggestive of apoptotic cell death, similarly treated DKO MEFs exhibited robust death

that modestly increased with co-treatment of zVAD-fmk suggestive of necroptotic cell

death (Figure 2B-C). When caspases are inhibited, the switch from DR-mediated

apoptosis to necroptosis is initiated by a RIPK1-RIPK3 interaction, which is regulated by

the ubiquitylation profile of RIPK1, resulting in RIPK3-mediated phosphorylation of MLKL,

which consequentially oligomerizes and permeabilizes the plasma membrane (Sun et al.,

2012). It has been reported that SV40-transformed MEFs do not express RIPK3

sufficiently to induce a strong necroptotic program and so we expanded our investigations

to different established cellular models (Moujalled et al., 2013). First, we utilized a

previously described model for studying necroptosis in which MEFs deficient in CYLD, a

RIPK1 deubiquintinase, were reconstituted with either CYLD or a vehicle control

(O'Donnell et al., 2011). $Cyld^{-/-}$ MEFs treated with TNFα+CHX exhibited cell death but

only the MEFs with reconstituted CYLD demonstrated increased or decreased death

following co-treatment with zVAD-fmk or the RIPK1 inhibitor, Necrostatin-1 (Nec1),

respectively, indicative of necroptosis (Degterev et al., 2005) (Figure 2D). Additionally,

MEFs dying by necroptosis exhibited similar labeling kinetics of Annexin V and YOYO3,

which parallels the labeling phenotype we observed in MEFs dying by ferroptosis.

Thus far, we relied on CHX co-treatment to reveal pro-death pathways by inhibiting

translation of induced pro-survival genes downstream of TNFα signaling. In order to better

study labeling kinetics of cells dying by necroptosis we utilized a SMAC mimetic,

birinapant (Brp), to inhibit RIPK1 ubiquitylation by cIAP1/2, which facilitates the pro-

survival signal cascade (Holler et al., 2000). As a consequence of inhibiting RIPK1

unbiquitylation, the requirement of CYLD to initiate necroptosis was greatly reduced and

therefore we moved to an MLKL knockout model for our necroptosis investigations (data

not shown). Primary $Mlkl^{+/+}$ adult ear fibroblasts (WT AEFs) were cultured from mice and

treated with TNFα+Brp with- or without co-treatment of zVAD-fmk or Nec1. While Annexin

V labeling kinetics appeared unchanged in WT AEFs instigated to undergo necroptosis

by addition of zVAD-fmk, YOYO3 demonstrated a dramatic and rapid labeling profile

specific to a necroptotic outcome (Figure 2E). Necroptosis is defined by the regulated

permeabilization of the plasma membrane by oligomeric MLKL and therefore the rapid

labeling by a viability dye is consistent with this phenotype (Wang et al., 2014). However,

the unchanged labeling kinetics of Annexin V was unexpected, particularly for cells which

became necroptotic hours prior. Inspection of the images collected during SPARKL

revealed a substantially distinct cellular morphology in necroptotic cells in which WT AEFs

did not contract, retained a topologically distinct and normal nucleus, and slowly exposed

PS resulting in a weak Annexin V signal over the large area of the cell (Figure 2F, upper

panel). Curiously, a subpopulation of the cells exhibited the same weak Annexin V signal

but demonstrated no YOYO3 labeling (Figure 2F, upper inset panel). It has been

observed that necroptotic cells are capable of exposing PS following MLKL activation but

prior to complete loss of plasma membrane integrity which could explain our observations

that WT AEFs slowly accumulate Annexin V labeling following induction of necroptosis

(Gong et al., 2017). Labeled WT AEFs co-treated with Nec1 resembled an apoptotic

phenotype characterized by cellular contraction and formation of apoptotic bodies;

however, there remained a few cells in the population which resembled necroptosis

(Figure 2F, bottom panel). To verify that our observations were indicative of necroptosis,

we repeated the experiment using $Mlkl^{-/-}$ primary adult ear fibroblasts (KO AEFs) and

observed that these cells were both less sensitive to TNFα+Brp and also completely

protected by addition of zVAD-fmk, suggestive of apoptotic cell death (Figure 2G). Finally,

we treated cocultured WT and KO AEFs with necroptotic inducers and analyzed the cells

with SPARKL to determine if the pattern of labeling would reflect the heterogeneity of the

population. As the ratio of WT to KO AEFs decreased, we observed a concomitant

decrease or absence of TNFα+Brp sensitivity and necroptotic labeling signatures (Figure

2H).

It is worth noting that minimal cell death was observed in TNFα+Brp treated WT

AEFs when co-treated with Nec1 (Figure 2E). Paradoxically, co-treating WT AEFs with

both zVAD-fmk and Nec1 (TBZN) did not abolish cell death but instead resembled the

TNFα+Brp only condition and exhibited apoptosis-like morphology (Figure 2F-H). One

explanation is that cells unable to undergo necroptosis eventually resolve the zVAD-

induced inhibition of caspases and revert to an apoptotic program. Alternatively, they may

be undergoing a necroptotic program that either does not require, or is revealed by the

inhibition of, RIPK1 (Lin et al., 2016; Newton et al., 2016). The KO AEFs do not parallel

these observations and while this could suggest that death in response to TBZN is MLKL-

dependent, data collected with SPARKL revealed that TBZN-induced cell death does not

resemble necroptosis by either labeling kinetics or morphology. Nec1 binds the kinase

domain of RIPK1, which is responsible for propagating the pro-death signal cascade, and

has been shown to selectively inhibit necroptosis in cells induced via a death receptor

and co-treatment with CHX$\pm$zVAD-fmk (Degterev et al., 2008). However, both apoptosis

and necroptosis are inhibited by Nec1 in cells which are induced via a co-treatment with

a SMAC mimetic, due to requirement of RIPK1 for the formation of the caspase-8

activating complex, and we observed this differential effect of Nec1 in our studies as well

(Wang et al., 2008) (Figures 2D-E).

We validated that the SPARKL workflow is not only high-throughput and less labor

intensive, but also versatile and capable of observing and characterizing cell death

following exposure to perturbagens. Specifically, we demonstrated that kinetic data

gathered by the SPARKL workflow can reveal deeper insights regarding the mechanism

of cell death when utilizing a dual-labeling approach. Cells dying by apoptosis sequentially

label with Annexin V and then with a viability dye while cells dying by ferroptosis exhibit

similar kinetics for both labels. In stark contrast, necroptotic cell death exhibited the

reverse phenotype in which cells robustly label with the viability dye and then slowly

accumulated Annexin V over time. It was recently reported that cells can exhibit Annexin

V labeling prior to viability dye labeling due to the formation of PS-containing "bubbles"

on the plasma membrane following MLKL activation (Gong et al., 2017). However, this

phenotype was demonstrated to be rapid, transient, and required highly-sensitive

technologies to detect. While we saw no evidence of this phenotype, we believe this is a

consequence of the difference in signal detection sensitivity and time resolution between

the methodologies and therefore do not believe the results to be inherently contradictory.

Furthermore, these differences exemplify the importance of utilizing multiple specialized

techniques to further understanding the molecular mechanisms of cell death.

Kinetic data gathered by SPARKL provides several quantitative metrics for investigative

and comparative analyses of cell death at both the single-cell and population level.

Cell death assays using flow cytometry provide only endpoint data and are typically

visualized as comparative bar graphs, quadrant plots of probe positivity, or histograms of

fluorescent intensity. SPARKL is less invasive to cultured cells and is more data-rich due

to both the collection of kinetic data and underlying cell imaging methodology. However,

while the utilization of real-time live-cell imaging is becoming more commonplace, most

users do not fully explore the depth of the technology or the data it generates when

analyzing their experiments. Furthermore, the advantages of these technologies are

further minimized by use of endpoint data, simplified comparisons, and labeling methods

repurposed from flow cytometry (Goodall et al., 2016). Therefore, we aimed to broaden

the landscape of experimental designs conducted with high-content imagers while

interrogating how data generated by SPARKL could best be utilized and analyzed in order

to examine phenomena currently overlooked by analogous workflows.

In the previous section, we characterized the labeling profiles for cells instigated

to die by several cell death pathways as observed by real-time live-cell imagers. Not only

did the labeling order differ between each pathway, but the kinetics and sensitivity of

those labeling events were also observably different. Therefore, we do not suggest that

either Annexin V or YOYO3 (or similar cell impermeable viability dye) is superior but rather

the in tandem use provides new methods to understand the kinetics of cell death in

response to a perturbagen. To demonstrate this, we investigated methods to quantify,

process, and express the labeling profiles for each cell death pathway. To articulate the

relative labeling kinetics for each pathway, we expressed kinetic data as the difference

between Annexin V- and YOYO3-positive events over time (Figure 3A). Representation

of data in this manner clearly identifies the period when differential labeling is most

observed and how this period is variable between treatments or pathways of cell death.

Interestingly, applying the same method to data collected from WT and KO AEFs revealed

substantially dissimilar sequential labeling, also in the TNFα+Brp alone treatment (Figure

3B). Thus, these data suggest that TNFα+Brp-induced cell death in KO AEFs was more

apoptosis-like but signaled by a different mechanism in the WT AEFs. Inspecting the

labeling data taken from a coculture of WT and KO AEFs clearly demonstrated how

labeling phenotypes are more difficult to identify in heterogeneous populations. The

cocultured AEFs retained similar labeling kinetics of Annexin V but exhibited notably

different YOYO3 labeling when compared to either homogeneous population, such as the

separation of the zVAD+Nec1 co-treatment (Figures 3C and 2G). When we compared

the labeling data by expressing data as the difference between relative Annexin V- and

YOYO3-positive events, we observed a trend concomitant with the combination of the

two cellular populations (Figures 3D).

It is important to note that the kinetic data collected by SPARKL represents

populations of dying cells labeling with fluorescent probes. For an individual cell, detection

of fluorescent labeling is a binary process and pixels are considered "positive" when they

exhibit fluorescence above a threshold value established during algorithm training (see

Materials and Methods). We utilized the AEF genetic model as an exaggerated example

to illustrate how data interpretation can be less clear in heterogeneous populations.

However, heterogeneity in response to a perturbagen, and therefore the kinetics of cell

death and labeling, is the culmination of many biological and stochastic factors and is not

necessarily so overt in isogenic populations (Roux et al., 2015; Inde and Dixon, 2018).

Therefore, we explored methods in which SPARKL could assess heterogeneity in a

population dying in response to a perturbagen. Instead of viewing data as cumulative,

kinetic data can be processed and expressed as the number of new events over time.

Comparing histograms of events per hour for both labels revealed a more rapid labeling

of Annexin V compared to YOYO3 in apoptotic models, overlapping histograms in

ferroptosis, and stark YOYO3-first sequential labeling in cells undergoing necroptosis

(Figure 3E). These data better illustrate labeling trends within the cell population and

therefore can be analyzed for population statistics (such as reporting mean, median, and

deviation of cell events within the population). Additionally, data expressed in this manner

can reveal subpopulations that label differentially and therefore could reflect different

biological mechanisms. For instance, we consistently observed "shelves" which were

most noticeable when the perturbagen resulted in rapid and robust cell death, such as

either Annexin V or YOYO3 for TNFα+CHX or YOYO3 for TNFα+Brp+zVAD.

High-content live-cell imagers observe the same cells over time and provide the

opportunity for investigators to track labeling kinetics on a single-cell level. We utilized

this technological advantage and applied it to our dual-labeling workflow in order to

assess and quantify sequential labeling for each cell within the imaging field. Image

stacks and analysis masks generated during the SPARKL workflow were processed using

ImageJ and Fiji (Schneider et al., 2012, Schindelin et al., 2012). Fluorescent events were

spatially and temporally defined as regions of interest (ROIs) and then cross-referenced

between images slices and fluorescent channels. Analyzed in this way, the time-to-signal

for both fluorophores was determined for each observed cell. WT MEFs and AEFs were

instigated to die, analyzed by this method, and graphed as the time to single-positive

followed by time to double-positive. Interestingly, while labeling kinetics were treatment-

specific, the time in which Annexin V-labeled cells sequentially labeled with YOYO3 was

highly variable (Figure 3F). For example, 75% of MEFs treated with TNFα+CHX+ABT

labeled with Annexin V by 5 hours and co-labeling with YOYO3 occurred, on average, 3.2

hours later but the slowest 10% were highly variable. This analysis observed cells once

they became Annexin V-positive and then assessed the time in which the cell became

double positive. As a consequence, ROIs which labeled either simultaneously or in

reverse order are visualized as a bar with no red phase and cells exhibiting this pattern

appeared more frequently and rapidly in models of non-apoptotic cell death. To

corroborate this observation, the same image stacks were reanalyzed to track the time to

YOYO3- and then double-positivity (Figure 3G). Very few cells instigated to undergo

apoptosis exhibited Annexin V labeling after YOYO3 (visualized by the lack of green

phase in most of the bars) but ferroptotic cells exhibited approximately 50% of YOYO3-

first sequential labeling. Interestingly, this analysis revealed a dramatic bifurcation within

the population of treated AEFs: while half of the population demonstrated a phenotype

consistent with necroptosis (i.e., rapid onset of YOYO3 followed by variable Annexin V

labeling), the other half exhibited a phenotype reminiscent of a mitochondrial pathway of

apoptosis (i.e., lengthy and highly variable labeling time, YOYO3 labeling after Annexin

V). This phenotype could be a consequence of birinapant-mediated inhibition within the

cell or the association of oligomeric MLKL with membranes other than the plasma

membrane, such as the mitochondria (Wang et al., 2014).

SPARKL data analyzed and graphed in this manner aids in visualization of

population trends by separately tracking the labeling kinetics of each individual cell.

Therefore, labeling metrics of the population can be statistically compared to further

characterize pathway- or treatment-specific patterns of cell death (Figure 3H). Sequential

labeling can be confirmed and quantified by comparing individual label kinetics to double-

positive kinetics. For example, YOYO3 and double-positive kinetics of MEFs treated with

VP16+ABT are not significantly different which indicated that double-positivity depended

on YOYO3 and that Annexin V labeling occurred first. Similarly, the reverse phenotype

was observed in AEFs undergoing necroptosis even though the population exhibited two

different labeling profiles. Interestingly, MEFs undergoing ferroptosis exhibited

significantly distinct labeling kinetics for Annexin V, YOYO3, and double-positivity, which

indicated no sequential labeling and likely reflects the biochemistry of the currently

unknown method by which lipid peroxidation results in cell death (Feng and Stockwell,

2018). Additionally, this method for comparing cellular populations provides a convenient

strategy to investigate the consequences of co-treatments and determine either

synergistic or differential biology (Figure 3I). For example, MEFs induced to undergo

apoptosis exhibited significantly different labeling kinetics in response to ABT-737 while

ferroptotic MEFs did not. Importantly, the pattern of distributions, while occurring earlier,

were conserved and therefore reflected a treatment-mediated synergy of the death

pathway, namely apoptosis. In stark contrast, the switch from apoptosis to necroptosis in

AEFs exhibited significantly different labeling kinetics as well as different patterns of

distribution. Collectively, these data demonstrate a method and rationale for analyzing

data collected by high-content live-cell imagers to characterize and compare differential

labeling kinetics of single cells in response to perturbagens.

While there are instances to do the deep analyses we described above, it does not

represent the common usage of high-content live-cell imagers for cell death analyses –

high-volume comparative studies. In order to take full advantage of the high-throughput

capacity of these technologies, complex kinetic data must to be expressed as simplified

key parameters. Here, we will review best practices for comparing SPARKL data as well

as several mathematical methods for rapid parameterization of collected data (Figure 4A).

Instinctually, investigators often use the maximum number of events to compare cell

death data and use language akin to "A had more or less death than B." However, the

amplitude of kinetic curves reflects extraneous factors such as the number of cells in the

population or the penetrance of a particular perturbagen. This is why graph axes are set

by control treatments and do not need to be the same scale between cell types (such as

data in Figure 1E). Additionally, using the maximal number of events (i.e.: death) is highly

reductionist and does not reflect metrics that are uniquely collected by real-time

methodologies such as the times of onset, 50% death, plateau, maximal rate of death,

etc. – collectively referred to as the "kinetics" of cell death. Therefore, we recommend

drawing comparisons from cell death kinetics.

The simplest method is to calculate the area under the curve (AUC) which

compresses kinetic data into a single parameter for convenient comparisons. WT and

DKO MEFs were instigated to die using a variety of perturbagens, co-treated with ABT-

737 or zVAD-fmk, and detected by Annexin V- and YOYO3-labling (Figure 4B). AUC

values were greater in cells co-treated with ABT-737 and reduced in cells co-treated with

zVAD-fmk for perturbagens activating apoptosis, but unchanged in cells undergoing

ferroptosis. AUC calculated from YOYO3 detection exhibit similar trends in response to

ABT-737 and zVAD-fmk albeit to a lesser degree (Figure 4B). These analyses retained

the observed trends of Annexin V and YOYO3 labeling within the raw kinetic data.

Alternative metrics of interest are the rate of cell death and the time to maximal rate of

death. The first derivative was calculated for data collected from MEFs treated with pro-

death perturbagens and labeled with Annexin V (Figure 4C). The derivative reports the

rate of cell death within the population, or the slope of kinetic curves, at each collected

time point. MEFs undergoing apoptosis demonstrated greater rates of cell death at earlier

time points in response to ABT-737 while exhibiting lower rates in response to zVAD-fmk

(Figure 4C, yellow and purple data). Additionally, data can be presented as the time-to-

maximal rate of death, however this parameter is only informative for samples exhibiting

significant death and labeling over the course of data collection (Figure 4D). These

mathematical analyses represent simple but robust methods to parameterize data

collected with SPARKL for convenient comparisons between experimental conditions.

In an effort to quantify data in a manner that better retains the richness of kinetic

data, we employed a non-linear fit analysis to the data collected with SPARKL. A Lag

One-Phase Exponential (LOPE) function provided the best fit for our data and has been

utilized successfully by other groups for modeling cell death kinetics (Forcina et al., 2017)

(Figure 4E). This mathematical function reflects the latency period prior to the onset of

cell death in which cells undergo biological signal transduction in response to

perturbagens. The LOPE function calculates two useful parameters during the fit analysis:

the duration of the lag phase (t') and the rate of cell death following the lag phase (R_D). It

is worth noting that this model assumes that R_D is the maximal rate of cell death which is

not always accurate. However, this model consistently performed better when fitting data

compared to other models (for example, sigmoidal functions which required more defined

asymptotes to model). Similar to graphs of raw kinetic data, the plateau amplitude is

dependent on the number of cells and therefore was not considered for parameterization

in comparative studies. Comparing t' values for WT MEFs labeled with Annexin V and

undergoing apoptosis demonstrated a clear trend in which ABT-737 co-treatment reduces

t', while the *t'* of MEFs undergoing ferroptosis was unchanged (Figure 4F). R_D trends

calculated from Annexin V data were harder to interpret since the metric only differed in

MEFs co-treated with CHX and ABT-737 (Figure 4G). We believe this may be an

indication that the selected perturbagen concentrations were excessive and therefore R_D

has plateaued, or an indication that the time of maximal death significantly differs from

the R_D calculated using LOPE. Interestingly, *t'* from YOYO3 data did not reflect changes

in response to ABT-737 and R_D values increased (Figures 4F-G, red data). Viewing these

data as two-parameter plots reveals treatment- or cell type-specific trends in *t'* and R_D

following co-treatment with ABT-737 (Figure 4H). This strategy for data visualization is

particularly effective when characterizing cell death mechanisms for perturbagens of

unknown biology and conducive to large-scale, multi-parametric drug panels (Forcina et

al., 2017; Fallahi-Sichani et al., 2013). While we have demonstrated methods that utilized

both Annexin V and YOYO3, this method for comparative studies is most effective when

utilizing a single labeling agent, whichever is most relevant to the experimental design.

The advent of high-content live-cell imagers provides investigators the opportunity

to collect and analyze cellular data in a high-volume, high-throughput manner in real time.

However, these technologies are not often used to their fullest potential and instead serve

as a surrogate for flow cytometric workflows. To capitalize on the advantages of these

technologies, we have provided several methods and best practices for utilizing and

interpreting the data in comparative studies. We described a multiplex labeling workflow

to demonstrate the variety of experimental designs compatible with these machines as

well as several methods to parameterize and appropriately compare collected data.

Collectively, these examples showcase the versatility, range of applications, breadth and

depth of collected data, and analyses investigators can apply to experimental paradigms

by using the SPARKL workflow.

Simultaneous collection of proliferation and cell death data to accurately characterize

perturbagen-induced cytostatic and cytotoxic effects.

The high-throughput capacity of SPARKL permits investigators to conduct

exhaustive optimizations and drug titrations with relative ease. However, perturbagens at

sub-lethal concentrations may still affect rates of cell proliferation, either directly or as a

consequence while mitigating the stress signal. Additionally, it has been shown that

growth rate inhibition metrics can correct otherwise confounding data in large-scale drug

screens (Hafner et al., 2016). Therefore, investigators require a method to properly

assess the cytostatic effects of perturbagens on cells in culture. This is commonly

accomplished by either measuring confluency or surface area metrics, which do not

accurately reflect cell numbers, or by requiring the investigator to generate cell lines

expressing fluorescent proteins, which is laborious, limiting to model systems, and not

conducive to high-throughput analyses (Artymovich and Appledorn, 2015). Previously,

we described a method utilizing non-toxic CFSE to label cells prior to treatment and data

acquisition (Gelles and Chipuk, 2016). This method required additional handling steps,

suffered mild photobleaching, and was not compatible with certain cell types. Therefore,

we sought to develop a label-and-go protocol of SPARKL that would simultaneously

collect proliferation and cell death data in real-time without requiring a separate cell-

labeling step (Figure 5A).

Probes labeling the nucleus provide robust and readily detectable fluorescent

events, which can be efficiently analyzed for accurate quantification of event number

within each sample. However, most nuclear labeling probes are either cell impermeable

or function as toxic DNA intercalators which are incompatible with time-course

experiments and the SPARKL workflow. Therefore, we investigated if cell-permeable,

low-affinity cyanine nucleic acid stains were suitable for prolonged incubation with cells.

MEFs were incubated with varying concentrations of SYTO fluorescent dyes and Annexin

V conjugated with AlexaFluor 594 (Annexin V-594) in order to assess the cytotoxic effects

of nucleotide labeling (Figure 5B). Each tested SYTO dyes exhibited toxicity at high

concentrations and degrees of cytostatic behavior at low- to mid-range concentrations.

Low-concentration ranges of SYTO16 and SYTO21 demonstrated no adverse effects to

cell proliferation or viability, and we ultimately selected SYTO21 for use within SPARKL

workflows. SYTO21 primarily labeled cells in the nucleus, was readily detected and

segmented, and capable of labeling daughter cells post-mitosis (Figure 5C). Additionally,

SYTO21 signal was retained in cells undergoing apoptosis and resulted in

SYTO21/Annexin V-594 double-positive events. MEFs incubated with SYTO21

demonstrated an increase in SYTO21-positive events over time, indicative of proliferation.

By contrast, MEFs undergoing apoptosis via treatment of TNFα+CHX exhibit no change

in the number of SYTO21-positive events and robustly labeled with Annexin V-594

(Figure 5D). Therefore, SYTO21 is suitable as a fluorescent probe to track cell

proliferation and population growth.

High-content live-cell imagers detect fluorescence and express data as a number

of positive events within a given area. However, this value is not normalized for total cell

number and therefore more events are detected in wells containing more cells.

Interpretations of data expressed in this manner may be obfuscated when cell numbers

vary between samples due to either treatment-induced or cell type-specific differences in

proliferation rates. Imperfect strategies to circumvent this issue undermine the

advantages of kinetic monitoring by either implementing tandem analysis via flow

cytometry, normalizing to end-point data collected using DNA-intercalating dyes or potent

inducers of cell death, or using confluency as a surrogate for cell number (Lopez et al.,

2016; Giampazolias et al., 2017). Therefore, we investigated whether SYTO21 labeling

could be used to normalize cell death data collected with SPARKL. MEFs were seeded

at several densities, incubated with SYTO21 and Annexin V-594, and treated with either

vehicle or TNFα+CHX. SYTO21 data accurately measured differences in cell number and

tracked cell proliferation in untreated MEFs; MEFs instigated to die exhibited constant

SYTO21 data and the concomitant Annexin V-positive data exemplified how cell number

can complicate data interpretation (Figure 5E). The number of Annexin V-positive events

at each timepoint was normalized by the corresponding SYTO21-positive events to

express data as the percent of the population exhibiting Annexin V-positivity (Figure 5F).

For comparison, Annexin V-positive endpoint data collected using SPARKL is shown both

as the raw number of events and the adjusted percentages normalized using total cell

number as reported by SYTO21 (Figure 5G). These data demonstrate that when coupled

with skilled cell culture practices, this dual-labeling method and analysis can control for

cell number differences due to differential cell seeding and cell type-inherent or treatment-

induced alterations to proliferation rates.

Cell death data normalization is particularly relevant when perturbagens do not

illicit a rapid death phenotype, which will result in more proliferation variability between

experimental conditions and therefore different maximal labeling. To demonstrate this,

we applied the normalization methodology to WT and DKO MEFs, which exhibit dramatic

differences in rates of proliferation, incubated with SYTO21 and Annexin V, and treated

with a panel of perturbagens. Annexin V data collected through SPARKL was normalized

by the SYTO21 data and provided values that can be directly compared between the two

cell types (Figure 6A). Perturbagens engaging BAX- and BAK-mediated apoptosis

demonstrated an attenuated lethal response in DKO MEFs (such as VP16 and calcium

ionophore, A23187) and perturbagens capable of engaging the intrinsic pathway of

apoptosis were synergized by co-treatment with ABT-737. When we replicated this

experiment using YOYO3 as the cell death probe and normalized for SYTO21 labeled

cells, we observed a significant loss of signal during longer experiments (Figure 6B).

When analyzed for longer than 24 hours, MEFs undergoing apoptosis by co-treatment

with TNFα+CHX exhibited sustained SYTO21- and Annexin V-positivity, but

demonstrated a loss of YOYO3 signal over time (Figure 6C). We analyzed the average

fluorescent intensity of positive events which revealed that YOYO3-positive events lost

fluorescence over time (Figure 6D). While photobleaching may contribute to this loss of

signal, we believe that the effect is indicative of changes to DNA content and subsequent

labeling by YOYO3, which would explain why the diminished signal is perturbagen-

specific and observed in cells undergoing apoptosis. Signal loss does not appear to be

due to cells leaving the focal plane as this would affect detection of all probes and the

data do not demonstrate this concomitant loss of signal in either SYTO21 or Annexin V

over time.

We capitalized on this method to normalize data and applied the same

mathematical analyses described in the previous section. Percentages of cell death

calculated by normalizing Annexin V data to SYTO21 were fit using the LOPE function

(Figure 6E). Samples co-treated with ABT-737 reveal a reduction in lag phase time (t'),

indicative of BCL-2 family contributions within the cell death pathway; by contrast, t' is

unchanged in response to ABT-737 in MEFs undergoing ferroptosis via erastin (Figure

6F). Consistent with our previous methods of analysis, receptor-mediated apoptosis by

either TNFα or TRAIL in conjunction with CHX demonstrated substantially different

kinetics when co-treated with ABT-737 and indicates a predominance for Type II signaling

in MEFs. When analyzing the initial rates of cell death (R_D), which became more

informative for data which had been cell number-normalized, we observed increased

rates in of death following co-treatment of ABT-737 in MEFs treated with CHX,

TNFα+CHX, TRAIL+CHX, or VP16 (Figure 6G). R_D was unchanged in MEFs treated with

erastin or A23187. Cell death data from MEFs treated with A23187 collectively indicates

a mechanism requiring BAX and BAK but only partially sensitized by inhibition of anti-

apoptotic BCL-2 proteins, which is a profile not shared amongst the other perturbagens

exampled within this work (Figure 6H). Control-treated MEFs demonstrated ambiguous

fit curves and therefore parameter extraction was highly variable and not informative

(Figures 6E-G, red panel). Similarly, DKO MEFs not exhibiting death in response to

particular perturbagens (such as VP16 and A23187) produced ambiguous fit curves and

were not analyzed further (data not shown). These data exemplify the utility of

mathematical parameterization on kinetic data which was normalized for differential cell

numbers as a consequence of cell-inherent or treatment-induced changes to proliferation.

Discussion

Here we described the SPARKL workflow which is capable of simultaneously gathering

proliferation and cell death data at single-cell and population-level resolutions. Using

multiplex fluorescent detection of cell death, investigators can dichotomize cells engaging

apoptotic or non-apoptotic signaling pathways in response to perturbagens. Additionally,

we described a novel method to quantify cell numbers and proliferation using a non-

disruptive fluorescent probe which is compatible with "label-and-go" workflows conducive

to high-throughput studies. Collection of these rich datasets provides in depth

comprehensive insight into the cytostatic effects of perturbagens as well as their cytotoxic

potential. Furthermore, we described and advanced robust and thorough mathematical

analyses that parameterize data for convenient comparative analyses and interpretations

of underlying cell death biology. The presented data demonstrate the versatility of the

technology using single concentration treatments, but these workflows and analyses are

well suited for experiments using perturbagen titrations and determining optimal

conditions for subsequent studies. Collectively, SPARKL is a versatile, data-rich, non-

intensive, and zero-handling method to visualize, detect, and quantify the kinetics of cell

death which, when coupled with mathematical analyses, enable investigators to conduct

thorough high-throughput comparative studies.

Limitations

The data presented herein have utilized either immortalized MEFs or primary AEFs to

demonstrate the workflows compatible with SPARKL. These adherent cells demonstrate

minimal PS exposure under normal growth conditions. In our method, cells are incubated

with labeled Annexin V which may bind stochastically-exposed PS on the outer leaflet of

the plasma membrane and result in fluorescent puncta. These puncta are easily excluded

while developing appropriate processing definitions, but may pose a difficulty for cell types

demonstrating significant basal PS localization on the outer leaflet. Additionally, while

MEFs tolerated the use of SYTO21 as a non-perturbing cell marker, other cell lines may

demonstrate increased sensitivity to the nucleotide label, varying its efficacy. During

analysis, accurate quantification of fluorescent events requires segmenting neighboring

fluorescent signals without inappropriately bifurcating individual events. Therefore, cell

lines that grow in colonies or exhibit a "cobble-stone" monolayer may require more

stringent optimization to adequately establish segmentation between neighboring cells.

Of note, apoptotic bodies demonstrated minimal detachment from the plate following cell

death and labeled sufficiently with our detection reagents. However, alternative cell

models or perturbagens may not remain adherent following cell death or may result in

complex morphology that could complicate the quantification or detection of fluorescent

events. While many of these applications can be modified for suspension cells adhered

to the plate by a binding agent, additional troubleshooting is necessary and event

segmentation is generally less representative due to cellular aggregates. Finally, high-

content live-cell imagers require more robust labeling for detection when compared to

flow cytometry and therefore certain methods, such as transient expression of fluorescent

proteins, may not be detectable in these high-throughput workflows.

References

Artymovich K., Appledorn D.M. (2015). A multiplexed method for kinetic measurements

of apoptosis and proliferation using live-content imaging. Methods Mol Biol. *1219*, 35-42.

Biton S., Ashkenazi A. (2011). NEMO and RIP1 control cell fate in response to extensive

DNA damage via TNF-α feedforward signaling. Cell *145*, 92-103.

Chan G.K., Kleinheinz T.L., Peterson D., Moffat J.G. (2013). A simple high-content cell

cycle assay reveals frequent discrepancies between cell number and ATP and MTS

proliferation assays. PLoS One *8*, e63583.

Chipuk J.E., Moldoveanu T., Llambi F., Parsons M.J., Green D.R. (2010). The BCL-2

family reunion. Mol Cell *37*, 299-310.

Degterev A., Huang Z., Boyce M., Li Y., Jagtap P., Mizushima N., … Yuan J. (2005).

Chemical inhibitor of nonapoptotic cell death with therapeutic potential for ischemic brain

injury. Nat Chem Biol *1*, 112–119.

Degterev, A., Hitomi, J., Germscheid, M., Ch'en, I. L., Korkina, O., Teng, X., … Yuan, J.

(2008). Identification of RIP1 kinase as a specific cellular target of necrostatins. Nat Chem

Biol *4*, 313–321.

Devaux P. (1991). Static and dynamic lipid asymmetry in cell membranes. Biochemistry

30, 1163–1173.

Dillon C, Weinlich R, Rodriguez D, Cripps J, Quarato G, Gurung P. (2014). RIPK1 blocks

early postnatal lethality mediated by caspase-8 and RIPK3. Cell *157*,1189–1202.

Dixon S.J., Lemberg K.M., Lamprecht M.R., Skouta R., Zaitsev E.M., Gleason C.E., Patel

D.N., Bauer A.J., Cantley A.M., Yang W.S., Morrison B., Stockwell B.R. (2012).

Ferroptosis: an iron-dependent form of nonapoptotic cell death. Cell *149*, 1060-72.

Fallahi-Sichani M., Honarnejad S., Heiser L.M., Gray J.W., Sorger P.K. (2013) Metrics

other than potency reveal systematic variation in responses to cancer drugs. Nat Chem

Biol *9*,708-14.

Feng H., Stockwell B.R. (2018). Unsolved mysteries: How does lipid peroxidation cause

ferroptosis? PLoS Biol *16*, e2006203.

Forcina G.C., Conlon M., Wells A., Cao J.Y., Dixon S.J. (2017). Systematic Quantification

of Population Cell Death Kinetics in Mammalian Cells. Cell Syst. *4*, 600-610.

Gaudet S., Spencer, S.L., Chen W.W., Sorger P.K. (2012). Exploring the contextual

sensitivity of factors that determine cell-to-cell variability in receptor-mediated apoptosis.

PLoS Comput Biol *8*, e1002482.

Gelles J.D., Chipuk J.E. (2016). Robust high-throughput kinetic analysis of apoptosis with

real-time high-content live-cell imaging. Cell Death Dis. *8*, e2758.

Giampazolias E., Zunino B., Dhayade S., Bock F., Cloix C., Cao K., … Tait S.W.G. (2017).

Mitochondrial permeabilization engages NF-κB-dependent anti-tumour activity under

caspase deficiency. Nat Cell Biol *19*, 1116–1129.

Gong Y.N., Guy C., Crawford J.C., Green D.R. (2017). Biological events and molecular

signaling following MLKL activation during necroptosis. Cell Cycle *16*, 1748-1760.

Gong Y.N., Guy C., Olauson H., Becker J.U., Yang M., Fitzgerald P., Linkerman A., Green,

D.R. (2017). ESCRT-III Acts Downstream of MLKL to Regulate Necroptotic Cell Death

and Its Consequences. Cell *169*, 286–300.

Goodall M.L., Fitzwalter B.E., Zahedi S., Wu M., Rodriguez D., Mulcahy-Levy J.M., …

Thorburn A. (2016). The Autophagy Machinery Controls Cell Death Switching between

Apoptosis and Necroptosis. Dev cell *37*, 337–349.

Hafner M., Niepel M., Chung M., & Sorger P.K. (2016). Growth rate inhibition metrics

correct for confounders in measuring sensitivity to cancer drugs. Nat methods *13*, 521–

527.

Holler, N., Zaru, R., Micheau, O., Thome, M., Attinger, A., Valitutti, S., … Tschopp, J.

(2000). Fas triggers an alternative, caspase-8–independent cell death pathway using the

kinase RIP as effector molecule. Nat Immunol *1*, 489–495.

Inde Z., Dixon S.J. (2018). The impact of non-genetic heterogeneity on cancer cell death.

Crit Rev Biochem Mol Biol *53*, 99–114.

Jiang L., Tixeira R., Caruso S., Atkin-Smith G., Baxter A., Paone S., Hulett M.D., Poon

I.K. (2016). Monitoring the progression of cell death and the disassembly of dying cells

by flow cytometry. Nat Protoc. *4*, 655-663.

Jost P.J., Grabow S., Gray D., McKenzie M.D., Nachbur U., Huang D.C., … Kaufmann T.

(2009). XIAP discriminates between type I and type II FAS-induced apoptosis. Nature,

460, 1035–1039.

Kawahara, A., Ohsawa, Y., Matsumura, H., Uchiyama, Y. & Nagata, S. (1998). Caspase-

independent cell killing by Fas-associated protein with death domain. J. Cell Biol. *143*,

1353–1360.

Kepp O., Galluzzi L., Lipinski M., Yuan J., Kroemer G. (2011). Cell death assays for drug

discovery. Nat Rev Drug Discov *10*, 221-37.

Koopman G., Reutelingsperger C., Kuijten G., Keehnen R., Pals S., van Oers M. (1994).

Annexin V for flow cytometric detection of phosphatidylserine expression on B cells

undergoing apoptosis. Blood *84*, 1415–1420.

Lin J., Kumari S., Kim C., Van T.M., Wachsmuth L., Polykratis A., Pasparakis M. (2016).

IPK1 counteracts ZBP1-mediated necroptosis to inhibit inflammation. Nature *540*, 124-

128.

Lopez J., Bessou M., Riley J.S., Giampazolias E., Todt F., Rochegüe T., … Tait S.W.G.

(2016). Mito-priming as a method to engineer Bcl-2 addiction. Nat Commun *7*, 10538.

Logue S., Elgendy M., Martin S. (2009). Expression, purification and use of recombinant

annexin V for the detection of apoptotic cells. Nat Protoc *4*, 1383-1395.

McStay G.P., Salvesen G.S., Green D.R. (2008). Overlapping cleavage motif selectivity

of caspases: implications for analysis of apoptotic pathways. Cell Death Differ. *15*, 322-

331.

Meers P., Mealy T. (1993). Calcium-dependent annexin V binding to phospholipids:

stoichiometry, specificity, and the role of negative charge. Biochemistry *32*: 11711–11721.

Moujalled D.M., Cook W.D., Okamoto T., Murphy J., Lawlor K.E., Vince J.E., Vaux D.L.

(2013). TNF can activate RIPK3 and cause programmed necrosis in the absence of

RIPK1. Cell Death Dis. *4*, e465.

Murphy J.M., Czabotar P.E., Hildebrand J.M., Lucet I.S., Zhang J.G., Alvarez-Diaz S., et

al. (2013). The pseudokinase MLKL mediates necroptosis via a molecular switch

mechanism. Immunity *39*: 443-53.

Newton K., Wickliffe K.E., Maltzman A., Dugger D.L., Strasser A., Pham V.C., … Dixit

V.M. (2016). RIPK1 inhibits ZBP1-driven necroptosis during development. Nature *540*,

129-133.

Oberst A., Bender C., Green D.R. (2008). Living with death: the evolution of the

mitochondrial pathway of apoptosis in animals. Cell Death Differ *15*, 1139-46.

O'Donnell M.A., Perez-Jimenez E., Oberst A., Ng A., Massoumi R., Xavier R., Green D.R.,

Ting A.T. (2011). Caspase 8 inhibits programmed necrosis by processing CYLD. Nat Cell

Bio *13*: 1437-42.

Oltersdorf, T., Elmore, S. W., Shoemaker, A. R., Armstrong, R. C., Augeri, D. J., Belli, B.

A., … Rosenberg, S. H. (2005). An inhibitor of Bcl-2 family proteins induces regression of

solid tumours. Nature, 435(7042), 677–681.

Onufriev M., Yakovlev A., Lyzhin A., Stepanichev M., Khaspekov L., Gulyaeva N. (2009).

A secreted caspase-3-substrate-cleaving activity at low pH belongs to cathepsin B: a

study on primary brain cell cultures. Biochemistry (Mosc) *74*, 281-287.

Peltzer N., Darding M., Walczak H. (2016). Holding RIPK1 on the Ubiquitin Leash in

TNFR1 Signaling. Trends Cell Biol. *26*, 445-461.

Roux J., Hafner M., Bandara S., Sims J.J., Hudson H., Chai D., Sorger, P.K. (2015).

Fractional killing arises from cell-to-cell variability in overcoming a caspase activity

threshold. Mol Syst Biol *11*, 803.

Schneider, C.A., Rasband, W.S., Eliceiri, K.W. (2012). NIH Image to ImageJ: 25 years of

image analysis. Nat Methods *9*, 671-675.

Schindelin, J., Arganda-Carreras, I., Frise, E. (2012). Fiji: an open-source platform for

biological-image analysis. Nat Methods *9*, 676-682.

Segawa K, Kurata S, Yanagihashi Y, Brummelkamp T, Matsuda F, Nagata S. (2014).

Caspase-mediated cleavage of phospholipid flippase for apoptotic phosphatidylserine

exposure. Science *344*, 1164–1168.

Spencer S.L., Gaudet S., Albeck J.G., Burke J.M., Sorger P.K. (2009). Non-genetic

origins of cell-to-cell variability in TRAIL-induced apoptosis. Nature *21*, 428-32.

Sun L., Wang H., Wang Z., He S., Chen S., Liao D., … Wang X. (2012). Mixed lineage

kinase domain-like protein mediates necrosis signaling downstream of RIP3 kinase. Cell

148, 213-27.

Thévenaz, P., Ruttimann, U.E., Unser, M. (1998). A Pyramid Approach to Subpixel

Registration Based on Intensity. IEEE Trans Image Process *7*, 727-41.

Vanden Berghe T., Vanlangenakker N., Parthoens E., Deckers W., Devos M., Festjens

N., Guerin C.J., Brunk U.T., Declercq W., Vandenabeele P. (2010). Necroptosis, necrosis

and secondary necrosis converge on similar cellular disintegration features. Cell Death

Differ. *17*, 922-30.

Vercammen D., Beyaert R., Denecker G., Goossens V., Van Loo G., Declercq W.,

Grooten J., Fiers W., Vandenabeele P. (1998). Inhibition of caspases increases the

sensitivity of L929 cells to necrosis mediated by tumor necrosis factor. J. Exp. Med. 187,

1477–1485.

Wang H., Sun L., Su L., Rizo J., Liu L., Wang L.F., Wang F.S., Wang X. (2014). Mixed

lineage kinase domain-like protein MLKL causes necrotic membrane disruption upon

phosphorylation by RIP3. Mol Cell *54*, 133-146.

Wang L., Du F., Wang X. (2008). TNF-alpha induces two distinct caspase-8 activation

pathways. Cell *133*, 693-703.

Wei M.C., Zong W.X., Cheng E.H., Lindsten T., Panoutsakopoulou V., Ross A.J., Roth

K.A., MacGregor G.R., Thompson C.B., Korsmeyer S.J. (2001). Proapoptotic BAX and

BAK: a requisite gateway to mitochondrial dysfunction and death. Science *292*, 727-30.

Weinlich R., Oberst A., Beere H.M., Green D.R. (2017). Necroptosis in development,

inflammation and disease. Nat Rev Mol Cell Biol *18*,127-136.

Yu K., Kennedy C.A., O'Neill M.M., Barton R.W., Tatake R.J. (2001). Disparate cleavage

of poly-(ADP-ribose)-polymerase (PARP) and a synthetic tetrapeptide, DEVD, by

apoptotic cells. Apoptosis *6*, 151-160.

Figure Legends

Figure 1. SPARKL provides data-rich kinetic analyses of cell death in real-time using live-cell imaging.

(**A**) Comparison of workflows for cell death detection assays using traditional flow cytometry methods or SPARKL live-cell imaging. SPARKL eliminates sample preparation and handling while coupling real-time detection and analysis into a simple, rapid, and less laborious protocol.

(**B**) MEFs were co-treated with CHX (50 µg/ml) and mTNFα (10 ng/ml) to induce cell death, incubated with titrations of Annexin V-FITC (67, 100, 133, 200, 400, 1000, 2000, and 4000 ng/ml) and YOYO3 (200, 250, 333, 400, 500, 667, 1000, and 2000 nM), and subjected to the SPARKL protocol.

(**C**) MEFs cultured in DMEM lacking or containing phenol red were treated as in **B**. Positive labeling events were analyzed for signal intensity as mean fluorescence intensity integrated over the event area. Relative fluorescence intensity is depicted as a heat map for each titration of Annexin V-FITC and YOYO3.

(**D**) $Bak^{+/+}Bax^{+/+}$ (WT) and $Bak^{-/-}Bax^{-/-}$ (DKO) MEFs were treated with DMSO (0.2%), mTNFα (10 ng/ml) + CHX (50 µg/ml), VP16 (25 µM), or Erastin (10 µM), and incubated with Annexin V-FITC (250 ng/ml) and YOYO3 (250 nM). Images were collected by an IncuCyte ZOOM with a 10× objective and the shown panels are representative images from the stack; 100 µm scale bar. Images were white-point corrected to normalize visualization of green and red pixel background in images lacking a significant fluorescent signal.

(**E**) Kinetic data analyzed from images stacks corresponding to cells from **D**. Dissimilar

range of axis reflects differences in number of total cells due to rates of proliferation. Axis

range is determined by internal controls within each experiment.

(**F**) Comparative endpoint data extracted from **E** at the indicated time-points.

Figure 2. SPARKL is versatile and capable at quantifying and interrogating different

pathways of cell death.

(**A**) Cells dying by apoptotic or non-apoptotic mechanisms exhibit differential labeling of

Annexin V or a cell viability dye (YOYO3) which are not clearly differentiated by flow

cytometric methodologies.

(**B**) WT MEFs were incubated in growth media containing Annexin V-FITC (250 ng/ml)

and YOYO3 (250 nM), treated with CHX (50 µg/ml), mTNFα (10 ng/ml) + CHX (50 µg/ml),

VP16 (25 µM), or Erastin (10 µM), co-treated with DMSO, ABT-737 (1 µM), or zVAD-fmk

(100 µM), and subjected to SPARKL.

(**C**) DKO MEFs were treated and incubated as in **B**.

(**D**) *Cyld*[-/-] MEFs reconstituted with either GST or CYLD were incubated in growth media

containing Annexin V-FITC (250 ng/ml) and YOYO3 (250 nM), mTNFα (20 ng/ml) + CHX

(50 µg/ml), zVAD-fmk (40 µM), and necrostatin-1 (10 µM), and subjected to SPARKL.

(**E**) Primary adult ear fibroblasts (AEFs) were cultured from *Mlkl*[+/+] and incubated in

growth media containing Annexin V-FITC (250 ng/ml) and YOYO3 (250 nM), mTNFα (50

ng/ml) + Birinapant (5 µM), zVAD-fmk (40 µM), and necrostatin-1 (10 µM), and subjected

to SPARKL.

(**F**) Representatives images collected by the IncuCyte ZOOM from *Mlkl*^{+/+} AEFs treated

with mTNFα (50 ng/ml), Birinapant (5 μM), zVAD-fmk (40 μM), with or without necrostatin-

1 (10 μM). Cells are labeled with Annexin-FITC and YOYO3 (green and red, respectively).

Stitched insets are a conserved magnification. Scale bars represent 100 μm.

(**G**) AEFs from either *Mlkl*^{+/+} or *Mlkl*^{-/-} mice were treated with mTNFα (50 ng/ml) +

Birinapant (10 μM), and co-treated with zVAD-fmk (40 μM), with or without necrostatin-1

(10 μM), and subjected to SPARKL in the presence of Annexin V-FITC (250 ng/ml) and

YOYO3 (250 nM).

(**H**) WT and *Mlkl* KO AEFs were cocultured in a variety of ratios and treated as in *G*.

Figure 3. A dual-reporter SPARKL workflow for characterizing modes of cell death
at population and single-cell level.

(**A**) Number of YOYO3-positive events was subtracted from the number of Annexin V-

positive events for cells analyzed in *Figure 2B*.

(**B**) Number of YOYO3-positive events was subtracted from the number of Annexin V-

positive events for cells analyzed in *Figure 2G*.

(**C**) WT and *Mlkl* KO AEFs were cocultured in a 1:1 ratio, treated with mTNFα (50 ng/ml)

+ Birinapant (10 μM), and co-treated with zVAD-fmk (40 μM), with or without necrostatin-

1 (10 μM), and subjected to SPARKL in the presence of Annexin V-FITC (250 ng/ml) and

YOYO3 (250 nM).

(**D**) Annexin V- and YOYO3-positive events for cells treated and analyzed in *C* were

normalized as a percent of total signal and the percent YOYO3-positive was subtracted

from the percent Annexin V-positive for each time point.

(**E**) Histogram data from cells treated and analyzed as in **A-D** shown as number of new

Annexin V (green) or YOYO3 (red) events occurring during each hour.

(**F**) WT MEFs were treated with CHX (50 µg/ml), mTNFα (10 ng/ml) + CHX (50 µg/ml),

VP16 (25 µM), or Erastin (10 µM), and co-treated with ABT-737 (1 µM); WT AEFs were

treated with mTNFα (50 ng/ml) + Birinapant (10 µM) and co-treated with zVAD-fmk (40

µM). Cells were subjected to SPARKL in the presence of Annexin V-FITC (250 ng/ml)

and YOYO3 (250 nM). Individual cells within the population imaged by the machine were

tracked and analyzed to identify the time at which a cell was single- and double-positive.

Panels show analysis where single-positive is defined as Annexin V-positive and double-

positive is defined by subsequent labeling with YOYO3.

(**G**) Same data as in **F** but panels show analysis where single-positive is defined as

YOYO3-positive and double-positive is defined by subsequent labeling with Annexin V.

(**H**) WT MEFs and WT AEFs treated and analyzed in **F**. Population distribution of time-to-

signal is shown for the following metrics: time to Annexin V (green), time to YOYO3 (red),

time to double-positive (yellow), time to YOYO3 after Annexin V (transparent red), and

time to Annexin V after YOYO3 (transparent green). Green, red, and yellow plots are

scaled to the left axis while the transparent plots are scaled to the right axis.

(**I**) WT MEFs and WT AEFs were treated and analyzed as in **F** and **G** without co-treatment

of ABT-737 or zVAD-fmk, respectively. Violin plots of collected data were compared with

data from **H** in which cells were co-treated as indicated. Green, red, and yellow plots are

scaled to the left axis while the transparent plots are scaled to the right axis.

Error bars are not shown for kinetic data to aid in data visualization. Violin plots display

the entire population of collected data and have been smoothened to aid in visualization.

Instances of flat ends are an artifact of the method used to create the frequency

distribution and do not reflect missing or truncated data points. Median and quartiles are

shown as a solid and dotted line, respectively. Population statistics were conducted by

performing a Mann-Whitney U test for each indicated comparison. Significance is noted

as follows: ****$P < 0.0001$, ***$P < 0.001$, **$P < 0.01$, *$P < 0.05$, ns = not significant.

Figure 4. Mathematical analysis of the SPARKL kinetic data provides several

parameters for high-throughput comparative assays.

(**A**) Example methods for mathematically analyzing kinetic data and extracting

parameters for cell death comparative studies. Area under the curve (AUC) provides a

single parameter for comparing kinetic data between treatments or cell types.

Mathematical conversion of kinetic data can provide additional comparative parameters

such as death rates via derivatization or death onset from non-linear fit analyses.

(**B**) WT and DKO MEFs were incubated in DMEM containing Annexin V-FITC (250 ng/ml)

and YOYO3 (250 nM), treated with CHX (50 µg/ml), mTNFα (10 ng/ml) + CHX (50 µg/ml),

VP16 (25 µM), or Erastin (10 µM), co-treated with DMSO, ABT-737 (1 µM), or zVAD-fmk

(100 µM), and subjected to SPARKL. Area under the curve (AUC) values were calculated

from kinetics of Annexin V- and YOYO3-labeling. Dissimilar range of axis reflects

differences in number of total cells due to rates of proliferation. Axis range is determined

by internal controls within each experiment.

(**C**) First derivative of data collected from cells treated as in **B** depicting the rates of

population cell death at each time point.

(**D**) Comparison of maximum rates of death as calculated from first derivatives in **C**.

(**E**) Fit curves using a Lag One-Phase Growth function (solid line) applied to data from

cells treated in **B** (dotted line). Curve for YOYO3 labeling kinetics in cells treated with

VP16 and zVAD-fmk was considered an ambiguous fit and therefore excluded from

parameter extraction and comparisons.

(**F**) Comparison of lag phase time (t') from Annexin V and YOYO3 data fits for cells treated

as indicated from **E**.

(**G**) Comparison of initial/maximal rate of death (R_D) from Annexin V and YOYO3 data fits

for cells treated as indicated from **E**.

(**H**) 2D graphs of parameters calculated in **E** and graphed in **F** and **G** provide a convenient

method for comparing treatment trends across labels and cell types.

Error bars are not shown in instances where they are smaller than the data symbol.

1023

Figure 5. Non-toxic label-and-go reagents used with SPARKL can normalize data

to account for changes in cell proliferation.

(**A**) Label-and-go method to collect proliferation data during Annexin V-binding assay for

composite cytostatic and cytotoxic effects of perturbagens.

(**B**) WT MEFs were incubated with titrations of SYTO nucleotide labeling reagents (0.03-

20 µM) in the presence of Annexin V-594 (1 µg/ml) and subjected to SPARKL in order to

assess toxicity of the SYTO dyes.

(**C**) WT MEFs were co-treated with CHX (50 µg/ml) and mTNFα (10 ng/ml) and incubated

with Annexin V-594 (1 µg/ml) and SYTO21 (300 nM). Images captured at indicated time-

points.

(**D**) Kinetic data analyzed from images stacks corresponding to cells from **C**.

(**E**) WT MEFs were seeded at the indicated numbers 24 hours prior to treatment with

mTNFα (10 ng/ml) + CHX (50 μg/ml) and incubation with Annexin V-594 (1 μg/ml) and

SYTO21 (300 nM). Cells were then subjected to SPARKL to analyze proliferation and cell

death kinetics.

(**F**) Cell death data from **E** was normalized using SYTO21 and expressed as percentage

of the population exhibiting death (% Death).

(**G**) Endpoint data of Annexin V-594-positive events from **E** and corresponding percent

death (% Death) normalization from **F**.

1043

Figure 6. Parameters extracted from non-linear fit analyses of normalized death

kinetics generate comparative profiles for studying mechanisms of cell death.

(**A**) WT and DKO MEFs were incubated in DMEM containing SYTO21 (300 nM), Annexin

V-549 (1 μg/ml), indicated treatments of CHX (50 μg/ml), TRAIL (25 ng/ml) + CHX (50

μg/ml), mTNFα (10 ng/ml) + CHX (50 μg/ml), VP16 (25 μM), Erastin (10 μM), or A23187

(5 μM), co-treated with either DMSO control or ABT-737 (1 μM), and subjected to

SPARKL workflow. Percent death (% Death) was calculated by Annexin V data

normalized by the SYTO21 data at each timepoint.

(**B**) Cells, treatments, and analysis as in **A** for cells cultured in DMEM containing SYTO21

(300 nM) and YOYO3 (250 nM).

(**C**) WT MEFs treated as in **A** were subjected to long-term, repetitive detection of

SYTO21-, Annexin V-594-, and YOYO3-positive events to assess signal permanence and

longevity.

(**D**) Average integrated intensity of events detected in **C** showing changes in fluorescent

intensity of reporters over the first 48 hours.

(**E**) Fit curves using a Lag One-Phase Growth function applied to data from WT cells in

A. Curve for DMSO-treated cells was considered an ambiguous fit and therefore its

parameterization is variable.

(**F**) Comparison of lag phase time (t') from fits applied to cells treated as indicated in **A**.

(**G**) Comparison of initial/maximal rate of death (R_D) from fits applied to cells treated as

indicated in **A**.

(**H**) 2D graph of parameters calculated in **E** and graphed in **F** and **G** provides a convenient

method for comparing trends across various cytotoxic treatments. Error bars are not

shown for kinetic data to aid in data visualization, and not shown in instances where they

are smaller than the data symbol.

Materials and Methods

Reagents and equipment

Cell culture reagents and chemicals were from Sigma Aldrich (St. Louis, MO, USA) or

Thermo Fisher Scientific (Waltham, MA, USA) unless otherwise stated. Drugs and

biologics as follows: ABT-737 (Abbott Pharmaceuticals, Lake Bluff, IL, USA); CHX

(Sigma); YOYO3 and SYTO reagents (Life Technologies/Thermo Fisher Scientific),

mTNFα and mTRAIL (Peprotech, Rocky Hill, NJ, USA); zVAD-fmk (ApexBio, Houston,

TX, USA). Recombinant Annexin V was purified, labeled with FITC or Alexa Fluor 594

(Thermo Fisher Scientific), and re-purified using Dye Removal Columns (Thermo Fisher

Scientific) following conjugation. Purity, stability, and fluorescence of labeled recombinant

Annexin V was verified by SDS-PAGE and Flow Cytometry as previously described

(Logue et al., 2009).

1081

Cell culture and treatments

Unless stated to the contrary, experiments were performed with $Bak^{+/+}Bax^{+/+}$ and

$Bak^{-/-}Bax^{-/-}$ (WT and DKO, respectively) SV40-transformed MEFs (ATCC, Manassas,

VA, USA). $Cyld^{-/-}$ MEFs were reconstituted with either a Flag tagged CYLD construct or

empty vector as described previously (O'Donnell et al. 2011). Primary adult ear fibroblasts

(AEFs) were isolated from $Mlkl^{+/+}$ and $Mlkl^{-/-}$ mice as follows: mouse ears were clipped,

minced, and incubated in trypsin at 37°C for 1 hour with repeated vortexing before being

cultured on tissue culture plates (Murphy et al., 2013). Cells were cultured in DMEM

containing 10% FBS, 2 mM L-glutamine, and antibiotics and maintained in humidified

incubators at 37°C with 5% CO_2. For kinetic studies, cells were seeded at approximately

$2 - 4 \times 10^3$ cells/well in a 96-well format and left to adhere for 24 hours. Prior to live-cell

imaging, growth media was replaced with complete phenol-red-free DMEM containing

indicated fluorescent reagents and perturbagens. DMEM contains sufficient calcium

concentrations for robust Annexin V labeling but other media formulations require calcium

supplementation for optimal Annexin V labeling (1.5 – 2 mM $CaCl_2$). Plates were pre-

warmed prior to data acquisition to avoid condensation and expansion of the plate, which

hinders auto-focusing during scan intervals. All tissue cultures were confirmed

mycoplasma free with the colorimetric PlasmoTest detection kit (Invivogen, San Diego,

CA, USA).

1101

SPARKL data acquisition and automated analysis

Kinetic experiments were performed with the IncuCyte ZOOM (Model 4459, Essen

Bioscience, Ann Arbor, MI, USA) residing in a tissue culture incubator maintained as

above, but other high-content live-cell fluorescent imagers are capable of replicating

these data (e.g., IN CELL analyzer). Experiments were conducted for 24 – 96 hours with

data collection every 1 – 2 hours to avoid photobleaching of fluorescent reporters. Using

the 10× objective, a single plane of view was collected per well for 96-well plate assays.

Phase contrast, green channel (Ex: 440/80 nm; Em: 504/44 nm; acquisition time: 400 ms),

and red channel (Ex: 565/05 nm; Em: 625/05 nm; acquisition time: 800 ms) were collected

for all experiments with spectral unmixing set as 3% of red removed from green. Images

collected were 1392 × 1040 pixels at 1.22 µm/pixel. Automated image analysis was

accomplished using the ZOOM software (V2018A) and defined by images collected using

the specific cell lines and fluorescent reporters pertaining to the experiment. Processing

definitions were designed using the settings for each fluorescent reporter as described

below (Table 1). Data are expressed as events per well. The y-axis scale was determined

for each experiment using internal controls to assess maximal death for a population.

Fluorescent intensity data were analyzed using the average integrated intensity (AU ×

μm^2) metric. Images of cell culture presented in this manuscript were white-point

corrected for clarity and normalization of automated background subtraction in the green

and red channel when no significant fluorescent event was detected. White-point

correction was accomplished by adjusting temperature and tint parameters to overlap

peak contributions of red, green, and blue pixels.

1124

1125 **Table 1: Channel settings for each fluorescent reporter used for processing**

1126 **definitions with the IncuCyte ZOOM**

	Channel	Parameters	Radius (μm)	Threshold (RFU)	Edge Sensitivity	Area (μm²)
Annexin V-AF594	Red	Top-Hat	10	5	-35	>150
Annexin V-FITC	Green	Top-Hat	25	3	-30	>100
Annexin V-FITC (low)	Green	Top-Hat	25	2	-30	>50
SYTO21	Green	Top-Hat	25	0.5	-25	>150
YOYO3	Red	Top-Hat	10	3	-46	>150

1127

Parameter extraction, mathematical processing, and data handling

Data exported from the IncuCyte ZOOM software were organized, handled, and analyzed

using Excel v16.16.9 (Microsoft) and Prism 8.1.1 (Graphpad Software Inc.). Axis range

for graphs was determined by internal negative and positive controls within each

experiment. Event count data were converted into histograms using Equation 1, where

#Signal[+] refers to the number of events using either fluorescently-labeled Annexin V or

YOYO3 for a given well "*i*" at time point "*j*":

1135
$$\#Events/hour_{well=i,time=j} = \left(\#Signal+_{i,j} - \#Signal+_{i,j-1}\right) \quad \text{(Equation 1)}$$

Area under the curve (AUC) values were calculated with the analysis tool in Prism and

utilized the trapezoid rule ignoring peaks that were less than 10% of the distance from

minimum to maximum Y values. The first derivative of collected data was calculated with

the analysis tool in Prism and smoothed the data using 4 neighbors and 2nd order

smoothing polynomial. Non-linear curve fits were calculated using the plateau followed

by one-phase association function in Prism using the least squares fitting method given

by Equation 2. Parameters extracted by Equation 2 are defined as follows: $D(t'')$ is the

amount of death for given time point t''; D_0 is the initial and minimum amount of death; D_M

is the plateau of maximal death; R_D is the initial and maximal rate of death; t' is the time

at which the lag plateau ends and exponential increase in death begins. All parameters

were left unconstrained for mathematical fitting. Fits listed as ambiguous by Prism only

occurred in conditions that did not result in death positivity and are highlighted in the text.

1148
$$D(t'') = D_0 + (D_M - D_0)\left(1 - e^{-R_D(t''-t')}\right) \quad \text{(Equation 2)}$$

1149   Experiments using SYTO21 to normalize data with cell counts was accomplished using

1150   Equation 3, where #Dead was determined by the event counts using either fluorescently-

1151   labeled Annexin V or YOYO3 for a given well "i" at time point "j":

1152
$$\%Death_{well=i,time=j} = \left(\frac{\#Dead_{i,j}}{\#SYTO+_{i,j}}\right) \quad \text{(Equation 3)}$$

1153   Unless otherwise noted, all experiments were conducted in triplicate and the presented

1154   data are the mean of replicates and representative of repeated experiments. When

1155   shown, error bars denote standard deviation of replicates; select graphs omit error bars

1156   for convenience of visualization.

1157

Single-cell two-hit kinetics

For experiments investigating sequential labeling of individual cells, analysis utilized

event masks generated by the ZOOM software which were exported for further

processing using external software packages. Specifically, masks for green and red

channels as well as the overlap masks were used for these analyses. Cell event masks

were recognized and spatio-temporally defined using ImageJ and Fiji software packages

(ImageJ, National Institutes of Health, Bethesda, ML, USA) (Schneider et al., 2012,

Schindelin et al., 2012). Images exported from the ZOOM software were imported into

ImageJ as stacks for each mask collection (green, red, and overlay), converted to 8-bit

grayscale and binary, and subjected to translation alignment using the StackReg and

TurboReg plugins (Thévenaz et al., 1998). Reference regions of interest (ROIs) were

autonomously defined by applying the Analyze Particles tool on the final image slice of

the overlay mask. These reference ROIs were applied to all slices within the image stacks

corresponding to the green, red, and overlay channels using the Multi-Measure tool.

Positive pixel intensity within ROIs defined the slice in which the signal occurred. These

data were then sorted and aligned in Excel by ROI reference number, thereby linking

identified signal slices from each channel to the same ROI for graphing and interpretation.

This workflow was used to quantify the progression of either green or red single-positivity

to double-positivity for single cells within in a population.

A

Flow Cytometry Workflow

Treatment → Incubation → Preparation → Analysis → End-point Data

%Death | t_1 t_2 t_3

SPARKL Workflow

Treatment → Incubation, Data Acquisition, & Analysis → Graph

Image Stacks | Kinetic Data | Timepoints

%Death vs Time — t_1 t_2 t_3

B Fluorescent Signal

TNFα+CHX — AV
Events (×10³) vs Time (hours)
UT

TNFα+CHX — Y3
Events (×10³) vs Time (hours)
UT

C Fluorescent Intensity

DMEM
− + : Phenol Red — AV
− + : Phenol Red — Y3

D

WT (*Bax⁺/⁺Bak⁺/⁺*) MEFs | DKO (*Bax⁻/⁻Bak⁻/⁻*) MEFs

DMSO: 0h 12h 24h | 0h 12h 24h
TNFα+CHX: 0h 12h 24h | 0h 12h 24h
VP16: 0h 12h 24h | 0h 12h 24h
Erastin: 0h 12h 24h | 0h 12h 24h

E

AV488+ or YOYO3+ Events per Well (×10³)

WT | DKO

DMSO | DMSO
TNFα+CHX | TNFα+CHX
VP16 | VP16
Erastin | Erastin

Time (hours)

F

AV488+ Events for WT or DKO MEFs (×10³)

■ WT ■ DKO

6h 12h 24h