Infectious disease phylodynamics with occurrence data

Leo A. Featherstone[1]*, Francesca Di Giallonardo[2], Edward C. Holmes[3,5,4], Timothy G. Vaughan[6,7], Sebastián Duchêne[1]

[1]Department of Microbiology and Immunology, Peter Doherty Institute for Infection and Immunity, University of Melbourne, Melbourne, VIC, Australia.

[2]The Kirby Institute, UNSW Sydney, Sydney, NSW, Australia.

[3]Marie Bashir Institute for Infectious Diseases and Biosecurity, The University of Sydney, Sydney, NSW, Australia.

[4]Charles Perkins Centre, School of Life and Environmental Sciences, The University of Sydney, Sydney, NSW, Australia.

[5]School of Medical Sciences, The University of Sydney, Sydney, NSW, Australia.

[6]Department of Biosystems Science and Engineering, ETH Zurich, Basel, Switzerland.

[7]Swiss Institute of Bioinformatics (SIB), Switzerland.

* Contact
Email: leo.featherstone@unimelb.edu.au

Abstract (350 words max. currently 173)

Point 1: Phylodynamic models use pathogen genome sequence data to infer epidemiological dynamics. With the increasing genomic surveillance of pathogens, especially amid the SARS-CoV-2 outbreak, new practical questions about their use are emerging.

Point 2: One such question focuses on the inclusion of un-sequenced case occurrence data alongside sequenced data to improve phylodynamic analyses. This approach can be particularly valuable if sequencing efforts vary over time.

Point 3: Using simulations, we demonstrate that birth-death phylodynamic models can employ occurrence data to eliminate bias in estimates of the basic reproductive number due to misspecification of the sampling process. In contrast, the coalescent exponential model is robust to such sampling biases, but in the absence of a sampling model it cannot

exploit occurrence data. Subsequent analysis of the SARS-CoV-2 epidemic in the

northwest USA supports these results.

Point 4: We conclude that occurrence data are a valuable source of information in

combination with birth-death models. These data should be used to bolster phylodynamic

analyses of infectious diseases and other rapidly spreading species in the future.

Key Words: Phylodynamics, pathogens, coalescent, birth-death, Bayesian statistics

Introduction

Outbreak investigations increasingly rely on genome sequencing of causative pathogens.

Phylodynamic methods take advantage of these data to infer epidemiological dynamics

(Rife et al., 2017). New sequencing technologies generate these data rapidly, such that

phylodynamic inferences can be conducted in actionable time frames (Gardy & Loman,

2018; Grubaugh et al., 2019; Hadfield et al., 2018). In this context, the main appeal of

phylodynamics is that it uses sequence data to infer epidemiological dynamics preceding

the earliest collected sample, or during periods without collected sequences, and offers

insight into transmission chains.

Phylodynamic models describe a branching process, modelling both how a branching

transmission chain and phylogenetic tree of the underlying pathogen evolve. These are

central to linking epidemiological dynamics to the evolution of a pathogen. In Bayesian

phylogenetic implementations the particular model of a branching process is part of the

prior and is sometimes referred to as the 'tree prior', such as the birth-death or coalescent

exponential. Internal nodes in the tree are associated with transmission events while the

tips of the tree represent sampling events (du Plessis & Stadler, 2015). The basic

reproductive number, R_0, is a key parameter that reflects the average number of secondary

infections in a fully susceptible population. The simplest tree priors that can infer R_0 posit

that the number of infected individuals increases exponentially over time. Although more

sophisticated methods now exist (Kühnert et al., 2014; Popinga et al., 2015; Rasmussen et

al., 2017; Vaughan et al., 2019; Volz & Siveroni, 2018), we focus here on tree priors

assuming simple exponential growth since they are appropriate for the early stages of an

outbreak and are increasingly used to assess the efficacy of public health interventions

(Geoghegan et al., 2020; Vasylyeva et al., 2019).

Two commonly used phylodynamic tree priors are the coalescent exponential and the birth-

death, both of which assume that the infected population size, N, grows at a rate r; $N(t)=e^{rt}$,

where t is time after the origin. From an epidemiological perspective, r is the difference

between the transmission rate, λ, and the become uninfectious rate, δ, ($r=\lambda-\delta$). $1/\delta$ is the

duration of infection. R_0 is estimated as $R_0=\lambda/\delta$. The exponential coalescent is a

generalisation of the Kingman-n coalescent where population size is a deterministic

function of time (Griffiths & Tavare, 1994; Volz et al., 2009, 2013). In contrast, the birth-

death tree prior assumes stochastic population growth with sampling through time

(Stadler, 2010; Stadler et al., 2012; Stadler & Yang, 2013). This is captured in the death

rate$\delta=\psi+\mu$, where μ is the recovery rate and ψ is the sampling rate such that the sampling

proportion, p, can be calculated as $p = \frac{\psi}{\psi+\mu}$.

Phylodynamic analyses draw from sequence data and sampling times (Biek et al., 2015;

Drummond et al., 2002, 2003; Rambaut, 2000; Rieux & Balloux, 2016). In the coalescent

exponential, sampling times are useful insofar as they influence the distribution of

coalescent events through time, influencing R_0 in turn. Coalescent models typically

condition upon sampling times instead of using them to infer sampling rates. Some

'augmented likelihood' approaches can combine the coalescent with a sampling process

(Volz & Frost, 2014), but they are not standard practice. For the birth-death tree prior, the

number of samples and their times are naturally informative because they are explicitly

modelled through the sampling rate (i.e. they inform ψ) (Boskova et al., 2018). This is a well

understood difference between the two tree priors, but its consequences remain to be

explored in the context of occurrence data. Although the amount of sequence data in

outbreak investigations has increased, a key consideration is that sequencing efforts are

often conducted only after relatively a large number of cases are reported. This latency in

sampling can bias estimates of epidemiological parameters. To visualise this, the trees in

Fig 1 were simulated under an R_0 of 2, a constant sampling effort, and over the course of 1

year. If sequencing were only conducted for samples collected after 0.75 years, samples

from the deep sections of the tree would be missed (*late sampling* in Fig 1). Such sampling

bias can mislead inferences of epidemiological dynamics because there is no sampling

data and very few branching events to inform inferences of the early stages of the outbreak.

Here we investigate bias in epidemiological parameters due to sampling heterogeneity and

present two approaches to reduce such bias using occurrence data. The first approach

involves using a birth-death skyline tree prior that requires an understanding of the

sampling effort (Stadler et al., 2013). If it is known that there was no attempt to collect

samples early in the outbreak, one can set two intervals for the ψ parameter where one is

zero. However, without knowledge of sampling effort this scenario is indistinguishable from

a constant sampling effort where initial prevalence was so low as to preclude obtaining any

sequence data early in an outbreak. The second approach consists of including early case

occurrences in analyses, where an occurrence is a laboratory confirmed case that was not

sequenced (*occurrences* scenario in Fig 1). Occurrence data are a relatively inexpensive

and often readily available source of information because they are traditionally used in

epidemiology and accurately identified via contact tracing and testing efforts. In a Bayesian

phylogenetic framework, topological uncertainty due to occurrence data is naturally

incorporated into the analysis through the posterior. An analogous approach can be used

to coherently specify fossil data for molecular clock calibration (Heath et al., 2014; Heath &

Moore, 2014). This approach and others have been modelled, but not applied in

phylodynamics hitherto (Gupta et al., 2020; Manceau et al., 2019).

Materials and Methods

Simulation study

We simulated phylogenetic trees under a birth-death process in MASTER v6.1 (Vaughan &

Drummond, 2013), with the following parameterisation; R_0=2 or 1.5, δ=91, p=0.05, and an

outbreak duration of one year (1/δ = 0.011 years, corresponding to an expected infectious

period of about 4 days). The number of tips and their ages are naturally variable (from 100

to 150 tips). We assumed a strict molecular clock with an evolutionary rate of 0.01

substitutions per site per year (subs/site/year) and the HKY+Γ substitution model to

produce alignments of 13,000 nucleotides using NELSI (Ho et al., 2015) and Phangorn v2.4

(Schliep, 2011). These settings are broadly similar to an influenza virus outbreak (Hedge et

al., 2013), but a rescaling of the epidemiological parameters could apply to many other

pathogens. We then assumed three sampling scenarios: (i) *constant* sampling with all

sequences from the simulation included (e.g. the sequence for every sample in the tree in

Fig 1 is included), (ii) *late sampling* only with samples after time T_s (e.g. only sequences for

samples after the dashed line in the tree in Fig 1), and (iii) *occurrences* in which sequence

data are available only after time T_s with those preceding recorded as occurrences. We set

T_s to 0.75 or 0.9 years. For each parameter configuration we simulated 100 sequence data

sets which were subsampled according to the three scenarios above. Occurrences were

emulated by replacing simulated DNA sequences with 'n' (i.e. missing data) in the

alignment. We analysed the data in BEAST v2.5 (Bouckaert et al., 2019) with coalescent

exponential and the birth-death tree priors. Our results focus on the birth-death, but the

coalescent exponential forms a valuable point of comparison through its robustness to

variation in sampling. For the *late sampling* scenario, we also considered the birth-death

skyline (BDSky in figures) with two intervals for the ψ parameter, with the interval time fixed

at T_s. We matched the substitution and clock model to those used to generate the data and

we used an informative prior on δ using a Γ distribution with mean fixed to the true value of

91 and standard deviation of 1.

We assessed the effectiveness of each analysis treatment using three statistics. First, we

considered the coverage as a measure of accuracy, or the number of times the 95%

highest posterior density (HPD) intervals covered the true value of a given parameter.

Second, we consider 'average bias', which is the difference between the posterior mean

and true mean for a given parameter averaged across the 100 simulations for each

sampling treatment. Third, we consider average 95% HPD width for each treatment, as a

measure of precision.

Empirical case study

To illustrate the accuracy of occurrence data relative to completely sequenced data sets we

analysed 821 whole genome sequences sampled from the SARS-CoV-2 pandemic from

Washington State, USA, and the adjacent Washington County, Oregon, downloaded from

GISAID (Supplementary material) and partially documented by (Bedford et al., 2020).

Accordingly, we downloaded 2,164 high-coverage genome sequences collected between

January 18 and June 30 2020, but selected the 821 sequences taken up to March 21 2020

to capture an exponential phase in the epidemic and sampling (Fig S1). We corroborated

exponential growth in the underlying population using an Epoch Sampling Proportion

Skyline Plot (Parag et al., 2020). We further divided this data set into five subsets as per our

simulation study: (i) 'complete sampling' including all 821 sequences; (ii) late sampling post

March 6 2020 (decimal date 2020.18) including 637 sequences; (iii) late sampling post

March 14 2020 (2020.20) including 340 sequences; (iv) late sampling post March 6 2020

(2020.18) including 637 sequences and 184 occurrences; and (v) late sampling post 2020.2

including 340 sequences and 481 occurrences. Including two late sampling data subsets

offers information about how inflation in R_0 varies with latency in sequences.

We then analysed each data set with each tree prior used in the simulation study with

BEASTv2.5. We first employed a birth-death model with serial sampling. We placed a

lognormal prior on R_0 with mean 0 and standard deviation of 1; fixed δ at 36.5 (i.e. 10-day

duration of infection as estimated recently (Price et al., 2020)); a β prior on sampling

proportion with shape and scale equal to 2 to penalise extreme values. Second, we used a

birth-death skyline with the same priors as the birth-death, but with two sampling rate

parameters. The first pertained to after the 2020.18 or 2020.2 cut-off, and the second to

before the cut-off. Both used the same beta prior for sampling proportion as for the birth-

death. Third, a coalescent exponential tree prior was used with a Laplace prior on growth

rate with mean 0 and scale 100 and an exponential prior with mean 100 on the coalescent

exponential effective population size (ϕ). For both tree priors, we assumed HKY+Γ

substitution model with a strict molecular clock rate fixed to 10^{-3} subs/site/year, following

recent estimates (Duchene et al., 2020). We ran a Markov chain Monte Carlo of 5×10^{8} steps,

sampling at every 1000^{th} step. We determined sufficient sampling from the posterior by

verifying that the effective sample size all parameters of interest was above 200.

Results

Simulation study

Analyses of data sets with late sampling using the birth-death model were least accurate in

estimating R_0. In only 12 of 100 simulations with R_0=2 did the 95% HPD include 2 (Table 1

and Fig 2a). The true value was never recovered for simulations with R_0=1.5 (Table S1 and

Fig S2). The birth-death skyline was more accurate with 95 and 92 of 100 simulations

covering R_0=2 and R_0=1.5 respectively. The coalescent exponential was also more accurate

with 100 and 80 simulations having HPD intervals that covered R_0=2 and R_0=1.5

respectively. However, this came at the cost of low precision as HPD width was the largest

for the coalescent out of all treatments.

In general, we observed that the birth-death model tended to overestimate R_0 while the

coalescent exponential underestimated it for data sets with late sampling (Fig 2). Estimates

of the evolutionary rate displayed an identical pattern to those of R_0, with the coalescent

exponential and the birth-death model being the most and least accurate respectively at

the expense of precision. However, the evolutionary rate appeared overall robust to the

choice of the tree prior, with the only treatment producing a less than 90% coverage being

the birth-death model with late sampling. This is a valuable consideration for analyses of

future outbreaks as considerable attention is initially devoted to estimating a reliable

evolutionary rate for a given pathogen because this is key to phylodynamic inference

(Duchene et al., 2020).

As expected, analyses of the data with constant sampling were accurate in a majority of

cases, with 94 and 89 out of 100 simulations covering R_0 alongside 94 and 92 for the

evolutionary rate under the birth-death and the coalescent exponential models,

respectively. The true model is the birth-death, and as such it is expected to perform better

than the coalescent. Estimates of R_0 including occurrence data were similar in accuracy to

those with complete sampling. A total of 94 analyses correctly estimated this parameter

under the birth-death model, and 96 analyses included the true value for the coalescent

exponential. Evolutionary rate estimates with occurrence data were similar, with 95

accurate estimates using the birth-death model and 91 using the coalescent exponential

(Table 1, Fig2a,d). These results are attributable to the fact that the birth-death model treats

sampling times as data, whereas the coalescent exponential model conditions on the

number of samples and their ages (Boskova et al., 2018; Stadler et al., 2015). In the birth-

death model, occurrence data improve accuracy when inferring R_0 and are also informative

about the age of the tree height under this tree prior, which can also improve the accuracy

of the evolutionary rate relative to the coalescent exponential model. But these estimates

are unlikely to be as accurate as those with complete sequence data because they include

less information.

The coalescent exponential model appears to be more robust to the sampling treatment,

with greater accuracy than the birth-death model across late sampling and occurrence

treatments. Our simulations suggest that this comes at the expense of less precise

estimates than those from the birth-death model (Table 1). In turn, birth-death and birth-

death skyline models tend to produce more precise estimates with less bias (Table 1, Fig

2). Together these results suggest that in a genomic-reporting scenario, the coalescent

exponential is suitable when sampling proportion is assumed to be low, when the sampling

process is otherwise poorly understood, or when no reliable occurrence data are available.

However, when increased precision is desirable and occurrence data are available, birth-

death tree priors may provide the sharper estimates with comparable accuracy. The choice

of tree prior could be optimised depending on prioritisation of precision and bias based on

the ordering of bars in Figure 2.

Empirical case study: SARS-CoV-2 from the northwest USA

Mirroring trends in our simulated data sets, the coalescent exponential returned consistent

estimates of R_0 across treatments which were generally lower than those inferred by the

birth-death tree prior (Fig 3a, Table 2). Coalescent exponential treatments again produced

wider HPD intervals than birth-death treatments, with the exception of late sampling which

was highly uncertain under the birth-death, as expected from simulations. Uncertainty in

posterior R_0 does not appear to change when substituting sequenced data for occurrence

data (Fig 3A), indicating that late samples are highly informative while occurrence data

contribute relatively little additional information to coalescent analyses. Moreover, we

observed a near perfect match between estimates from analyses with only late sampling

and those that included occurrences. This pattern can be explained because occurrence

data have no influence on marginal posterior estimates under the coalescent. By contrast,

our simulations show small differences in performance between coalescent analyses with

late sampling and those with occurrence data, which we attribute to noise in the simulation

study.

The results of the birth-death analyses recapitulate our observation from simulations that

later sampling inflates estimates of R_0, and that occurrence data rectify this (Figure 3B table

2). Complete sampling gave a mean R_0 of 1.96 (95% HPD: 1.85, 2.07) and late sampling

with occurrence data estimated mean R_0 of 1.95 and 2.00 (95% HPDs: 1.8, 2.11 and 1.9,

2.12 for post - 2020.18 and 2020.2 respectively). These estimates are slightly lower than

those from earlier work to estimate R_0 in the Washington state epidemic (Vaughan et al.,

2020). This discrepancy may be due to the former being conducted earlier when the virus

may have been spreading more rapidly. Late sampling alone inferred a mean R_0 of 2.44 and

3.53 for post 2020.18 and 2020.2 (2.31, 2.58 and 3.24, 3.82 95% HPDs respectively). The

way in which the latest sampling data set inferred the highest values of R_0 further suggests

that upward bias increases with lateness in sampling.

In both late sampling treatments, the birth-death skyline posterior distributions of R_0 were

lower than their equivalents under the standard birth-death model, with later sampling

corresponding to lower estimates (Fig 3). This is consistent with the simulated data (Fig S2),

and suggests that including occurrence data is a preferential strategy to rectify posterior R_0

estimates amid late genome sequence sampling. Furthermore, the entropy of each birth-

death based posterior R_0 distribution, a measure of uncertainty, is comparable at 3.68-3.78

as calculated with the mlf R package (Peterson, 2018). This further suggests that the

topological uncertainty induced by occurrence data does not considerably increase

uncertainty in posterior R_0 (Fig 3).

Discussion

Occurrence data in empirical phylodynamic studies

Occurrence data represent an extreme case of when genome coverage in samples is poor. Herein we show that low-coverage samples can be useful in phylodynamics so long as the sequences analysed are accurate. An outstanding task is to characterise an upper-bound on the relative proportion of occurrence to genomic samples from which genomic samples can still inform tree topology for epidemiological dynamics. To this end, we caution against over-inflating occurrence among genomic data sets without comparing to results obtained with genomic samples alone.

Our simulations and empirical data analyses reveal that occurrence data are a rich source of information for birth-death tree priors that can dramatically improve the accuracy and precision in estimates of epidemiological parameters. A key consideration is that occurrences should represent confirmed cases that would have been sequenced if sequencing effort had been constant, and which are known to belong to a particular outbreak, such as via contact tracing. Combining occurrence and sequence data can be particularly useful in situations where it is unknown if sequence sampling has been constant over time or where there exist several confirmed cases but a smaller number of sequences. This is valuable amid recently emerging outbreaks where combining both sources of data can provide sharper and more timely insight into the recent evolution of the pathogen in question.

Acknowledgements

We are grateful to Trevor Bedford, authors and groups, originating laboratories, and submitting laboratories of sequences downloaded from GISAID. We provide a full acknowledgement table in the supplementary information. SD and LAF were supported by a Discovery Early Career Fellowship from the Australian Research Council (DE190100805), awarded to SD. ECH is supported by an Australian Research Council Laureate Fellowship (FL170100022).

Authors' contributions:

All authors contributed to the design of experiments and writing of the manuscript. LF conducted analyses of empirical data and lead writing of the manuscript. FG contributed initial datasets, writing, and guidance with figures. TV contributed to writing the manuscript and mathematical concepts. EH contributed to writing of the manuscript and original ideas.

SD conceived of fundamental concepts in the manuscript, conducted simulations, and

contributed to writing.

Data availability:

Input files to generate trees in MASTER and to analyse sequence data in BEAST according

to the birth-death skyline, birth-death, and the coalescent exponential tree priors, and

accession numbers for empirical SARS-CoV-2 virus data. Available at:

github.com/sebastianduchene/birth-death-sampling. Accession numbers for empirical

SARS-CoV-2 virus data and the GISAID acknowledgements table are available as

supplementary data online.

References

Bedford, T., Greninger, A. L., Roychoudhury, P., Starita, L. M., Famulare, M., Huang, M.-L.,

Nalla, A., Pepper, G., Reinhardt, A., Xie, H., Shrestha, L., Nguyen, T. N., Adler, A.,

Brandstetter, E., Cho, S., Giroux, D., Han, P. D., Fay, K., Frazar, C. D., … Jerome, K.

R. (2020). *Cryptic transmission of SARS-CoV-2 in Washington State* [Preprint].

Epidemiology. https://doi.org/10.1101/2020.04.02.20051417

Biek, R., Pybus, O. G., Lloyd-Smith, J. O., & Didelot, X. (2015). Measurably evolving

pathogens in the genomic era. *Trends in Ecology and Evolution*, *30*(6), 306–313.

Boskova, V., Stadler, T., & Magnus, C. (2018). The influence of phylodynamic model

specifications on parameter estimates of the Zika virus epidemic. *Virus Evolution*,

4(1), vex044.

Bouckaert, R., Vaughan, T. G., Barido-Sottani, J., Duchêne, S., Fourment, M.,

Gavryushkina, A., Heled, J., Jones, G., Kühnert, D., Maio, N. D., Matschiner, M.,

Mendes, F. K., Müller, N. F., Ogilvie, H. A., Plessis, L. du, Popinga, A., Rambaut, A.,

Rasmussen, D., Siveroni, I., … Drummond, A. J. (2019). BEAST 2.5: An advanced

software platform for Bayesian evolutionary analysis. *PLOS Computational Biology*,

15(4), e1006650. https://doi.org/10.1371/journal.pcbi.1006650

Drummond, A. J., Nicholls, G. K., Rodrigo, A. G., & Solomon, W. (2002). Estimating

mutation parameters, population history and genealogy simultaneously from

temporally spaced sequence data. *Genetics*, *161*(3), 1307–1320.

Drummond, A. J., Pybus, O. G., Rambaut, A., Forsberg, R., & Rodrigo, A. G. (2003).

Measurably evolving populations. *Trends in Ecology & Evolution*, *18*(9), 481–488.

du Plessis, L., & Stadler, T. (2015). Getting to the root of epidemic spread with

phylodynamic analysis of genomic data. *Trends in Microbiology*, *23*(7), 383–386.

Duchene, S., Featherstone, L., Haritopoulou-Sinanidou, M., Rambaut, A., Lemey, P., &

Baele, G. (2020). Temporal signal and the phylodynamic threshold of SARS-CoV-2.

BioRxiv, 2020.05.04.077735. https://doi.org/10.1101/2020.05.04.077735

Gardy, J. L., & Loman, N. J. (2018). Towards a genomics-informed, real-time, global

pathogen surveillance system. *Nature Reviews Genetics*, *19*(1), 9.

Geoghegan, J. L., Ren, X., Storey, M., Hadfield, J., Jelley, L., Jefferies, S., Sherwood, J.,

Paine, S., Huang, S., Douglas, J., Mendes, F. K. L., Sporle, A., Baker, M. G.,

Murdoch, D. R., French, N., Simpson, C. R., Welch, D., Drummond, A. J., Holmes, E.

C., … de Ligt, J. (2020). *Genomic epidemiology reveals transmission patterns and*

dynamics of SARS-CoV-2 in Aotearoa New Zealand [Preprint]. Infectious Diseases

(except HIV/AIDS). https://doi.org/10.1101/2020.08.05.20168930

Griffiths, R. C., & Tavare, S. (1994). Sampling theory for neutral alleles in a varying

environment. *Philosophical Transactions of the Royal Society of London. Series B:*

Biological Sciences, *344*(1310), 403–410. https://doi.org/10.1098/rstb.1994.0079

Grubaugh, N. D., Ladner, J. T., Lemey, P., Pybus, O. G., Rambaut, A., Holmes, E. C., &

Andersen, K. G. (2019). Tracking virus outbreaks in the twenty-first century. *Nature*

Microbiology, *4*(1), 10.

Gupta, A., Manceau, M., Vaughan, T., Khammash, M., & Stadler, T. (2020). The probability

distribution of the reconstructed phylogenetic tree with occurrence data. *Journal of*

Theoretical Biology, *488*, 110115. https://doi.org/10.1016/j.jtbi.2019.110115

Hadfield, J., Megill, C., Bell, S. M., Huddleston, J., Potter, B., Callender, C., Sagulenko, P.,

Bedford, T., & Neher, R. A. (2018). Nextstrain: Real-time tracking of pathogen

evolution. *Bioinformatics*, *34*(23), 4121–4123.

Heath, T. A., Huelsenbeck, J. P., & Stadler, T. (2014). The fossilized birth–death process for

coherent calibration of divergence-time estimates. *Proceedings of the National*

Academy of Sciences, *111*(29), E2957–E2966.

Heath, T. A., & Moore, B. R. (2014). Bayesian inference of species divergence times. In M.-

H. Chen, L. Kuo, & P. O. Lewis (Eds.), *Bayesian Phylogenetics, Methods, Algorithms,*

and Applications (pp. 277–318). CRC Press.

Hedge, J., Lycett, S. J., & Rambaut, A. (2013). Real-time characterization of the molecular

epidemiology of an influenza pandemic. *Biology Letters*, *9*(5), 20130331.

Ho, S. S., Duchêne, S., & Duchêne, D. A. (2015). Simulating and detecting autocorrelation

of molecular evolutionary rates among lineages. *Molecular Ecology Resources*,

15(4), 688–696. https://doi.org/10.1111/1755-0998.12320

Kühnert, D., Stadler, T., Vaughan, T. G., & Drummond, A. J. (2014). Simultaneous

reconstruction of evolutionary history and epidemiological dynamics from viral

sequences with the birth–death SIR model. *Journal of The Royal Society Interface*,

11(94), 20131106. https://doi.org/10.1098/rsif.2013.1106

Manceau, M., Gupta, A., Vaughan, T., & Stadler, T. (2019). *The ancestral population size*

conditioned on the reconstructed phylogenetic tree with occurrence data [Preprint].

Evolutionary Biology. https://doi.org/10.1101/755561

Parag, K. V. (n.d.). *Jointly Inferring the Dynamics of Population Size and Sampling Intensity*

from Molecular Sequences. 16.

Peterson, K. (2018). *mlf: Machine Learning Foundations* (1.2.1) [Computer software].

https://CRAN.R-project.org/package=mlf

Popinga, A., Vaughan, T., Stadler, T., & Drummond, A. J. (2015). Inferring Epidemiological

Dynamics with Bayesian Coalescent Inference: The Merits of Deterministic and

Stochastic Models. *Genetics*, *199*(2), 595–607.

https://doi.org/10.1534/genetics.114.172791

Price, D. J., Shearer, F. M., Meehan, M. T., McBryde, E., Moss, R., Golding, N., Conway, E.

J., Dawson, P., Cromer, D., Wood, J., Abbott, S., McVernon, J., & McCaw, J. M.

(2020). *Early analysis of the Australian COVID-19 epidemic* [Preprint]. Epidemiology.

https://doi.org/10.1101/2020.04.25.20080127

Rambaut, A. (2000). Estimating the rate of molecular evolution: Incorporating non-

contemporaneous sequences into maximum likelihood phylogenies. *Bioinformatics*,

16(4), 395–399. https://doi.org/10.1093/bioinformatics/16.4.395

Rasmussen, D. A., Kouyos, R., Günthard, H. F., & Stadler, T. (2017). Phylodynamics on

local sexual contact networks. *PLOS Computational Biology*, *13*(3), e1005448.

https://doi.org/10.1371/journal.pcbi.1005448

Rieux, A., & Balloux, F. (2016). Inferences from tip-calibrated phylogenies: A review and a

practical guide. *Molecular Ecology*, *25*(9), 1911–1924.

Rife, B. D., Mavian, C., Chen, X., Ciccozzi, M., Salemi, M., Min, J., & Prosperi, M. C. (2017).

Phylodynamic applications in 21st century global infectious disease research.

Global Health Research and Policy, *2*(1), 13. https://doi.org/10.1186/s41256-017-

0034-y

Schliep, K. P. (2011). phangorn: Phylogenetic analysis in R. *Bioinformatics*, *27*(4), 592–593.

Stadler, T. (2010). Sampling-through-time in birth-death trees. *Journal of Theoretical*

Biology, *167*(3), 696–404.

Stadler, T., Kouyos, R., von Wyl, V., Yerly, S., Böni, J., Bürgisser, P., Klimkait, T., Joos, B.,

Rieder, P., & Xie, D. (2012). Estimating the basic reproductive number from viral

sequence data. *Molecular Biology and Evolution*, *29*(1), 347–357.

Stadler, T., Kühnert, D., Bonhoeffer, S., & Drummond, A. J. (2013). Birth–death skyline plot

reveals temporal changes of epidemic spread in HIV and hepatitis C virus (HCV).

Proceedings of the National Academy of Sciences, *110*(1), 228–233.

Stadler, T., Vaughan, T. G., Gavryushkin, A., Guindon, S., Kühnert, D., Leventhal, G. E., &

Drummond, A. J. (2015). How well can the exponential-growth coalescent

approximate constant-rate birth–death population dynamics? *Proceedings of the*

Royal Society B: Biological Sciences, *282*(1806), 20150420.

Stadler, T., & Yang, Z. (2013). Dating phylogenies with sequentially sampled tips.

Systematic Biology, *62*(5), 674–688.

Vasylyeva, T. I., du Plessis, L., Pineda-Peña, A. C., Kühnert, D., Lemey, P., Vandamme, A.-

M., Gomes, P., Camacho, R. J., Pybus, O. G., Abecasis, A. B., & Faria, N. R. (2019).

Tracing the Impact of Public Health Interventions on HIV-1 Transmission in Portugal

Using Molecular Epidemiology. *The Journal of Infectious Diseases*, *220*(2), 233–243.

https://doi.org/10.1093/infdis/jiz085

Vaughan, T. G., & Drummond, A. J. (2013). A stochastic simulator of birth–death master

equations with application to phylodynamics. *Molecular Biology and Evolution*,

30(6), 1480–1493.

Vaughan, T. G., Leventhal, G. E., Rasmussen, D. A., Drummond, A. J., Welch, D., & Stadler,

T. (2019). Estimating Epidemic Incidence and Prevalence from Genomic Data.

Molecular Biology and Evolution, *36*(8), 1804–1816.

https://doi.org/10.1093/molbev/msz106

Vaughan, T. G., Nadeau, S. A., Sciré, J., & Stadler, T. (2020, March 13). Phylodynamic

Analyses of outbreaks in China, Italy, Washington State (USA), and the Diamond

Princess. *Virological.Org*. https://virological.org/t/phylodynamic-analyses-of-

outbreaks-in-china-italy-washington-state-usa-and-the-diamond-princess/439

Volz, E. M., & Frost, S. D. W. (2014). Sampling through time and phylodynamic inference

with coalescent and birth–death models. *Journal of the Royal Society Interface*,

11(101). https://doi.org/10.1098/rsif.2014.0945

Volz, E. M., Koelle, K., & Bedford, T. (2013). Viral phylodynamics. *PLOS Computational*

Biology, *9*(3), e1002947.

Volz, E. M., Pond, S. L. K., Ward, M. J., Brown, A. J. L., & Frost, S. D. W. (2009).

Phylodynamics of infectious disease epidemics. *Genetics*, *183*(4), 1421–1430.

Volz, E. M., & Siveroni, I. (2018). Bayesian phylodynamic inference with complex models.

PLOS Computational Biology, *14*(11), e1006546.

https://doi.org/10.1371/journal.pcbi.1006546

Figure legends

Fig 1. Example of a phylogenetic trees generated under a birth-death process with a basic reproductive number (R_0) of 2, and a becoming uninfectious rate(δ) of 100 for three analysis scenarios. The solid line denotes the number of samples collected over time. In *constant sampling* samples are collected and sequenced at a rate ψ=5 (i.e. sampling probability, *p*, of 0.05). In *late sampling* samples are collected and sequenced after time T_s shown with the dashed line. In *occurrence data* samples are collected constantly over time, but only sequenced after time T_s, such that before T_s only occurrences (sampling times with no sequence data) are included. Blue circles represent samples with sequence data, whereas those in orange correspond to occurrences. In the *occurrence data* scenario, a Bayesian analysis would integrate over their phylogenetic uncertainty. The solid line represents the number of samples collected over time. In *late sampling* there are no samples collected before T_s, such that assuming constant sampling can produce a bias in estimates of epidemiological dynamics.

Fig 2. Bar ordering varies across plots to reflect preferential performance in each statistic such that those in red are most preferential. A) The number of simulations (out of 100) for which HPDs for R_0 captured 2, the value simulated under. B) Mean bias in R_0 across simulation treatments. C) Mean HPD width in R_0 across simulation treatments. D) The number of simulations (out of 100) for which HPDs for evolutionary rate captured 0.01, the value simulated under. E) Mean bias in Rate across simulation treatments. F) Mean HPD width in rate across simulation treatments.

Fig 3. Posterior estimates of R₀ for SARS-CoV-2 genome data. Constant sampling refers to using all 821 genomes in the empirical dataset. Post 2020.18 refers to only including sequences from 2020-03-04 and afterwards. Post 2020.2 refers to the same from 2020-03-14 and afterwards. A) Posterior densities of the basic reproductive number, R_0 under the coalescent exponential. B) Posterior densities for estimates of the basic reproductive number, R_0 under the birth death. In B, birth-death and birth-death skyline posteriors for R_0 and post cut-off sampling proportions are overlapping.

Tables

Table 1. Results of the simulation study with R_0 of 2 and evolutionary rate of 0.01 subs/site/year. The rows correspond to the seven treatments. For R_0 and evolutionary rate (subs/site/year), columns denote the number of simulations (out of 100) where the value

used to generate the data was contained within the 95% highest posterior density (HPD),
also referred to as coverage and reflecting accuracy; average bias measured the average
difference between posterior mean R_0 and 2; and the average HPD width. BD stands for
birth-death, CE for coalescent exponential, and BDSky to the birth-death skyline model
with two sampling intervals.

Treatment	R_0 Within HPD	R_0 Mean Bias	R_0 Mean HPD Width	Rate Within HPD	Rate Mean Bias	Rate Mean HPD Width
BD Constant Sampling	94	0.072	0.364	94	0.00057	0.00285
CE Constant Sampling	89	0.115	0.481	92	0.00067	0.00298
BD Late Sampling	12	0.364	0.515	58	0.00233	0.00511
BDSky Late Sampling	95	0.125	0.591	90	0.00122	0.00559
CE Late Sampling	100	0.156	0.786	97	0.00138	0.00646
BD + Occurrences	94	0.076	0.384	95	0.00097	0.00450
CE + Occurrences	96	0.163	0.748	91	0.00140	0.00605

Table 2. Posterior estimates of R_0 and p using the birth-death for the SARS-CoV-2
empirical dataset. Rows correspond to the 12 treatments.

Sampling Treatment	Mean R_0	95% HPD
BD Constant Sampling	1.96	(1.85, 2.07)
BD Post 2020.18	2.44	(2.31, 2.58)
BD Post 2020.18 + Occurrences	1.97	(1.87, 2.08)
BD Post 2020.2	3.53	(3.24, 3.82)
BD Post 2020.2 + Occurrences	1.96	(1.83, 2.09)
BDSky Post 2020.18	1.57	(1.43, 1.71)
BDSky Post 2020.2	1.48	(1.35, 1.63)
CE Constant Sampling	1.52	(1.4, 1.65)
CE Post 2020.18	1.51	(1.39, 1.64)
CE Post 2020.18 + Occurrences	1.50	(1.38, 1.62)
CE Post 2020.2	1.43	(1.3, 1.58)
CE Post 2020.2 + Occurrences	1.43	(1.29, 1.57)

Supplementary material

Table S1. Results of the simulation study with R_0=1.5, evolutionary rate of 0.01
subs/site/year, and late sampling starting at 0.9 years of a total time of 1 year. The rows

correspond to the seven treatments. The first two columns denote the number of
simulations (out of 100) where the value used to generate the data was contained within the
95% highest posterior density (HPD). The last two columns are a measure of precision of
the estimates calculated as the estimated mean estimate of R_0 and the evolutionary rate
divided by the 95% HPD width, such that large values imply low precision. Here we report
the mean value over 100 simulations.

	R_0 within 95% HPD	Evol. rate within 95% HPD	Mean R_0 / HPD width	Mean evol. rate / HPD width
Late sampling BD const.	0	37	0.20	0.39
Late sampling BD skyline	92	93	0.25	0.54
Late sampling Coal. exp.	80	87	0.27	0.59
Constant sampling BD const.	97	94	0.16	0.23
Constant sampling Coal. exp.	96	92	0.16	0.23
Birth Death + Occurrences.	92	86	0.16	0.35
Coalescent Exponential + Occurrences	66	69	0.23	0.43

Fig S1. The temporal distribution of SARS-CoV-2 samples taken from Washington State
and Washington County, Oregon, during the COVID-19 pandemic downloaded from

GISAID. Colouring represents the subset of these data that we analysed and vertical lines

show our two cut-offs for late sampling.

Fig S2. Posterior densities for estimates of the basic reproductive number, R_0, and the

evolutionary rate for 100 simulations with true R_0 of 1.5 and an evolutionary rate of 0.01

subs/site/year. The bars represent the 95% highest posterior density (HPD) and the points

are the mean. Estimates are ordered from lowest to highest mean. We analysed the data by

sampling late in the outbreak only (i.e. after 0.75 of the tree height), with a constant

sampling effort (with all samples sequenced), and by including occurrence data. The

colours represent four different tree priors; red for the coalescent exponential, blue for the

birth-death skyline, and orange for the birth-death with constant sampling. For the data

with sampling late in the outbreak only we use the birth-death skyline tree prior with

constant R_0 and two intervals for the sampling rate, ψ, before time 0.75. This tree priori not

applicable to analyses with complete sampling or with occurrence data where sampling is

constant. The dashed horizontal lines correspond to the true parameter value used to

generate the data.