NAVIP: Unraveling the Influence of Neighboring Small Sequence Variants on Functional Impact Prediction

Jan-Simon Baasner[1], Andreas Rempel[2,3], Dakota Howard[4] and Boas Pucker[1,5,*]

1 Genetics and Genomics of Plants, Faculty of Biology & Center for Biotechnology, Bielefeld University, 33615 Bielefeld, Germany

2 Genome Informatics, Faculty of Technology & Center for Biotechnology, Bielefeld University, 33615 Bielefeld, Germany

3 Graduate School "Digital Infrastructure for the Life Sciences" (DILS), Bielefeld Institute for Bioinformatics Infrastructure (BIBI), Bielefeld University, 33615 Bielefeld, Germany

4 Biology and Computer Science Department, Furman University, Greenville, South Carolina, USA

5 Plant Biotechnology and Bioinformatics, Institute of Plant Biology & BRICS, TU Braunschweig, 38106 Braunschweig, Germany

* Correspondence: b.pucker@tu-braunschweig.de

Abstract

Once a suitable reference sequence has been generated, intraspecific variation is often assessed by re-sequencing. Variant calling processes can reveal all differences between strains, accessions, genotypes, or individuals. These variants can be enriched with predictions about their functional implications based on available structural annotations, i.e. gene models. Although these functional impact predictions on a per-variant basis are often accurate, some challenging cases require the simultaneous incorporation of multiple adjacent variants into this prediction process. Examples include neighboring variants which modify each other's functional impact. The Neighborhood-Aware Variant Impact Predictor (NAVIP) considers all variants within a given protein coding sequence when predicting the functional consequences. As a proof of concept, variants between the *Arabidopsis thaliana* accessions Columbia-0 and Niederzenz-1 were annotated. NAVIP is freely available on GitHub (https://github.com/bpucker/NAVIP) and accessible through a web server (https://pbb-tools.de).

Author Summary

Intraspecific variation gains increasing relevance as reference genome sequences are available for many investigated (plant) species. Understanding the functional consequences of sequence variants between individuals of a population is a challenge. SnpEff (Cigolnai et al., 2012) is the current standard tool to predict the functional impact of sequence variants, but does only consider one sequence variant at the time. We developed NAVIP to properly handle cases in which multiple sequence variants are clustering together and influence each other's functional impact. A comparison of two *Arabidopsis thaliana* accessions demonstrates the relevance of considering multiple sequence variants simultaneously for the prediction of changes in encoded

proteins. NAVIP is universally applicable to non-model organisms. All underlying code is freely available from GitHub and we operate a web server for users' convenience.

Keywords: sequence variants, variant annotation, SNPs, SNVs, InDels, mutations

Introduction

Re-sequencing projects, examining many individuals or accessions of a species [1–3], are becoming increasingly important in plant research. Approaches similar to genome-wide association studies (GWAS) which are based on mapping-by-sequencing (MBS) are frequently applied in a wide range of crop species [4–7]. They are boosted by a rapidly increasing availability of high-quality reference genome sequences for crops [8–12], technological advances in long-read sequencing [13], and low sequencing costs [14,15]. *De novo* assemblies are still beneficial for the detection of large structural variants [16–21] and especially to reveal novel sequences [17,18,20,22], but the reliable detection of modifying single nucleotide variants (SNVs) can be achieved based on (short) read mappings. Nevertheless, long-read sequencing is also frequently used to explore the intraspecific diversity, as more sequence variants can be detected in previously inaccessible genomic regions [23,24].

Once identified, the annotation of sequence variants is performed by predicting their functional implications based on the available gene models (structural annotation). Leading tools such as ANNOVAR [25] and SnpEff [26] currently perform this prediction efficiently by focusing on a single variant at a time. An impact prediction facilitates the identification of targets for

post-GWAS analyses and can lead to the identification of small sequence variants that form the molecular basis of economically relevant phenotypic differences [6,27,28]. Although the effect prediction for single variants is computationally efficient and usually correct, there is a minority of challenging cases in which predictions based on a single variant alone cannot be accurate. Multiple InDels could either lead to frameshifts or they could compensate for each other's effect leaving the sequence with minimal modifications [29–31]. Two SNVs occurring in the same codon could lead to a different amino acid substitution compared to the apparent effects resulting from an isolated analysis of each of these SNVs.

Here we present a computational tool for accurately predicting the combined effect of phased variants on annotated coding sequences. The Neighborhood-Aware Variant Impact Predictor (NAVIP) was developed to investigate large variant data sets of plant re-sequencing projects, but is not limited to the annotation of variants in plants. As a proof of concept, NAVIP was used to identify cases between the *A. thaliana* accessions Columbia-0 (Col-0) and Niederzenz-1 (Nd-1) where an accurate impact prediction needs to consider multiple variants at a time.

Results

Features of NAVIP

NAVIP predicts the functional impact of sequence variants by considering all sequence variants affecting the coding sequence of a gene simultaneously. Users need to supply a set of sequence variants (VCF), a reference genome sequence (FASTA), and a structural annotation (GFF3). NAVIP returns an annotated VCF file and FASTA files with corrected coding and

polypeptide sequences. If phased sequence variants are provided in the VCF file, NAVIP

performs separate analyses for the different haplophases.

There are multiple installation options described in the GitHub repository

(https://github.com/bpucker/NAVIP) and NAVIP is also available free of charge through a web

server (https://pbb-tools.de/NAVIP). This makes NAVIP accessible to a wide range of users and

applicable to data sets of various sizes. Uploaded files are only used for the intended analysis

and deleted 48 hours after offering the results for download. The web server is able to send

notification emails upon completion of a job which can serve as documentation and facilitates

the analysis of large data sets.

Relevance of NAVIP for predicting premature stop codons

Running NAVIP on an *A. thaliana* Nd-1 data set with 644,261 SNVs (S1 File, S2 File) took about

5 minutes with a single core and a peak memory usage of about 3 GB RAM. To the best of our

knowledge, SnpEff is the most frequently used tool for the annotation of variants and also

universally applicable. Consequently, the NAVIP output was compared with the SnpEff

predictions produced for the same data set and structural annotation. The results are largely

congruent, but interesting cases for comparison are predictions of premature stop codons as

these have severe biological consequences. While a single SNV would cause a premature stop

codon, the simultaneous presence of two SNVs can result in an amino acid encoding codon

(**Figure 1a**). Of 600 premature stop codons predicted by SnpEff, 144 were identified as amino

acid substitutions when considering multiple SNVs in the same codon via NAVIP (**Figure 1b**).

Given the total of 600 predicted premature stop codons in this Nd-1 data set, 24% were false

positive predictions. NAVIP revealed that tyrosine occurs frequently instead of a premature stop

codon, because the tyrosine codons are very similar to two of the three stop codons. There are

also additional 17 premature stop codons predicted by NAVIP, which are the consequence of

two sequence variants influencing the same codon. Despite the surprisingly large difference between the SnpEff and NAVIP results when it comes to predicting premature stop codons, the differences in affected genes are smaller. Many genes with a predicted premature stop codon have multiple downstream premature stop codons. While an individual prediction of a premature stop codon might be wrong, the gene can still be correctly identified by both tools (S3 File). When a premature stop codon results in a loss-of-function event, the accumulation of additional variants is likely due to a lack of purifying selection. To support the assumption that genes with premature stop codons lost their function, the rate of amino acid changing variants in these genes was compared to all other genes (**Figure 1c, Figure 1d**). The number of variants changing amino acids (dN) to those resulting in the same amino acid (dS) were calculated for all genes. A significantly higher proportion of amino acid changing variants were observed in genes with predicted premature stop codons compared to all other genes (Mann-Whitney U test, p-value=10^{-167}). An additional comparison of the average expression of genes with a premature stop codon against all other protein encoding genes (**Figure 1e and f**) revealed also a significant difference (Mann-Whitney U text, p-value=10^{-70}).

Figure 1: (a) This illustration shows the concept of two SNVs affecting the same codon resulting in different prediction outcomes. (b) Second site variants within the same codon turn premature stop codons predicted by SnpEff into amino acid substitutions. In 144 cases, premature stop codons are substitutions by the displayed amino acids.(c) The proportion of amino acid changing variants is significantly higher in genes with premature stop codons predicted by NAVIP (blue) compared to all other genes (red). (d) The proportion of amino acid changing variants is significantly higher in genes with premature stop codons predicted by SnpEff (blue)

compared to all other genes (red). Data underlying these visualizations are available in S3 File. (e) Comparison of the average expression of genes with a premature stop codon predicted by NAVIP against all other protein encoding genes with available expression data. (f) Comparison of the average expression of genes with a premature stop codon predicted by SnpEff against all other protein encoding genes with available expression data.

Role of compensating InDels (cInDels)

InDels can compensate for each others' frameshift when occurring together (**Figure 2a**). While the first InDel can alter the reading frame, the second one could revert the reading frame back to the original one thus resulting in only a few altered codons enclosed by the two events. Since premature stop codons can emerge in the novel codons following the first frameshift, the distance between such InDels was is expected to be very small. An analysis of the distance distribution of the InDels between Nd-1 and Col-0 (S4 File) revealed that most compensating InDels (cInDels) occur within a very short distance of 2-8 bp (**Figure 2b**). Multiples of three are more frequent than other distances of a similar size, which might be connected to the length of codons.

Figure 2: (a) Theoretical concept of two InDels compensating each others' frameshift. The first insertion changes the reading frame, while the second insertion shifts the reading frame back to the original one. While each individual variant would suggest a loss-of-function due to a frameshift mutation, the combination of both results in 'only' two additional amino acids in the gene product. (b) Distribution of distances between compensating InDels (cInDels). As the second InDel can compensate for the frameshift caused by the upstream InDel, distances between such cInDels are short and frequently multiples of three. In total, 484 genes were identified to contain cInDels in the Nd-1 data set.

Discussion

This study demonstrates features of NAVIP by utilizing a previously generated set of high confidence sequence variants [35]. There is always a trade-off between sensitivity and specificity in the variant calling process [35,40] and the aim in this study was to obtain a high specificity data set (see S1 File for details). The benchmarking of NAVIP through comparison with SnpEff neutralizes the influence of the sequence variant data set quality on the outcome.

As an additional validation of the outcome, NAVIP results were analyzed for additional amino acid substitutions in genes with premature stop codons. The frequency of such variants was higher in genes with premature stop codons compared to others suggesting a lack of purification selection in these genes which could point to pseudogenization. The comparison against all other genes also clearly revealed the increased frequency of amino acid substitutions in genes with premature stop codons. Additionally, a low expression of genes with premature stop codons compared to other genes suggests a pseudogenization. In summary, the properties observed for genes with premature stop codons match the expectations thus supporting the biological validity of the data set.

We developed NAVIP to simultaneously assess the impact of all neighboring sequence variants in protein encoding sequences and to be universally applicable. The described cases in the comparison of two *A. thaliana* accessions demonstrate the necessity to have such a tool at hand. NAVIP revealed the presence of second site mutations that compensate for other variants e.g. turning a presumed premature stop codon into an amino acid substitution or vice versa. Another example are frameshifts resulting from InDels that are compensated by downstream InDels, which shift the reading frame back to the original pattern. Neglecting these interactions of sequence variants during the functional impact prediction can lead to mis-annotation. While NAVIP was developed to accurately predict changes in the polypeptide sequence based on DNA sequence variants, downstream tools are needed to predict consequences of these changes on the function of proteins. Tools like SIFT [45], PolyPhen-2 [49], or SNAP2 [50] could be applied for this next step.Many computational tools for the assessment of DNA sequence variant impact focus on human data sets [41–44]. The objective is often to identify pathogenic variants [45,46]. Universally applicable tools like SnpEff [26], which are also suitable to analyze plant data sets, predict the impact of isolated sequence variants. The purpose of NAVIP is to offer novel functionalities to the plant science community and other communities working on

non-model organisms. NAVIP could boost the power of re-sequencing studies by opening up the field of compensating or in general mutually influencing variants. Such variants have the potential to reveal new insights into patterns of molecular evolution and especially co-evolution of sites. The consideration of multiple variants during the effect prediction could reveal novel targets in GWAS-like approaches. The availability through a web server enables a large community of scientists without computational skills to benefit from NAVIP.

The remaining challenge is now the reliable detection of sequence variants prior to the application of NAVIP. A range of tools is available for the mapping of short reads and the following identification of sequence variants [35]. There is also rapid progress in the development of long read mapping tools [51,52] and the following variant identification [53–56]. For heterozygous and polyploid species, phasing of these variants is another task that needs to be addressed in the future. The correct prediction of functional implications relies on the correct assignment of variants to respective haplophases. If provided with accurately phased variants, NAVIP can perform predictions for highly heterozygous and even polyploid species. Previous studies demonstrated that sequence variants might only affect individual isoforms in a negative way [46]. NAVIP analyzes all annotated transcript isoforms and would be able to discover such cases. Currently, a major limitation is the lack of isoform-resolved annotation for non-model plant species. Given the rapid progress in long read sequencing [13,47,48], it is likely that highly accurate structural annotation will become available for most plant species in the next few years.

Materials and Methods

Implementation of the Neighborhood-Aware Variant Impact Predictor (NAVIP)

The Neighborhood-Aware Variant Impact Predictor (NAVIP) (https://github.com/bpucker/NAVIP) is implemented in Python3. NAVIP requires a VCF file containing sequence variants, a FASTA file containing the reference sequence, and a GFF3 file containing the structural annotation (gene models) as input. The variants provided must be homozygous or in a phased state to allow an accurate impact prediction per allele. If no information about the phasing is provided, all variants are assumed to be in the same haplophase. Effects on all annotated transcripts are evaluated per gene by taking into account the presence of all given variants simultaneously. NAVIP generates a new VCF file with an additional annotation field and additional report files. One annotation string in the VCF output file matches the SnpEff result format, but also has a NAVIP-specific string with additional information (see the manual for details: https://github.com/bpucker/NAVIP/wiki). NAVIP also produces FASTA files with sequences harboring all variants. An automatic assessment of complementing InDels (cInDels) reveals the relevance of simultaneously considering all InDels within a coding sequence when predicting their impact. All NAVIP scripts can be downloaded from the above-mentioned GitHub repository and do not require the installation of any dependencies other than the Python packages. NAVIP is also available through a web server (https://pbb-tools.de/NAVIP) free of charge. Files are kept confidential and will be deleted 48 h after offering the results for download.

Identification and validation of sequence variants

Illumina sequencing reads of *A. thaliana* Nd-1 [16] were mapped to the *A. thaliana* Col-0 reference genome sequence (TAIR10) [32] via BWA MEM v.0.7.13 [33] using the –m option to avoid spurious hits. Variant calling was performed via GATK v3.8 [34] based on the developers' recommendation. This combination of BWA MEM and GATK was previously identified as a reliable approach for this particular data set [35]. All processes were wrapped into Python scripts (https://github.com/bpucker/variant_calling) to facilitate automatic execution on a high-performance compute cluster. An initial variant set was generated based on hard filtering criteria recommended by the GATK developers. The two following variant calling runs considered the set of surviving variants from the previous round as the gold standard to avoid the need for hard filtering.

Since a high-quality genome sequence assembly of Nd-1 was previously generated [17], we harnessed this sequence to validate all variants identified by short-read mapping. Starting at the north end of each chromosome sequence, variants sorted by genomic position were successively tested by taking the upstream sequence from Col-0, modifying it according to all upstream *bona fide* variants, and searching for it in the Nd-1 assembly (S5 File). Variants were admitted to the following analysis if the assembly supported them. This consecutive inspection of all variants enabled a reliable removal of false positives, leading to a set of high-confidence variants. The genome-wide distribution of the sequence variants was assessed using a previously developed Python script [16].

An independent confirmation of randomly selected sequence variants was performed using Sanger sequencing. *A. thaliana* Nd-1 plants were grown as previously described [16] to extract DNA from leaf tissue using a cetyltrimethylammonium bromide (CTAB)-based method [36]. Oligonucleotides flanking the regions that harbor the variants of interest were designed manually (S6 File). Amplification via PCR, analysis of PCR products via agarose gel

electrophoresis, purification of PCR products, Sanger sequencing, and evaluation of results were following previously established protocols [37].

Comparison of NAVIP and SnpEff stop gain prediction

To the best of our knowledge, SnpEff [26] is the most widely used tool for predicting the functional consequences of sequence variants, thus it was selected for comparison. NAVIP can only provide more accurate predictions about functional consequences if multiple sequence variants interfere, e.g. if multiple SNVs are located within the same codon. Otherwise, the predictions of NAVIP and SnpEff would be the same. Consequently, the following comparison focuses only on cases of multiple sequence variants that might interfere with each other.

SnpEff v4.1f [26] was applied with default parameters to the *A. thaliana* Nd-1 variant data set to predict the effects of SNVs based on the Araport11 [38] structural annotation of the TAIR10 genome sequence of *A. thaliana* Columbia-0. NAVIP was also applied to the same data set for benchmarking. Predictions of premature stop codons were compared between NAVIP and SnpEff results, as these cases have the potential to show biologically important differences. This analysis was performed exclusively on SNVs to avoid the influence of frameshifts that would be caused by InDels. Only the most upstream predicted premature stop codon within any gene was considered in this analysis. To support the loss of function of the affected genes, the frequency of amino acid changing variants (dN) was compared to the number of variants that did not alter the encoded amino acid (dS). This ratio was compared between genes with premature stop codons and all other genes, expecting a higher ratio of variants that change the encoded amino acids if the gene undergoes pseudogenization. The Python package plotly was used to visualize these data distributions in violin plots. If no dS events were discovered in an analyzed gene, a pseudocount was added to both dN and dS to enable the ratio calculation. dN/dS ratios greater than 10 were set to this maximum value to enable visualization. A Mann-Withney U test was

performed using Python to test for significant differences between the two groups. When genes with a premature stop codon undergo pseudogenization, they may show lower than average gene expression. Therefore, a comparison of the expression of genes with a premature stop codon against all other protein-coding genes was performed. A previously compiled count table based on all publicly available paired-end RNA-seq data sets of *A. thaliana* [39] was harnessed for this analysis. Differences were visualized using the Python package plotly as described above, with the expression values clipped at 50 to enable an informative visualization. All Python scripts developed for these analyses are freely available on GitHub (https://github.com/bpucker/variant_calling).

Assessment of compensating InDels (cInDels)

An independent analysis of insertions/deletions (InDels) was performed by NAVIP to understand the relevance of considering all InDels within a CDS simultaneously. Transcripts with predicted frameshifts were analyzed to identify downstream insertions/deletions which are compensating each other's effect, i.e. the second frameshift is reverting an upstream frameshift. The distance between these events was analyzed by NAVIP and is included in the standard output. This analysis is not restricted to pairs of cInDels, but can also handle multiple InDels compensating each other's frameshifts.

Availability and requirements

Project name: NAVIP

Project home page: https://github.com/bpucker/NAVIP

Operating system(s): Linux (website is platform independent)

Programming language: Python3

Other requirements: Python3

License: GNU General Public License v3.0

RRID: SCR_024838

Data availability

The data sets supporting the results of this article are publicly available or included within the article and its additional files. Python scripts developed and applied for this study are available on GitHub: https://github.com/bpucker/NAVIP (https://doi.org/10.5281/zenodo.10613052) and https://github.com/bpucker/variant_calling (https://doi.org/10.5281/zenodo.10613055).

Declarations

Authors' contributions

BP designed research. JSB wrote the NAVIP code. AR updated the NAVIP code and added NAVIP to the pbb-tools web server. JSB, DH, and BP conducted bioinformatic analyses. DH and BP performed experimental validation. JSB, AR, and BP wrote the manuscript. All authors read and approved the final manuscript version and agreed to its submission.

Financial Disclosure Statement

The authors received no specific funding for this work.

Competing Interests

JSB, AR, and DH have no competing interests. BP is head of the technology transfer center Plant Genomics and Applied Bioinformatics at iTUBS. This does not alter our adherence to PLOS policies on sharing data and materials.

Acknowledgments

We acknowledge support by members of Genetics and Genomics of Plants, Bioinformatics Resource Facility, and Sequencing Core Facility at the Center of Biotechnology. We thank Hanna Schilbert for critical reading of the manuscript. We thank the Center for Biotechnology (CeBiTec) at Bielefeld University for providing an environment to perform the computational analyses. Many thanks to the German network for bioinformatics infrastructure (de.NBI, grant 031A533A) and the Bioinformatics Resource Facility (BRF) at the Center for Biotechnology (CeBiTec) at Bielefeld University for providing an environment to perform the computational analyses. bioRender.com was used to construct some of the figures. We acknowledge support by the Open Access Publication Funds of Technische Universität Braunschweig.

References

1.   Alonso-Blanco C, Andrade J, Becker C, Bemm F, Bergelson J, Borgwardt KM, et al. 1,135 Genomes Reveal the Global Pattern of Polymorphism in Arabidopsis thaliana. Cell. 2016;166: 481–491. doi:10.1016/j.cell.2016.05.063

2.   Duan N, Bai Y, Sun H, Wang N, Ma Y, Li M, et al. Genome re-sequencing reveals the history of apple and supports a two-stage model for fruit enlargement. Nat Commun. 2017;8: 249. doi:10.1038/s41467-017-00336-7

3. Lobaton JD, Miller T, Gil J, Ariza D, de la Hoz JF, Soler A, et al. Resequencing of Common Bean Identifies Regions of Inter–Gene Pool Introgression and Provides Comprehensive Resources for Molecular Breeding. Plant Genome. 2018;11: 170068. doi:10.3835/plantgenome2017.08.0068

4. James GV, Patel V, Nordström KJV, Klasen JR, Salomé PA, Weigel D, et al. User guide for mapping-by-sequencing in Arabidopsis. Genome Biol. 2013;14: R61. doi:10.1186/gb-2013-14-6-r61

5. Mascher M, Jost M, Kuon J-E, Himmelbach A, Aßfalg A, Beier S, et al. Mapping-by-sequencing accelerates forward genetics in barley. Genome Biol. 2014;15: R78. doi:10.1186/gb-2014-15-6-r78

6. Schilbert HM, Pucker B, Ries D, Viehöver P, Micic Z, Dreyer F, et al. Mapping-by-Sequencing Reveals Genomic Regions Associated with Seed Quality Parameters in Brassica napus. Genes. 2022;13: 1131. doi:10.3390/genes13071131

7. Sielemann K, Pucker B, Orsini E, Elashry A, Schulte L, Viehöver P, et al. Genomic characterization of a nematode tolerance locus in sugar beet. BMC Genomics. 2023;24: 748. doi:10.1186/s12864-023-09823-2

8. Dohm JC, Minoche AE, Holtgräwe D, Capella-Gutiérrez S, Zakrzewski F, Tafer H, et al. The genome of the recently domesticated crop plant sugar beet (Beta vulgaris). Nature. 2014;505: 546–549. doi:10.1038/nature12817

9. Jarvis DE, Ho YS, Lightfoot DJ, Schmöckel SM, Li B, Borm TJA, et al. The genome of Chenopodium quinoa. Nature. 2017;542: 307–312. doi:10.1038/nature21370

10. Lightfoot DJ, Jarvis DE, Ramaraj T, Lee R, Jellen EN, Maughan PJ. Single-molecule sequencing and Hi-C-based proximity-guided assembly of amaranth (Amaranthus hypochondriacus) chromosomes provide insights into genome evolution. BMC Biol. 2017;15: 74. doi:10.1186/s12915-017-0412-4

11. Siadjeu C, Pucker B, Viehöver P, Albach DC, Weisshaar B. High Contiguity de novo

Genome Sequence Assembly of Trifoliate Yam (Dioscorea dumetorum) Using Long Read
Sequencing. Genes. 2020;11: 274. doi:10.3390/genes11030274

12. Marks RA, Hotaling S, Frandsen PB, VanBuren R. Representation and participation across
    20 years of plant genome sequencing. Nat Plants. 2021;7: 1571–1578.
    doi:10.1038/s41477-021-01031-8

13. Pucker B, Irisarri I, Vries J de, Xu B. Plant genome sequence assembly in the era of long
    reads: Progress, challenges and future directions. Quant Plant Biol. 2022;3: e5.
    doi:10.1017/qpb.2021.18

14. Stein LD. The case for cloud computing in genome informatics. Genome Biol. 2010;11:
    207. doi:10.1186/gb-2010-11-5-207

15. Christensen KD, Dukhovny D, Siebert U, Green RC. Assessing the Costs and
    Cost-Effectiveness of Genomic Sequencing. J Pers Med. 2015;5: 470–486.
    doi:10.3390/jpm5040470

16. Pucker B, Holtgräwe D, Sörensen TR, Stracke R, Viehöver P, Weisshaar B. A De Novo
    Genome Sequence Assembly of the Arabidopsis thaliana Accession Niederzenz-1
    Displays Presence/Absence Variation and Strong Synteny. PLOS ONE. 2016;11:
    e0164321. doi:10.1371/journal.pone.0164321

17. Pucker B, Holtgräwe D, Stadermann KB, Frey K, Huettel B, Reinhardt R, et al. A
    chromosome-level sequence assembly reveals the structure of the Arabidopsis thaliana
    Nd-1 genome and its gene set. PLOS ONE. 2019;14: e0216233.
    doi:10.1371/journal.pone.0216233

18. Zapata L, Ding J, Willing E-M, Hartwig B, Bezdan D, Jiao W-B, et al. Chromosome-level
    assembly of Arabidopsis thaliana Ler reveals the extent of translocation and inversion
    polymorphisms. Proc Natl Acad Sci. 2016;113: E4052–E4060.
    doi:10.1073/pnas.1607532113

19. Fan X, Chaisson M, Nakhleh L, Chen K. HySA: a Hybrid Structural variant Assembly

approach using next-generation and single-molecule sequencing technologies. Genome Res. 2017;27: 793–800. doi:10.1101/gr.214767.116

20. Michael TP, Jupe F, Bemm F, Motley ST, Sandoval JP, Lanz C, et al. High contiguity Arabidopsis thaliana genome assembly with a single nanopore flow cell. Nat Commun. 2018;9: 541. doi:10.1038/s41467-018-03016-2

21. Wala JA, Bandopadhayay P, Greenwald NF, O'Rourke R, Sharpe T, Stewart C, et al. SvABA: genome-wide detection of structural variants and indels by local assembly. Genome Res. 2018;28: 581–591. doi:10.1101/gr.221028.117

22. Zhou Y, Massonnet M, Sanjak JS, Cantu D, Gaut BS. Evolutionary genomics of grape (Vitis vinifera ssp. vinifera) domestication. Proc Natl Acad Sci U S A. 2017;114: 11715–11720. doi:10.1073/pnas.1709257114

23. De Coster W, Weissensteiner MH, Sedlazeck FJ. Towards population-scale long-read sequencing. Nat Rev Genet. 2021;22: 572–587. doi:10.1038/s41576-021-00367-3

24. Olson ND, Wagner J, McDaniel J, Stephens SH, Westreich ST, Prasanna AG, et al. PrecisionFDA Truth Challenge V2: Calling variants from short and long reads in difficult-to-map regions. Cell Genomics. 2022;2: 100129. doi:10.1016/j.xgen.2022.100129

25. Wang K, Li M, Hakonarson H. ANNOVAR: functional annotation of genetic variants from high-throughput sequencing data. Nucleic Acids Res. 2010;38: e164. doi:10.1093/nar/gkq603

26. Cingolani P, Platts A, Wang LL, Coon M, Nguyen T, Wang L, et al. A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff. Fly (Austin). 2012;6: 80–92. doi:10.4161/fly.19695

27. Hou L, Zhao H. A review of post-GWAS prioritization approaches. Front Genet. 2013;4: 280. doi:10.3389/fgene.2013.00280

28. Ries D, Holtgräwe D, Viehöver P, Weisshaar B. Rapid gene identification in sugar beet using deep sequencing of DNA from phenotypic pools selected from breeding panels.

BMC Genomics. 2016;17: 236. doi:10.1186/s12864-016-2566-9

29. Schneeberger K, Ossowski S, Ott F, Klein JD, Wang X, Lanz C, et al. Reference-guided assembly of four diverse Arabidopsis thaliana genomes. Proc Natl Acad Sci U S A. 2011;108: 10249–10254. doi:10.1073/pnas.1107739108

30. Cao J, Schneeberger K, Ossowski S, Günther T, Bender S, Fitz J, et al. Whole-genome sequencing of multiple Arabidopsis thaliana populations. Nat Genet. 2011;43: 956–963. doi:10.1038/ng.911

31. Gan X, Stegle O, Behr J, Steffen JG, Drewe P, Hildebrand KL, et al. Multiple reference genomes and transcriptomes for Arabidopsis thaliana. Nature. 2011;477: 419–423. doi:10.1038/nature10414

32. Lamesch P, Berardini TZ, Li D, Swarbreck D, Wilks C, Sasidharan R, et al. The Arabidopsis Information Resource (TAIR): improved gene annotation and new tools. Nucleic Acids Res. 2012;40: D1202–D1210. doi:10.1093/nar/gkr1090

33. Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. ArXiv13033997 Q-Bio. 2013 [cited 20 Oct 2020]. Available: http://arxiv.org/abs/1303.3997

34. Van der Auwera GA, Carneiro MO, Hartl C, Poplin R, del Angel G, Levy-Moonshine A, et al. From FastQ data to high confidence variant calls: the Genome Analysis Toolkit best practices pipeline. Curr Protoc Bioinforma Ed Board Andreas Baxevanis Al. 2013;11: 11.10.1-11.10.33. doi:10.1002/0471250953.bi1110s43

35. Schilbert HM, Rempel A, Pucker B. Comparison of Read Mapping and Variant Calling Tools for the Analysis of Plant NGS Data. Plants. 2020;9: 439. doi:10.3390/plants9040439

36. Rosso MG, Li Y, Strizhov N, Reiss B, Dekker K, Weisshaar B. An Arabidopsis thaliana T-DNA mutagenized population (GABI-Kat) for flanking sequence tag-based reverse genetics. Plant Mol Biol. 2003;53: 247–259. doi:10.1023/B:PLAN.0000009297.37235.4a

37. Pucker B, Holtgräwe D, Weisshaar B. Consideration of non-canonical splice sites improves gene prediction on the Arabidopsis thaliana Niederzenz-1 genome sequence. BMC Res

Notes. 2017;10: 667. doi:10.1186/s13104-017-2985-y

38. Cheng C-Y, Krishnakumar V, Chan AP, Thibaud‑Nissen F, Schobel S, Town CD. Araport11: a complete reannotation of the Arabidopsis thaliana reference genome. Plant J. 2017;89: 789–804. doi:10.1111/tpj.13415

39. Choudhary N, Pucker B. Conserved amino acid residues and gene expression patterns associated with the substrate preferences of the competing enzymes FLS and DFR. bioRxiv; 2023. p. 2023.11.05.565693. doi:10.1101/2023.11.05.565693

40. Olson ND, Lund SP, Colman RE, Foster JT, Sahl JW, Schupp JM, et al. Best practices for evaluating single nucleotide variant calling methods for microbial genomics. Front Genet. 2015;6: 235. doi:10.3389/fgene.2015.00235

41. Holcomb D, Hamasaki-Katagiri N, Laurie K, Katneni U, Kames J, Alexaki A, et al. New approaches to predict the effect of co-occurring variants on protein characteristics. Am J Hum Genet. 2021;108: 1502–1511. doi:10.1016/j.ajhg.2021.06.011

42. Liu Y, Yeung WSB, Chiu PCN, Cao D. Computational approaches for predicting variant impact: An overview from resources, principles to applications. Front Genet. 2022;13. Available: https://www.frontiersin.org/articles/10.3389/fgene.2022.981005

43. Wang D, Li J, Wang Y, Wang E. A comparison on predicting functional impact of genomic variants. NAR Genomics Bioinforma. 2022;4: lqab122. doi:10.1093/nargab/lqab122

44. Katsonis P, Wilhelm K, Williams A, Lichtarge O. Genome interpretation using in silico predictors of variant impact. Hum Genet. 2022;141: 1549–1577. doi:10.1007/s00439-022-02457-6

45. Ng PC, Henikoff S. SIFT: predicting amino acid changes that affect protein function. Nucleic Acids Res. 2003;31: 3812–3814. doi:10.1093/nar/gkg509

46. Brandes N, Goldman G, Wang CH, Ye CJ, Ntranos V. Genome-wide prediction of disease variant effects with a deep protein language model. Nat Genet. 2023;55: 1512–1522. doi:10.1038/s41588-023-01465-0

47. Marx V. Method of the year: long-read sequencing. Nat Methods. 2023;20: 6–11. doi:10.1038/s41592-022-01730-w

48. Al-Dossary O, Furtado A, KharabianMasouleh A, Alsubaie B, Al-Mssallem I, Henry RJ. Long read sequencing to reveal the full complexity of a plant transcriptome by targeting both standard and long workflows. Plant Methods. 2023;19: 112. doi:10.1186/s13007-023-01091-1

49. Adzhubei IA, Schmidt S, Peshkin L, Ramensky VE, Gerasimova A, Bork P, et al. A method and server for predicting damaging missense mutations. Nat Methods. 2010;7: 248–249. doi:10.1038/nmeth0410-248

50. Hecht M, Bromberg Y, Rost B. Better prediction of functional effects for sequence variants. BMC Genomics. 2015;16: S1. doi:10.1186/1471-2164-16-S8-S1

51. Amarasinghe SL, Ritchie ME, Gouil Q. long-read-tools.org: an interactive catalogue of analysis methods for long-read sequencing data. GigaScience. 2021;10: giab003. doi:10.1093/gigascience/giab003

52. Sahlin K, Baudeau T, Cazaux B, Marchet C. A survey of mapping algorithms in the long-reads era. Genome Biol. 2023;24: 133. doi:10.1186/s13059-023-02972-3

53. Ahsan MU, Liu Q, Fang L, Wang K. NanoCaller for accurate detection of SNPs and indels in difficult-to-map regions from long-read sequencing by haplotype-aware deep neural networks. Genome Biol. 2021;22: 261. doi:10.1186/s13059-021-02472-2

54. Shafin K, Pesout T, Chang P-C, Nattestad M, Kolesnikov A, Goel S, et al. Haplotype-aware variant calling with PEPPER-Margin-DeepVariant enables high accuracy in nanopore long-reads. Nat Methods. 2021;18: 1322–1332. doi:10.1038/s41592-021-01299-w

55. Cleal K, Baird DM. Dysgu: efficient structural variant calling using short or long reads. Nucleic Acids Res. 2022;50: e53. doi:10.1093/nar/gkac039

56. Huang N, Xu M, Nie F, Ni P, Xiao C-L, Luo F, et al. NanoSNP: a progressive and haplotype-aware SNP caller on low-coverage nanopore sequencing data. Bioinformatics.

Supporting Information

S1 File: Detailed description of the variant calling process, the validation process, and the resulting sequence variant data set.

S2 File: VCF file containing SNVs between Nd-1 and Col-0.

S3 File: Detailed information about premature stop codons predicted by NAVIP and/or SnpEff.

S4 File: VCF file containing InDels between Nd-1 and Col-0.

S5 File: Schematic illustration of the variant validation process.

S6 File: FASTA file containing oligonucleotide sequences used for the generation and sequencing of amplicons to validate randomly selected sequence variants.