Title

Comparison of prokaryotic communities between the fields showing different

disinfestation effects by anaerobic soil disinfestation

Authors

Chol Gyu Lee[1,2], Toshiya Iida[1], Eriko Matsuda[3], Kayo Yoshida[3], Masato Kawabe[4],

Masayuki Maeda[5], Yasunori Muramoto[6], Hideki Watanabe[6], Yoko Otani[7], Kazhiro

Nakaho[8], Moriya Ohkuma[1]

Affiliations

[1]*Japan Collection of Microorganisms, RIKEN BioResource Center, Tsukuba, Ibaraki*

305-0074, Japan.

[2]*Graduate School of Bio-Applications and Systems Engineering, Tokyo University of*

Agriculture and Technology, Koganei, Tokyo, 184-8588, Japan

[3]*Ishikawa Agriculture and Forestry Research Center, Kanazawa, Ishikawa 920-3198,*

Japan.

⁴Horticultural Research Institute, Toyama Prefectural Agricultural, Forestry and

Fisheries Research Center, Tonami, Toyama 939-1327, Japan.

⁵Niigata Agricultural Research Institute, Niigata, Nagaoka, Niigata 940-0826, Japan.

⁶Gifu Prefectural Agricultural Technology Center, Matamaru, Gifu 501-1152, Japan.

⁷Wakayama Agricultural Experiment Station, Kinokawa, Wakayama, 640-0423, Japan.

⁸Institute of Vegetable and Floriculture Science, National Agriculture and Food

Research Organization, Tsu, Mie 514-2392, Japan.

Abbreviations

ASD: Anaerobic soil disinfestation, OTU: Operation taxonomic units, PCR: Polymerase

chain reaction,

Keywords

Bacterial wilt, Betaproteobacteria, Clostridia, Indicator species analysis, Multiple

fields

Abstract

Soil-borne pathogens are serious problems for crop production. Chemical-dependent

soil fumigation method has been an effective method for controlling soil-borne pathogens,

but it is escaped for food safety and environmental pollution. Anaerobic soil

disinfestation (ASD) is a chemical-independent fumigation method and can reduce

pathogens under 40 cm soil depth. However, their effect is not stable under the field

condition. Soil microbes have important roles for ASD. Microbial community reflects

the soil environment and some unique microbes are be detected under different

condition. If common microbes are shown in well-disinfested multiple fields, they will

be a good indicator for the success of ASD. For this purpose, we conducted the same

ASD treatment with 17 tomato greenhouses where tomato bacterial wilt occurred in the

geographically different areas. In this study, we compared soil prokaryotic communities

in the field which indicate different disinfestation effects after ASD treatment among

several fields using the next-generation sequencer. Weighted-UniFrac analysis showed

that prokaryotic communities in the fields showing different disinfestation effect were

not different and they were roughly separated to sampling fields. The relative abundance

of *Betaproteobacteria* and *Clostridia* were significantly increased in well- disinfested

fields. Indicator species analysis revealed that twenty-five operational taxonomic units

(OTUs) were specifically detected from various well-disinfested soil and 18 OTUs

belonged to phylogenetically diversified *Clostridia*. Other OTUs belonged to aerobic

bacteria and were not previously detected from ASD treated fields. Our results showed

that the prokaryotic communities were not largely differed by ASD efficiency but small

changing will be important for ASD.

1. Introduction

Soil-borne pathogens cause various plant diseases, including take-all and damping-off, crown rots, and wilting. Bacterial wilt caused by *Ralstonia solanacearum* has a host range exceeding 200 species and over 50 families (Kelman 1998). Disinfestation of this pathogen is difficult because the pathogen is distributed evenly in a deep layer soil below 40 cm (Graham and Lloyd 1979). Several approaches have been attempted for bacterial wilt control, including soil amendments, crop rotation, and field sanitation (Wang and Lin 2005). Soil fumigation with chemical pesticides is an effective method for killing the bacterial wilt pathogens but its efficacy tends to be unstable in deep soil and it is escaped for food safety and environmental pollution.

Anaerobic soil disinfestation (ASD) is an effective method to reduce soil-borne pathogens (Shinmura 2000). This method consists of incorporation of labile organic matter in the soil, irrigation, and covering soil surface with polyethylene film. Organic matter increases microbial respiration and irrigation purges soil air, polyethylene film prevents an inflow of oxygen from the atmosphere, then reductive soil condition is induced (Block et al. 2000; Momma et al. 2006). It is effective for the soil in the lower

layer under 40 cm, which uses water-soluble organic such as low concentration ethanol

or molasses as a carbon source (Momma et al. 2013). Therefore, ASD using

water-soluble carbon sources is useful for the disinfection of *R. solanacearum* in

deep-layer soils and is environmentally-friendly. However, the disinfestation effects by

ASD is unstable under field condition (Shrestha et al. 2016). Enough soil temperature,

incubation period, and amount of carbon amendments are needed to success of

disinfestation.

Soil microbes have important roles for ASD. Several microbes increase after the ASD

treatment and they may be involved in suppression of pathogens (Momma et al. 2007;

Mowlick et al. 2013, Huang et al. 2015, 2016; Strauss et al. 2017; Mazzola et al. 2018).

Soil microbes reflect soil environment and they are considered as one of the indicators

for soil healthiness (van Burggen and Semenov 2000; Mendes et al. 2011). Therefore,

different disinfestation efficiency fields can have the different soil microbial

communities. Recently, Mazzola et al. (2018) suggested that microbial community

changing by ASD is different among sampling fields. If the microbes which is increased

in well-disinfested soil are commonly detected among several fields, they will be a good

indicator for ASD treatment. For this purpose, we conducted the ASD treatment with

same substrate in 17 geographically different greenhouses and they showed different

disinfestation efficiency. We compared the prokaryotic communities among the fields

showing different disinfestation effect. This multi-fields study will show that 1)

microbial community may differ depending on disinfestation efficiency and 2) the

well-disinfested soil may have unique microbes compare with not-well-disinfested soil

among various fields.

2. Materials and Methods

2.1. Sampling field and ASD treatment

Field experiments were established in 17 greenhouses in 8 fields in Japan in which

occurrences of bacterial wilt were observed (Table 1). Sugar-containing diatoms,

defined as a sugar-rich byproduct, are a waste discharged from the food-processing

industry that 40 % sugar in weight. The sugar-containing diatoms were mixed into the

soil at a ratio of 15 t ha^{-1} with the rototiller in 30 cm soil layer. The field was covered by

polyethylene and flooded with water more about 150 L m^{-2}. Disinfestation was

conducted 21 days excepted for Ha field (17 days), because soil temperature in this field

were reached more than 35 °C during disinfestation period. ASD were treated in the

greenhouse (15 m × 6 m) and replicates were conducted among different greenhouses.

There were three replications in the Ha, Ni, Ts, and To fields, two replications in the Sa

field, and no replication in the IS, Gi, and Wa fields. The samples were collected from

each greenhouse before and after ASD treatment from two different depths (20-30 cm

and 40-50 cm) in each greenhouse using a Gauge Auger DIK-106B core sampler (Daiki

Rika Kogyo Co., LTD, Saitama, Japan) in Is, Ha, To, Gi, and Wa fields (9 greenhouse ×

4 soil samples, Totally 36 soil samples). In Ni, Ts and Sa field, only two soil samples

(before and after ASD treatment in the upper layer soil) were collected from each

greenhouse in these fields (8 greenhouse × 2 soil samples, Totally 16 soil samples). A

total of 52 soil samples were collected for analysis. Soil samples were collected from 5

randomly chosen points in each greenhouse and mixed well.

2.2 Quantification of *R. solanacearum* in the field

The most probable number (MPN)-PCR method, which is semi-quantitive *R*.

solanacearum counting method, was conducted as described by Inoue and Nakaho

(2014). Briefly, 10 g of soil was eluted in cultivation buffer and the soil extract was

diluted with buffer 10, 100, and 1000 times. Each aliquot was incubated at 35 °C for

approximately 24 h, and nested-PCR targeting *phcA* gene, which is a key gene for the

appearance of wilt disease, was conducted on the culture sample using the primer pairs

phcA2981f (5′-TGGATATCGGGCTGGCAA-3′) and phcA4741r

(5′-CGCTTTTGCGCAAAGGGA-3′) for the first step and phcA3538f

(5′-GTGCCACAGCATGTTCAGG-3′) and phcA4209r

(5′-CCTAAAGCGCTTGAGCTCG-3′) for the second step. The number of samples that

appeared in the PCR amplification around 700 bp length was collectively checked

against the MPN table. The detection limit of *R.solanacearum* by this method is 3 to

2400 colony forming units (cfu) g^{-1} dry soil and a detection indicates the risk of

outbreaking of bacterial wilt in the field.

2.3 Tag-encoded amplicon sequencing targeted by 16S rRNA gene for prokaryotes

DNA was extracted from 0.5 g soil with an ISOIL kit for bead beating (Nippongene,

Tokyo, Japan) following the manufacturer's instructions. DNA quantification and

integrity were measured using a Nanodrop spectrophotometer (Thermo Fisher Scientific,

Waltham, MA, USA) and by gel visualization (0.8% agarose in TAE). PCR of each

sample was performed to amplify the V4 variable region of the 16S rRNA gene using

the bacterial and archaeal universal primers, 515F

(5'-GTGCCAGCMGCCGCGGTAA-3') and 806R

(5'-GGAC-TACVSGGGTATCTAA-3') (Camporaso et al. 2011). The library was

prepared by adaptor ligation and PCR primer pairs using the TruSeq Nano DNA Library

Prep Kit (Illumina, San Diego, CA, USA). When two or more bands were detected by

1.5%-agarose gel electrophoresis, PCR products of approximately 300 bp in length were

excised from the gel and purified using a MonoFas DNA purification kit (GL Sciences,

Tokyo, Japan) for prokaryotes. One soil sample (the upper layer soil in To3 after ASD

treatment) was not amplified after PCR amplification and was excluded from the analyses.

Each PCR amplicon was cleaned twice using an Agencourt AMPure XP system

(Beckman Coulter, Inc, Palo Alto, CA, USA) to remove primers and short DNA

fragments, and was quantified using a Qubit Fluorometer (Invitrogen, Carlsbad, CA,

USA). After successful amplification, the PCR products were adjusted to equimolar

concentrations and subjected to unidirectional pyrosequencing at Bioengineering Lab.

Co., Ltd. (Kanagawa, Japan) on a MiSeq instrument (Illumina). A total of 3,114,333

sequences in the 51 samples were obtained after sequencing (Supplemental Table 1).

Sequencing data were deposited in the DNA Data Base of Japan (DDBJ) Sequence

Read Archive under accession number DRA006673.

2.4 Data analyses

The analysis of the sequencing data was conducted according to Lee et al. (2017). Raw

fasta files were pre-processed in the Quantitative Insights into Microbial Ecology

(QIIME) bioinformatic pipeline (Caporaso et al. 2010). Data from the read sequences,

quality, flows, and ancillary metadata were analyzed using the QIIME pipeline. Quality

filtering consisted of discarding reads < 200 bp or > 1000 bp in length, excluding

homopolymer runs of > 6 bp and continuous ambiguous bases of > 6, and accepting one

barcode correction and two primer mismatches. Moreover, a mean quality score < 25

was also removed. Finally, singleton operational taxonomic units (OTUs) and chimeric

sequences were removed for statistical analyses. Denoising was performed using the

built-in Denoiser algorithm, and chimera removal and OTU picking were accomplished

with USEARCH 61 considering a pairwise identity percentage of 0.97. Taxonomy

assignment was performed using the Ribosomal Database Project (RDP) naïve Bayesian

classifier with a minimum confidence of 0.8 against the Greengenes database, October

2012 release and Basic Local Alignment Search Tool (BLAST). OTU-based analysis

was performed on pyrotag-based datasets to calculate the richness and diversity using

the phyloseq R package, version 1.7.24 (McMurdie and Holmes 2013). The diversity

within each individual sample was estimated using the non-parametric Shannon

diversity index. The Chao1 estimator was calculated to estimate the richness of each

sample. Multivariate analysis of community structure and diversity was performed on

pyrotag-based datasets using a weighted Unifrac dissimilarity matrix calculated in

QIIME, jackknifing (1000 reiterations) read abundance data at the deepest level

possible (41,012 reads), and using unconstrained ordination by a Principal Coordinate

Analysis (PCoA) for prokaryotes in each soil layer. *K*-means clustering was used to

verify the effects of sampling fields and ASD treatment on prokaryotic community.

Finally, indicator value analysis was done using the indicspecies R package (De Cáceres

2009) with the aim of identifying OTUs associated with suppressive soil, rather than the

conducive soil or vice versa ($p < 0.05$). We created two patterns of soil groups. One

group was divided into samples from the soil before or after ASD treatment. The other

group was divided based on the efficiency of disinfestation. The datasets were merged

and the abundant OTUs that were associated with the pathogen non-detected soil after

ASD treatment were evaluated in each layer ($p < 0.001$). The number of random

permutation tests for the calculation of the indicator values was 999.

2.5 Construction of phylogenetic tree

The phylogenetic tree of clostridial members was constructed by a 1000-fold bootstrap

analysis using neighbor-joining methods with the Clustal W program (version 2.0

Developments) software (Larkin et al. 2007) and the NJ plot software (Perrière and

Gouy 1996). The phylogenetic tree contained 16S rRNA sequences of several clusters

of clostridial strains.

3. Results

3.1. Effectiveness of ASD treatment with sugar-containing diatoms

R. solanacearum in soil was not detected after treated ASD with sugar-contained

diatoms in 14 of the 17 greenhouses and 7 of the 9 greenhouses in the upper and lower

layers, respectively, (Table 2). Therefore, the ASD treatments were appropriate in each

fields. Some fields (in IS, Ha1, and Ha2 field) were not-well-disinfested by ASD. We

classified the soil samples from greenhouses into two groups based on the effect of

ASD: not-well-disinfested (disinfestation failure) soil as F-soil and well-disinfested

(disinfestation success) soil as S-soil.

3.2 Changes of prokaryotic communities after ASD treatment

The sequences were clustered into 74,539 OTUs at 97% similarity. Among S- and F-

soil, the DNA concentrations were significantly decreased in the upper layer of the

S-soil and the lower layer of the F-soil (Figure 1, Supplemental Table 2). Shannon

indices and Chao1 were decreased after ASD in the upper. OTU numbers were

significantly decreased in F-soil regardless of soil layers.

PCoAs based on weighted-UniFrac analysis showed that prokaryotic communities

were roughly classified by sampling fields (Figure 2). In the upper layer, Ha fields

contained one cluster regardless of ASD treatment and disinfestation efficiency (Figure

2A). IS, Wa and Ts fields were separated by before and after ASD treatment but not

affected by disinfestation efficiency. In the lower layer soil, IS, Ha, and Gi were also

different among sampling fields and ASD treatment, but not different by S- and F-soil

(Figure 2B). The microbes collected from same fields comprised same clusters before

and after ASD treatment in spite of ASD efficiency was different.

After ASD treatment, the relative abundance of *Firmicutes* was increased by more

than 2-times in 8 greenhouses (Supplemental Table 3). In the upper layer soil,

Firmicutes was significantly increased, while *Crenarchaeota* was decreased after ASD

treatement (Figure 3). *Acidobacteria* and *Verrcomicrobia* in the upper layer of the

S-soil, *Acidobacteria*, and *Crenarchaeota* in the lower layer of the F-soil were

significantly decreased. In the class level, *Clostridia* that belonged to the *Firmicutes*

phylum were increased more than 2-times after ASD in 15 of 16 and 8 of 9 greenhouses

in the upper and lower layers, respectively (Supplemental Table 4). Moreover,

Betaproteobacteria and *Clostridia* were significantly increased after ASD treatment in

the upper layer of the S-soil (Figure 4). In the S-soil, *Sphingobacteria* and *Cytophagia*

were significantly decreased in the upper layer, and *Thermoleophilia* and *Actinobacteria*

were decreased in the lower layer.

3.4 Specifically increased OTUs in S-soil

We successfully detected 25 OTUs specifically increased in the S-soil by ASD from

more than 8 of the 16 sites or 4 of the 6 sites from the upper and lower layers,

respectively (Table 3). They belonged to several classes, including *Clostridia*, *Bacillus*,

Betaproteobacteria, *Gammaproteobacteria*, and *Mollicutes*. Eighteen out of the 25

OTUs belonged to *Clostridia*. The phylogenetic trees consisting of OTUs related to

clostridial members showed that diversified *Clostridia* were specifically increased after

ASD treatment in the S-soil (Figure 5). Three OTUs belonged to the *Caloramator* group,

Clostridium cluster III and two OTUs belonged to the *Christensenella* group. OTU 50

248 (*Caloramator*) was commonly detected in 11 upper layer and five lower layer soil

249 samples. OTU311169 (*Bacillus*), OTU199301 and OTU624373 (*Christensenella*), OTU

250 112 (*Clostridium*), and OTU56 (*Rhodothermaceae*) were also increased in 10 of the 14

251 S-soils, while OTU934369 and OTU304 (*Caloramator*) and OTU212333

252 (*Symbiobacterium*) were commonly detected in more than five of the six fields of S-soil.

253 Prokaryotes decreased by ASD treatment common to the various fields in this criterion

254 were not detected (data not shown).

256 **4. Discussion**

257  In this study, we showed that microbial communities were not different even if the

258 fields indicated different disinfestation effect by ASD. Especially, in Ha field, we used

259 three contiguous greenhouses and they treated the same field managements for more

260 than 10 years. They showed a different disinfestation effect but were included in the

261 same cluster. Our previous study also showed that prokaryotic community structures

262 were affected more by indigenous communities than the disease suppression effect (Lee

263 et al. 2017). These results suggested that the difference of indigenous soil communities

are not affected to controlling of soil-borne pathogen. Runia et al. (2014) showed

Firmicutes are increased by ASD treatment among 5 soil types using the real-time PCR

method. We also showed that *Clostridia* belonged to Firmimutes have increased in

S-soil. *Clostridia*, are gram-positive anaerobic bacteria. During ASD, decreases in

oxygen availability along with increases in pH occur, which promotes the increased

prevalence of anaerobic microbes (Momma et al. 2006; Mowlick et al. 2013, 2014).

Several studies indicated that *Clostridia* increase in number and become dominant

bacteria following ASD treatment using plant material or low concentration ethanol as a

substrate (Huang et al. 2015, 2016; Mowlick et al. 2012; Strauss et al. 2017). The prior

and present results indicate that *Clostridia* increased after ASD treatment regardless of

soil types, field locations, and substrate types and they may be key microbes for

disinfestation by ASD. *Caloramator*-like OTUs were the most commonly found OTUs

in the upper and lower layers of various fields. They are rod-shaped, obligate anaerobic,

thermophilic endospore producers that can ferment several sugars and produce acetate,

isobutyrate, lactate, and other VFAs, while they do not degrade cellulose (Collins et al.

1994; Ogg and Patel 2011). Mowlick et al. (2012) reported that 63% of clostridial

clones that were increased after ASD treatment belonged to clostridium cluster I,

including the *Oxobacter* and *Caloramator* groups. *Symbiobacterium* is rod-shaped

thermophiles that are syntrophic with *Bacillus* spp., which possess a glucose

degradation pathway (Ohno et al. 2000). *Christensenella* is a non-spore forming, short,

straight rod with tapered ends; it can use various sugars and produce VFAs during

fermentation (Morotomi et al. 2012). *Christensenella* were detected in an anaerobic

cellulolytic microbial consortium from mangrove soil (Gao et al. 2014). These results

suggest that phylogenetically diverse clostridial members might be responsible for

suppression of pathogens through their production of VFAs during anaerobic

decomposition of sugar-containing diatoms.

Strauss et al. (2017) reported the increased of *Betaproteobacteria* by ASD treatment

with organic bran at soil depths of 15.2 cm and 45.7 cm. We showed

Noviherbaspirillum, belonged to *Betaproteobacteria*, which were specifically detected

from S-soil. They are gram-negative, aerobic, non-spore forming rods capable of

utilizing several carbohydrates as carbon sources (Lin et al. 2013). Moreover, several

Bacillus species have been described as biological control agents for bacterial wilt and

have been detected from soil after ASD treatment (Mowlick et al. 2012; Yamamoto et al.

2014; Yuliar et al. 2015). The family *Rhodothermaceae* of the class Cytophagia

comprises six genera; most are isolated from extreme environments, such as hypersaline

crystallized ponds in the Dead Sea and submarine hot spring (Xia et al. 2015). They

exhibit extreme thermophilic or halophilic growth characteristics. Mazzola et al (2018)

showed that these *Cytophagaceae* bacteria were significantly increased after ASD

treatment with rice bran. However, their role for disinfestation does not appear.

Symbiobacterium, *Christensenella*, *Noviherbaspirillum*, and *Rhodothermaceae* were not

previously detected from ASD treated soil. They might have important roles for

disinfestation of *R. solanacearum*. *Noviherbaspirillum*, *Bacillus*, and *Rhodothermaceae*

are aerobic bacteria that were also increased by ASD, as with anaerobic bacteria. In the

soil environment, some *Bacillus* might contribute to the rapid decrease of soil redox

potential at the initial stage of ASD treatment by consuming oxygen (Mowlick et al.

2012). Therefore, they might have important roles in maintaining anoxic soil conditions

during ASD treatment.

In conclusion, we revealed prokaryotic communities were not strongly affected by

disinfestation efficiency compare with sampling fields, but *Clostridia* and

Betaproteobacteria were significantly increased in well-disinfested soil. Twenty-five

OTUs were specifically detected in S-soil and most of them were affiliated to *Clostridia*,

which was well-known bacteria for ASD effect. However, some microbes were not

previously detected from ASD treated fields. These microbes will be a good indicator

for well-disinfested soil and future study will appear the role of these microbes for

disinfestation.

Acknowledgements

This work was supported by the Cabinet Office, Government of Japan, Cross-ministerial

Strategic Innovation Promotion Program (SIP), and the "Technologies for creating

next-generation agriculture, forestry and fisheries" (funding agency: Bio-oriented

Technology Research Advancement Institution, NARO).

This work was also partially supported by the RIKEN Competitive Program for

Creative Science and Technology, a Grant-in-Aid for Scientific Research from the Japan

Society for Promotion of Science, No. 17K01447 (to M.O.), and by the Sasakawa

Scientific Research Grant from The Japan Science Society (to C.G.L.). We would like to

thank Editage (www.editage.jp) for English language editing.

References

Blok, W.J., Lamers, J.G., Termorshuizen, A.J., Bollen, G.J., 2000. Control of soilborne

plant pathogens by incorporating fresh organic amendments followed by tarping.

Phytopathol. 90, 253-259.

Caporaso, J.G., J. Kuczynski, J. Stombaugh, *et al*. 2010. QIIME allows analysis of

high-throughput community sequencing data. Nature Methods. 7, 335–336.

Caporaso, J.G., C.L. Lauber, W.A., Walters, D., Berg-Lyons, C.A., Lozupone, P.J.,

Turnbaugh, N., Fierer and R. Knight. 2011. Global patterns of 16S rRNA diversity

at a depth of millions of sequence per sample. Proc. of Natl. Acad. Sci. U. S. A. 108,

S4516–4522.

Collins, M.D., Lawson, P.A., Willems, A., Cordoba, J. J., Fernandez-Garayzabal, J.,

Garcia, P., Cai, J., Hippe, H., Farrow, J.A.E. 1994. The phylogeny of the genus

Clostridium: proposal of five new genera and eleven new species combinations. Int.

J. Syst. Bacteriol. 44, 812-826.

De Cáceres, M., Legendre. P., 2009. Associations between species and groups of sites:

indices and statistical inference. Ecology. 90, 3566–3574.

Gao, Z.M., Xu, X., Ruan, L.W., 2014. Enrichment and characterization of an anaerobic

cellulolytic microbial consortium SQD-1.1 from mangrove soil. Appl. Microbiol.

Biotechnol. 98, 465-474.

Graham, J. Lloyd, A.B., 1979. Survival of potato strain (race 3) of *Pseudomonas*

solanacearum in the deeper soil layers. Aust. J. Agric. Res. 30, 489–496.

Huang, X., Liu, L., Wen, T., Zhang, J., Wang, F., Cai, Z., 2016. Changes in the soil

microbial community after reductive soil disinfestation and cucumber seedling

cultivation. Appl. Microbiol. Biotechnol. 100, 5581-5593.

Huang, X., Liu, L., Wen, T., Zhu, R., Zhang, J., Cai, Z., 2015. Illumina Miseq

investigations on the changes of microbial community in the *Fusarium oxyporum*

f.sp. *cubense* infested soil during and after reductive soil disinfestation. Microbiol.

Res. 181, 33-42.

Inoue, Y., Nakaho, K., 2014. Sensitive quantitative detection of *Ralstonia*

solanacearum in soil by the most probable number-polymerase chain reaction

(MPN-PCR) method. Appl. Microbiol. Biotechnol. 98, 4169–4177.

Kelman, A., 1998. One hundred and one years of research on bacterial wilt, p. 1–5. *In*

P.H. Prior, C. Allen, J. Elphinstone (ed.), Bacterial Wilt Disease: Molecular and

Ecological Aspects, Springer, Heidelberg.

Lee, C.G., Iida, T., Uwagaki, Y., Otani, Y., Nakaho, K., Ohkuma, M., 2017.

Comparison of prokaryotic and eukaryotic communities in soil samples with and

without tomato bacterial wilt collected from different fields. Microb. Environ. 32,

376-385.

Lin, S.Y., Hameed, A., Arun, A.B., Liu, Y.C., Hsu, Y.H., Lai, W.A., Rekha, P.D.,

Young, C,C., 2013. Description of *Noviherbaspirillum malthae* gen. nov., sp. nov.,

isolated from an oil-contaminated soil, and proposal to reclassify *Herbaspirillum soli*,

Herbaspirillum aurantiacum, *Herbaspirillum canariense* and *Herbaspirillum*

psychrotolerans as *Noviherbaspirillum soli* comb. nov., *Noviherbaspirillum*

aurantiacum comb. nov., *Noviherbaspirillum canariense* comb. nov. and

Noviherbaspirillum psychrotolerans comb. nov. based on polyphasic analysis. Int. J.

Syst. Microbiol. 63, 4000-4007.

Liu, L., Kong., J., Cui, H., Zhang, J., Wang, F., Cai, Z., Huang, X., 2016. Relationships

of decomposability and C/N ratio in different types of organic matter with

suppression of *Fusarium oxysporum* and microbial communities during reductive

soil disinfestation. Biol. Control. 101, 103-113.

Mazzola, M., Hewacitharana, S.S., Strauss, S.L., Shennan, C., Muramoto, J., 2016.

Anaerobic soil disinfestation and *Brassica* seed meal amendment alter soil

microbiology and system resistance. Int. J. Fruit. Sci. 16, 47-58.

Mazzola, M., Muramoto, J., Shennan, C., 2018. Anaerobic disinfestation induced

changes to soil microbiome disease incidence and strawberry fruit yields in

California field trials. Appl. Soil. Ecol. 127, 74-86.

McMurdie, P.J., Holmes. S., 2013. phyloseq: An R package for reproducible interactive

analysis and graphics of microbiome census data. PLoS One. 8, e61217.

Mendes, R., Kruijt, M., de Bruijin I., et al. 2011. Deciphering the rhizosphere

microbiome for disease-suppressive bacteria. Science. 3321097-1100.

Momma, N., Momma, M., Kobara, Y., 2010. Biological soil disinfestation using

ethanol: effect on *Fusarium oxyporum* f. sp. <u>*lycoperisici*</u> and soil microorganisms. J.

Gen. Plant Pathol. 76, 336-344.

Momma, N., 2008. Biological soil disinfestation (BSD) of soilborne pathogens and its

possible mechanisms. Jpn. Agr. Res. Q. 7-12.

Momma, N., Kobara, Y., Uematsu, S., Kita, N., Shinmura, A., 2013. Development of

biological soil disinfestaion in Japan. Appl. Microbiol. Biotechnol. 97, 3801-3809.

Momma, N., Yamamoto, K., Shimandi, P., Shishido, M., 2006. Role of organic acids in

the mechanisms of biological soil disinfestation (BSD). J. Gen. Plant Pathol. 72,

247-252.

Momma, N., Usami, T., Shishido, M,. 2007. Detection of *Clostridium* sp. including

biological soil disinfestation (BSD) and suppression of pathogens causing Fusarium

wilt and bacterial wilt of tomato by gases evolved during BSD (in Japanese with

English summary). Soil Microorg. 61, 3-9.

Morotomi, M., Nagai, F., Watanabe, Y., 2012. Description of *Christensenella minuta*

gen. nov., sp. nov., isolated from human feces, which forms a distinct btranch in the

order *Clostridiales*, and proposal of *Christensenellaceae* fam. nov. Int. J. Syst.

Microbiol. 62, 144-149.

Mowlick, S., Yasukawa, H., Inoue, T., Takehara, T., Kaku, N., Ueki, K., Ueki, A., 2013.

Suppression of spinach wilt disease by biological soil disinfestation incorporated

with *Brassica juncea* plants in association with changes in soil bacterial

communities. Crop Prot. 54, 185-193.

Mowlick, S, Takehara, T., Kaku, N,, Ueki, K., Ueki, A., 2013. Proliferation of

diversified clostridial species during biological soil disinfestation incorporated with

plant biomass under various conditions. Appl. Microbiol. Biotechnol. 97,

8365-8379.

Mowlick, S, Hirota, K., Takehara, T., Kaku, N., Ueki, K., Ueki, A., 2012 Development

of anaerobic bacterial community consisted of diverse clostridial species during

biological soil disinfestation amended with plant biomass. Soil. Sci. Plant. Nutr. 58,

273-287.

Ogg, C.D., Patel, B.K.C., 2011. *Caloramator mitchellensis* sp. nov., a thermoanaerobe

isolated from the geothermal waters of the Great Artesian Basin of Australia, and

emended description of the genus *Caloramator*. Int. J. Syst. Microbiol. 61,

644-653.

Ohno, M., Shiratori, H., Park, M.J., Saitoh, Y., Kumon, Y., Yamahista, N., Hirata, A.,

Nishida, H., Ueda, K., Beppu, T., 2000. *Symbiobacterium thermophilum* gen. nov.,

sp. nov., a symbiotic thermophile that depends on co-culture with a *Bacillus* strain

for growth. Int. J. Syst. Microbiol. 50, 1829-1832.

Perrière, G., Gouy, M., 1996. WWW-Query: an online retrieval system for biological

sequence banks. Biochime. 78, 364-369.

Runia, W.T., Thoden, T.C., Molendijk, L.P.G., van den Berg, W., Termorshuizen, A.J.,

Streminska, M.A., van der Wurff, A.W.G., Feli, H., Meints, H., 2014. Unravelling

the mechanism of pathogen inactivation during anaerobic soil disinfestation. Acta.

Hortic. 1044, 177-193.

Shinmura, A., 2000 Causal agent and control of root rot of welsh onion. PSJ Soilborne

Disease Work Report. 20, 133-143.

Shrestha, U., Auge, R.M., Butler, D.M., 2016. A meta-analysis of the impact of

anaerobic soil disinfestation on pest suppression and yield of horticultural crops.

Front. Plant. Sci. 7, 1254.

Straus, S.L., Kluepfel, D.A., 2015. Anaerobic soil disinfestation: A

chemical-indipendent approach to pre-plant control of plant pathogens. J. Integr.

Agr. 14, 2309-2318.

Strauss, S.L., Greenhut, R.F., McClean, A.E., Kluepfel, D.A., 2017. Effect of anaerobic

soil disinfestation on the bacterial community and key soilborne phytopathogenic

agents under walnut tree-crop nursery conditions. Plant Soil. 415, 493-506.

van Agtmaal, M., van OS, G.J., Hol, W.H.G., Hundscheid, M.P.J., Runia, W.T.,

Hordijk, C.A., de Boer, W., 2015. Legacy effects of anaerobic soil disinfestation on

soil bacterial community composition and production of pathogen-suppression

volatiles. Front. Microbiol. 6, 701.

Van Bruggen, A.H.C., Semenov, A. M., 2000. In search of biological indicators for soil

health and disease suppression. Appl. Soil. Ecol. 15, 13-24.

Wang, J.F., Lin, C.H., 2005. Integrated management of tomato bacterial wilt.

AVRDC-The world vegetable center, Taiwan.

Xia, J., Zhou, Y.X., Zhao, L.H., Chen, G.J., Du, Z.J., *Longimonas halophile* gen. nov.,

isolated from marine solar saltern. Int. J. Syst. Microbiol. 65, 2272-2276.

Yamamoto, S., Shiraishi, S., Kawagoe, Y., Mochizuki, M., Suzuki, S., 2015. Impact of

Bacillus amyloliquefaciens S13-3 on control of bacterial wilt and powdery mildew

in tomato. Pest Manag. Sci. 71, 722-727.

Yuliar, Nion, Y.A., Toyota, K., 2015. Recent trends in control methods for bacterial

wilt disease caused by *Ralstonia solanacearum*. Microb. Environ. 30, 1-11.

Figure legend

Figure 1. Comparison of DNA concentration, prokaryotic diversity and richness

between the F- and S-soil.

Asterisk represents a pair of means that show significant difference between the S- and

F- soil (t-test, $p < 0.05$).

Figure 2. A UniFrac-weighted principal component analysis of the soil prokaryotic

communities in the upper layer (A) and lower layer (B) of the field before and after

ASD treatment.

Closed circle: S-soil before ASD treatment, Open circle: S-soil after ASD treatment,

Closed triangle: F-soil before ASD treatment, Open triangle: F-soil after ASD

treatment.

The clustering was conducted by *k*-means analysis.

Figure 3. Comparison of the prokaryotic phyla which observed significant difference

between the S- and F-soil

Asterisk represents a pair of means that show a significant difference between the S-

and C- soil (Tukey's test, $p < 0.05$).

Figure 4. Comparison of the prokaryotic class that observed a significant difference

between the S- and F-soil

Asterisk represents a pair of means that show a significant difference between the S-

and C- soil (Tukey's test, $p < 0.05$).

Figure 5. Neighbor-joining tree showing the phylogenetic relationships of the

clostridial OTUs based on 16S rRNA gene sequencing (According to the clostridial

cluster analysis by Collins et al. 1994). The OTUs obtained from this study are shown

in gothic letters. *Bacillus subtilis* was used as the outgroup. Bold letters indicate OTUs

that specifically appeared in the S-soil.

Figure1

Fig. 2A

Fig. 2B

Fig. 3

Fig. 4

Fig. 5

Table 1. Detailed information on the sampling fields

Field	Ishikawa	Hakusan	Niigata	Tsubame	Sanjo	Toyama	Gifu	Wakayama
Code	IS	Ha	Ni	Ts	Sa	To	Gi	Wa
latitude and longitude	36° 38′ N, 136° 42′ E	36° 29′ N, 136° 30′ E	37° 55′ N, 139° 9′ E	37° 38′ N, 138° 55′ E	37° 39′ N, 139° 57′ E	36° 61′ N, 136° 96′ E	35° 44′ N, 136° 70′ E	33° 27′ N, 135° 14′ E
Soil types	Gleysol	Gleysol	Fluvisol	Fluvisol	Fluvisol	Gleysol	Gleysol	Cambisols
Replication	1	3	3	3	2	3	1	1
pH	5.54	7.22	7.11	7.03	7.24	6.07	6.08	7.49
EC (mS cm-1)	0.03	0.31	0.32	0.24	0.20	0.29	0.12	0.88
ASD conducted schedule	7/27/2016 - 8/18/2016	7/11/2016 - 7/28/2016	8/18/2016 - 9/9/2016	7/31/2016 - 8/21/2016	9/1/2016 - 9/22/2016	8/2/2016 - 8/26/2016	8/11/2016 - 9/1/2016	7/29/2016 - 8/19/2016

Table 2. The number of *R.solanacearum* in each field before and after ASD treatment

	Soil layer	ASD treatment	IS	Ha1	Ha2	Ha3	Ni1	Ni2	Ni3	Ts1	Ts2	Ts3	San1	San2	To1	To2	To3	Gi	Wa
Number of *R.solanacearum* in soil (cfu soil^{-1})	Upper	Before	> 2400	1100	46	23	95	210	460	95	75	120	160	43	64	43	23	> 2400	> 2400
		After	75	21	15	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
	Lower	Before	240	1100	39	13	-	-	-	-	-	-	-	-	11	53	12	1100	1100
		After	3	21	11	N.D.	-	-	-	-	-	-	-	-	N.D.	N.D.	N.D.	N.D.	N.D.

The number of each fields indicate replication number
N.D.; *R. solanacearum* were not detected with this method.
-; No samples

Supplemental Table 1. Read numbers of the 16S rRNA gene in each soil sample

			Before	After
	IS	Upper	48,338	56,750
		Lower	51,336	66,408
IS	Ha-1	Upper	50,451	41,012
		Lower	54,223	65,242
	Ha-2	Upper	50,052	50,403
		Lower	54,362	65,817
	Ha-3	Upper	44,053	158,609
		Lower	50,653	56,204
To	1	Upper	49,838	52,458
		Lower	42,379	44,913
	2	Upper	120,447	106,275
		Lower	46,493	50,754
	3	Upper	-	55,666
		Lower	47,673	56,354
Ni	Shi1	Upper	70,925	58,473
	Shi2	Upper	65,166	65,462
	Shi3	Upper	63,392	63,830
	Tsu1	Upper	67,593	69,027
	Tsu2	Upper	60,868	58,135
	Tsu3	Upper	59,287	60,065
	San1	Upper	79,424	55,336
	San2	Upper	64,800	60,657
Gi		Upper	54,337	62,183
		Lower	58,254	47,825
Wa		Upper	57,969	61,009
		Lower	55,401	57,752

Supplemental Table2
[Click here to download high resolution image](#)

Supplemental Table 2. The change rate of the soil chemical properties or prokaryotic diversity, richness and DNA concentration before to after ASD treatment.

		IS	IS	Ha-1	Ha-2	Ha-3	1	To	2	3	Shi1	Shi2	Shi3	Ni	Tsu1	Tsu2	Tsu3	San1	San2	Gi	Wa
DNA concentration (µg g soil⁻¹)	Upper	0.20		0.26	0.49	0.31	0.32		0.32	0.30	0.50	0.59	0.58		0.35	0.64	0.23	1.35	1.66	0.27	0.45
	Lower	0.34		0.23	0.53	0.44	0.32		0.21	0.24	-	-	-		-	-	-			3.5	0.89
Shannon index	Upper	0.96		0.92	0.84	1.00	0.99		1.00	-	0.99	1.01	0.92		0.86	0.88	0.87	0.97	0.97	0.97	0.90
	Lower	0.79		0.90	0.83	0.95	0.99		0.98	0.98	-	-	-		-	-	-			0.80	0.87
Observed OTU	Upper	0.77		0.62	0.58	0.98	0.9		0.59	-	0.98	0.96	0.82		0.52	0.51	0.53	0.94	0.88	0.99	0.76
	Lower	0.65		0.78	0.65	0.80	0.91		1.03	0.77	-	-	-		-	-	-			0.32	0.66
Chao 1 estimator	Upper	0.08		0.62	0.64	0.98	0.88		0.84	-	0.96	0.99	0.82		0.55	0.54	0.58	0.99	0.89	1.01	0.72
	Lower	0.59		0.68	0.59	0.82	0.91		0.94	0.88	-	-	-		-	-	-			0.44	0.67

The red and blue cells indicate increasing or decreasing factors after ASD treatment compared with the before

treatment. The color intensities indicates the rate of each difference.

-; No samples

Supplemental Table 3. The change rate of the soil prokaryotic phylum before to after ASD treatment.

		IS				To			Ni								Gi	Wa
		IS	Ha-1	Ha-2	Ha-3	1	2	3	Shi1	Shi2	Shi3	Tsu1	Tsu2	Tsu3	San1	San2		
Proteobacteria	Upper	1.00	0.85	1.01	0.81	1.26	0.93	-	1.09	1.08	1.46	0.58	0.62	0.72	1.24	1.26	0.94	0.39
	Lower	1.79	1.07	1.52	1.07	1.28	1.30	1.18	-	-	-	-	-	-	-	-	1.18	0.79
Firmicutes	Upper	1.54	1.84	0.82	2.39	2.11	1.42	-	1.09	0.80	1.34	2.64	2.83	2.95	0.98	0.71	1.43	3.85
	Lower	0.87	3.45	2.02	1.93	0.59	0.73	0.28	-	-	-	-	-	-	-	-	0.54	0.12
Acidobacteria	Upper	0.66	1.06	1.66	0.64	0.68	1.03	-	0.65	0.71	0.35	0.59	0.65	0.70	0.53	0.49	1.17	1.22
	Lower	0.42	0.54	0.37	0.48	1.05	0.88	1.90	-	-	-	-	-	-	-	-	1.17	0.34
Bacteroidetes	Upper	0.56	3.51	0.87	2.07	1.17	0.60	-	0.86	0.79	1.05	0.10	0.17	0.07	1.13	1.86	0.23	0.40
	Lower	0.46	4.15	5.94	2.04	1.42	1.39	0.55	-	-	-	-	-	-	-	-	0.30	0.47
Chloroflexi	Upper	1.39	0.94	1.03	0.78	0.70	1.77	-	1.35	1.44	0.79	2.53	2.03	2.68	1.30	1.36	0.92	0.72
	Lower	0.45	0.73	0.64	1.16	0.75	1.05	1.23	-	-	-	-	-	-	-	-	0.90	0.52
Gemmatimonadetes	Upper	0.80	0.83	1.54	1.18	0.93	0.84	-	0.74	0.79	0.52	2.30	2.33	1.63	0.92	1.10	0.55	0.90
	Lower	0.45	0.44	0.33	0.62	1.07	0.89	1.02	-	-	-	-	-	-	-	-	1.36	0.59
Actinobacteria	Upper	1.05	1.30	1.03	1.08	1.30	0.98	-	0.68	0.76	0.64	1.52	1.09	0.88	0.70	0.71	0.41	0.42
	Lower	0.75	0.87	0.78	0.68	1.02	0.52	0.62	-	-	-	-	-	-	-	-	0.67	0.20
Crenarchaeota	Upper	0.29	0.19	9.36	0.00	0.50	1.24	-	0.53	1.21	0.27	0.23	0.20	0.96	0.17	0.27	0.50	0.14
	Lower	0.12	0.19	0.04	0.59	1.02	1.02	0.55	-	-	-	-	-	-	-	-	1.17	5.82
Verrucomicrobia	Upper	0.47	0.89	0.64	1.35	0.53	0.78	-	0.79	1.01	0.60	0.27	0.40	0.16	0.49	0.38	0.20	0.26
	Lower	0.32	1.29	0.82	0.65	1.58	0.61	1.84	-	-	-	-	-	-	-	-	0.36	0.85
Planctomycetes	Upper	0.81	0.59	1.10	0.78	0.44	1.15	-	0.78	0.79	0.59	0.58	0.49	0.98	0.66	0.83	0.74	0.31
	Lower	0.32	0.37	0.39	0.58	1.11	0.67	1.43	-	-	-	-	-	-	-	-	1.29	0.07

The red and blue cells indicate increasing or decreasing factors after ASD treatment compared with the before treatment. The color intensities indicates the rate of each difference.

-; No samples

Supplemental Table 4. The change rate of the soil prokaryotic class before to after ASD treatment.

Phylum	Class		IS	IS	IS	IS	To	To	To	Ni	Ni	Ni	Ni	Ni	Ni	Ni	Ni	Gi	Wa
			IS	Ha-1	Ha-2	Ha-3	1	2	3	Shi1	Shi2	Shi3	Tsu1	Tsu2	Tsu3	San1	San2	Gi	Wa
Proteobacteria	Alphaproteobacteria	Upper	0.88	1.273	0.85	0.945	1.242	0.928	-	0.803	0.833	0.955	0.346	0.376	0.408	0.844	0.761	0.518	0.405
		Lower	0.52	1.435	1.452	0.563	0.997	0.754	0.678	-	-	-	-	-	-	-	-	0.129	0.324
	Betaproteobacteria	Upper	1.34	0.459	1.229	1.329	2.182	1.295	-	1.491	1.309	3.709	0.89	1.205	1.822	3.358	3.412	1.009	0.491
		Lower	3.16	1.026	0.695	1.957	2.069	1.463	1.36	-	-	-	-	-	-	-	-	0.45	0.433
	Deltaproteobacteria	Upper	0.86	0.66	1.045	1.307	0.936	0.758	-	1.163	1.581	0.797	1.676	1.555	1.25	0.755	0.873	2.116	0.449
		Lower	1.02	0.953	1.036	2.279	1.335	0.924	1.117	-	-	-	-	-	-	-	-	1.383	0.329
	Gammaproteobacteria	Upper	1.00	0.772	0.596	0.375	0.755	0.78	-	1.242	1.067	1.071	0.245	0.393	0.345	0.808	1.172	0.56	0.205
		Lower	0.30	0.88	4.755	0.411	0.919	0.832	0.342	-	-	-	-	-	-	-	-	0.284	0.274
Firmicutes	Bacilli	Upper	0.96	1.798	1.865	1.934	1.048	1.077	-	0.718	0.663	0.646	1.826	1.981	2.518	0.821	0.458	1.096	1.773
		Lower	0.47	1.736	1.291	0.868	0.542	1.11	0.688	-	-	-	-	-	-	-	-	0.2	2.731
	Clostridia	Upper	5.79	2.375	6.151	5.619	11.921	4.588	-	2.266	1.354	3.242	7.663	7.569	5.503	2.804	3.574	3.132	21.45
		Lower	3.017	11.363	3.987	3.901	5.254	7.113	2.289	-	-	-	-	-	-	-	-	1.46	16.45
Acidobacteria	Acidobacteria-6	Upper	0.66	0.216	0.444	0.639	0.561	1.706	-	0.729	0.736	0.422	0.594	0.676	0.637	0.727	0.701	0.628	0.285
		Lower	0.251	0.246	0.257	0.589	0.531	0.515	0.818	-	-	-	-	-	-	-	-	0.382	0.216
	Acidobacteriia	Upper	0.466	1.995	0.637	0.826	0.834	0.612	-	0.726	0.798	0.373	0.299	0.494	0.877	5.479	3.111	1.75	0.293
		Lower	0.44	1.385	0.81	0.436	2.013	1.525	3.538	-	-	-	-	-	-	-	-	1.08	0.129
	Solibacteres	Upper	2.782	1.361	0.882	1.372	1.292	0.808	-	0.733	0.527	0.742	2.317	1.496	1.581	1.118	1.231	0.701	5.314
		Lower	1.068	0.841	0.778	0.587	0.974	1.05	1.949	-	-	-	-	-	-	-	-	0.895	3.996
	[Chloracidobacteria]	Upper	0.329	0.06	0.037	0.926	0.735	1.022	-	0.38	0.384	0.346	0.053	0.231	0.041	0.33	0.267	0.669	0.352
		Lower	0.204	0.259	0.401	0.74	0.868	0.349	0.366	-	-	-	-	-	-	-	-	0.069	0.185
Bacteroidetes	Sphingobacteria	Upper	0.698	4.267	3.244	1.958	0.613	0.57	-	0.991	0.606	0.282	0.183	0.324	0.063	0.298	0.364	0.521	0.82
		Lower	0.644	6.449	10.178	1.27	0.582	0.526	0.292	-	-	-	-	-	-	-	-	0.188	0.609
	Cytophagia	Upper	0.051	1.157	2.666	2.625	0.163	0.325	-	0.401	0.412	0.379	0.001	0.003	0.018	0.139	0.271	0.041	0.011
		Lower	0.139	1.044	2.852	0.691	0.287	0.212	0.212	-	-	-	-	-	-	-	-	0.05	0.012
Chloroflexi	Anaerolineae	Upper	2.26	0.072	0.064	1.752	0.476	2.129	-	1.884	1.859	0.919	0.851	1.12	3.981	1.56	2.082	1.365	0.302
		Lower	0.47	0.389	0.606	1.719	0.853	1.309	2.35	-	-	-	-	-	-	-	-	1.293	0.149
Gemmatimonadetes	Gemmatimonadetes	Upper	0.94	1.829	0.782	1.442	1.018	0.824	-	0.647	0.744	0.559	4.01	4.075	2.878	0.654	0.654	0.387	0.874
		Lower	0.89	1.118	0.524	0.629	1.134	0.856	0.487	-	-	-	-	-	-	-	-	0.075	0.627
	Gemm-1	Upper	0.596	0.198	0.051	0.424	0.728	0.761	-	0.83	0.683	0.427	1.102	0.96	0.754	1.325	1.807	0.654	0.791
		Lower	0.284	0.236	0.274	0.621	1.005	0.827	1.452	-	-	-	-	-	-	-	-	0.516	0.716
Actinobacteria	Actinobacteria	Upper	0.88	1.057	0.921	0.742	1.4	0.922	-	0.699	0.738	0.56	1.913	1.395	0.858	0.753	0.706	0.306	0.312
		Lower	1.279	1.334	0.407	0.414	0.362	1.05	0.208	-	-	-	-	-	-	-	-	0.145	0.406
	Thermoleophilia	Upper	1.484	1.776	0.943	1.359	1.343	0.916	-	0.805	0.89	0.958	1.077	0.652	1.096	0.737	0.779	0.466	0.565
		Lower	0.397	0.728	0.902	0.462	0.707	0.537	0.613	-	-	-	-	-	-	-	-	0.559	0.632
Crenarchaeota	Thaumarchaeota	Upper	0.284	0.209	0.027	0.003	0.5	1.237	-	0.529	1.226	0.253	0.194	0.193	0.902	0.13	0.233	0.451	0.139
		Lower	0.093	0.239	0.034	0.581	0.527	0.616	0.281	-	-	-	-	-	-	-	-	1.326	0.112
Nitrospirae	Nitrospira	Upper	0.417	0.127	0.19	0.241	0.609	1.283	-	1.422	1.748	0.747	0.438	0.926	0.873	0.53	0.377	1.773	0.288
		Lower	0.413	0.184	0.317	0.695	0.98	0.813	1.872	-	-	-	-	-	-	-	-	2.201	0.186