Article

Comparison of prokaryotic communities among fields exhibiting different disinfestation effects by anaerobic soil disinfestation

Chol Gyu Lee[1,2], Toshiya Iida[1], Eriko Matsuda[3], Kayo Yoshida[3], Masato Kawabe[4], Masayuki Maeda[5], Yasunori Muramoto[6], Hideki Watanabe[6], Yoko Otani[7], Kazhiro Nakaho[8], and Moriya Ohkuma[1]

[1] Japan Collection of Microorganisms, RIKEN BioResource Center, Tsukuba, Ibaraki 305-0074, Japan; cglee@go.tuat.ac.jp (C. G. L.); tiida@riken.jp (T. I); mohkuma@riken.jp (M. O.)

[2] Graduate School of Bio-Applications and Systems Engineering, Tokyo University of Agriculture and Technology, Koganei, Tokyo, 184-8588, Japan;

[3] Ishikawa Agriculture and Forestry Research Center, Kanazawa, Ishikawa 920-3198, Japan; eriko-m@pref.ishikawa.lg.jp (E. M.), kyoshida@pref.ishikawa.lg.jp (K. Y.)

[4] Horticultural Research Institute, Toyama Prefectural Agricultural, Forestry and Fisheries Research Center, Tonami, Toyama 939-1327, Japan; masato.kawabe@pref.toyama.lg.jp (M. K.)

[5] Niigata Agricultural Research Institute, Niigata, Nagaoka, Niigata 940-0826, Japan; maeda.masayuki@pref.niigata.lg.jp (M. M.)

[6] Gifu Prefectural Agricultural Technology Center, Matamaru, Gifu 501-1152, Japan; watanabe-hideki@pref.gifu.lg.jp (H. W.), muramoto-yasunori@pref.gifu.lg.jp (Y. M.)

[7] Wakayama Agricultural Experiment Station, Kinokawa, Wakayama, 640-0423, Japan; otani_y0005@pref.wakayama.lg.jp (Y. O.)

[8] Institute of Vegetable and Floriculture Science, National Agriculture and Food Research Organization, Tsu, Mie 514-2392, Japan. nakaho@affrc.go.jp (K. N.)

Correspondence: cglee@go.tuat.ac.jp (C.G. L)

Abstract: Anaerobic soil disinfestation (ASD) is a chemical-independent fumigation method used for reducing the abundance of pathogens at soil depths of <40 cm. However, its disinfestation efficiency is unstable under field conditions. The microbial community reflects the soil environment and is a good indicator of soil health. Therefore, soil with a good disinfestation efficiency may have a unique microbial community. The aim of the present study was to compare the prokaryotic communities among soils obtained from 17 geographically different greenhouses that experienced tomato bacterial wilt but exhibited different disinfestation efficiencies after ASD treatment with the same substrate. In the present study, soil prokaryotic communities in the field, which indicate difference in disinfestation effects after ASD treatment among several fields, were compared using next-generation sequencing. The prokaryotic communities in the fields showing different disinfestation effects were roughly separated into sampling fields. The relative abundances of *Betaproteobacteria* and *Clostridia* were significantly increased in well-disinfested fields. Overall, 25 operational taxonomic units (OTUs) were specifically increased in various well-disinfested soils and 18 OTUs belonged to phylogenetically diversified *Clostridia.* Other OTUs belonged to aerobic bacteria and were not previously detected in sample collected from ASD-treated fields. The results showed that the changes to the prokaryotic communities did not affect ASD efficiency, whereas changes in the abundance of specific microbes in the community were related to disinfestation.

Keywords: Bacterial wilt; Betaproteobacteria; Clostridia; Indicator species analysis; Multiple fields; Sugar-containing diatoms

1. Introduction

Soil-borne pathogens cause various plant diseases, including take-all, damping-off, crown rot, and wilting. Bacterial wilt caused by *Ralstonia solanacearum* has a host range exceeding 200 species from >50 families [1]. Soil disinfestation is challenging because this pathogen is distributed evenly at depths of >40 cm [2]. Several approaches have been attempted to control for bacterial wilt, including soil amendment, crop rotation, and field sanitation [3]. Although soil fumigation with chemical pesticides is an effective method for killing the pathogen causing bacterial wilt, the efficacy tends to be unstable in deep soil and the chemicals must escape to ensure food safety and prevent environmental pollution.

Anaerobic soil disinfestation (ASD) is an effective method to reduce the abundance of soil-borne pathogens [4]. This method comprises the incorporation of labile organic matter in the soil, irrigation, and covering the soil surface with polyethylene film. Organic matter increases microbial respiration, irrigation purges soil air, and polyethylene film prevents oxygen inflow from the atmosphere, which collectively induce reductive soil conditions [5,6]. Moreover, ASD using water-soluble organics, such as low-concentration ethanol or molasses as the carbon source, is effective for soil at depths of <40 cm [7]. Therefore, ASD using water-soluble carbon sources is suitable for the disinfection of *R. solanacearum* in deep-layer soils and is environmentally friendly. However, the disinfestation effects of ASD are unstable under field conditions [8]. A sufficient soil temperature, incubation period, and amount of carbon amendments are needed for the success of disinfestation.

Soil microbes reflect the soil environment and are considered an index of soil health [9,10]. ASD increases the abundance of several microbes that may be involved in the suppression of pathogens [11–16]. Therefore, fields with different disinfestation effects may have different soil microbial communities. Microbes that increase in abundance in well-disinfested soil that are commonly detected in several fields may be good candidate indicators for the efficiency of ASD treatment. The aim of the present study was to compare the prokaryotic communities among soils obtained from 17 geographically different greenhouses that had different disinfestation efficiencies after ASD treatment with the same substrate. The results of this multi-fields study showed that the soil microbial communities differed with the disinfestation efficiency of particular field and the well-disinfested fields had unique soil microbes, as compared with those not well-disinfested.

2. Materials and Methods

2.1. Sampling field and ASD treatment

Field experiments were established in 17 greenhouses situated on 8 fields in Japan that experienced bacterial wilt (Table 1).

Table 1. Characteristics of the sampled fields

Table 1. Detailed information on the sampling fields

Field	Ishikawa	Hakusan	Niigata	Tsubame	Sanjo	Toyama	Gifu	Wakayama
Code	Is	Ha	Ni	Ts	Sa	To	Gi	Wa
latitude and longitude	36° 38′ N, 136° 42′ E	36° 29′ N, 136° 30′ E	37° 55′ N, 139° 9′ E	37° 38′ N, 138° 55′ E	37° 39′ N, 139° 57′ E	36° 61′ N, 136° 96′ E	35° 44′ N, 136° 70′ E	33° 27′ N, 135° 14′ E
Soil types	Gleysol	Gleysol	Fluvisol	Fluvisol	Fluvisol	Gleysol	Gleysol	Cambisols
Replication	1	3	3	3	2	3	1	1
pH	5.54	7.22	7.11	7.03	7.24	6.07	6.08	7.49
EC (mS cm-1)	0.03	0.31	0.32	0.24	0.20	0.29	0.12	0.88
ASD conducted schedule	7/27-8/18	7/11-7/28	8/18-9/9	7/31-8/21	9/1-9/22	8/2-8/26	8/11-9/1	7/29-8/19

Sugar-containing diatoms are discharged from food-processing facilities as by-products of the
filtration of saccharified liquids. The main components of such by-products are sugars derived from
the saccharified solution of tapioca starch and diatoms used as a filtering aid. These by-products,
containing 40% by weight, were powdered and mixed into the soil with a rototiller at a ratio of 15 t
ha^{-1} (approximately 6.0 g carbon kg soil^{-1}) at a depth 30 cm. Thereafter, the field was covered with
transparent polyethylene film (thickness, 0.1 mm) and flooded with more than approximately 150 L
of water m^{-2}. Each site was flooded at the time of disinfestation, and no irrigation was conducted
afterward. Disinfestation was conducted for 21 days with the exception of field Ha (17 days)
because the soil temperature of this field was >35°C during disinfestation. Each greenhouse (15 × 6
m) was subjected to ASD treatment. There were three replicates from fields Ha, Ni, Ts, and To; two
from the field Sa; and none from fields Is, Gi, and Wa. Soil samples were collected from each
greenhouse on the fields Is, Ha, To, Gi, and Wa before and after ASD treatment from two different
depths—20–30 and 40–50 cm—using a core sampler (Gauge Auger DIK-106B; Daiki Rika Kogyo Co.,
Ltd, Saitama, Japan); i.e., 9 greenhouses × 2 depths × 2 sampling times = 36 soil samples in total.
Only two soil samples (before and after ASD treatment in the upper layer soil) were collected from
each greenhouse on fields Ni, Ts, and Sa (8 greenhouse × 2 sampling times = 16 soil samples).
Overall, 52 soil samples were collected for analysis. Soil samples were collected from five randomly
chosen points in each greenhouse and were mixed well.

2.2. Quantification of R. solanacearum in the field

The most probable number–polymerase chain reaction (MPN–PCR) method, which is a
semi-quantitative *R. solanacearum* counting method [17], was conducted. Briefly, 10 g of soil was
eluted into cultivation buffer and the soil extract was diluted with buffer to 10-, 100-, and 1000-fold.
Each sample was incubated at 35°C for approximately 24 h. Thereafter, nested-PCR was performed
using the samples as templates. The primer pair phcA2981f (5′-TGGATATCGGGCTGGCAA-3′) and
phcA4741r (5′-CGCTTTTGCGCAAAGGGA-3′) was used in the first step of the PCR reaction and the
primer pair phcA3538f (5′-GTGCCACAGCATGTTCAGG-3′) and phcA4209r
(5′-CCTAAAGCGCTTGAGCTCG-3′) was used in the second step to target the *phcA*, which is key for
the appearance of wilt disease. The number of samples that resulted in PCR products of ~700-bp
length was collectively checked against the MPN table. The detection limit of *R. solanacearum* using
this method is 3–2400 colony forming units g^{-1} dry soil and a detection indicates the risk of
outbreaks of bacterial wilt in the field.
DNA was extracted from 0.5 g of soil using the ISOIL for bead beating kit (Nippon Gene Co.,
Ltd., Tokyo, Japan), according to the manufacturer's instructions. DNA quantification and integrity
were measured using a NanoDrop spectrophotometer (Thermo Fisher Scientific, Waltham, MA,
USA) and gel visualization (0.8% agarose in Tris/acetic acid/ethylenediaminetetraacetic acid buffer),
respectively. The V4 region of the 16S rRNA gene of each sample was amplified by PCR using the
bacterial and archaeal universal primers 515F (5′-GTGCCAGCMGCCGCGGTAA-3′) and 806R
(5′-GGAC-TACVSGGGTATCTAA-3′) [18]. A library was prepared by adaptor ligation with the PCR
primer pairs using the TruSeq Nano DNA Library Prep Kit (Illumina, Inc., San Diego, CA, USA).
When two or more bands were detected using 1.5%-agarose gel electrophoresis, PCR products of
approximately 300 bp in length were excised from the gel, non-specific amplicons were removed,
and the products were purified using a MonoFas DNA purification kit for prokaryotes (GL Sciences,
Inc., Tokyo, Japan). One soil sample (the upper layer of soil of To3 after ASD treatment) was not
amplified after PCR amplification and was excluded from analysis. Each PCR amplicon was cleaned
twice to remove the primers and short DNA fragments using the Agencourt AMPure XP system
(Beckman Coulter, Inc., Brea, CA, USA) and quantified using a Qubit Fluorometer (Invitrogen
Corporation, Carlsbad, CA, USA). Following successful amplification, the PCR products were

adjusted to equimolar concentrations and subjected to unidirectional pyrosequencing, which was
performed by Bioengineering Lab. Co., Ltd. (Kanagawa, Japan) using a MiSeq instrument (Illumina,
Inc.). Overall, 3,114,333 sequences were obtained from the 51 samples (Supplemental Table 1).
Sequencing data were deposited in the DNA Database of Japan Sequence Read Archive under the
accession number DRA006673.

2.4. Data analyses

The sequencing data were analyzed as previously described [19]. Raw FASTA files were
pre-processed with the Quantitative Insights into Microbial Ecology (QIIME) bioinformatics pipeline
[20]. Data from the read sequences, quality, flows, and ancillary metadata were analyzed using the
QIIME pipeline. Quality filtering consisted of discarding reads of <200 or >1000 bp in length,
excluding homopolymer runs of >6 bp and continuous ambiguous bases of >6 bp, and accepting one
barcode correction and two primer mismatches. Moreover, reads with a mean quality score of <25
were removed. Finally, singleton operational taxonomic units (OTUs) and chimeric sequences were
removed for statistical analyses. Denoising was performed using the built-in denoiser algorithm,
and chimera removal and OTU picking were accomplished using the USEARCH 61 sequence
analysis tool (http://www.drive5.com/usearch/download.html) and a pairwise identity percentage
of 0.97. Taxonomy assignment was performed using the Ribosomal Database Project naïve Bayesian
classifier with a minimum confidence of 0.8 against the Greengenes database (October 2012 release;
http://greengenes.secondgenome.com/) and the Basic Local Alignment Search Tool
(https://blast.ncbi.nlm.nih.gov/Blast.cgi). The richness and diversity of pyrotag-based datasets were
determined by OTU-based analysis using the phyloseq R package, version 1.7.24 [21]. The alpha
diversity within each individual sample was estimated using the non-parametric Shannon diversity
index. The Chao1 estimator was used to estimate the richness of each sample. A multivariate
analysis of the community structure and diversity of the pyrotag-based datasets was performed
using a weighted UniFrac dissimilarity matrix calculated in the QIIME, jackknifing (1000
reiterations) read abundance data at the deepest possible level (41,012 reads), and using
unconstrained ordination by a principal coordinate analysis (PCoA) for prokaryotes in each soil
layer. *K*-means clustering was used to verify the effects of the sampling fields and ASD treatment on
prokaryotic communities. Finally, an indicator value analysis was conducted using the indicspecies
R package [22] to identify OTUs associated with suppressive soil rather than conducive soil or vice
versa ($p < 0.05$). Two soil group patterns were created. One group was divided into samples from the
soil before and after ASD treatment. The other group was divided based on disinfestation efficiency.
The datasets were merged and the abundant OTUs that were associated with the pathogen-free soil
after ASD treatment were evaluated in each layer ($p < 0.001$). The number of random permutation
tests for the calculation of the indicator values was 999.

2.5. Construction of phylogenetic tree

A phylogenetic tree of clostridial members was constructed by a 1000-fold bootstrap analysis
using the neighbor-joining method with the Clustal W multiple sequence alignment algorithm
(version 2.0) [23] and NJ plot software [24]. The phylogenetic tree contained the 16S rRNA sequences
of several clusters of clostridial strains.

3. Results

3.1. Effectiveness of ASD treatment with sugar-containing diatoms

R. solanacearum in soil was not detected after ASD treatment in in the upper layers of 14 (82.4%)
of 17 greenhouses and lower layers of 7 (77.8%) of 9 greenhouses (Table 2). Therefore, ASD treatment
was appropriate in each field. These soils were defined as well-disinfested (disinfestation success
soil: S-soil). However, some fields (Is, Ha1, and Ha2) were not well disinfested (disinfestation failure
soil: F-soil).

Table 2. The number of *R. solanacearum* in each field before and after ASD treatment

	Soil layer	ASD treatment	IS	Ha1	Ha2	Ha3	Ni1	Ni2	Ni3	Ts1	Ts2	Ts3	San1	San2	To1	To2	To3	Gi	Wa
Number of *R.solanacearum* in soil (cfu soil^{-1})	Upper	Before	> 2400	1100	46	23	95	210	460	95	75	120	160	43	64	43	23	> 2400	> 2400
		After	75	21	15	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
	Lower	Before	240	1100	39	13	-	-	-	-	-	-	-	-	11	53	12	1100	1100
		After	3	21	11	N.D.	-	-	-	-	-	-	-	-	N.D.	N.D.	N.D.	N.D.	N.D.

3.2. Changes of prokaryotic communities after ASD treatment

The sequences were clustered into 74,539 OTUs at 97% similarity. Among the S- and F- soils, the DNA concentrations were significantly decreased in the upper layer of the S-soils and the lower layer of the F-soils (Figure 1, Supplemental Table 2).

Figure 1. Comparison of DNA concentrations and prokaryotic diversity and richness between the F- and S-soils. Asterisks represent a significant difference in the mean values before and after ASD treatment (Tukey's test, p < 0.05). F-soil: not well-disinfested soil, S-soil: well-disinfested soil

The Shannon index and Chao1 were significantly decreased in the S- and F-soils after ASD treatment of the upper layer. OTU numbers before and after soil sampling were significantly decreased in the F-soils regardless of the soil layer.

PCoAs based on weighted UniFrac analysis showed that the prokaryotic communities were roughly classified by sampling fields before and after ASD treatment (Figure 2).

Fig. 2A

Fig. 2B

Figure 2. A UniFrac-weighted principal component analysis of the soil prokaryotic communities in
the upper layer (A) and lower layer (B) of the field before and after ASD treatment. Closed circle:
S-soil before ASD treatment, Open circle: S-soil after ASD treatment, Closed triangle: F-soil before

ASD treatment, Open triangle: F-soil after ASD treatment. The clustering was conducted using
K-means analysis.

The microbes collected from the same field comprised the same cluster before and after ASD
treatment, despite the difference in ASD efficiency.
In the upper layer, field Ha contained one cluster regardless of ASD treatment and
disinfestation efficiency (Figure 2A). Despite separation on the basis of before and after ASD
treatment in fields Is, Wa, and Ts, the disinfestation efficiency had no effect. There were differences
in the microbial communities of the lower layer soils of fields Is, Ha, and Gi after ASD treatment;
however, no differences were detected between the S- and F-soils (Figure 2B). After ASD treatment,
a more than two-fold increase in the relative abundance of Firmicutes was observed in eight
greenhouses (Supplemental Table 3). In the upper layer soil, the relative abundance of Firmicutes
was significantly increased after ASD treatment, whereas that of Crenarchaeota was decreased
(Figure 3).
210

Figure 3. Comparison of the prokaryotic phyla with significant differences in abundance between the
S- and F-soils. Asterisks represent a significant difference in the mean value of before and after ASD
treatment (Tukey's test, $p < 0.05$). F-soil: not well-disinfested soil, S-soil: well-disinfested soil

The relative abundance of *Acidobacteria* and *Verrucomicrobia* in the upper layer of the S-soil
remained unchanged, whereas that of *Acidobacteria* and *Crenarchaeota* in the lower layer of the F-soil
was significantly decreased. At the class level, a more than two-fold increase in the relative
abundance of *Clostridia* of the phylum Firmicutes was observed after ASD treatment in the
upper and lower layers of 15 (93.8%) of 16 and 8 (88.9%) of nine greenhouses, respectively
(Supplemental Table 4). Moreover, the relative abundances of *Betaproteobacteria* and *Clostridia*
were significantly increased after ASD treatment in the upper layer of the S-soil (Figure 4). In
the S-soil, *Sphingobacteria* and *Cytophagia* were significantly decreased in the upper layer, and
Thermoleophilia and *Actinobacteria* were significantly decreased in the lower layer.

Figure 4. Comparison of the prokaryotic class with significant differences in abundance between the S- and F-soils. Asterisks represent a significant difference in the mean values before and after ASD treatment (Tukey's test, p < 0.05). F-soil: not well-disinfested soil, S-soil: well-disinfested soil

3.4. Specific increased OTUs in various S-soils

Indicspecies analysis detected 25 OTUs from the upper and lower layers that were specifically increased in the S-soil after ASD treatment in >8 (50%) of 16 sites and 4 (66.7%) of 6 sites, respectively (Table 3).

Table 3. Shared prokaryotic OTUs with significantly increased abundance in the S-soils of different fields

OTU number	Upper	Lower	Upper + Lower	Closest relatives	Class	Identity
OTU50	11	5	16	*Caloramator sp. NST1-2*	Clostridia	88%
OTU199301	10	4	14	*Christensenella sp. strain 2NS-PRS3-s1*	Clostridia	95%
OTU311169	11	3	14	*Bacillus sp. strain FJAT-25547*	Bacillus	86%
OTU624373	10	3	13	*Christensenella sp. strain 2NS-PRS3-s1*	Clostridia	95%
OTU112	10	3	13	*Clostridium cellobioparum strain JCM 1422*	Clostridia	93%
OTU999898	9	3	12	*Acetivibrio cellulolyticus strain HL-2*	Clostridia	94%
OTU113750	9	3	12	*Ruminococcaceae bacterium HZ254R*	Clostridia	99%
OTU572575	9	3	12	*Clostridium sp. 6-31*	Clostridia	90%
OTU151400	8	4	12	*Ruminococcaceae bacterium Otu096*	Clostridia	91%
OTU56	10	2	12	*Rhodothermaceae bacterium strain MEBiC09517*	Bacteroidetes	89%
OTU107205	8	4	12	*Gracilibacter thermotolerans strain JW/YJL-S1*	Clostridia	91%
OTU445172	9	3	12	*Moorella humiferrea strain 64-FGQ*	Clostridia	89%
OTU184099	8	4	12	*Carboxydocella manganica strain SLM 61*	Clostridia	88%
OTU568807	9	2	11	*Thiomicrospira sp. SL-1*	Gammaproteobacteria	87%
OTU533198	8	3	11	*Noviherbaspirillum agri strain K-1-15*	Betaproteobacteria	98%
OTU273540	8	3	11	*Desulfotomaculum sp. MP104*	Clostridia	91%
OTU212333	6	5	11	*Symbiobacterium thermophilum strain IAM 14863*	Clostridia	92%
OTU365547	8	3	11	*Dendrosporobacter sp. LW3.5*	Negativicutes	94%
OTU88528	8	3	11	*Acholeplasmatales bacterium canine oral taxon 375*	Mollicutes	86%
OTU304	6	5	11	*Caloramator sp. strain Qq2*	Clostridia	91%
OTU141173	8	2	10	*Mangrovibacterium diazotrophicum strain SCSIO N0430*	Bacteroidetes	89%
OTU221127	8	2	10	*Peptococcaceae bacterium DRI-13*	Clostridia	93%
OTU934369	4	6	10	*Caloramator mitchellensis strain VF08*	Clostridia	99%
OTU395236	8	2	10	*Clostridium cellulosi*	Clostridia	85%
OTU539	8	2	10	*Bacillus boroniphilus strain S8-01*	Bacillus	86%

The number indicates the field number where specific OTUs were detected.

The identified microbes belonged to several classes, including Clostridia, Bacillus, Betaproteobacteria, Gammaproteobacteria, and Mollicutes. Clostridia accounted for 18 (72%) of the 25 OTUs. In a phylogenetic tree of OTUs related to clostridial members, various Clostridia species were found to be specifically increased after ASD treatment in the S-soil (Figure 5).

Figure 5. Neighbor-joining tree showing the phylogenetic relationships of the clostridial OTUs based on 16S rRNA gene sequencing (According to the clostridial cluster analysis [25]). OTUs obtained from this study are shown in gothic letters. Bacillus subtilis was used as the outgroup. Gothic letters indicate OTUs that specifically appeared in the S-soil.

Three OTUs belonged to the *Caloramator* group, *Clostridium* cluster III, whereas two OTUs
belonged to the *Christensenella* group. OTU 50 (*Caloramator*) was commonly detected in 11 upper
layer and 5 lower layer soil samples. Furthermore, OTU311169 (*Bacillus*), OTU199301 and
OTU624373 (*Christensenella*), OTU 112 (*Clostridium*), and OTU56 (*Rhodothermaceae*) were increased in
10 (71.4%) of the 14 S-soils, whereas OTU934369 and OTU304 (*Caloramator*) and OTU212333
(*Symbiobacterium*) were commonly detected in >5 of the 6 fields of S-soils. On the other hand, no
prokaryotes reduced by ASD were found to be common in more than half of the fields (data not
shown).

4. Discussion

The results of the present study showed no significant differences in the abundances of
microbial communities among fields with different disinfestation effects. In our previous study,
prokaryotic communities were more affected by indigenous communities than the disease
suppression effect [19]. The soil from a specific site typically contains microbial communities of
greater similarity irrespective of soil treatment [26]. In the Ha field, three contiguous greenhouses
received the same field management regimen for >10 years. Although there was a difference in
the disinfestation effects, these samples were classified into the same cluster. These results
suggest that the differences among indigenous soil communities are unaffected by controlling
soil-borne pathogens.
Several OTUs that mainly belonged to *Clostridia* were specifically increased in several S-soils
despite differences in indigenous prokaryotic communities. *Clostridia* are gram-positive anaerobic
bacteria. Several studies have reported that the abundance of Clostridia species increases and it
becomes the dominant bacteria after ASD treatment using plant material or low-concentration
ethanol as a substrate [13,14,27]. During ASD, decreased oxygen utilization promotes the increased
prevalence of anaerobic microbes [6,12,27]. Effective ASD requires sufficient soil reduction [28].
Therefore, well-disinfested soil is a good reduction condition, which may promote the growth of
several anaerobic bacteria, such as *Clostridia*. Moreover, some *Clostridia* can produce spores that are
resistant to high temperatures during the disinfestation period. Therefore, environmental changes
might have promoted the increased abundance of Clostridia, some of which play important roles in
the suppression of pathogen growth and proliferation. *Caloramator* are rod-shaped, obligate
anaerobic, thermophilic endospore-producers that ferment several sugars and produce acetate,
isobutyrate, lactate, and other volatile fatty acids (VFAs) but do not degrade cellulose [25,29]. More
than half of clostridial clones that were increased after ASD treatment belonged to Clostridium
cluster I, which included the *Oxobacter* and *Caloramator* groups [30]. *Symbiobacterium* spp. is a
rod-shaped thermophiles that are syntrophic with *Bacillus* spp.; which possess a glucose degradation
pathway [31]. *Christensenella* is a non-spore forming, short, straight rod with tapered ends and can
use various sugars and produce VFAs during fermentation [32]. These results suggest that
phylogenetically diverse Clostridia members might be responsible for the suppression of pathogens
via the production of VFAs during the anaerobic decomposition of sugar-containing diatoms.
Moreover, other bacteria were increased after ASD in several S-soils. For example,
Noviherbaspirillum, belonging to the class Betaproteobacteria, were specifically detected in the S-soils.
Noviherbaspirillum are gram-negative, aerobic, non-spore forming rods capable of using several
carbohydrates as carbon sources [33]. Betaproteobacteria increased after ASD treatment with wheat
bran application in soil depths of 15.2 and 45.7 cm [34]. Moreover, several *Bacillus* species have been
described as biological control agents against bacterial wilt and have been detected in soil after ASD
treatment [30,35,36]. Cytophagaceae bacteria were significantly increased after ASD treatment with
rice bran [16]. Although not previously detected in ASD-treated soil, *Symbiobacterium*, *Christensenella*,
Noviherbaspirillum, and *Rhodothermaceae* might play important roles in the disinfestation of *R.*
solanacearum. *Noviherbaspirillum*, *Bacillus*, and *Rhodothermaceae* are aerobic bacteria, but were
increased by ASD, just as the anaerobic bacteria. In the soil environment, some *Bacillus* might
contribute to the rapid decrease of the soil redox potential at the initial stage of ASD treatment via
the consumption of oxygen [30]. Therefore, they might have important roles in maintaining anoxic

soil conditions during ASD treatment. The abundances of Bacteroidetes, Acidobacteria, Planctomycetes, and Gammaproteobacteria increased after ASD treatment and some have been associated with the improvement of crop yields and suppression of plant diseases [26,37–39]. The fungi of the genus *Zopfiella*, which have been found to cause disease after ASD treatment, can prevent damping-off disease in cucumber [40]. Therefore, the aerobes detected in this study have the potential to directly suppress the growth and proliferation of plant pathogens.

The results of this study showed that *Clostridia* members were increased after effective ASD treatment in several fields. Moreover, some may be involved in the suppression of pathogen proliferation during ASD treatment. However, the killing mechanism of pathogens by ASD is not well -known. Some studies have shown that VFAs or ferrous iron can directly kill pathogens [15,41]. Future isolation studies are required to clarify the roles of the microbes detected in this study. These microbes may be involved in the suppression of pathogen proliferation; therefore, they could be good indicators of successful disinfestation by ASD. By revealing the ecology, there may be several strategies to increase the abundances of beneficial microbes.

In conclusion, prokaryotic communities were not strongly affected by the disinfestation efficiency among the sampling fields. However, the relative abundance of *Clostridia* and *Betaproteobacteria* were significantly increased in well-disinfested soil. Further, 25 OTUs were specifically detected in S-soils and most were affiliated to *Clostridia*, which is well-known to promote the effects of ASD. However, some microbes were not previously detected in ASD-treated fields. These microbes present good indicators of well-disinfested soil. Nonetheless, future studies are required to further elucidate the role of these microbes in disinfestation.

Funding Information

This work was supported by the Cabinet Office, Government of Japan, Cross-ministerial Strategic Innovation Promotion Program (SIP), and the "Technologies for creating next-generation agriculture, forestry and fisheries" (funding agency: Bio-oriented Technology Research Advancement Institution, NARO). This work was also partially supported by the RIKEN Competitive Program for Creative Science and Technology, a Grant-in-Aid for Scientific Research from the Japan Society for Promotion of Science, No. 17K01447 (to M.O.), and by the Sasakawa Scientific Research Grant from The Japan Science Society (to C.G.L.). We would like to thank Editage (www.editage.jp) for English language editing.

Supplementary Materials: The following are available online at www.mdpi.com/link, Table S1: Read numbers of the 16S rRNA gene in each soil sample, Table S2: The ratio of the DNA concentrations, prokaryotic diversity, and richness before to those after ASD treatment, Table S3: The ratio of the prokaryotic phylum before to that after ASD treatment, Table S4: The ratio of the prokaryotic class before to that after ASD treatment.

Acknowledgments: We would like to thank Enago (www.enago.jp) for the English language editing.

Author Contributions: All authors contributed to the intellectual input and provided assistance to this study. Each author contributed to conducting the ASD in each prefecture (E. M. and K. Y. in Ishikawa, M. K. in Toyama, M. M. in Niigata, Y. M. and H. W. in Gifu, and Y. O. in Wakayama prefecture). T. I., K. N., and M. O. have supervised the work and approved the manuscript for publication.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Kelman, A. One hundred and one years of research on bacterial wilt. In *Bacterial Wilt Disease: Molecular and Ecological Aspects*; Prior, P., Allen, C., Elphinstone, J., Eds.; Springer Berlin Heidelberg: Berlin, Heidelberg, 1998; pp. 1–5 ISBN 978-3-662-03592-4.

2. Graham, J.; Lloyd, A.B. Survival of potato strain (race 3) of *Pseudomonas*
solanacearum in the deeper soil layers. *Aust. J. Agric. Res.* **1979**, *30*, 489–496.
https://www.publish.csiro.au/cp/AR9790489 (access on 15 May 2019).
3. Weller, D.M. Biological control of soilborne plant pathogens in the rhizosphere
with Bacteria. *Annu. Rev. Phytopathol.* **1988**, *26*, 379–407.
https://doi.org/10.1146/annurev.py.26.090188.002115 (access on 15 May 2019).
4. Shinmura, A. Causal agent and control of root rot of welsh onion. In
Proceedings of the PSJ soil-borne disease workshop report; 2000; Vol. 20, pp.
133–143.
5. Blok, W.J.; Lamers, J.G.; Termorshuizen, A.J.; Bollen, G.J. Control of soilborne
plant pathogens by incorporating fresh organic amendments followed by
tarping. *Phytopathology* **2000**, *90*, 253–259.
https://www.ncbi.nlm.nih.gov/pubmed/18944617 (access on 15 May 2019).
6. Momma, N.; Yamamoto, K.; Simandi, P.; Shishido, M. Role of organic acids in
the mechanisms of biological soil disinfestation (BSD). *J. Gen. Plant Pathol.* **2006**,
72, 247–252. https://link.springer.com/article/10.1007/s10327-006-0274-z (access
on 15 May 2019).
7. Momma, N.; Kobara, Y.; Uematsu, S.; Kita, N.; Shinmura, A. Development of
biological soil disinfestations in Japan. *Appl. Microbiol. Biotechnol.* **2013**, *97*,
3801–3809. https://www.ncbi.nlm.nih.gov/pubmed/23549745 (access on 15
May 2019).
8. Shrestha, U.; Augé, R.M.; Butler, D.M. A meta-analysis of the impact of
anaerobic soil disinfestation on pest suppression and yield of horticultural
crops. *Front. Plant Sci.* **2016**, *7*.
https://www.ncbi.nlm.nih.gov/pubmed/27617017 (access on 15 May 2019).
9. van Bruggen, A.H.C.; Semenov, A.M. In search of biological indicators for soil
health and disease suppression. *Appl. Soil Ecol.* **2000**, *15*, 13–24.
https://www.sciencedirect.com/science/article/pii/S0929139300000688 (access
on 15 May 2019).
10. Mendes, R.; Kruijt, M.; de Bruijn, I.; Dekkers, E.; van der Voort, M.; Schneider,
J.H.M.; Piceno, Y.M.; DeSantis, T.Z.; Andersen, G.L.; Bakker, P.A.H.M.; et al.
Deciphering the rhizosphere microbiome for disease-suppressive bacteria.
Science **2011**, *332*, 1097–1100. https://www.ncbi.nlm.nih.gov/pubmed/21551032
(access on 15 May 2019).
11. Momma, N.; Usami, T.; Shishido, M. Detection of *Clostridium* sp. inducing
biological soil disinfestation (BSD) and suppression of pathogens causing
fusarium wilt and bacterial wilt of tomato (*Lycopersicon esculentum*) by gases
evolved during BSD. *Soil Microorg. (Japan)* **2007**.
https://ci.nii.ac.jp/naid/110009468481 (access on 15 May 2019).
12. Mowlick, S.; Takehara, T.; Kaku, N.; Ueki, K.; Ueki, A. Proliferation of
diversified clostridial species during biological soil disinfestation incorporated
with plant biomass under various conditions. *Appl. Microbiol. Biotechnol.* **2013**,
97, 8365–8379.

https://www.sciencedirect.com/science/article/pii/S0929139318301422 (access
on 15 May 2019)
13. Huang, X.; Liu, L.; Wen, T.; Zhang, J.; Wang, F.; Cai, Z. Changes in the soil
microbial community after reductive soil disinfestation and cucumber seedling
cultivation. *Appl. Microbiol. Biotechnol.* **2016**, *100*, 5581–5593.
https://www.ncbi.nlm.nih.gov/pubmed/26875875 (access on 15 May 2019).
14. Huang, X.; Liu, L.; Wen, T.; Zhu, R.; Zhang, J.; Cai, Z. Illumina MiSeq
investigations on the changes of microbial community in the *Fusarium*
oxysporum f.sp. cubense infected soil during and after reductive soil
disinfestation. *Microbiol. Res.* **2015**, *181*, 33–42.
https://www.ncbi.nlm.nih.gov/pubmed/26640050 (access on 15 May 2019).
15. Strauss, S.L.; Kluepfel, D.A. Anaerobic soil disinfestation: A
chemical-independent approach to pre-plant control of plant pathogens. *J.*
Integr. Agr. **2015**, *14*, 2309–2318.
https://www.sciencedirect.com/science/article/pii/S2095311915611182 (access
on 15 May 2019).
16. Mazzola, M.; Hewavitharana, S.S.; Strauss, S.L.; Shennan, C.; Muramoto, J.
Anaerobic soil disinfestation and *Brassica* seed meal amendment alter soil
microbiology and system resistance. *Int. J. Fruit Sci* **2016**, *16*, 47–58.
https://www.tandfonline.com/doi/abs/10.1080/15538362.2016.1195310 (access
on 15 May 2019)
17. Inoue, Y.; Nakaho, K. Sensitive quantitative detection of Ralstonia
solanacearum in soil by the most probable number-polymerase chain reaction
(MPN-PCR) method. *Appl. Microbiol. Biotechnol.* **2014**, *98*, 4169–4177.
https://www.ncbi.nlm.nih.gov/pubmed/24584461 (access on 15 May 2019).
18. Caporaso, J.G.; Lauber, C.L.; Walters, W.A.; Berg-Lyons, D.; Lozupone, C.A.;
Turnbaugh, P.J.; Fierer, N.; Knight, R. Global patterns of 16S rRNA diversity at
a depth of millions of sequences per sample. *Proc. Natl. Acad. Sci. U. S. A.* **2011**,
108, 4516–4522. https://www.ncbi.nlm.nih.gov/pubmed/20534432 (access on
15 May 2019).
19. Lee, C.G.; Iida, T.; Uwagaki, Y.; Otani, Y.; Nakaho, K.; Ohkuma, M.
Comparison of prokaryotic and eukaryotic communities in soil samples with
and without tomato bacterial wilt collected from different fields. *Microb.*
Environ. **2017**, *32*, 376–385.
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5745023/ (access on 15 May
2019).
20. Caporaso, J.G.; Kuczynski, J.; Stombaugh, J.; Bittinger, K.; Bushman, F.D.;
Costello, E.K.; Fierer, N.; Pena, A.G.; Goodrich, J.K.; Gordon, J.I. QIIME allows
analysis of high-throughput community sequencing data. *Nature Methods* **2010**,
7, 335. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3156573/ (access on 15
May 2019).

21. McMurdie, P.J.; Holmes, S. phyloseq: an R package for reproducible interactive
analysis and graphics of microbiome census data. *PloS one* **2013**, *8*, e61217.
https://www.ncbi.nlm.nih.gov/pubmed/23630581 (access on 15 May 2019)
22. Cáceres, M.D.; Legendre, P. Associations between species and groups of sites:
indices and statistical inference. *Ecology* **2009**, *90*, 3566–3574.
https://esajournals.onlinelibrary.wiley.com/doi/full/10.1890/08-1823.1 (access
on 15 May 2019).
23. Larkin, M.A.; Blackshields, G.; Brown, N.P.; Chenna, R.; McGettigan, P.A.;
McWilliam, H.; Valentin, F.; Wallace, I.M.; Wilm, A.; Lopez, R.; et al. Clustal W
and Clustal X version 2.0. *Bioinformatics* **2007**, *23*, 2947–2948.
https://www.ncbi.nlm.nih.gov/pubmed/17846036 (access on 15 May 2019).
24. Perrière, G.; Gouy, M. WWW-Query: an online retrieval system for biological
sequence banks. *Biochime.* **1996,** *78*, 364-369.
https://www.ncbi.nlm.nih.gov/pubmed/8905155   (access on 15 May 2019).
25. Collins, M.D.; Lawson, P.A.; Willems, A.; Cordoba, J.J.; Fernandez-Garayzabal,
J.; Garcia, P.; Cai, J.; Hippe, H.; Farrow, J.A. The phylogeny of the genus
Clostridium: proposal of five new genera and eleven new species combinations.
Int. J. Syst. Bacteriol. **1994**, *44*, 812–826.
https://www.ncbi.nlm.nih.gov/pubmed/7981107 (access on 15 May 2019).
26. Mazzola, M.; Muramoto, J.; Shennan, C. Anaerobic disinfestation induced
changes to the soil microbiome, disease incidence and strawberry fruit yields
in California field trials. *Appl. Soil Ecol.* **2018**, *127*, 74–86.
https://www.sciencedirect.com/science/article/pii/S0929139318301422 (access
on 15 May 2019)
27. Mowlick, S.; Yasukawa, H.; Inoue, T.; Takehara, T.; Kaku, N.; Ueki, K.; Ueki, A.
Suppression of spinach wilt disease by biological soil disinfestation
incorporated with *Brassica juncea* plants in association with changes in soil
bacterial communities. *Crop Protect.* **2013**, *54*, 185–193.
https://www.sciencedirect.com/science/article/pii/S0261219413002111   (access
on 15 May 2019).
28. Shennan, C.; Muramoto, J.; Koike, S.; Baird, G.; Fennimore, S.; Samtani, J.;
Bolda, M.; Dara, S.; Daugovish, O.; Lazarovits, G. Anaerobic soil disinfestation
is an alternative to soil fumigation for control of some soilborne pathogens in
strawberry production. *Plant Pathol.* **2018**, *67*, 51–66.
29. Ogg, C.D.; Patel, B.K.C. Caloramator mitchellensis sp. nov., a thermoanaerobe
isolated from the geothermal waters of the Great Artesian Basin of Australia,
and emended description of the genus *Caloramator. Int. J. Syst. Evol. Microbiol.*
2011, *61*, 644–653. https://www.ncbi.nlm.nih.gov/pubmed/20400665   (access on
15 May 2019).
30. Mowlick, S.; Hirota, K.; Takehara, T.; Kaku, N.; Ueki, K.; Ueki, A. Development
of anaerobic bacterial community consisted of diverse clostridial species
during biological soil disinfestation amended with plant biomass. *Soil Sci. Plant*
Nutr. **2012**, *58*, 273–287.

https://www.tandfonline.com/doi/abs/10.1080/00380768.2012.682045 (access on 15 May 2019).

31. Ohno, M.; Shiratori, H.; Park, M.J.; Saitoh, Y.; Kumon, Y.; Yamashita, N.; Hirata, A.; Nishida, H.; Ueda, K.; Beppu, T. *Symbiobacterium thermophilum* gen. nov., sp. nov., a symbiotic thermophile that depends on co-culture with a *Bacillus* strain for growth. *Int. J. Syst. Evol. Microbiol.* **2000**, *50 Pt 5*, 1829–1832. https://www.ncbi.nlm.nih.gov/pubmed/11034494 (access on 15 May 2019).

32. Morotomi, M.; Nagai, F.; Watanabe, Y. Description of *Christensenella minuta* gen. nov., sp. nov., isolated from human faeces, which forms a distinct branch in the order *Clostridiales*, and proposal of *Christensenellaceae* fam. nov. *Int. J. Syst. Microbiol.* **2012**, *62*, 144–149. https://www.ncbi.nlm.nih.gov/pubmed/21357455 (access on 15 May 2019).

33. Lin, S.-Y.; Hameed, A.; Arun, A.B.; Liu, Y.-C.; Hsu, Y.-H.; Lai, W.-A.; Rekha, P.D.; Young, C.-C. Description of *Noviherbaspirillum malthae* gen. nov., sp. nov., isolated from an oil-contaminated soil, and proposal to reclassify *Herbaspirillum soli*, *Herbaspirillum aurantiacum*, *Herbaspirillum canariense* and *Herbaspirillum psychrotolerans* as *Noviherbaspirillum soli* comb. nov., *Noviherbaspirillum aurantiacum* comb. nov., *Noviherbaspirillum canariense* comb. nov. and *Noviherbaspirillum psychrotolerans* comb. nov. based on polyphasic analysis. *Int. J. Syst. Microbiol.* **2013**, *63*, 4100–4107. https://www.ncbi.nlm.nih.gov/pubmed/23749276 (access on 15 May 2019).

34. Strauss, S.L.; Greenhut, R.F.; McClean, A.E.; Kluepfel, D.A. Effect of anaerobic soil disinfestation on the bacterial community and key soilborne phytopathogenic agents under walnut tree-crop nursery conditions. *Plant Soil* **2017**, *415*, 493–506. https://link.springer.com/article/10.1007/s11104-016-3126-4 (access on 15 May 2019).

35. Yamamoto, S.; Shiraishi, S.; Kawagoe, Y.; Mochizuki, M.; Suzuki, S. Impact of *Bacillus amyloliquefaciens* S13-3 on control of bacterial wilt and powdery mildew in tomato. *Pest Manag. Sci.* **2015**, *71*, 722–727. https://www.ncbi.nlm.nih.gov/pubmed/24889125 (access on 15 May 2019).

36. Yuliar; Nion, Y.A.; Toyota, K. Recent trends in control methods for bacterial wilt diseases caused by *Ralstonia solanacearum*. *Microb. Environ.* **2015**, *30*, 1–11. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4356456/ (access on 15 May 2019).

37. Meng, T.; Yang, Y.; Cai, Z.; Ma, Y. The control of *Fusarium oxysporum* in soil treated with organic material under anaerobic condition is affected by liming and sulfate content. *Biol. Fertil. Soils* **2018**, *54*, 295–307. https://link.springer.com/article/10.1007/s00374-017-1260-7 (access on 01 July 2019).

38. van Agtmaal, M.; van Os, G.J.; Hol, W.H.G.; Hundscheid, M.P.J.; Runia, W.T.; Hordijk, C.A.; de Boer, W. Legacy effects of anaerobic soil disinfestation on soil bacterial community composition and production of pathogen-suppressing

513  volatiles. *Front. Microbiol.* **2015**, *6*.
514  https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4498103/ (access on 01 July
515  2019).
516 39. Hewavitharana, S.S.; Ruddell, D.; Mazzola, M. Carbon source-dependent
517  antifungal and nematicidal volatiles derived during anaerobic soil
518  disinfestation. *Eur. J. Plant Pathol.* **2014**, *140*, 39–52.
519  https://link.springer.com/article/10.1007/s10658-014-0442-5 (access on 01 July
520  2019).
521 40. Liu, L.; Huang, X.; Zhao, J.; Zhang, J.; Cai, Z. Characterizing the key agents in a
522  disease-suppressed soil managed by reductive soil disinfestation. *Appl. Environ.*
523  *Microbiol.* **2019**, *85*, e02992-18. https://aem.asm.org/content/85/7/e02992-18
524  (access on 01 July 2019).
525 41. Momma, N. Biological Soil Disinfestation (BSD) of soilborne pathogens and its
526  possible mechanisms. *JARQ* **2008**, *42*, 7–12.
527  https://www.jstage.jst.go.jp/article/jarq/42/1/42_7/_article (access on 01 July
528  2019).
529
530