Insights from Fisher's geometric model on the likelihood of speciation under different

histories of environmental change

Ryo Yamaguchi[1, 2] and Sarah P. Otto[3]

[1]Department of Advanced Transdisciplinary Sciences, Hokkaido University, Sapporo,

Hokkaido, Japan

[2]Department of Biological Sciences, Tokyo Metropolitan University, Hachioji, Tokyo, Japan

[3]Department of Zoology & Biodiversity Research Centre, University of British Columbia,

Vancouver, British Columbia, Canada, V6T 1Z4

Corresponding author: Ryo Yamaguchi

Email: ryamaguchi@sci.hokudai.ac.jp

Abstract

The formation of new species via the accumulation of incompatible genetic changes is thought
to result either from ecologically-based divergent natural selection or the order by which
mutations happen to arise, leading to different evolutionary trajectories even under similar
selection pressures. There is growing evidence in support of both ecological speciation and
mutation-order speciation, but how different environmental scenarios affect the rate of species
formation remains underexplored. We use a simple model of optimizing selection on multiple
traits ("Fisher's geometric model") to determine the conditions that generate genetic
incompatibilities in a changing environment. We find that incompatibilities are likely to
accumulate in isolated populations adapting to different environments, consistent with
ecological speciation. Incompatibilities also arise when isolated populations face a similar
novel environment; these cases of mutation-order speciation are particularly likely when the
environment changes rapidly and favors the accumulation of large-effect mutations. In addition,
we find that homoploid hybrid speciation is likely to occur either when new environments arise
in between the parental environments or when parental populations have accumulated large-
effect mutations following a period of rapid adaptation. Our results indicate that periods of
rapid environmental change are particularly conducive to speciation, especially mutation-order
or hybrid speciation.

35
36

37 **Keywords**

38 Ecological speciation, mutation-order speciation, homoploid hybrid speciation, Fisher's
39 geometric model.

Introduction

According to the ecological theory of speciation, populations diverge when exposed to different

environmental conditions (Schluter 2000). Metaphorically, populations in different

environments are thought to evolve phenotypically across different fitness landscapes,

diverging over time as they approach different fitness peaks. This theory can be tested using

reciprocal transplant experiments in which the performance of phenotypically divergent

populations and their hybrid progeny are measured in both parental environments. The results

of such studies often indicate that the environment matters, with phenotypic differentiation and

the performance of hybrids depending on the environment (Table 5.1 in Schluter 2000).

An alternative theory of speciation – "mutation-order speciation" – proposes that

populations diverging under similar selection pressures may ultimately become reproductively

isolated because they accumulate different sets of mutations, some of which are incompatible

with one another (Mani and Clarke 1990; Schluter 2009; Schluter and Conte 2009; Mendelson

et al. 2014). The focus here is on the reduction in fitness caused by combining genes that have

independently spread in different populations, so-called "Bateson-Dobzhansky-Muller"

incompatibilities (BDMI), regardless of the environment in which hybrids are assayed.

Empirical studies have suggested that both divergent selection and parallel

selection are relatively common (Table 5.2 in Schluter 2000), suggesting that both ecological

and mutation-order speciation may underlie the formation of new species in nature.

Ecological speciation is often inferred based on local adaptation of parental populations to

different abiotic (e.g., soil chemistry) or biotic conditions (e.g., pollinator communities) (see

Waser and Campbell 2004 for a review of evidence in plants). While such data provides

evidence for ecology-dependent selection, it does not prove that ecological differences

underly reproductive isolation, rather than intrinsic genetic incompatibilities (BDMIs).

Additional evidence for ecological speciation comes from reciprocal transplant experiments

where hybrids formed by backcrossing to the foreign parent are less fit than backcrosses to

the local parent (Rundle and Whitlock 2001), as seen in sticklebacks (Rundle 2002).

Alternative evidence for ecological speciation is provided when populations that have

independently adapted to a novel environment and that are isolated from their ancestral

populations remain interfertile with other such populations from similar environments

(Schluter 2000). Such "parallel ecological speciation" has been found in a number of animal

taxa, including sticklebacks and other fish, Phasmatodea walking-stick insects, and *Littorina*

snails, but evidence in plants remains weak (Ostevik *et al.* 2012).

Empirical evidence for mutation-order speciation includes cases of reciprocal gene

loss of duplicated genes in *Drosophila*, *Arabidopsis*, and yeast, where the loss of one gene

copy or the other is thought to have occurred under similar selection pressures, reducing

fitness in hybrids that lack the gene (see discussion and review of evidence by Muir and

Hahn 2015). Genetic conflict, such as meiotic drive or cytonuclear conflicts, has also been

shown to generate reproductive isolation in a manner consistent with mutation-order

speciation in a wide variety of animal and plant species (Crespi and Nosil 2013).

Experimental evolution studies in the lab have also provided evidence for both

ecological speciation and mutation-order speciation. In such studies, multiple independent

cultures of organisms are allowed to evolve under either different environmental conditions or

identical conditions, following which genetic incompatibilities are measured between the

replicate populations. Evidence for ecological speciation has been obtained in studies that have

adapted lines to different conditions. For example, Dettman *et al*. (2007) evolved multiple

populations of the yeast *Saccharomyces cerevisiae* in either rich media with high salt or in

minimal media with low glucose. In most replicates, hybrid diploids performed significantly

worse than the strain that had adapted to a particular local environment. Similarly, Dettman *et*

al. (2008) adapted *Neurospora crassa* to either high salinity or low temperature conditions,

finding evidence for the first stages of parallel ecological speciation. When crossed,

reproductive success was lower among hybrids formed from strains adapted to different

environments than among hybrids from strains adapted to the current environment being tested.

Evidence for mutation-order speciation includes an experimental evolution study by

Kvitek and Sherlock (2011) in *S. cerevisiae* involving multiple replicate populations

propagated under glucose limitation. Genomic sequencing revealed that mutations in two genes

(MTH1 and HXT6/HXT7) arose repeatedly in different lines but were never found together.

Subsequent crosses established that adaptive mutations at each of these two genes were

deleterious when combined, reducing the fitness of hybrids. Similarly, a study of *S. cerevisiae*

adapting in the presence of the fungicide nystatin found that even the first-step mutations that

arise during adaptation can be incompatible (Ono *et al*. 2017). When lines that had accumulated

different resistance mutations were crossed, the double mutants performed less well than single

mutants in many cases ("sign epistasis"). Sign epistasis was also observed in an experiment

with the bacterium, *Methylobacterium extorquens*, evolving to tolerate a genetically-

engineered change to how formaldehyde is oxidized (Chou *et al*. 2014). Again, adaptive

mutations tended to work poorly when combined.

Previous theoretical studies have modeled the fitness of recombinant hybrids under

divergent and parallel adaptation following a period of adaptation with *de novo* mutations

(Barton 2001; Chevin *et al*. 2014; Fraïsse *et al*. 2016). These models have clarified how

segregation of different adaptive alleles reduces hybrid fitness ("segregation load"), via the

separation of coadapted alleles. Using Fisher's geometric model to track fitness variability

among hybrids between independently adapting populations, Chevin *et al*. (2014) found that

segregation load (isolated from extrinsic, environment-dependent, fitness effects) is insensitive

to whether adaptation occurred to the same or to different environmental optima (e.g., Figure

2 in Chevin *et al*. 2014). This work is a launching point for the current study, where we explore

how the distribution of potential mutations and the speed of adaptation influence the mean

fitness of hybrids, combining the segregation load and the mismatch to the environment in

which hybrid and parental fitnesses are measured (i.e., both intrinsic and extrinsic

incompatibilities; their Equation 1, see also Equation 3 of Barton 2001). Fraïsse *et al*. (2016)

further found that Fisher's geometric model can generate many of the empirical patterns of

speciation studies, from Haldane's rule to asymmetrical introgression.

As noted in these two studies, large-effect mutations are more likely to contribute to

genetic incompatibilities. This prediction is consistent with the observation that

incompatibilities arise often in short-term experimental evolution studies, because those studies

typically induce strong selection favoring large-effect mutations. Large-effect mutations are

more likely to cause hybrid breakdown both because large-effect mutations can have large

pleiotropic side effects and because combining such mutants can overshoot the optimal fitness

(Figure 1). Our work thus aims to explore the environmental conditions that allow such strong

genetic incompatibilities to arise, measuring how overall reproductive isolation, including both

intrinsic and extrinsic isolation, depends on the speed and direction of environmental change,

as well as the average effect size of new mutations.

In another recent study, Simon *et al.* (2018) looked not at the processes generating

divergence between parental populations but at the fitness relationships between different

hybrid genotypes. Given a set of divergent alleles between the parents, hybrid fitness depends

on two main quantities, the hybrid index, H (the total proportion of alleles from one parent),

and heterozygosity, p_{12} (the proportion of divergent sites that carry one allele from each parent

population). Using a Brownian bridge approximation to walk from hybrids with only alleles

from one parent ($H = 0$) to hybrids with only alleles from the other parent ($H = 1$) and using

Fisher's geometrical model to describe fitness, the authors predict the average fitness of all

hybrid combinations (H, p_{12}) given remarkably few summary statistics (the breakdown scores

of the two parents and one additional fitted parameter; see their Equation 6), matching

empirical data sets well. Importantly, Simon *et al.* (2018) take the divergence between the

parents as a given (the number of diverged sites and the average trait variance contributed by

each site). By contrast, we focus here on how the history of environmental change influences

the genetic differences observed between the parents and the amount of hybrid breakdown (i.e.,

the fitted parameter in Simon *et al*. (2018)). In particular, we explore the conditions under which large effect mutations are likely to accumulate, leading to greater trait variance per locus and greater hybrid breakdown, even among parental populations experiencing the same environments. We then use the Brownian bridge approximation of Simon *et al.* (2018) to fit fitness data from a variety of hybrid types.

Specifically, using simulations, we assess how the history of adaptation to environmental change and the degree of environmental similarity between two populations impact the mean fitness of a variety of hybrid crosses (e.g., F_1, F_2, backcross to local, backcross to foreign parents) for both haploid and diploid species. We illustrate how the degree of reproductive isolation depends on the history of adaptation, the mean mutational size, and the cross considered. Our results emphasize how the nature of past selection – both the direction and the speed of selection in each population – has a substantial effect on the amount of hybrid breakdown predicted.

We also consider how the modularity of the genotype-phenotype relationship impacts the development of reproductive isolation under mutation-order speciation. Modularity refers to the observation that genes cluster according to which traits are affected by genetic perturbations. Such clustering is common, for example in the genotype-phenotype maps measured using large gene deletion collections in yeast, nematodes, and mice (Wang *et al*. 2010). Modularity may arise when genes in the same module are subject to similar regulatory networks or have phenotypic effects that are spatially or temporally confined within a particular tissue or cellular component. To vary modularity, we investigate a generalized form of Fisher's geometric model (FGM), in which the level of pleiotropy can be tuned. The original FGM assumes that a mutation affects all traits under consideration (i.e., universal pleiotropy; fitness landscape is a single hypersphere). Here, we also consider a "multi-sphere model," where each mutation affects only one module or cluster of traits (modular pleiotropy, Welch and Waxman 2003; Chevin *et al*. 2010; Wagner and Zhang 2011; Hill and Zhang 2012; Martin 2014). Given that we simulate a changing environment affecting the optimum in only one sphere, we predicted that there would be a higher fraction of compatible genetic changes in organisms with greater modularity, because the pleiotropic effects of adaptive mutations and any compensatory mutations would be restricted to the same subset of traits. Thus, we expected a higher degree of parallel evolution, leading to less reproductive isolation, in cases where the environmental change affected the same module. Unexpectedly we found a non-monotonic relationship between the extent of reproductive isolation and the degree of modularity, which we were able to ascribe to a counteracting tendency for mutations to arise more often in different modules, especially late in the divergence process when there are many modules.

Finally, we explore the potential for hybrids to colonize new sites beyond the range

of the parental species. Crosses between individuals from two adapted populations can produce

a wide variety of hybrid genotypes, potentially allowing hybrids to survive beyond the

environmental conditions in which parental genotypes can survive (Albertson and Kocher

2005). We thus explored the range of environments in which hybrids were more likely to

establish than parental individuals, potentially contributing to hybrid speciation (i.e., without a

change in ploidy: "homoploid hybrid speciation"; Rieseberg 1997).

Overall, our work explores how the nature of environmental change and the

modularity of the genetic architecture influence the development of reproductive isolation and

the potential for hybrid speciation.

Model

We first describe the fitness landscape in our model and then describe how we simulate the

evolution of reproductive isolation.

Fitness landscape

Fisher's model assumes multivariate stabilizing selection around a single optimum. We

extend this model by defining an individual's phenotype as m sets of n-dimensional vectors,

where m is the number of spheres and n is the number of traits in each sphere. Specifically,

the phenotype of an individual in the ith sphere is $\mathbf{z}_i = \left\{ z_{i,1}, z_{i,2}, ..., z_{i,n} \right\}$, whose components

$z_{i,j}$ are the phenotypic values for trait j; we denote the individual's phenotype across all

spheres as $\mathbf{Z} = \{ \mathbf{z_1}, \mathbf{z_2}, ..., \mathbf{z_m} \}$.

The fitness component based on the set of traits described by the ith sphere depends

on the distance of the phenotype from the optimum $\mathbf{o}_i = \left\{ o_{i,1}, o_{i,2}, ..., o_{i,n} \right\}$, measured as the

Euclidean distance $\|\mathbf{z}_i - \mathbf{o}_i\| \equiv \sqrt{\sum_{j=1}^{n} \left(z_{i,j} - o_{i,j} \right)^2}$. Assuming that an individual's relative fitness

is calculated by considering fitness across all spheres in a multiplicative fashion, fitness is

given by:

$$w(\mathbf{Z}) = \exp(-q \sum_{i=1}^{m} \|\mathbf{z}_i - \mathbf{o}_i\|^k), \qquad \text{(Eq. 1)}$$

where q sets the overall strength of selection and k determines the curvature of the fitness

function. The parameter k corresponds to the degree of epistasis (Tenaillon 2014). Throughout,

we assume that fitness declines from a peak height at the optimum, according to a Gaussian

fitness surface ($k = 2$). It is worth noting that selection is assumed independent and equally

strong on all $m \times n$ traits, for a given distance to the optimum (isotropic selection).

Each mutation additively shifts the phenotypic value relative to its present value. 214   While each mutation has an additive effect on the phenotype, the effect on fitness is not additive 215   but depends on the height of the fitness function for those phenotypes (Eq. 1). In the multi- 216   sphere model, we assume that a mutation affects one, and only one, sphere, chosen at random. 217   Once a sphere is chosen, the mutation affects all of the n traits in the sphere under consideration, 218   in a random direction (isotropic mutation). While the trait axes can be defined to allow isotropy 219   with respect to either selection or mutation (assuming a single optimum), it is not possible to 220   do so for both. Nevertheless, previous work exploring a non-isotropic FGM model has shown 221   that the isotropic FGM serves as a good approximation, in many cases, if we consider n to be 222   an effective number of dimensions (Waxman and Welch 2005; Martin and Lenormand 2006). 223   Martin (2014) also argued for an isotropic model with universal pleiotropy as an approximation 224   to a broader class of functions.

225   For both haploids and homozygous diploids, we denote the effect of a mutation

226   occurring within sphere i by $\Delta\mathbf{z}_i = \left\{\Delta z_{i,1}, \Delta z_{i,2}, ..., \Delta z_{i,n}\right\}$, with $\Delta z_{i,j}$ describing the effect on

227   trait j within that sphere. We assume that a mutation affects each trait within the sphere 228   independently, with $\Delta z_{i,j}$ proportional to a draw from a standard normal distribution, denoted 229   as ξ_j (Hartl and Taubes 1998; Fraïsse *et al.* 2016). The total effect size of the mutation was 230   then set to r, drawn from an exponential distribution with mean λ, adjusting the mutational 231   effect on each component trait j according to $\Delta z_{i,j} = r\xi_j / \sqrt{\sum_{l=1}^{n} \xi_l^2}$ (for alternative 232   distributions, see discussion in Poon & Otto 2000). The phenotypic value of the mutant 233   individual was then set to $z_{i,j} + \Delta z_{i,j}$ for all traits j within sphere i and left unchanged for 234   traits in other modules. Denoting the resulting mutant phenotypic vector as $\mathbf{Z} + \Delta\mathbf{Z}$, the

selection coefficient of the mutation is $s \equiv \dfrac{w(\mathbf{Z} + \Delta\mathbf{Z})}{w(\mathbf{Z})} - 1$. In diploids, the above describes the

properties of homozygous mutations. Heterozygous diploids are assumed to be shifted half as 237   far in phenotype space, $\Delta\mathbf{Z}/2$ (i.e., we assume additivity on a phenotypic scale). The

dominance coefficient for fitness is then defined as $h \equiv \left(\dfrac{w(\mathbf{Z} + \Delta\mathbf{Z}/2)}{w(\mathbf{Z})} - 1\right) \Big/ s$.

Simulation methods

We consider a species that is sexually reproducing with non-overlapping generations. The 242   species is sub-divided into two finite populations, each comprising N hermaphrodites with no 243   migration between them (i.e., allopatry). Initially, these populations are genetically identical,

having just separated geographically. Genomes may be haploid or diploid and are assumed to

be large enough for each mutation to occur at a unique site, with free recombination among

sites (i.e., assuming an infinite sites model).

247   Mutations that affect fitness occur stochastically every generation, but we assume

that the mutation rate is quite low to keep the product of the rate and population size much

smaller than one. This condition allows us to assume that a population is monomorphic most

of the time (Kimura and Ohta 1969; Taylor *et al.* 2006). Thus, we consider the fate of mutations

one at a time (Orr 1998; Chevin *et al.* 2014). Each mutation that arises reaches fixation within

a single population with probability $\dfrac{1-\exp(-2s)}{1-\exp(-2Ns)}$ in haploids and $\dfrac{\int_{0}^{\frac{1}{2N}} e^{-2Ns(2h-1)x(1-x)-2Nsx}dx}{\int_{0}^{1} e^{-2Ns(2h-1)x(1-x)-2Nsx}dx}$

in diploids (Crow and Kimura 1970). Note that overdominance naturally arises, at least

transiently, in diploid models of adaptation to a new optimum (Sellis *et al.* 2011). We assume

that the population size is small enough that polymorphisms are sufficiently transient to be

ignored, although this assumption is more tenuous in diploids with overdominance. To simplify

the simulations, we track time in units of fixation events, where we randomly choose mutations

and assign them to one or the other population.

259   As the two populations accumulate mutations, they independently traverse the fitness

landscapes in the two patches. While this necessarily leads to genetic divergence, the

populations may remain similar phenotypically or may diverge initially and subsequently

converge (or vice versa), depending on the optima in the two patches. To explore the impact of

parallel versus divergent selection on reproductive isolation, we study the following two

environmental scenarios:

(1) Adaptation to a common optimum, shifted from the position of the initial population, (to

investigate mutation-order speciation).

(2) Adaptation to different optima (to investigate ecological speciation).

For each of these scenarios, we explore how the extent of reproductive isolation depends on

whether the environment shifts rapidly (in one step) or gradually (in multiple steps).

Specifically, we define the vector of optima across all spheres as $\mathbf{O}=\left\{\mathbf{o}_1,\mathbf{o}_2,...,\mathbf{o}_m\right\}$. By

assumption, the initial phenotype $\mathbf{Z}_0$ and the initial optimum, $\mathbf{O}_0$, are the same in the two

populations and set to the origin in all *m* spheres. We then allow environmental change by

shifting the optimum phenotype, which induces directional selection until the population

approaches the new optimum, at which point the population experiences stabilizing selection.

Under scenario (1), the optimum starts at $\mathbf{O}_0$ and moves to a final optimum of $\mathbf{O}_E$, either in

one step (rapid environmental change) or in τ time steps, with each step occurring after a

fixation event (gradual environmental change). Under scenario (2), adaptation proceeds

towards $\mathbf{O}_A$ in patch A and towards $\mathbf{O}_B$ in patch B, again where the optima move either

rapidly once or gradually over τ fixation events.

We measure the level of reproductive isolation between the two populations through

crosses, generating F_1, F_2, and backcross (BC) hybrids. To generate F_1 individuals in the

haploid case, for example, when there are x fixed mutants in patch A and y fixed mutations in

patch B, we randomly transmit each mutation to offspring with a 50% probability, so that there

are, on average, $(x+y)/2$ mutations in each hybrid. In the diploid case, all F_1 hybrids are identical

and are heterozygous for all x and y mutations. The phenotype of a hybrid individual is then

calculated as the sum of the mutational vectors, and its fitness is assessed in both parental

environments. This is repeated 1000 times for each hybrid type. FGM simulations for

adaptation and for producing hybrids were written in C++ and *Mathematica*, respectively

(hosted on Github, https://github.com/ryamaguchi0731/FGM-master).

We were also interested in the time course over which reproductive incompatibilities

developed. Whereas some models have assumed that all divergent pairs of loci are equally

likely to give rise to speciation barriers (Orr and Turelli 2001), we speculated that large-effect

mutations that accumulate during periods of rapid adaptation to a changing environment may

contribute disproportionately. For each population, we thus accumulate an equal number of

fixation events in two phases: the phase during which adaptation occurs ("adaptation phase")

and the phase during which the population moves randomly around the optimum ("stationary

phase"), with drift as well as selection affecting the fate of a mutation. For simplicity, we define

the initial adaptation phase as the time until the first mutation fixes with a negative selection

coefficient in each population. With a moving optimum, the adaptation phase is defined as the

time until the first mutation fixed with a negative selection coefficient after the optimum stops

moving, as sometimes deleterious alleles fixed when the phenotype happens to be near the

current optimum. In addition, for modular pleiotropy, we assume that the adaptation phase

continues until the first allele with a negative selection coefficient fixes within the focal sphere

in which the optimum has moved.

To translate the number of fixation events into a more biologically meaningful time

scale, we calculated the expected number of generations until appearance and fixation of the

next mutation and summed this quantity across the series of mutations fixed. Given the

mutation rate u, we first calculated the waiting time until the appearance of the next "successful"

mutation that would survive stochastic loss while rare. This waiting time was assumed to be

exponentially distributed with a rate of events equal to the mutation rate, u (set as 10^{-5}

throughout this paper) multiplied by the number of haplotypes in the population ($c\,N$, where c

= 1 for haploids and 2 for diploids) times the average fixation probability, P. The latter varies

depending on the current distance of the population from the optimum and was determined by

integrating P over the distribution of possible mutations for each state of the population. The

mean waiting time of this exponential distribution is $1/(cNuP)$ generations. Then, we calculated

the fixation time of the successful allele that was drawn, using the expected waiting time until

fixation for a new mutation (Ewens 2004; Otto and Day 2007). Given the selection coefficient

and the dominance coefficient of a mutation, the expected waiting time, conditional upon

fixation, is approximately (Kimura 1980):

322
$$T = \int_0^1 \int_0^y \frac{2}{V(x)} e^{-2\int_x^y \frac{M(z)}{V(z)}dz} dxdy \,, \tag{Eq. 2}$$

where M and V are the mean and variance of the change of mutation allele frequency p per

generation. Specifically, $M(p) = sp(1-p)$ and $V(p) = \frac{p(1-p)}{N}$ in haploids and

$M(p) = p(1-p)[sp+h(1-2p)]$ and $V(p) = \frac{p(1-p)}{2N}$ in diploids. The expected waiting

time until the next mutation appears and fixes then becomes $1/(cNuP)$ plus T.

Results

Evolutionary trajectory of adaptation

Under Fisher's geometric model (FGM), mutations have nearly a 50% chance of being

beneficial when a population starts far from the optimum relative to the effect size of the

mutation, but they are more likely to overshoot the optimum and less likely to fix once the

population approaches the optimum (Fisher 1930; Kimura 1983). As a consequence, during the

process of adaptation, the average phenotypic effect size of fixed mutations is typically larger

at first and declines as populations become increasingly adapted (Figure S1a; Orr 1998). When

the distribution of mutations has a larger average magnitude, we find that the optimum is

reached faster, as expected (Figure S1c). After nearing the optimum, populations accumulate a

combination of slightly deleterious and beneficial mutations (inset to Figure S1b), resulting in

a phase of mutation-selection-drift where the populations stochastically move around the

optimum through successive fixation events.

We then accumulated the same total number of fixation events in the initial

adaptation phase and in the second stationary phase in each parental population (allowing

each parental population to accumulate a different number of fixed mutations). This

procedure ensured that we allowed enough time to pass in the stationary phase that the level

of divergence contributed by the two phases is the same (in terms of substitutions). Table S2

gives the number of fixation events in each parental population, the corresponding expected

number of generations, and the proportion of time in the adaptation phase. The adaptation

phase was typically much faster than the stationary phase. After the adaptation and stationary

phase had finished, we performed crosses between populations to determine how hybrid

fitness depended on the nature of environmental change, the mutation effect size, ploidy, and

the modularity of pleiotropy.

Adaptation to a common optimum – Figure 2 shows hybrid fitness after a period of adaptation

towards a common optimum in haploids and diploids (for an exploration of introgression

effects, see Fraïsse *et al*., 2016), assuming that the optimum shifted in one step (rapid

environmental change). For the parameters considered, substantial intrinsic reproductive

isolation is observed in all hybrids, with the exception of diploid F_1s. Diploid F_1s are not yet

recombinant but heterozygous at all sites that differ between the parents. When parents near

the same optimum are crossed in the FGM, averaging their phenotype in the diploid F_1

reduces the distance to the peak, as discussed by Barton (2001), Fraïsse et al. (2016), and

Simon et al. (2018). Indeed, we find that the fitness of F_1 hybrids is about three times closer

to the optimum than the average parental fitness, although this difference is too small to see

in Figure 2 (parental and F_1 hybrid fitnesses of 0.999273 and 0.999784, respectively, in panel

(b) with $\lambda = 0.04$ and of 0.997224 and 0.999152 in panel (d) with $\lambda = 0.12$).

By contrast, all recombinant hybrids have reduced average fitness, relative to the

parents, reflecting the segregation load studied by Chevin *et al*. (2014). Importantly, the

incompatibility results primarily from the large-effect mutations that accumulate during

adaptation, which combined can generate hybrids that move far off the peak fitness and that

exhibit substantial pleiotropic side effects (Figure 1). Such incompatibilities are not observed

to any substantial degree when the optimum remains at the origin, even with the same

number of fixation events (Figure S2). That is, reproductive isolation is not simply a function

of the number of divergent alleles. Rather, the vast majority of reproductive isolation arose

during the first adaptation phase when large-effect mutations were likely to fix. Considering

only these mutations, F_2 hybrid fitness, for example, was reduced below the optimum by, on

average, 0.72 in haploids and 0.74 in diploids. By contrast, F_2 hybrid fitness remained much

higher, 0.99 in haploids and 0.98 in diploids, when we add the same number of mutations that

occurred only during the stationary phase to the optimum (see Table S2 for the number of

fixation events in each population).

Hybrid breakdown did not accumulate linearly over time during the adaptation

phase (Figure S3a). Indeed, early on hybrids were slightly fitter than the average of the

parental populations when fitness was measured in their new environment, presumably due to

the large advantage of combining mutations and approaching the distant optimum. Only after

several mutations had fixed did hybrid fitness decline, on average, relative to the parents.

We next asked how much of the hybrid breakdown was caused by mutational

effects along the trait axis of environmental change (e.g., hybrids overshooting the optimum)

versus pleiotropy (e.g., hybrids breaking apart compensatory gene combinations). To do this,

we measured hybrid fitness only along the axis of environmental change, setting all other trait

values to the optimum (black slice in pie chart). The remainder of the hybrid breakdown is

due to pleiotropy. We find that the majority of the observed breakdown was due to pleiotropy

(Figure 2).

While we have illustrated the fitness drop in several hybrid forms, the Brownian

bridge approximation of Simon et al. (2018) allows us to connect the fitness of any hybrid

genotype, given the parental fitness, the hybrid index (H), the heterozygosity of the hybrid

(p_{12}), and one free parameter, which measures the extent of hybrid breakdown. As illustrated

in Figure S4 (a,b), the method of Simon et al. (2018) works well to predict the fitness of each

class of hybrids, once we fit the free parameter by minimizing the sum of squared deviations

between the observed and expected fitnesses for the different hybrids.

During the period of adaptation, large mutations can fix, even if they shift the

phenotype substantially away from the origin in some directions, as long as the average

phenotypic distance to the optimum is shorter. In subsequent generations, these negative

pleiotropic side effects favor the fixation of compensatory mutations that move the phenotype

back toward the optimum (Figure 1). In recombinant hybrids, however, these compensatory

mutations are separated from the mutant backgrounds in which they are favored, reducing

fitness. Consequently, the fitness reduction in hybrids is more severe when mutational effect

sizes (and hence pleiotropic side effects) are larger, on average (see Figure 3a,b with a 0°

angle for populations experiencing a common optimum, with dots from left to right

representing smaller to larger average effects). It is also worth highlighting the substantial

amount of variation in reproductive isolation observed across replicates, depending on the

exact adaptive path taken, with more variability occurring when the average mutational effect

is larger (for Figure 3a, for example, the standard deviation in the average F_1 haploid fitness

among replicates rose from 7.41×10^{-3} when $\lambda=0.04$ to 1.26×10^{-2} when $\lambda=0.12$).

The above simulations were repeated but allowing the optimum to move gradually

towards its final position, $\mathbf{O}_E$. With a history of slower environmental change, large-effect

mutations have a lower probability of fixation, reducing the accumulation of negative

pleiotropic side-effects. Consequently, we find that much weaker reproductive isolation

develops, with higher hybrid fitness, after a period of slow environmental change, even when

ultimately reaching the same optimum (see Figure 3c,d with a 0° angle for populations

experiencing a common optimum, with dots from left to right representing slower to faster

environmental change). We thus conclude that rapid environmental change, resulting in the

accumulation of large-effect mutations, is particularly conducive to mutation-order

speciation.

Adaptation to different optima – In the next scenario, we simulated isolated populations

adapting to different fixed optima, assessing hybrid fitness in both parental environments.

The results are qualitatively different in diploids, and now all hybrids, including F_1 hybrids,

have lower fitness within each environment, as their phenotype falls between the optima of

the two environments (extrinsic reproductive incompatibility). The larger the angle between

the two optima, the lower the fitness of hybrids (Figure 3), as well as the lower the average

fitness of the two parents (grey dots). The average fitness of the two parents is very close to

that expected by the respective optima to which they have adapted (Table S1), with one

parent having fitness near one and the other having a fitness that declines to zero as the angle

of divergence rises. Also shown is the expected fitness if the hybrid were phenotypically mid-

way between the two parents (dashed line). When the new environmental optima are at 30° or

45° relative to the ancestral population, the recombinant hybrids are less fit than either the

average of the parents or the parental mid-point, indicating intrinsic incompatibilities. At 90°,

some of the recombinant hybrids have a high proportion of local alleles, causing the hybrid

offspring to rise above that expected for the mid-parent phenotype (less extrinsic

incompatibility). Hybrid breakdown is more severe when mutational effect sizes are larger

(see sets of three adjacent points in Figure 3a,b), indicating stronger intrinsic

incompatibilities, but this effect is weak when parents are highly diverged (90°) because of

the counteracting advantage of recombinants being more likely to carry large effect alleles

adapted to patch A. The temporal dynamics of hybrid breakdown is illustrated in Figure S3.

Again, divergence accumulates slowly at first, especially when the new optima are similar

(30° or 45°).

We next explored the sensitivity of ecological speciation to the speed of

environmental change. As observed with mutation-order speciation, greater reproductive

isolation arises when the environment changes rapidly (see sets of four adjacent points in

Figure 3c,d), but again the effect is smaller than the effect of the angle between the two

environments. Essentially, the strong extrinsic reproductive incompatibility caused when

parents are adapted to different environmental optima masks the intrinsic incompatibilities

that arise from accumulating incompatible mutations (Figure 1). The more similar the current

parental environments (the lower the angle between them), the greater the difference in

hybrid breakdown observed between a history of slow versus rapid environmental change.

Again, the Brownian bridge approximation of Simon et al. (2018) allows us to

correctly predict the fitness of any hybrid genotype, by fitting one parameter along with the

parental fitnesses, H, and p_{12} (Figure S4). The fitted parameter, a measure of the extent of

hybrid breakdown, depends on the current parental environments and is greater when

populations have been selected to more divergent optima (compare Figure S4 panels c,d to

e,f). Consistent with the conclusions drawn above, the breakdown parameter is also greater

when fit to simulation data that have allowed larger-effect mutations to accumulate, either

because the average effect of mutations is larger (as in Figure 3a,b) or because the

environmental change occurred faster (as in Figure 3c,d).

When parental population have experienced divergent selection toward different

optima, backcross hybrids to the local parent, BC_L, show much higher fitness than backcross

hybrids to the foreign parent, BC_F (Figure S5a,b), a hallmark of ecological speciation (Rundle

and Whitlock 2001). The relative performance of the various hybrids depends, however, on

the angle between the two optima (compare Figure S5c,d with an angle of 90° to Figure S5a,b

with 30°). As a measure of the importance of ecological speciation, Rundle and Whitlock

(2001) suggest calculating the fitness of BC_L minus the fitness of BC_F, averaged over the two

local environments, as this estimates the strength of additive-by-environment interactions

(AE). Indeed, AE rises from zero when parental environments are identical to ~0.4-0.6 for

environments separated by an angle of 90° (Figure 4a,b). Across the full range of angles,

however, AE behaves non-monotonically (Figure 4a,b). For very dissimilar environments, the

fitness of the local BC_L falls to zero, severely limiting the maximum value of AE.

The degree to which ecological speciation contributes to reproductive isolation

may be better captured by measuring AE relative to the maximum value it could take (i.e., AE

divided by the fitness of BC_L). This rescaled value measures the degree to which additive-by-

environment interactions reduce the fitness of hybrids backcrossed to the foreign parent

compared to the fitness expected based on backcrosses with the local parent. When AE/BC_L

is zero, fitness is the same regardless of which parental population is used in the backcrosses

(as expected when the parental environments are the same), whereas an AE/BC_L of one

indicates that the genetic differences between the parents reduces the fitness of backcross

hybrids to zero when raised in the foreign environment (as expected following strong

divergent selection). The rescaled measure AE/BC_L does rise monotonically with the angle

between the optima and hence the degree of divergent selection (Figure 4c,d), although

sample sizes would need to be sufficient to avoid bias caused by measurement error in the

denominator.

Impact of modularity

We next investigated the degree to which modularity of the phenotype would affect the

development of reproductive incompatibilities between adapting populations. To investigate

different degrees of modular pleiotropy, we held the total number of phenotypic dimensions

constant (the number of spheres times the number of traits per sphere = *m n*) but varied the

number of spheres. To simplify the presentation, we consider only F_1 hybrids in haploid

simulations. Our prediction was that we would observe less negative pleiotropy when the

phenotype was more modular, leading to less hybrid breakdown. While we did see that, as

expected, pleiotropy contributed less to hybrid breakdown in more modular simulations (pie

charts in Figure 5a), hybrid fitness actually declined with high levels of modularity (Figure 5a).

Specifically, reproductive isolation showed a non-monotonic pattern, with more hybrid

breakdown in the fully pleiotropic model (*m* = 1) and in the model with no pleiotropy (*n* = 1)

than with intermediate levels of pleiotropy.

To understand this counter-intuitive result, we investigated how the genes contributing to

hybrid breakdown accumulated over time. Relatively little contribution came from mutations

accumulating during the stationary phase, so we sub-divided the mutations accumulated during

the adaptation phase into two groups: the first 50% of mutations that fixed and the second 50%

of mutations that fixed. With more modular genetic networks (*m* larger; Figure 5b), mutations

that fixed during the first half of adaptation were more likely to work well together, improving

fitness along the axis of environmental change and increasing mean hybrid fitness, as we had

expected. When the environmental change occurs in a module with few traits (low *n*),

mutations that fix in one population are more likely to point towards the optimum and still be

beneficial in another population. By contrast, as the population approached the optimum during

the second half of the adaptation phase, small-effect mutations in different modules made up

an increasingly large fraction of the mutations that fixed in different populations (recall that

the stationary phase was defined to start only after a deleterious mutation arose in the module

experiencing environmental change). Consequently, the more modules, the less likely

mutations in different populations would compensate for one another, so that decreased hybrid

fitness was observed with more modularity (Figure 5c). Thus, the role of modularity in

speciation had opposite effects during rapid early periods of adaptation, where reproductive

isolation developed faster in less modular systems, compared to later periods of relatively slow

adaptation, where reproductive isolation developed faster in more modular systems.

As an alternative explanation, we asked whether there was a substantial difference in the

magnitude of mutations as a function of modularity. The magnitude of the largest mutation that

fixed in each population was, however, similar across the combinations of *m* and *n* explored

(Figure S6), and so this explanation does not appear to contribute substantially to the non-

monotonic pattern with modularity.

Homoploid hybrid speciation

As discussed above, the history of environmental change can have a strong impact on the

degree of reproductive isolation exhibited among populations. Here, we briefly explore how

this history can also impact the potential to produce hybrids that are able to colonize new

environments, leading to hybrid speciation. Specifically, we investigated the possibility of

homoploid hybrid speciation by crossing the two parental populations generated in our

simulations and measuring the fitnesses of hybrids across a range of possible new environments.

For each environment, representing a potential site for colonization, the fitness of F_1 hybrid

haploids was assessed given the optimal phenotype in that environment, $\mathbf{O}_P$. The probability

of establishment in a site, P_e, was determined for any genotype by the branching process

approximation $1 - P_e = e^{-wbP_e}$, assuming a Poisson number of offspring with mean equal to the

genotypic fitness *w* times the average reproductive capacity, *b*, of an individual with full fitness

of *w*=1 (*b* was set to 5 in the simulations). We repeated this calculation for 1000 hybrids and

averaged P_e to get the expected establishment probability. We then measured the establishment

probability of hybrids relative to the average establishment probability of the two parental

genotypes.

Figure 6 illustrates the colonization success of hybrid haploids versus their parents,

as a function of the phenotypic optimum in the colonized environment. Whether hybrids are

more likely to colonize a given environment depends strongly on the history of adaptation in

the parental populations. When the parental populations diverged in a constant environment,

hybrids are almost never better colonists than the parental populations (top panels; optimum

constant at zero as in Figure S2). By contrast, hybrids are more variable following a period of

adaptation to a common environment (mutation-order speciation), and so at least some

individual hybrids are able to colonize substantially different environments than the parental

populations (using parents from simulations reported in Figure 3 with a 0° angle between

environments). With ecological speciation between parental populations adapted to different

environments (> 0° angle in Figure 3), hybrids are much better at colonizing new environments

that fall between the optima of the parental populations (bottom panels). Homozygous diploid

F_2s behave similarly, but the establishment probability of diploids in general would have to

account for segregation variance after colonization, which was not explored.

557   Thus, the history of the parental populations has a large influence on the fate of

hybrids and their ability to survive in and colonize new environments. Even when the parental

populations are phenotypically similar, distinct hybrids that can survive quite different

environmental conditions are more likely to be produced if the parents had recently undergone

a period of adaptation and accumulated different large effect mutations (compare middle panels

to top panels of Figure 6).

Discussion

We used a simple model of optimizing selection on multiple traits ("Fisher's geometric model")

to explore speciation in the face of environmental change. Because the path of adaptation in

Fisher's geometric model varies among populations evolving in allopatry, genetic crosses

between populations can yield mis-matched combinations of adaptive mutations, producing

hybrid offspring of lower fitness (Chevin *et al*. 2014; Fraïsse *et al*. 2016; Simon *et al*. 2018).

Our main conclusion from this model is that periods of environmental change are likely engines

of speciation for allopatric populations, allowing populations to develop strong intrinsic

reproductive incompatibilities even when they are adapting to similar environmental changes

("mutation-order speciation"; Figure 3), as well as contributing additional intrinsic

incompatibilities to the extrinsic hybrid breakdown that occurs when parents are adapted to

different environments ("ecological speciation"; Figure 3). Reproductive incompatibilities

were strongest when large-effect mutations could accumulate, i.e., during times of rapid

environmental change, which challenges the notion that all genetic substitutions are equally

likely to generate reproductive incompatibilities.

579   Using the metaphor of a "speciation clock" (e.g., Coyne & Orr 1989), these results

580   imply that the clock will vary in rate, sometimes speeding up and sometimes slowing,

581   depending on the local environmental circumstances and the chance of experiencing a period

582   of adaptation. Averaged over these circumstances, however, we would still predict a constant

583   rate of ticking, but our work predicts greater variation in the amount of BDMIs for a given

584   period of divergence time, depending on the history of stasis or change in the environment.

585   We also explored the relative importance of mutation-order speciation versus

ecological speciation by comparing the fitness of different hybrid crosses. Rundle and Whitlock

(2001) showed that the contribution of ecological speciation could be captured by the additive-

by-environment interaction (AE), estimated from the fitness of individuals backcrossed to the

local population (BC$_L$) minus the fitness of individuals backcrossed to a foreign population

(BC$_F$). Only under ecological speciation are these backcross fitnesses expected to differ. We

find that this AE measure does indeed rise with the angle of environmental optima between

two evolving populations (measured here relative to their ancestral origin), but only when that

angle was small. With larger environmental differences, the fitness of even the local

backcrossed individuals (BC$_L$) could be very low, leading AE to drop (Figure 4a,b). We thus

caution against using low AE values to distinguish cases of mutation-order from ecological

speciation, especially when backcross fitnesses are all low. As a correction for this non-

monotonic behavior, we instead investigated a new measure of the importance of ecological

speciation, dividing AE by its maximum expected value (BC$_L$), which we found does rise with

the angle of environmental differences in our study and hence is a better indicator of the extent

of divergent selection (Figure 4c,d). It would be valuable for future studies to determine the

sampling properties of this new measure and its ability to distinguish mutation-order and

ecological speciation in simulated and empirical studies.

In addition, we considered how the modularity of the genotype-phenotype

relationship impacted the development of reproductive isolation under mutation-order

speciation. With a more modular genetic architecture, allopatric populations experiencing the

same environmental change are more likely to accumulate mutations that are compatible,

because the negative pleiotropic effects of adaptive mutations are constrained within the

module affected by the environmental change. Thus, we expected a higher degree of parallel

evolution, leading to less reproductive isolation, in cases where the environment changed in

the same direction in different populations. Unexpectedly, we found a non-monotonic

relationship between the extent of reproductive isolation and the degree of modularity (Figure

5). By separating out the first half of mutations fixed during the rapid early phase of adaptation

from the latter half of mutations fixed during the adaptation phase, we were able to verify that

reproductive isolation is less likely to develop in more modular systems early on during

adaptation when selection is strong, but this is counteracted by a greater chance that mutations

generate reproductive isolation during later phases of adaptation as populations more slowly

approach the optimum. The key difference between these phases was whether fixed mutations

tended to fix primarily in the same module – the sphere with the shifting environment – or in

different modules. Once the populations were near the optimum, different populations tended

to fix mutations and compensatory mutations in different modules, substantially increasing the

chance of hybrid breakdown when crossed.

Finally, we explored how the history of adaptation of the parental populations

influenced the potential to produce hybrid offspring that could colonize a new environment and

lead to homoploid hybrid speciation. Because hybrids can vary in phenotype beyond the range
of parental populations, hybrids can successfully colonize environments inaccessible to
parental populations, potentially contributing to homoploid hybrid speciation (Gross and
Rieseberg 2005). Importantly, we found that hybrid speciation was especially likely when the
parental populations had undergone a period of rapid adaptation in allopatry (Figure 6). The
parents were then more likely to differ by large-effect mutations, allowing hybrid genotypes to
vary more broadly in phenotype and producing at least some individuals that could succeed in
more extreme environments. The potential for colonization success by hybrids was observed
whether the parental populations had adapted to a common environment (mutation-order
speciation, Figure 6f) or different environments (ecological speciation, Figure 6i), although the
environment where hybrids were most successful differed (in a halo around the common
parental environment versus midway between them, respectively). We thus predict that periods
of adaptation by the parental forms may be particularly conducive to subsequent homoploid
hybrid speciation.

Recombining pre-existing variation is a powerful way of generating new species over
short periods of time. Indeed, hybrid speciation is thought to underlie adaptive radiation in a
number of systems (Marques *et al.* 2019; Martin and Richards 2019). Based on our simulations,
we suggest that large effect mutations following a period of adaptation may also contribute
disproportionately to admixture variation, contributing to intrinsic and extrinsic
incompatibilities, transgressive traits, and novel trait combinations in hybrid swarms (compare,
for example, figure 6f,i to 6c).

In this study, we have made several simplifying assumptions that warrant more
exploration in future work. For one, we assumed that populations are largely monomorphic,
punctuated in time by the fixation of new mutations. This assumption breaks down in larger
populations (N) or in environments with a larger number of potential mutations (higher u),
either of which increases the flux of mutations into a population. A higher flux of mutations
can increase the rate at which large-effect mutations appear, increasing the probability that
intrinsic incompatibilities arise, but more mutations also increase the standing genetic variation
of the parental populations, which can mean that the same alleles are present and fix in the
parental populations when the environment changes (Thompson *et al*. 2019). Which effect is
more important depends on the exact details, e.g., when the parental populations became
isolated relative to the change in environment. Future work is needed to explore the
development of reproductive isolation allowing for polymorphic alleles at multiple loci
throughout the genome. Relaxing the assumption of monomorphism is particularly important
for diploid populations, because large effect mutations often arise that overshoot the optimum,

leading to overdominance that maintains polymorphism (Sellis *et al*. 2011).

Another simplifying assumption in our model is the Gaussian fitness landscape ($k =$
2). Fraïsse *et al*. (2016) investigated various k values with divergence via genetic drift around
an optimum and found that speciation genes could fix with a nonnegligible probability only
when the number of traits (n) was sufficiently small and k was sufficiently large (their Figure
2a). This is consistent with our finding that BDMIs do not develop to any appreciable extent
in populations that remain near the optimum. With a history of adaptation allowing large-effect
mutations, however, BDMIs do arise even with a quadratic fitness surface (Figure 2 here, see
also Barton *et al*. 2001), although the results of Fraïsse *et al*. (2016) suggest that even more
hybrid breakdown would be observed with $k > 2$.

In addition, it would be worthwhile exploring other forms of selection beyond the
single-peaked surface considered here. With multiple optima or other complex fitness
landscapes (e.g., the "holey adaptive landscape" of Gavrilets 1997), mutation-order speciation
is thought to be even more likely, as populations move to different points on the fitness
landscape, depending on the order in which mutations arise (Gavrilets 2004). Nevertheless, we
conjecture that the degree of reproductive isolation may still be stronger when initially further
away from the peaks of a rugged fitness surface because large-effect mutations may then be
more likely to fix in allopatric populations, relative to the small-effect mutations providing
slight improvements in fitness near a local optimum. Future work would be valuable that
explores how the shape of the fitness landscape affects the likelihood of accumulating the large
effect mutations that contribute disproportionately to hybrid breakdown.

As an example of speciation across a more complex fitness landscape, Benkman
(2003) showed that the fitness surface of red crossbills feeding on cones from different
coniferous tree species displayed different peaks, associated with distinct call types, with both
ridges of relatively high fitness and deep fitness valleys. Although not explored in this paper,
the combination of fitness ridges with deep valleys might facilitate reproductive isolation by
allowing large-effect mutations that reach different points along the ridge and that, when
combined, cause hybrid phenotypes to exhibit low fitness.

Fluctuating selection could also contribute to the accumulation of large-effect
mutations and be a potent driver of speciation (see, e.g, Barton 2001; Chevin *et al*. 2014;
Fraïsse *et al*. 2016). For example, beak phenotypes of the medium ground finch *Geospiza fortis*
on Galapagos have experienced selection that fluctuates between life stages, years and
generations (Price and Grant 1984; Grant and Grant 2008). Stabilizing selection with a
fluctuating optimum allows the population to accumulate relatively large mutations, which
contribute disproportionately to the accumulation of reproductive isolation.

In conclusion, our work shows that the tempo of speciation does not tick at a steady
rate. Rather, reproductive isolation between allopatric populations should rise faster after bouts
of rapid environmental change. A rapidly changing environment is particularly important when
selection acts in parallel among populations (mutation-order speciation), as extrinsic
reproductive incompatibilities are lacking. In our simulations, crossing the few large-effect
mutations that arose during bouts of rapid adaptation caused more hybrid breakdown than
combining many small-effect mutations, whose effects tended to cancel (Figure 1; see also eq.
3 in Chevin *et al.* 2014). These results help explain the surprisingly rapid appearance of genetic
incompatibilities during short-term experimental evolution studies in the lab (e.g., Dettman *et*
al. 2007, 2008; Kvitek and Sherlock 2011; Chou *et al.* 2014; Ono *et al.* 2017). Periods of strong
selection are unusually powerful drivers of speciation, helping to account for why reproductive
isolation arises so rapidly in experimental evolution studies and yet the biological world has
not diversified into an infinite variety of species.

Author contributions

RY and SPO designed the study; RY developed the computer simulation code and conducted analyses in consultation with SPO; RY and SPO wrote the manuscript.

Data accessibility

All code necessary to repeat the analysis described in this study have been made available. C++ source codes of our simulation models for adaptation process, *Mathematica* (version 11.2.0.0) codes for calculating hybrids genotypes and fitnesses, R (version 3.5.1) scripts that we used to produce figures, and simulation results in this paper are hosted on Github (https://github.com/ryamaguchi0731/FGM-master).

Acknowledgements

This work was supported by funding from the Japanese Society for the Promotion of Science (Research Fellowship for Young Scientists, DC1-14J02775 and PD-17J01380, to RY) and from the Natural Sciences and Engineering Research Council of Canada (RGPIN-2016-03711 to SPO). We thank the following people for their helpful comments on previous drafts of the manuscript: Yoh Iwasa, Matt Osmond. Ken Thompson, Sam Flaxman, Mohamad Noor, Bryn Wiley, and three anonymous reviewers. We are grateful to Matt Osmond for suggesting the normalization of AE by BC_L as a more informative measure of ecological speciation.

Literature Cited

1.   Albertson, R. C., & Kocher, T. D. (2005). Genetic architecture sets limits on transgressive
segregation in hybrid cichlid fishes. *Evolution*, *59*(3), 686-690.

2.   Barton, N. H. (2001). The role of hybridization in evolution. *Molecular Ecology*, *10*(3),
551-568.

3.   Benkman, C. W. (2003). Divergent selection drives the adaptive radiation of
crossbills. *Evolution*, *57*(5), 1176-1181.

4.   Chevin, L. M., Decorzent, G., & Lenormand, T. (2014). Niche dimensionality and the
genetics of ecological speciation. *Evolution*, *68*(5), 1244-1256.

5.   Chevin, L. M., Martin, G., & Lenormand, T. (2010). Fisher's model and the genomics of
adaptation: restricted   pleiotropy,   heterogenous   mutation,   and   parallel
evolution. *Evolution*, *64*(11), 3213-3231.

6.   Chou, H. H., Delaney, N. F., Draghi, J. A., & Marx, C. J. (2014). Mapping the fitness
landscape of gene expression uncovers the cause of antagonism and sign epistasis between
adaptive mutations. *PLoS Genetics*, *10*(2), e1004149.

7.   Crespi, B., & Nosil, P. (2013). Conflictual speciation: species formation via genomic
conflict. *Trends in Ecology & Evolution*, *28*(1), 48-57.

8.   Crow, J. F., & Kimura, M. (1970). An introduction to population genetics theory. *An
Introduction to Population Genetics Theory*. Harper and Row, New York USA.

9.   Dettman, J. R., Anderson, J. B., & Kohn, L. M. (2008). Divergent adaptation promotes
reproductive isolation among experimental populations of the filamentous fungus
Neurospora. *BMC Evolutionary Biology*, *8*(1), 35.

10. Dettman, J. R., Sirjusingh, C., Kohn, L. M., & Anderson, J. B. (2007). Incipient speciation
by divergent adaptation and antagonistic epistasis in yeast. *Nature*, *447*(7144), 585-588.

11. Ewens, W. J. (2004). *Mathematical Population Genetics 1: Theoretical Introduction*. 2nd
edition. Springer-Verlag New York, Inc. New York, USA.

12. Fisher, R. A. (1930). *The Genetical Theory of Natural Selection*. Clarendon Press, Oxford,
UK.

13. Fraïsse, C., Gunnarsson, P. A., Roze, D., Bierne, N., & Welch, J. J. (2016). The genetics
of speciation: Insights from Fisher's geometric model. *Evolution*, *70*(7), 1450-1464.

14. Gavrilets, S. (1997). Evolution and speciation on holey adaptive landscapes. *Trends in
Ecology & Evolution*, *12*(8), 307-312.

15. Gavrilets, S. (2004). *Fitness landscapes and the origin of species (MPB-41)* (Vol. 41).
Princeton University Press, Princeton NJ.

16. Grant, P. R., & Grant, B. R. (2008). *How and why species multiply: the radiation of*

Darwin's finches. Princeton University Press, Princeton NJ.

17. Gross, B. L., & Rieseberg, L. H. (2005). The ecological genetics of homoploid hybrid speciation. *Journal of Heredity*, *96*(3), 241-252.

18. Hartl, D. L., & Taubes, C. H. (1998). Towards a theory of evolutionary adaptation. *Genetica*, *102*, 525-533.

19. Hill, W. G., & Zhang, X. S. (2012). On the pleiotropic structure of the genotype–phenotype map and the evolvability of complex organisms. *Genetics*, *190*(3), 1131-1137.

20. Kimura, M. (1980). Average time until fixation of a mutant allele in a finite population under continued mutation pressure: Studies by analytical, numerical, and pseudo-sampling methods. *Proceedings of the National Academy of Sciences*, *77*(1), 522-526.

21. Kimura, M. (1983). *The neutral theory of molecular evolution*. Cambridge University Press, Cambridge UK.

22. Kimura, M., & Ohta, T. (1969). The average number of generations until fixation of a mutant gene in a finite population. *Genetics*, *61*(3), 763.

23. Kvitek, D. J., & Sherlock, G. (2011). Reciprocal sign epistasis between frequently experimentally evolved adaptive mutations causes a rugged fitness landscape. *PLoS Genetics*, *7*(4), e1002056.

24. Mani, G. S., & Clarke, B. C. (1990). Mutational order: a major stochastic process in evolution. *Proceedings of the Royal Society of London B: Biological Sciences*, *240*(1297), 29-37.

25. Marques, D. A., Meier, J. I., & Seehausen, O. (2019). A combinatorial view on speciation and adaptive radiation. *Trends in Ecology & Evolution*, *34*(6), 531-544.

26. Martin, G. (2014). Fisher's geometrical model emerges as a property of complex integrated phenotypic networks. *Genetics*, *197*(1), 237-255.

27. Martin, G., & Lenormand, T. (2006). A general multivariate extension of Fisher's geometrical model and the distribution of mutation fitness effects across species. *Evolution*, *60*(5), 893-907.

28. Martin, C. H., & Richards, E. J. (2019). The paradox behind the pattern of rapid adaptive radiation: How can the speciation process sustain itself through an early burst? *Annual Review of Ecology, Evolution, and Systematics*, *50*, 569-593.

29. Mendelson, T. C., Martin, M. D., & Flaxman, S. M. (2014). Mutation-order divergence by sexual selection: diversification of sexual signals in similar environments as a first step in speciation. *Ecology Letters*, *17*(9), 1053-1066.

30. Muir, C. D., & Hahn, N. W. (2015). The limited contribution of reciprocal gene loss to

increased speciation rates following whole-genome duplication. *The American Naturalist*,
185(1), 70-86.

31. Ostevik, K. L., Moyers, B. T., Owens, G. L., & Rieseberg, L. H. (2012). Parallel
ecological speciation in plants? *International Journal of Ecology*, *2012*.

32. Ono, J., Gerstein, A. C., & Otto, S. P. (2017). Widespread genetic incompatibilities
between first-step mutations during parallel adaptation of Saccharomyces cerevisiae to a
common environment. *PLoS Biology*, *15*(1), e1002591.

33. Orr, H. A. (1998). The population genetics of adaptation: the distribution of factors fixed
during adaptive evolution. *Evolution*, *52*(4), 935-949.

34. Orr, H. A., & Turelli M. (2001). The evolution of postzygotic isolation: Accumulating
Dobzhansky-Muller incompatibilities. *Evolution*, *55*(6), 1085–1094.

35. Otto, S. P., & Day, T. (2007). *A Biologist's Guide to Mathematical Modeling in Ecology*
and Evolution. Princeton University Press, Princeton NJ.

36. Poon, A., & Otto, S. P. (2000). Compensating for our load of mutations: freezing the
meltdown of small populations. *Evolution*, *54*(5), 1467-1479.

37. Price, T. D., & Grant, P. R. (1984). Life history traits and natural selection for small body
size in a population of Darwin's finches. *Evolution*, *38*(3), 483-494.

38. Rieseberg, L. H. (1997). Hybrid origins of plant species. *Annual Review of Ecology and*
Systematics, *28*(1), 359-389.

39. Rundle, H. D. (2002). A test of ecologically dependent postmating isolation between
sympatric sticklebacks. *Evolution*, *56*(2), 322-329.

40. Rundle, H. D., & Whitlock, M. C. (2001). A genetic interpretation of ecologically
dependent isolation. *Evolution*, *55*(1), 198-201.

41. Schluter, D. (2000). *The ecology of adaptive radiation*. Oxford University Press, Oxford
UK.

42. Schluter,   D.   (2009).   Evidence   for   ecological   speciation   and   its
alternative. *Science*, *323*(5915), 737-741.

43. Schluter, D., & Conte, G. L. (2009). Genetics and ecological speciation. *Proceedings of*
the National Academy of Sciences, *106*(Supplement 1), 9955-9962.

44. Sellis, D., Callahan, B. J., Petrov, D. A., & Messer, P. W. (2011). Heterozygote advantage
as a natural consequence of adaptation in diploids. *Proceedings of the National Academy*
of Sciences, *108*(51), 20666-20671.

45. Simon, A., Bierne, N., & Welch, J. J. (2018). Coadapted genomes and selection on
hybrids: Fisher's geometric model explains a variety of empirical patterns. *Evolution*
Letters, *2*(5), 472-498.

46. Taylor, C., Iwasa, Y., & Nowak, M. A. (2006). A symmetry of fixation times in
evoultionary dynamics. *Journal of Theoretical Biology*, *243*(2), 245-251.

47. Tenaillon, O. (2014). The utility of Fisher's geometric model in evolutionary
genetics. *Annual Review of Ecology, Evolution, and Systematics*, *45*, 179-201.

48. Thompson, K. A., Osmond, M. M. and Schluter, D. (2019). Parallel genetic evolution and
speciation from standing variation. *Evolution Letters*, *3*(2), 129-141.

49. Wagner, G. P., & Zhang, J. (2011). The pleiotropic structure of the genotype–phenotype
map: the evolvability of complex organisms. *Nature Reviews Genetics*, *12*(3), 204-213.

50. Wang Z, Liao B-Y, Zhang J. 2010. Genomic patterns of pleiotropy and the evolution of
complexity. *Proceedings of the National Academy of Sciences*, 107(42), 18034–18039.

51. Waser N. M. & Campbell D. R. (2004). Ecological speciation in flowering plants. In:
Adaptive Speciation, eds. Dieckmann U., Doebeli M., Metz J.A.J. & Tautz D., pp. 264–
277. Cambridge University Press, Cambridge UK.

52. Waxman, D., & Welch, J. J. (2005). Fisher's microscope and Haldane's ellipse. *The*
American Naturalist, *166*(4), 447-457.

53. Welch, J. J., & Waxman, D. (2003). Modularity and the cost of
complexity. *Evolution*, 57(8), 1723-1734.

Table 1. Notation used. Default values are given in square brackets, except where noted in the

text.

Notation	Description
m	Number of spheres [1]
n	Number of traits in each sphere [10]
$\mathbf{z}_i = \left\{ z_{i,1}, z_{i,2}, ..., z_{i,n} \right\}$	Phenotype in the ith sphere
$\mathbf{o}_i = \left\{ o_{i,1}, o_{i,2}, ..., o_{i,n} \right\}$	Optimum phenotype in the ith sphere
$w\left(\mathbf{z}_1, \mathbf{z}_2, ..., \mathbf{z}_m \right)$	Individual's relative fitness
q	Parameter translating the phenotypic distance from the optimum into the strength of selection [1]
k	Degree of epistasis [2]
$\mathbf{Z} = \left\{ \mathbf{z}_1, \mathbf{z}_2, ..., \mathbf{z}_m \right\}$	Vector of phenotype sets of all spheres. $\mathbf{Z}_I$ is an initial condition
r	Mutation size, drawn from an exponential distribution with mean λ
ξ_j	Random number drawn from a standard normal distribution
$\Delta \mathbf{z}_i = \left\{ \Delta z_{i,1}, \Delta z_{i,2}, ..., \Delta z_{i,n} \right\}$	Phenotypic change due to mutation in the ith sphere
$\Delta \mathbf{Z} = \left\{ \Delta \mathbf{z}_1, \Delta \mathbf{z}_2, ..., \Delta \mathbf{z}_m \right\}$	Vector of mutational effects across all spheres
s	Selection coefficient
h	Measure of dominance
N	Number of individuals in a population [2000]
c	Ploidy level [1 for haploids, 2 for diploids]
u	Mutation rate [10^{-5}]
b	Number of offspring per individual of maximum fitness when calculating establishment probability [5]
$\mathbf{O} = \left\{ \mathbf{o}_1, \mathbf{o}_2, ..., \mathbf{o}_m \right\}$	Vector describing the optimal phenotype across all sets of spheres

851   Table S1. The location of the environmental optima.

Angle between two optima	Population 1	Population 2	Distance from Z_0 to O
0º	{2,0,0,0,0,0,0,0,0,0}	{2,0,0,0,0,0,0,0,0,0}	2
30º	$\{-\sqrt{2-\sqrt{3}}, -\sqrt{2+\sqrt{3}},$ 0,0,0,0,0,0,0,0\}	$\{\sqrt{2-\sqrt{3}}, -\sqrt{2+\sqrt{3}},$ 0,0,0,0,0,0,0,0\}	2
45º	$\{-\sqrt{2-\sqrt{2}}, -\sqrt{2+\sqrt{2}},$ 0,0,0,0,0,0,0,0\}	$\{\sqrt{2-\sqrt{2}}, -\sqrt{2+\sqrt{2}},$ 0,0,0,0,0,0,0,0\}	2
90º	$\{-\sqrt{2}, -\sqrt{2},$ 0,0,0,0,0,0,0,0\}	$\{\sqrt{2}, -\sqrt{2},$ 0,0,0,0,0,0,0,0\}	2
180º	{-2,0,0,0,0,0,0,0,0,0}	{2,0,0,0,0,0,0,0,0,0}	2

Vector angle is defined as $\text{Arccos}\left[\dfrac{a \cdot b}{\|a\| \, \|b\|}\right]$ where a and b are different vectors.

Table S2. Number of fixation events tracked within the parental populations P_1 and P_2 and the
corresponding number of generations, based on the expected waiting time until fixation. In
each population, we tracked the same total number of fixations events during the adaptation
phase and the stationary phase (e.g., 92 for P_1 and 88 for P_2 in each phase for the first set of
simulations).

Figure number & Population		Total fixations	Number of generations	% of time in adaptation phase
Figure 2a, 6a-f, S2a, S4a (Haploid)	P_1	184	2.31×10^7	8.32
	P_2	176	1.86×10^7	3.89
Figure 2b, S2b, S4b (Diploid)	P_1	204	6.73×10^7	4.99
	P_2	184	5.72×10^7	2.75
Figure 2c (Haploid)	P_1	68	2.62×10^7	6.59
	P_2	76	2.81×10^7	8.44
Figure 2d (Diploid)	P_1	124	1.19×10^8	3.29
	P_2	122	1.65×10^8	4.52
Figure 3a,c, 4a,c, S3a, S4a (Haploid, $\lambda = 0.04$)	P_1	159.52 ± 2.95	$1.85\times10^7 \pm 6.14\times10^5$	5.94 ± 0.38
	P_2	153.12 ± 4.45	$1.74\times10^7 \pm 7.54\times10^5$	7.15 ± 1.12
Figure 3a (Haploid, $\lambda = 0.08$)	P_1	99.20 ± 1.98	$1.94\times10^7 \pm 6.17\times10^5$	5.99 ± 0.41
	P_2	98.44 ± 2.38	$1.86\times10^7 \pm 8.81\times10^5$	6.21 ± 0.46
Figure 3a (Haploid, $\lambda = 0.12$)	P_1	83.44 ± 1.87	$2.25\times10^7 \pm 8.37\times10^5$	5.79 ± 0.43
	P_2	83.40 ± 1.44	$2.36\times10^7 \pm 6.63\times10^5$	7.03 ± 0.48
Figure 3b,d, 4b,d, S4b (Diploid, $\lambda = 0.04$)	P_1	183.04 ± 2.93	$5.69\times10^7 \pm 1.73\times10^6$	3.86 ± 0.25
	P_2	181.48 ± 4.79	$5.36\times10^7 \pm 2.73\times10^6$	5.55 ± 1.01
Figure 3b (Diploid, $\lambda = 0.08$)	P_1	133.08 ± 2.99	$1.11\times10^8 \pm 8.62\times10^6$	6.07 ± 1.97
	P_2	128.60 ± 4.91	$1.06\times10^8 \pm 5.79\times10^6$	6.40 ± 2.12
Figure 3b (Diploid, $\lambda = 0.12$)	P_1	111.88 ± 2.31	$1.26\times10^8 \pm 1.28\times10^6$	5.68 ± 1.99
	P_2	112.56 ± 1.61	$1.30\times10^8 \pm 3.71\times10^6$	5.97 ± 1.98
Figure 3c (Haploid, slow)	P_1	222.36 ± 1.88	$3.05\times10^7 \pm 4.08\times10^5$	17.8 ± 0.51
	P_2	216.20 ± 1.68	$2.97\times10^7 \pm 4.41\times10^5$	17.4 ± 0.48
Figure 3c (Haploid, medium)	P_1	166.32 ± 2.67	$2.02\times10^7 \pm 4.94\times10^5$	9.46 ± 0.27
	P_2	166.12 ± 2.05	$2.02\times10^7 \pm 4.40\times10^5$	9.87 ± 0.33

Figure 3c	P_1	155.24 ± 3.52	$1.79\times10^7 \pm 6.42\times10^5$	7.64 ± 1.31
(Haploid, fast)	P_2	158.22 ± 3.23	$1.78\times10^7 \pm 6.32\times10^5$	6.34 ± 0.31
Figure 3d	P_1	215.12 ± 1.82	$8.02\times10^7 \pm 1.05\times10^6$	18.1 ± 0.70
(Diploid, slow)	P_2	216.64 ± 1.78	$8.11\times10^7 \pm 1.16\times10^6$	17.0 ± 0.67
Figure 3d	P_1	180.04 ± 3.08	$5.83\times10^7 \pm 1.87\times10^6$	8.06 ± 0.29
(Diploid, medium)	P_2	170.12 ± 2.88	$5.41\times10^7 \pm 1.61\times10^6$	7.91 ± 0.34
Figure 3d	P_1	177.68 ± 4.70	$5.47\times10^7 \pm 2.38\times10^6$	7.52 ± 1.74
(Diploid, fast)	P_2	177.88 ± 4.11	$5.66\times10^7 \pm 2.34\times10^6$	6.69 ± 1.12
Figure 3a,c, 4a,c, S3b,	P_1	154.24 ± 3.84	$1.71\times10^7 \pm 7.37\times10^5$	5.83 ± 0.64
S4c	P_2	158.28 ± 3.48	$1.78\times10^7 \pm 6.54\times10^5$	5.90 ± 0.40
(Haploid, 30º, $\lambda = 0.04$)				
Figure 3a	P_1	101.36 ± 1.70	$1.92\times10^7 \pm 5.28\times10^5$	5.46 ± 0.41
(Haploid, 30º,	P_2	100.36 ± 2.75	$1.91\times10^7 \pm 7.88\times10^5$	6.30 ± 0.85
$\lambda = 0.08$)				
Figure 3a	P_1	80.28 ± 1.52	$2.24\times10^7 \pm 6.07\times10^5$	7.36 ± 0.51
(Haploid, 30º,	P_2	82.48 ± 1.70	$2.19\times10^7 \pm 7.84\times10^5$	6.29 ± 0.63
$\lambda = 0.12$)				
Figure 3a,c, 4a,c, S3c	P_1	155.20 ± 2.81	$1.78\times10^7 \pm 5.35\times10^5$	5.48 ± 0.33
(Haploid, 45º,	P_2	153.26 ± 3.85	$1.77\times10^7 \pm 6.03\times10^5$	6.70 ± 0.88
$\lambda = 0.04$)				
Figure 3a	P_1	103.24 ± 2.21	$2.06\times10^7 \pm 7.01\times10^5$	6.15 ± 0.42
(Haploid, 45º,	P_2	106.96 ± 2.31	$2.14\times10^7 \pm 6.42\times10^5$	6.61 ± 0.47
$\lambda = 0.08$)				
Figure 3a	P_1	81.72 ± 1.84	$2.26\times10^7 \pm 8.27\times10^5$	6.77 ± 0.48
(Haploid, 45º,	P_2	85.20 ± 1.78	$2.46\times10^7 \pm 7.65\times10^5$	7.93 ± 0.58
$\lambda = 0.12$)				
Figure 3a,c, 4a,c, 6g-i,	P_1	156.88 ± 2.45	$1.80\times10^7 \pm 4.57\times10^5$	6.05 ± 0.36
S3d, S4e,	P_2	149.96 ± 3.17	$1.66\times10^7 \pm 6.31\times10^5$	5.10 ± 0.26
(Haploid, 90º, $\lambda = 0.04$)				
Figure 3a	P_1	101.60 ± 2.10	$1.99\times10^7 \pm 6.31\times10^5$	6.75 ± 0.39
(Haploid, 90º,	P_2	101.44 ± 2.68	$1.99\times10^7 \pm 8.18\times10^5$	7.24 ± 0.63
$\lambda = 0.08$)				

Figure 3a (Haploid, 90°, $\lambda = 0.12$)	P_1	83.52 ± 1.76	$2.38\times10^7 \pm 8.53\times10^5$	7.97 ± 0.59
	P_2	81.64 ± 1.76	$2.29\times10^7 \pm 7.79\times10^5$	7.05 ± 0.50
Figure 3b,d, 4b,d, S4d (Diploid, 30°, $\lambda = 0.04$)	P_1	189.32 ± 2.88	$6.02\times10^7 \pm 1.57\times10^6$	4.70 ± 0.30
	P_2	175.08 ± 5.62	$5.40\times10^7 \pm 2.53\times10^6$	6.67 ± 1.58
Figure 3b (Diploid, 30°, $\lambda = 0.08$)	P_1	134.12 ± 2.56	$1.14\times10^8 \pm 3.81\times10^6$	4.09 ± 0.73
	P_2	133.56 ± 2.30	$1.24\times10^8 \pm 3.42\times10^6$	3.45 ± 0.44
Figure 3b (Diploid, 30°, $\lambda = 0.12$)	P_1	111.92 ± 2.19	$1.29\times10^8 \pm 4.47\times10^6$	3.65 ± 0.93
	P_2	109.80 ± 2.25	$1.23\times10^8 \pm 4.31\times10^6$	4.79 ± 0.79
Figure 3b,d, 4b,d (Diploid, 45°, $\lambda = 0.04$)	P_1	188.16 ± 4.86	$5.90\times10^7 \pm 2.23\times10^6$	5.34 ± 0.82
	P_2	185.28 ± 3.94	$5.78\times10^7 \pm 1.79\times10^6$	5.05 ± 1.01
Figure 3b (Diploid, 45°, $\lambda = 0.08$)	P_1	138.08 ± 2.61	$1.19\times10^8 \pm 3.99\times10^6$	3.84 ± 0.40
	P_2	138.28 ± 2.31	$1.20\times10^8 \pm 3.45\times10^6$	4.31 ± 0.45
Figure 3b (Diploid, 45°, $\lambda = 0.12$)	P_1	110.88 ± 2.75	$1.22\times10^8 \pm 5.33\times10^6$	3.49 ± 0.71
	P_2	115.72 ± 1.60	$1.31\times10^8 \pm 4.04\times10^6$	3.13 ± 0.49
Figure 3b,d, 4b,d, S4f (Diploid, 90°, $\lambda = 0.04$)	P_1	186.06 ± 3.25	$5.82\times10^7 \pm 1.75\times10^6$	3.92 ± 0.26
	P_2	190.88 ± 3.69	$5.80\times10^7 \pm 1.99\times10^6$	4.08 ± 0.25
Figure 3b (Diploid, 90°, $\lambda = 0.08$)	P_1	133.56 ± 2.68	$1.15\times10^8 \pm 4.08\times10^6$	3.79 ± 0.47
	P_2	138.88 ± 3.16	$1.15\times10^8 \pm 4.30\times10^6$	4.73 ± 0.88
Figure 3b (Diploid, 90°, $\lambda = 0.12$)	P_1	116.92 ± 1.83	$1.38\times10^8 \pm 4.19\times10^6$	3.84 ± 0.45
	P_2	112.16 ± 1.87	$1.27\times10^8 \pm 4.25\times10^6$	3.55 ± 0.55
Figure 3c (Haploid, 30°, slow)	P_1	218.76 ± 1.22	$3.28\times10^7 \pm 3.47\times10^5$	19.88 ± 4.58
	P_2	213.56 ± 1.12	$2.92\times10^7 \pm 2.47\times10^5$	11.52 ± 1.55
Figure 3c (Haploid, 30°,	P_1	165.96 ± 2.20	$2.21\times10^7 \pm 4.62\times10^5$	10.70 ± 2.17
	P_2	166.48 ± 2.01	$2.18\times10^7 \pm 4.33\times10^5$	9.94 ± 2.43

medium)

Figure 3c (Haploid, 30º, fast)	P_1	162.68 ± 1.90	$2.08{\times}10^7 \pm 4.23{\times}10^5$	7.86 ± 2.67
	P_2	154.52 ± 2.41	$1.96{\times}10^7 \pm 5.02{\times}10^5$	8.29 ± 3.52
Figure 3c (Haploid, 45º, slow)	P_1	219.16 ± 1.30	$3.37{\times}10^7 \pm 3.69{\times}10^5$	20.86 ± 4.49
	P_2	215.68 ± 1.31	$2.89{\times}10^7 \pm 2.36{\times}10^5$	10.43 ± 1.82
Figure 3c (Haploid, 45º, medium)	P_1	166.24 ± 2.06	$2.18{\times}10^7 \pm 4.85{\times}10^5$	11.63 ± 2.84
	P_2	164.88 ± 2.18	$2.17{\times}10^7 \pm 5.18{\times}10^5$	10.56 ± 5.83
Figure 3c (Haploid, 45º, fast)	P_1	160.44 ± 2.06	$2.05{\times}10^7 \pm 3.97{\times}10^5$	7.96 ± 2.29
	P_2	155.24 ± 2.19	$1.93{\times}10^7 \pm 4.86{\times}10^5$	7.19 ± 2.77
Figure 3c (Haploid, 90º, slow)	P_1	217.48 ± 1.25	$3.27{\times}10^7 \pm 2.78{\times}10^5$	20.33 ± 4.91
	P_2	215.72 ± 1.13	$2.87{\times}10^7 \pm 2.71{\times}10^5$	8.57 ± 2.03
Figure 3c (Haploid, 90º, medium)	P_1	164.92 ± 1.55	$2.17{\times}10^7 \pm 3.59{\times}10^5$	10.60 ± 1.82
	P_2	168.68 ± 2.08	$2.15{\times}10^7 \pm 4.43{\times}10^5$	8.25 ± 2.00
Figure 3c (Haploid, 90º, fast)	P_1	156.08 ± 2.25	$1.93{\times}10^7 \pm 5.06{\times}10^5$	8.95 ± 8.51
	P_2	160.44 ± 2.69	$2.00{\times}10^7 \pm 5.46{\times}10^5$	7.51 ± 2.76
Figure 3d (Diploid, 30º, slow)	P_1	214.32 ± 1.14	$8.01{\times}10^7 \pm 7.08{\times}10^5$	18.43 ± 4.68
	P_2	217.00 ± 1.21	$8.05{\times}10^7 \pm 7.60{\times}10^5$	17.45 ± 4.15
Figure 3d (Diploid, 30º, medium)	P_1	173.20 ± 1.80	$5.61{\times}10^7 \pm 1.08{\times}10^6$	8.00 ± 2.11
	P_2	174.92 ± 1.88	$5.61{\times}10^7 \pm 1.10{\times}10^6$	7.89 ± 1.81
Figure 3d (Diploid, 30º, fast)	P_1	175.84 ± 2.65	$5.39{\times}10^7 \pm 1.42{\times}10^6$	6.64 ± 8.76
	P_2	184.92 ± 2.05	$5.75{\times}10^7 \pm 1.00{\times}10^6$	5.46 ± 2.01
Figure 3d (Diploid, 45º, slow)	P_1	213.16 ± 1.27	$7.98{\times}10^7 \pm 9.08{\times}10^5$	18.76 ± 4.32
	P_2	215.08 ± 1.24	$8.00{\times}10^7 \pm 8.54{\times}10^5$	17.63 ± 4.79
Figure 3d (Diploid, 45º, medium)	P_1	169.32 ± 1.79	$5.42{\times}10^7 \pm 1.08{\times}10^6$	8.12 ± 2.11
	P_2	168.36 ± 2.40	$5.26{\times}10^7 \pm 1.33{\times}10^6$	8.90 ± 2.37
Figure 3d (Diploid, 45º, fast)	P_1	167.56 ± 3.52	$5.02{\times}10^7 \pm 1.93{\times}10^6$	7.25 ± 10.39
	P_2	178.92 ± 2.40	$5.54{\times}10^7 \pm 1.41{\times}10^6$	5.26 ± 1.84
Figure 3d (Diploid, 90º, slow)	P_1	215.20 ± 1.19	$8.08{\times}10^7 \pm 6.94{\times}10^5$	18.93 ± 4.75
	P_2	216.56 ± 1.29	$8.02{\times}10^7 \pm 8.52{\times}10^5$	17.45 ± 4.31

Figure 3d (Diploid, 90º, medium)	P_1	169.60 ± 2.03	$5.68\times10^7 \pm 1.02\times10^6$	8.84 ± 2.29
	P_2	171.76 ± 2.39	$5.34\times10^7 \pm 1.48\times10^6$	9.06 ± 4.76
Figure 3d (Diploid, 90º, fast)	P_1	173.12 ± 2.66	$5.28\times10^7 \pm 1.59\times10^6$	6.55 ± 2.92
	P_2	182.48 ± 2.40	$5.77\times10^7 \pm 1.42\times10^6$	5.71 ± 2.75
Figure 4a,c (Haploid, 180º)	P_1	147.32 ± 3.41	$1.75\times10^7 \pm 5.24\times10^5$	6.00 ± 0.83
	P_2	151.13 ± 4.42	$1.78\times10^7 \pm 6.19\times10^5$	7.02 ± 1.06
Figure 4b,d (Diploid, 180º)	P_1	187.72 ± 3.91	$5.81\times10^7 \pm 2.76\times10^6$	4.68 ± 0.98
	P_2	182.84 ± 3.80	$5.33\times10^7 \pm 1.62\times10^6$	4.19 ± 0.37
Figure 5, S6 (1, 10)	P_1	168.17 ± 4.56	$1.67\times10^7 \pm 2.99\times10^5$	9.81 ± 3.48
	P_2	166.64 ± 3.07	$1.77\times10^7 \pm 5.15\times10^5$	4.40 ± 1.74
Figure 5, S6 (2, 5)	P_1	133.56 ± 3.47	$2.97\times10^7 \pm 1.59\times10^6$	26.44 ± 1.42
	P_2	128.76 ± 3.06	$2.73\times10^7 \pm 1.15\times10^6$	25.09 ± 1.05
Figure 5, S6 (5, 2)	P_1	119.88 ± 5.84	$3.33\times10^7 \pm 2.54\times10^6$	43.17 ± 0.99
	P_2	114.92 ± 5.11	$3.04\times10^7 \pm 1.81\times10^6$	43.47 ± 1.05
Figure 5, S6 (10, 1)	P_1	138.36 ± 8.12	$4.09\times10^7 \pm 2.81\times10^6$	47.76 ± 0.90
	P_2	133.84 ± 8.35	$4.64\times10^7 \pm 5.36\times10^6$	46.31 ± 1.44
Figure S4c, S5a (Haploid)	P_1	122	1.24×10^7	2.84
	P_2	138	1.52×10^7	3.44
Figure S4d, S5b (Diploid)	P_1	166	4.26×10^7	1.80
	P_2	196	5.64×10^7	4.77
Figure S4e, S5c (Haploid)	P_1	144	1.19×10^7	2.59
	P_2	168	2.02×10^7	4.07
Figure S4f, S5d (Diploid)	P_1	188	5.23×10^7	2.98
	P_2	202	6.14×10^7	4.45

Figure 1: Conceptual figure illustrating why rapid environmental change is more likely to lead
to reproductive isolation. Both panels show a two-dimensional phenotype with a new optimum
at the center of the sphere, illustrating the adaptive paths taken by two populations (red and
white). In (a), the environment shifted rapidly and favored large-effect mutations; in (b), the
environment shifted slowly and favored small-effect mutations. Rapid environmental change
(a) is more likely to generate genetic incompatibilities that reduce hybrid fitness because
combining large-effect mutations can shift the offspring off of the fitness peak and because
such mutations can have substantial negative pleiotropic effects, here measured as
displacements along the y-axis (see dashed black arrow for one example hybrid combination).
By contrast, slow environmental change (b) is more likely to lead to many small-effect
mutations, which when combined create hybrids whose phenotypes remain near the peak.

Figure 2. Mutation-order speciation and fitness of hybrids in (a, c) haploids and (b, d) diploids.
We arbitrarily consider population 1 to be the "local" population used as the backcross parent,
BC_L (backcross results using the "foreign" population 2, BC_F, are similar). The phenotypic
optimum was displaced from the origin along the x-axis by 2 units, after which the appearance
and fixation of mutations was tracked for a total of (a) 360, (b) 388, (c) 144, (d) 246
substitutions (see Table S2). The majority of fitness loss accumulates during the first adaptation
phase (open circles). Pie charts illustrate the fraction of the hybrid breakdown due to
phenotypic change along the single axis that has undergone environmental change (black),
while the remainder is due to departures from the optimum along other trait dimensions (due
to pleiotropy, red in haploids, blue in diploids). Top panels show $\lambda = 0.04$ and bottom $\lambda = 0.12$.
Parameters: $m=1$, $n=10$, $N=2000$, $q=1.0$, $k = 2.0$, with distributions and solid points
representing the mean $\pm$ SD calculated from one simulation. 1000 individuals were generated
for each hybrid type.
884

Figure 3. Stronger reproductive isolation arises via ecological speciation when populations are selected in more divergent directions, with more isolation arising when mutations have a larger average effect size (top panels) and the environment changes rapidly (bottom panels) in haploids (a, c) and diploids (b, d). For top panels, mutations are drawn from an exponential distribution with $\lambda = 0.04$, 0.08, and 0.12 (three adjacent points from left to right). For bottom panels, the optimum is gradually moved from the origin in one trait dimension over $\tau = 100$ (slow), 50 (medium), 20 (fast) or 0 (immediate) fixation events until the new optimum is reached (four adjacent points from left to right), assuming $\lambda = 0.04$. The x-axis gives the angle between the optima $\mathbf{O}_A$ in patch A and $\mathbf{O}_B$ in patch B relative to the ancestral optimum $\mathbf{O}_0$ (as illustrated on the right bottom). Grey points show the average fitness of the two parental genotypes (we arbitrarily measure all fitnesses in patch A, to which population 1 is adapted). Dashed lines show the expected fitness of the hybrids at the parental mid-point. All other parameters are as in Figure 2 except that this figure is calculated from 50 simulations.

Figure 4. Estimating the degree of ecologically-dependent reproductive isolation. Top panels give $BC_L - BC_F$ for (a) haploids and (b) diploids, as a function of the angle between the optima toward which the parental populations were evolving. Bottom panels give $(BC_L - BC_F)/BC_L$ for (c) haploids and (d) diploids. While $BC_L - BC_F$ shows a non-monotonic relationship, $(BC_L - BC_F)/BC_L$ rises monotonically with the degree of ecological difference between the two environments. Mutations are drawn from an exponential distribution with $\lambda = 0.04$. All other parameters are as in Figure 2, with distributions and solid points representing the mean $\pm$ SE calculated from 50 simulations.

Figure 5. Effect of modularity on the accumulation of reproductive isolation in haploid F_1 hybrids evolving to a common optimum. Panel (a) includes all mutations fixed, while (b) includes only the first 50% of mutations fixed during the adaptation phase and (c) includes only the second 50% of mutations fixed during the adaptation phase; panel (d) includes only mutations fixed during the stationary phase. In each case, the phenotypic effects were added to parents whose phenotypes were at the origin. In (a), pie charts illustrate the fraction of the total hybrid breakdown along the single phenotypic axis experiencing environmental change (black), versus fitness effects on all other trait dimensions. All other parameters are as in Figure 2 with $\lambda = 0.04$, except that this figure changes the number of spheres along the x-axis (from 1 on left to 10 on right) and is calculated from 50 simulations.

Figure 6. Establishment probability of haploid hybrids relative to parents when colonizing a new environment. The optimal phenotype in the new environment is varied along two dimensions (x and y axes), holding the optimum at zero for all other phenotypic dimensions ($n=10$). Left panels show the establishment probability measured in each environment for the best of the two parental genotypes (red dots). Middle panels show the establishment probability of F_1 recombinant hybrids (black dots). Shading indicates establishment probability, from low (blue) to high (orange), as indicated at top. Right panels show the colonization success of hybrids relative to the parents, where the degree of orange indicates the amount by which hybrids have a mean establishment probability greater than the mean of the two parents (fixation probabilities $<10^{-6}$ were set to 10^{-6} to avoid division by small numbers). The potential for hybrid speciation depends strongly on the history of the parental populations: (a-c) diverged in a constant environment (parents from Figure S2a), (d-f) diverged through parallel adaptation to a new optimum at 2 along the x-axis (mutation-order speciation, parents from Figure 2a), (g-i) diverged through adaptation to different environments (ecological speciation with an angle between the two optima of 90°, parents from Figure S5c). All other parameters are as in Figure 2, with $b = 5$ and $\lambda = 0.04$.

Figure S1. Trajectories within haploid simulations of (a) phenotypic effect size among fixed
mutations, (b) selection coefficient, (c) distance to the optimum, and (d) time in units of
generations, all plotted against the number of fixation events. Blue, red, and black represent
maximum, minimum, and mean, respectively. Selection coefficients are positive in the first
"adaptation phase" (taking, on average, 86.9 fixation events) but are more often negative in the
second "stationary phase" (inset to b). In (c), mutations are drawn from an exponential
distribution with $\lambda = 0.04$ (right solid line), 0.08 (center dashed line), and 0.12 (left dashed-
dotted line). All other panels set $\lambda = 0.04$. 100 simulations were run for each parameter set.
The other parameters are $m=1$, $n=10$, $N=2000$, $q=1$ and $k=2$.
941

942   Figure S2. Little reproductive isolation arises during mutation-order speciation during the

943   stationary phase in (a) haploids or (b) diploids. Note that the y-axis range does not start at 0.

944   Data based on the mutations accumulated in Figure 2 (see Table S2 for the number of events).

945   All other parameters are as in Figure 2 with $\lambda = 0.04$.

Figure S3. Trajectories of F_1 fitness relative to the average parental fitness in haploid populations evolving under different environmental scenarios, plotted against the number of fixation events. The trajectories depend on the history of the parental populations: (a) diverged in a constant environment (mutation-order speciation with an angle between the two optima of $0°$), (b-d) diverged through adaptation to different environments (ecological speciation with an angle between the two optima of $30°$, $45°$, and $90°$, respectively). Blue, red, and black represent maximum, minimum, and mean, respectively. F_1 fitness values are shown over time relative to the average fitness of the two parental populations, all measured in patch A at the same time (F_1 hybrids and parents are equal in fitness, on average, along dashed line at one). Inset figures show the slight heterosis that develops early on. All other parameters are as in Figure S1 except that 50 simulations were run for each parameter set.

Figure S4. The fitness of hybrids can be predicted using the Brownian bridge approximation of Simon *et al.* (2018). Left and right panel represent haploid and diploid simulations respectively. Top panels show hybrid fitnesses measured in patch A following mutation-order speciation in the same parental environments (as in Figure 2a,b). Remaining panels show hybrid fitnesses measured in patch A following ecological speciation in populations adapting to optima that differ by 30° (middle panels as in Figure S5a,b) or 90° (bottom panels; as in Figure S5c,d). The dots represent data generated data in the current paper. The dashed line is predicted by fitting the free parameter $E(S_{\dagger})$ of Simon *et al.* (2018) by minimizing the sum of squared deviations between the observed and predicted fitnesses. Three different colors represent three different simulation runs. Choosing $E(S_{\dagger})$ based on the class of hybrid with the lowest fitness, as suggested by Simon et al. (2018), does not always provide a good fit, especially when the parents differ in fitness, but fitting the $E(S_{\dagger})$ parameter provides a robust predictor to the full set of hybrid fitnesses.

Figure S5. Ecological speciation and the fitness of hybrids in (a, c) haploids and (b, d) diploids. We illustrate the case where the "local" environment is that of population 1 (P_1), with fitnesses measured in its patch A and using $\lambda = 0.04$. The angle between the optima of the two environments is set to 30° in top panels and 90° in bottom panels, relative to the origin (see Table S1 for details). All other parameters are as in Figure 2, with distributions and solid points representing the mean $\pm$ SD calculated from one simulation. 1000 individuals were generated for each hybrid type.

977   Figure S6. The magnitude of the largest mutation fixed in each of the 50 simulations illustrated

978   in Figure 5 (one-way ANOVA, p>0.05). All parameters are the same as in Figure 5, with

979   distributions and solid points representing the mean $\pm$ SD.