1

Predicting gait adaptations due to ankle plantarflexor muscle weakness and contracture using physics-based musculoskeletal simulations

10   Carmichael F. Ong[1], Thomas Geijtenbeek[2], Jennifer L. Hicks[1], Scott L. Delp[1,3,4*]

14   [1] Department of Bioengineering, Stanford University, Stanford, California, United States of
15   America
16
17   [2] Department of Biomechatronics & Human-Machine Control, Delft University of Technology,
18   Delft, The Netherlands
19
20   [3] Department of Mechanical Engineering, Stanford University, Stanford, California, United
21   States of America
22
23   [4] Department of Orthopaedic Surgery, Stanford University, Stanford, California, United States of
24   America
25

26   * Corresponding author

27   Email: delp@stanford.edu (SLD)

Abstract

Deficits in the ankle plantarflexor muscles, such as weakness and contracture, occur commonly in conditions such as cerebral palsy, stroke, muscular dystrophy, and Charcot-Marie-Tooth disease. While these deficits likely contribute to observed gait pathologies, elucidating cause-effect relationships is difficult due to the often co-occurring biomechanical and neural deficits. To elucidate the effects of weakness and contracture, we systematically introduced isolated deficits into a musculoskeletal model and generated simulations of walking to predict gait adaptations due to these deficits. We developed a planar model containing 9 degrees of freedom and 18 musculotendon actuators, and an optimization framework through which we imposed simple objectives, such as minimizing cost of transport while avoiding falling and injury, and maintaining head stability. We first validated that our model could generate gaits that reproduced experimentally observed kinematic, kinetic, and metabolic trends for two cases: 1) walking at prescribed speeds between 0.50 m/s and 2.00 m/s and 2) walking at self-selected speed. We then applied mild, moderate, and severe levels of muscle weakness or contracture to either the soleus (SOL) or gastrocnemius (GAS) or both of these major plantarflexors (PF) and retrained the model to walk at a self-selected speed. The model was robust to all deficits, finding a stable gait in all cases. Severe PF weakness caused the model to adopt a slower, "heel-walking" gait. Severe contracture of only SOL or both PF yielded similar results: the model adopted a "toe-walking" gait with excessive hip and knee flexion during stance. These results highlight how plantarflexor weakness and contracture may contribute to observed gait patterns. Our model, simulation and optimization framework, and results are freely shared so that others can reproduce and build upon our work.

Author summary

Deficits in the muscles that extend the ankle are thought to contribute to abnormal walking patterns in conditions such as cerebral palsy, stroke, muscular dystrophy, and Charcot-Marie-Tooth disease. To study how deficits in these muscles contribute to abnormal walking patterns, we used computer simulations to systematically introduce muscle deficits into a biomechanically accurate model. We first showed that our model could discover realistic walking patterns over a wide range of speeds when we posed a simple objective: walking while consuming a minimum amount of energy per distance, maintaining head stability, and avoiding injury. We then used the model to study the effect of two commonly observed problems: muscle weakness and muscle tightness. We found that severe weakness of the ankle extensors caused the model to adopt a slower, "heel-walking" gait, and severe tightness caused the model to adopt a crouched, "toe-walking" gait. These results highlight how deficits in the ankle extensor muscles may contribute to abnormal walking patterns commonly seen in pathological populations.

Introduction

The ankle plantarflexor muscles play an important role in human walking, and deficits in these muscles are thought to contribute to gait pathologies. Previous work has shown that the plantarflexor muscles are important for generating forward acceleration during the push-off phase of walking [1,2]. Individuals with cerebral palsy, stroke, muscular dystrophy, or Charcot-Marie-Tooth (CMT) disease often exhibit plantarflexor muscle weakness [3–9] and contracture [3,9–15]. However, these muscular deficits often co-occur with other abnormalities, such as skeletal deformities and neural deficits, making it difficult to assess the independent contributions of these muscular deficits to gait abnormalities.

Previous experimental work has sought to better understand the cause-effect relationship

between plantarflexor weakness and gait pathologies by performing a tibial-nerve block in one leg

in healthy subjects [16]. Adaptations on the affected side included a reduced ability to transfer

weight to the contralateral leading limb during terminal stance and a shortened single-limb stance

duration. Although this experiment helps to establish a cause-effect relationship, nerve blocks are

invasive and thus difficult to repeat for many muscles. Further, these experiments cannot reveal

the effects of contracture, which produces abnormally high passive stiffness of muscles.

Musculoskeletal simulations built from experimental gait data have been used to study gait

pathologies. For example, simulations of individuals with cerebral palsy have quantified individual

muscle contributions to body weight support and forward propulsion [17], the minimum muscle

strength required to walk in a crouch gait [18], and the contributions of contracture and spasticity

to increased hamstring resistance [19]. These studies suggest strong links between muscle deficits

and the observed gait adaptations; however, since these studies tracked experimental data from

patients with a combination of muscular, skeletal, and neural deficits, the independent effects of

muscular weakness and contracture on the observed gait adaptations cannot be assessed.

Simulations in which kinematics are generated *de novo* (i.e., without tracking experimental

data) can help reveal cause-effect relationships between muscular deficits and gait abnormalities.

Researchers have created controllers that could generate various gait patterns, including level

walking [20–23] and running [21,23], inclined walking [22,23], loaded walking [22], stair ascent,

and turning [23]. More recently, Song and colleagues sought to understand which factors

contributed to decreased walking performance in the elderly population [24]. By systematically

introducing the neural, muscular, or skeletal deficits seen in this population and training each

model to walk, they determined that decreased muscle strength and mass of all muscles contributed

most to typically observed gait adaptations. Although simulations of this type are valuable, they

are not yet broadly used because it is challenging to generate simulations *de novo* and to reproduce

previous work. Overcoming these challenges has been difficult because of the sensitivity of

forward simulations to a software environment and limited sharing of software and models. More

work is also needed to validate that simulations of this type capture the adaptations observed in

experiments (e.g., due to varying walking speeds), before adjusting the model to generate

simulations under new conditions that cannot be observed in controlled experiments.

The goal of our work was to determine which gait adaptations arise from weakness or

contracture of the plantarflexor muscles. To this end, we first created and validated an optimization

framework and musculoskeletal model that could generate realistic motions *de novo*. Our

controller followed the previously described reflex-based controllers [20,21,23], and the

parameters of this controller were iteratively updated within the optimization framework. We

validated our results over a wide range of prescribed walking speeds and at self-selected speed;

this step was necessary as low gait speed is commonly observed in individuals with gait

impairments. We then introduced weakness or contracture in the plantarflexor muscles, generated

new gait patterns, and analyzed subsequent changes in the kinematics and kinetics of gait. Our

model and results are freely available at https://simtk.org/projects/pfdeficitsgait along with a

Docker build file and setup files so others can reproduce our work. We also provide downloads

for the control and optimization framework (SCONE, Simulation and CONtrol Environment) and

accompanying documentation at http://scone.software to enable others to use and build upon our

simulation framework.

Results and Discussion

We generated our simulations using a single shooting method (Fig 1), which is discussed

in more detail in the Methods section. We first compare our simulations of unimpaired walking to

experimental kinematic, kinetic, and metabolic data. We then detail how we modeled plantarflexor

weakness and contracture and discuss the kinematic and kinetic adaptations caused by introducing

these deficits into our musculoskeletal model. Video highlights of our results are provided in S1

Video, and videos of all simulations discussed here are provided in S2 Video.

Fig 1. Single shooting framework for dynamic optimization. We trained a planar,

musculoskeletal model actuated by 18 Hill-type musculotendon actuators by optimizing the

parameters of a gait controller based on an objective function that sought to minimize metabolic

cost, avoid falling and injury, and stabilize the head. A gait controller computed muscle

excitations, u(t), for a musculoskeletal model to generate a forward simulation. Sensory feedback,

based on the model's muscle and joint states, was used in a feedback loop with the gait controller.

The objective function quantified the performance of each simulation, and a Covariance Matrix

Adaptation Evolutionary Strategy (CMA-ES) optimization method updated the values of the

variables in the optimization problem.

Validating the model's gait over a range of speeds

We validated the ability of our model and optimization framework to capture trends in

kinematics, kinetics, and spatio-temporal measures when walking at different speeds by generating

seven simulations of gait at prescribed speeds between 0.50 m/s and 2.00 m/s, at intervals of 0.25

m/s. All simulations found a steady gait pattern at a speed within 0.05 m/s of the speed prescribed

by the optimization's objective function (S2 Video). Individual comparisons of each speed to

experimental data [25] are provided in S1 Fig.

Simulated kinematic and kinetic adaptations, including joint angles, joint moments, and

ground reaction forces (GRFs), matched trends observed in experimental data [25] with increasing

speeds (Fig 1). Hip and ankle joint angle excursions increased with increasing speeds. Kinetic data,

such as peak flexion and extension moments about the hip, knee, and ankle, and the peak horizontal

and vertical GRFs, also increased with increasing speeds. Some experimental trends, however,

were not captured in our simulations. The knee joint angle range and peak flexion during swing

did not increase at higher speeds. At the slowest prescribed speeds (0.50 m/s and 0.75 m/s), the

GRFs show a slow resonance not seen in experimental data. This was likely due to our choice for

contact parameters, which yielded a softer contact than previous work [22]. While this choice

likely contributed to resonance, it also sped simulations up by roughly 2- to 3-fold. We note one

aberrant point for the simulation at the slowest speed (0.50 m/s) in the knee joint moment plot that

shows a spike of knee extension in early stance, but it does not appear to affect other kinematic or

kinetic measures. Overall, while these results show that our simulations capture many salient

kinematic and kinetic trends, we must be cautious in interpreting GRF results at speeds at or below

0.75 m/s.

Fig 2. Kinematics and kinetics of simulated walking over a range of speeds. Seven prescribed

speeds between 0.50 m/s (blue) and 2.00 m/s (red) at intervals of 0.25 m/s were analyzed. Joint

angles (left column) and joint moments (middle column) are plotted for the hip (top row), knee

(middle row), and ankle (bottom row). Positive joint angles indicate flexion, and positive joint

moments indicate extension. Ground reaction forces (GRFs) (right column) are shown for both

horizontal (top row) and vertical (middle row) directions. With exceptions at the knee joint, we

observed trends that occur in experimental data [25], including greater joint angle ranges, joint

moments, and ground reaction forces at higher speeds.

Percent stance phase (i.e., the percent of time each foot is in stance during a gait cycle)

decreases, and step length and cadence increased with increasing speeds as observed

experimentally [25–28] (Fig 2). Percent stance phase was within 2 standard deviations (SD) of

experimental data at all speeds. Although the trends of step length and cadence follow

experimental data, the simulations generated gaits with a longer step length and a slower cadence

than observed experimentally. Our simulations were also able to reproduce experimentally

observed trends in metabolic cost of transport over the range of speeds we tested (Fig 2, right

panel). In particular, a greater increase in cost of transport was observed when walking slower than

optimal as compared to faster than optimal.

Fig 3. Spatio-temporal and metabolic parameters over a range of speeds. Simulated (dots) and

experimental (lines) data are plotted for percent stance phase (left), step length (middle left),

cadence (middle right), and cost of transport (right). Experimental data for the left three panels are

shown with 2 standard deviations [25]. In the right panel, the magnitude and characteristic shape

of the cost of transport bowl were similar between the simulated data and experimental data sets

[26–28].

Validating self-selected gait of the model

We tested the ability of our model and optimization framework to generate a realistic gait

at self-selected speed by generating simulations that did not have a prescribed speed within the

optimization's objective function. Instead, we imposed a minimum speed of 0.75 m/s, which

helped guide the optimizer to better solutions but was sufficiently low that it did not affect the final

optimized solutions. We generated 3 simulations by initializing the optimization framework with

the optimized parameters for 3 of the prescribed speeds from the previous section: slowest (0.50

m/s), middle (1.25 m/s), and fastest (2.00 m/s) speeds. The optimization yielded 3 simulations with

similar speeds of 1.23 m/s, 1.21 m/s, and 1.29 m/s, respectively. Each speed was within 2 SDs of

the expected self-selected speed (1.25 ± 0.15 m/s) for an individual of our model's leg length [25].

This suggests that our optimization framework can robustly find solutions and is not sensitive to

initial guess. For simplicity, only the simulation initialized with the middle speed is described in

the remainder of this section.

Our simulated kinematic and kinetic trajectories, including joint angles, joint moments,

and GRFs, were similar both in value and shape to experimental data for self-selected gait (Fig 4).

For the majority of the gait cycle, simulated trajectories were within 2 SDs of experimental data

describing self-selected gait. For each of the trajectories, the root-mean-squared error (RMSE)

between simulated and experimental data was no more than 1.55 SD, and for each trajectory except

the knee extension moment, the normalized cross-correlation (NCC) [29], which measures shape

similarity between the simulated and experimental data sets, was at least 0.82 (Table 1).

Fig 4. Kinematics and kinetics of self-selected gait. Simulated (black line) kinematics and

kinetics are compared to experimental data (gray area) [25]. Joint angles (left column) and joint

moments (middle column) are plotted for the hip (top row), knee (middle row), and ankle (bottom

row). Ground reaction forces (right column) are shown for both horizontal (top row) and vertical

(middle row) directions. Positive angles indicate flexion, and positive moments indicate extension.

Note that the experimental data for hip flexion angle was shifted by 11.6° to account for the

difference in pelvis orientation definitions between the experimental data set and the

musculoskeletal model.

Table 1. Similarity metrics between simulated and experimental data for self-selected gait.

	Angles			Moments			GRFs (stance only)	
	Hip	**Knee**	**Ankle**	**Hip**	**Knee**	**Ankle**	**Horizontal**	**Vertical**
Root-mean-squared error (SD)	1.55	1.48	0.82	1.08	1.46	1.17	1.04	0.78
Normalized cross-correlation	0.95	0.97	0.82	0.83	-0.04	0.95	0.95	0.99

Root-mean-squared error compares simulation mean trajectories to experimental mean and

standard deviation, reported in units of standard deviation (SD). Given the experimental mean and

standard deviation over the gait cycle, we compute a Z-score for the simulation data at every 1%

of the gait cycle. Then, we take the square root of the sum of squares of all the Z-scores.

Normalized cross-correlation compares mean trajectories between simulation and experimental

data [25].

The three trajectories with the highest RMSEs, hip joint angle (1.55 SD), knee joint angle

(1.48 SD), and knee joint moment (1.46 SD), each contained times when the simulation deviated

by more than 2 SDs from the experimental mean. The hip and knee joint angles showed excessive

flexion during the swing phase. These were likely compensations due to the simplifications in the

model, as degrees of freedom that contributed to foot clearance, such as pelvic list and hip

abduction, were not included in the model. Despite the larger RMSEs, the high NCCs of the

simulated hip (0.95) and knee (0.97) joint angles to their corresponding experimental trajectories

indicate the shape of the simulated and experimental trajectories matched well. The simulated and

experimental knee extension moments did not have a similar shape, with a NCC of -0.04. However,

there were large standard deviations in the experimental data during this part of the gait cycle,

which is not considered when computing NCC. Differences during early stance explain this

discrepancy in knee joint moment trajectories, as the model generated a knee flexion moment when

experimental data exhibits a knee extension moment. The model took long steps and landed with

an extended knee, causing the GRF to generate a knee extension moment, rather than a knee flexion

moment in early stance. Thus, the muscles had to generate a knee flexion moment, rather than a

knee extension moment, to prevent hyperextension.

Simulating walking with plantarflexor weakness or contracture

The results of the previous two sections gave us confidence that this model and

optimization framework could generate realistic gaits over a wide range of speeds, and could self-

select a realistic gait and speed. We then used the framework to study how gait would adapt to

deficits in the plantarflexors by adjusting the model's parameters and using the same optimization

framework.

We modeled mild, moderate, and severe weakness by reducing a muscle's maximum

isometric force (F_o^m) to 25%, 12.5%, and 6.25%, respectively, of its original value. The resulting

values for F_o^m were similar to those used in a previous study that computed the minimum

plantarflexor muscle strength needed to walk either normally or in a crouch [18]. We modeled

mild, moderate, and severe contracture by reducing a muscle's optimal fiber length (l_o^m) to 85%,

70%, and 55%, respectively, of its original value. The severe case was based on experimental data

[30] and previously used as a model of contracture [31]. We separately applied each of these 6

deficits to the soleus only (SOL), gastrocnemius only (GAS), or both plantarflexors (PF), to yield

18 total deficit cases.

The optimization framework generated a stable gait in all deficit cases, highlighting its

ability to change neural control in response to weakness and contracture (S2 Video). Furthermore,

the model's gait was robust to most deficits, and many cases had kinematic and kinetic measures

within 2 SDs of walking at a self-selected speed (Fig 5, S2 Fig). By count, the most affected cases

were moderate PF weakness, severe PF weakness, severe SOL contracture, and severe PF

contracture. In the following sections, we detail the weakness and contracture cases that caused

the model to adopt a gait with kinematic or kinetic features outside of 2 SDs of normal walking at

a self-selected speed. We provide complete kinematic and kinetic trajectories for all simulations

of muscle weakness and contracture in S3 Fig.

Fig 5. Z-scores of key kinematic and kinetic measures of gait for all deficit cases. Z-scores

were computed using previous experimental data of individuals walking at a self-selected speed

[25]. All 18 cases (dots) are plotted for each measure. A black dot indicates any case within 2 SDs

of normal (gray band). A blue or orange dot indicates either a weakness or contracture case,

respectively, which is outside of 2 SDs. All of these cases are labeled with the affected muscle or

muscles: soleus (SOL), gastrocnemius (GAS), or both (PF); and severity level: mild (Mild),

moderate (Mod), or severe (Sev). One other case is labeled (see Peak Plantarflexion Moment, PF

Mod*) as it was a clear outlier compared to other deficit cases. The count above each line (e.g., *n*

= 13) indicates the number of unlabeled dots.

Walking with plantarflexor weakness

Plantarflexor weakness decreases the capacity of muscles to generate ankle plantarflexion

moments, which are critical for generating forward propulsion during walking [1,2,16]. Our

simulations supported this notion as plantarflexion moments during walking were most affected

by moderate and severe PF weakness and minimally affected by SOL or GAS weakness only (Fig

5, third row). This suggests that in cases of weakness to only the SOL or GAS, the unimpaired

plantarflexor had sufficient capacity to compensate for weakness in the other muscle. For moderate

and severe PF weakness, peak ankle plantarflexion moment decreased by 43% and 69%,

respectively, when compared to simulated unimpaired walking (Fig 6, bottom row). This

highlights the reduced moment generating capacity in these cases.

Fig 6. Ankle kinematics and kinetics for moderate and severe PF weakness. Simulated ankle

kinematics and kinetics for unimpaired walking (black lines) and walking with PF weakness (blue

lines) are compared to experimental data (gray area) [25]. Ankle dorsiflexion (top) and ankle

plantarflexion moment (bottom) are plotted for cases of moderate (left) and severe (right) PF

weakness.

Other gait parameters, such as peak horizontal GRF, self-selected walking speed, and

percent stance were abnormal for moderate and severe PF weakness as well (Fig 5). Compared to

simulated unimpaired walking, peak horizontal GRF decreased by 55% and 70% for moderate and

severe PF, respectively, which led to slower walking speeds of 1.01 m/s and 0.87 m/s, respectively.

These decreases in gait speed in response to weakness are unsurprising given the direct relationship

between the push-off phase and walking speed. Reduced walking speed is commonly observed in

individuals with cerebral palsy [32], stroke [33], muscular dystrophy [34], and CMT disease [35],

where plantarflexor weakness is common. After accounting for these reduced walking speeds, the

model spent an increased percentage of the gait cycle in stance phase, 69% and 74% with moderate

and severe PF weakness, respectively. For comparison, simulated unimpaired walking at a slower

speed of 0.75 m/s had a percent stance phase of only 63% (Fig 3). These large increases in percent

stance phase are consistent with trends seen in the pathological population. For example, CMT

patients who had a "plantar flexor strength deficit" were found to adopt a slower gait with an

increased time spent in stance phase compared to the gait chosen by patients without the strength

deficit [36].

Dramatic kinematic compensations at the ankle were observed at the ankle in cases of

moderate and severe PF weakness. These models adopted a calcaneal, or "heel-walking", gait,

landing with substantially increased ankle dorsiflexion that was maintained throughout stance (Fig

6, top row). When compared with simulated unimpaired walking, the model with moderate or

severe PF weakness had an increased ankle dorsiflexion at initial contact of 15° and 16°,

respectively, and an increased average ankle dorsiflexion during stance of 7° and 11°, respectively.

This type of gait is often observed in patients who have received either an intramuscular

aponeurotic recession of the plantarflexors or Achilles tendon lengthening to correct contracture

[37–39], and it is commonly thought that overlengthening during these procedures can lead to

plantarflexor weakness [40,41]. Our simulations support the notion that plantarflexor weakness

alone could cause calcaneal gait.

While it has been previously hypothesized that plantarflexor weakness may contribute to

individuals adopting a crouch gait [18,38,39,41], our simulations did not support this hypothesis.

Crouch gait is characterized by increased knee flexion during stance and commonly occurs with

increased hip flexion during stance, but our optimization framework found gaits that did not

require substantial plantarflexion moment and did not have increased knee and hip flexion during

stance (Fig 5, bottom two rows). Our results also differ from previous work that observed a mild

crouch with weakened plantarflexors [21], but a key difference was that our simulations did not

have a constrained speed whereas the previous work had a prescribed speed of 1.25 m/s. Although

our work does not support the notion that plantarflexor weakness alone will lead to crouch gait, it

does not rule out that it could be a contributing factor. For example, it might be that plantarflexor

weakness is often associated with weakness and contracture in other muscle groups, and a

combination of these deficits along with impaired neural control may lead to crouch gait.

331　Walking with plantarflexor contracture

Plantarflexor contracture increases passive ankle plantarflexion moments, shifting the

ankle's resting position to a more plantarflexed state and limiting ankle range of motion [11,13].

All cases of severe contracture (i.e., SOL, GAS, and PF) had significantly decreased peak values

of dorsiflexion in stance (Fig 5, fifth row). One case, mild SOL contracture, appeared to have an

opposite effect, but we believe this is an anomaly as the overall ankle position trajectory did not

differ much from the unimpaired case (S3D Fig). Models with severe contracture landed on their

toes as the ankle was plantarflexed beyond 0° at initial contact, and stayed on their toes throughout

stance (Fig 7, top row). Equinus, or "toe-walking," gait is commonly observed in individuals with

cerebral palsy [42,43]. Plantarflexor contracture is thought to be a major contributing factor

[11,44], and our results support this theory.

343　Fig 7. Ankle kinematics and kinetics for severe contracture. Simulated ankle kinematics and

kinetics for unimpaired walking (black lines) and walking with severe contracture (orange lines)

are compared to experimental data (gray area) [25]. Ankle dorsiflexion (top) and ankle

plantarflexion moment (bottom) are plotted for cases of SOL (left), GAS (middle), and PF (right)

contracture.

All cases of severe contracture had significantly increased plantarflexion moments during

early stance (Fig 7, bottom row), which was expected since an equinus gait necessitates large

plantarflexion moments at initial contact to prevent excessive dorsiflexion. In particular, severe

SOL or PF contracture had a peak plantarflexion moment during early stance rather than during

the push-off phase. Although peak values were largely unaffected, plantarflexion moments during

the push-off phase did decrease which led to a subsequent decrease in peak horizontal GRF; all

contracture cases had a peak horizontal GRF that was between 2% and 23% lower than our

simulated unimpaired case (Fig S2, fourth row).

As severity of contracture increased, models adapted by walking slower and with a shorter

stance phase. The three slowest conditions, which included severe SOL, moderate PF, and severe

PF contracture, all walked at a significantly slower speed of 1.08 m/s with a slightly smaller percent

stance phase of 59% (Fig 5, top two rows). This small decrease in percent stance phase agrees well

with experimental data of individuals walking in equinus [45].

In cases of severe SOL, GAS, and PF contracture, the model adopted gaits that were more

crouched throughout the whole stance phase, as measured by increased knee and hip flexion, when

compared to simulated unimpaired walking (Fig 8). Cases of SOL and PF contracture were more

crouched than cases of GAS contracture (S2 Fig, bottom two rows). These results highlight two

conclusions. First, contracture in the plantarflexor muscles can lead to dramatic compensations at

joints that they do not span. Second, because contracture of only GAS had the smallest effect, our

results suggest that the crouched postures adopted by the model were affected less by the GAS

directly flexing the knee and more by walking in equinus.

Fig 8. Hip and knee kinematics for severe contracture. Simulated hip and knee kinematics for

unimpaired walking (black lines) and walking with severe contracture (orange lines) are compared

to experimental data (gray area) [25]. Hip flexion (top) and knee flexion (bottom) are plotted for

cases of SOL (left), GAS (middle), and PF (right) contracture.

Limitations and future work

Simplifications in the model may have affected the observed kinematic trajectories. The

model was constrained to sagittal plane motion and consequently lacked degrees of freedom, such

as hip abduction, pelvis list, and pelvis rotation, which allow for out-of-plane motions. This made

foot clearance more difficult and was likely the cause of compensations in the sagittal plane that

increased knee and hip flexion during swing. For the models with deficits, these planar constraints

did not allow for commonly observed compensations that are out of the sagittal plane, such as hip

circumduction [46,47].

Although we chose to model weakness by decreasing F_o^m and contracture by decreasing l_o^m,

there are other ways to model these deficits. For instance, weakness could also be modeled as

decreased maximum muscle excitation, which may be important in cases such as stroke [48], and

contracture could also be modeled as decreased tendon slack length [49] or increased passive

muscle stiffness [50]. These alternate modeling approaches may have led to different gait

compensations, and future work to test sensitivity of our results to variations in modeling choices

would be valuable.

391   Although our objective function contained terms that are thought to capture high-level

392 goals of walking, such as minimizing cost of transport [51] while maintaining head stability [52]

393 (see Methods), it may not comprehensively represent goals of human walking, especially amongst

394 individuals with gait pathologies. For example, some individuals may choose a trade-off between

395 metabolic efficiency and joint loading or stability to perturbations. Furthermore, the relative trade-

396 off between these competing objectives likely varies between individuals.

397   Due to the non-convex nature of the optimization problem and the stochastic nature of our

398 optimizer, the solutions provided here are likely to be local minima rather than global minima. To

399 gain confidence in our solutions, we followed previous work that used multiple optimizations from

400 the same seed and restarted optimizations from the previous best result [23,53], and furthermore,

401 we demonstrated how our solution was robust to different initial seeds (see section on "Validating

402 self-selected gait of the model"). This strategy, however, was computationally expensive, and

403 further work is needed to better search this complex, high-dimensional space. Since a good starting

404 guess can be important for finding reasonable results, we provide our results to assist other

405 researchers in future studies.

406   Although this work presents a validated framework for studying how isolated changes in

407 muscle parameters may lead to changes in gait, future work on how gait adapts with changes in

408 neural control will further our understanding of pathologic gait. Reflex-based controllers, like the

409 one used here, were useful to ensure that the optimization problem was tractable, but a controller

410 like this cannot predict changes in the underlying control structure. Recent advances in the field of

411 reinforcement learning may be able to generate robust simulations with control patterns that do

412 not have a specified and simplified control structure [54,55].

Conclusion

We present a model and optimization framework suitable for studying sagittal plane adaptations during walking due to muscular deficits. We first validated that, over a wide range of speeds, our model and framework captured many salient features of gait. This framework was then used to study how the model would adapt its gait when plantarflexor weakness or contracture was introduced. Severe plantarflexor weakness caused the model to adopt a calcaneal gait without crouching, while severe plantarflexor contracture caused the model to adopt an equinus gait. Our simulations also showed that contracture of only SOL or both PF caused the model to walk in a more crouched posture than if contracture was applied to only GAS, which suggests that walking in equinus is the major contributor to crouch gait rather than force from the GAS directly flexing the knee. We provide our results freely at https://simtk.org/projects/pfdeficitsgait and software at http://scone.software so that others can easily extend this work.

Methods

Our optimization framework used a single shooting method to solve the dynamic optimization problem of generating a simulation of gait (Fig 1). We implemented our model in OpenSim 3.3 [56,57] and used an optimization and control framework (SCONE) to implement the gait controller, perform the simulation using OpenSim as the plant, and optimize the parameters of our problem.

Musculoskeletal model

The musculoskeletal model (Fig 9) was a planar model based on a previous model used to study the lower limb [58]. The musculoskeletal model had 9 degrees of freedom (dof): a 3-dof

planar joint connected the pelvis to ground; a 1-dof pin joint represented each hip and ankle; and

a 1-dof joint with coupled rotation and translation represented each knee. The lumbar joint was

locked at 5° of flexion based on previous walking data [2].

Fig 9. A planar musculoskeletal model for walking. The musculoskeletal model had 9 degrees

of freedom (dof): the pelvis-to-ground joint was a 3-dof planar joint, each hip and ankle was a 1-

dof pin joint, and each knee was a 1-dof coupled joint. The model's 18 musclulotendon actuators

(red lines) represented the 9 major uniarticular and biarticular muscle groups per leg that drive

sagittal plane motion: iliopsoas (ILPSO), gluteus maximus (GMAX), rectus femoris (RF),

biarticular hamstrings (HAMS), vasti (VAS), biceps femoris short head (BFSH), gastrocnemius

(GAS), tibialis anterior (TA), and soleus (SOL). A compliant contact model was used to generate

forces between the spheres at the heel and toes of the feet and the ground plane.

The model was actuated using eighteen Hill-type muscle-tendon units [59], nine per leg.

Muscles with similar functions in the sagittal plane were combined into single muscle-tendon units

with combined peak isometric forces to represent 9 major muscle groups per leg: iliopsoas

(ILPSO), gluteus maximus (GMAX), rectus femoris (RF), biarticular hamstrings (HAMS), vasti

(VAS), biceps femoris short head (BFSH), gastrocnemius (GAS), tibialis anterior (TA), and soleus

(SOL). Peak isometric forces were based on a previous musculoskeletal model [60], whose muscle

volumes were based on young, healthy subjects [3]. The tendon strain at peak isometric force was

4.9% [60,61] for all muscles except for the plantarflexors, whose values were set to 10% [62]. To

better match passive muscle forces during walking [63], we adjusted the passive force-length

curves of the HAMS, VAS, and RF to decrease their passive forces (S1 Table). The tendon slack

length of each muscle was calculated based on experimental data [64], as done previously in other

musculoskeletal models [60,65]. We set maximum muscle fiber contraction velocity to 15 l_o^m/s

[62,66]. All muscle-tendon parameters are summarized in S1 Table.

Ligaments were modeled as nonlinear, rotational springs that generated torques when a

joint was hyperflexed or hyperextended. In our model, ligaments generated torques when the hip

was flexed beyond 120° or extended beyond 30°; the knee was flexed beyond 140° or extended

beyond 0°; and the ankle was dorsiflexed beyond 20° or plantarflexed beyond 40°.

A compliant contact model [67,68] was used to generate forces between the feet and the

ground. Each foot had three contact spheres: one sphere with a radius of 5 cm represented the heel,

and two spheres, each with a radius of 2.5cm, represented the toes. All contact spheres had the

following parameters: stiffness coefficient of 500,000 N/m$^{3/2}$, dissipation coefficient of 1.0 s/m,

static and dynamic friction coefficients of 0.8, viscous friction coefficient of 0, and transition

velocity of 0.1 m/s.

As described in the Results and Discussion, we introduced two types of deficits into the

musculoskeletal model: weakness and contracture. We modeled mild, moderate, and severe

weakness by reducing the peak isometric force of a muscle (F_o^m) by 25%, 12.5%, and 6.25%,

respectively, of the value in the unimpaired model. We modeled mild, moderate, and severe

contracture by reducing the optimal fiber length of a muscle (l_o^m) by 85%, 70%, and 55%,

respectively, of the value in the unimpaired model. All 6 of these muscle deficits were applied to

the SOL, the GAS, or both the SOL and GAS, for a total of 18 cases.

Gait controller

The gait controller was based on previous reflex-based controllers for human locomotion

[20,21,23] and used a combination of a high-level state machine and low-level control laws to

calculate muscle excitations. Our controller had 5 high-level states that represented key points in

the gait cycle for each leg: early stance (ES), mid-stance (MS), pre-swing (PS), swing (S), and

landing preparation (LP). Of the 5 transitions between these high-level states, transitions between

4 of the states were controlled by 4 thresholds that were free parameters in the optimization: 1) ES

to MS: horizontal distance between the ipsilateral foot and pelvis was lower than a threshold; 2)

PS to S: GRF on the ipsilateral foot was lower than a threshold; 3) S to LP: horizontal distance

between the ipsilateral foot and pelvis was greater than a threshold; and 4) LP to ES: GRF on

ipsilateral foot was greater than a threshold. The fifth transition, MS to PS, was not controlled by

a free parameter and occurred when the contralateral leg entered ES state.

Low-level control laws were active or inactive depending on the high-level controller (Fig

10). There were 5 types of control laws: constant (C), length feedback (L), velocity feedback (V),

force feedback (F), and proportional-derivative (PD). All muscle-based feedback laws were

positive feedback (i.e., L+, V+, F+) onto the same muscle, except for a negative force feedback

(F-) from the soleus to the tibialis anterior. PD control was only used to control the pelvis tilt angle

(θ) and velocity ($\dot{\theta}$) with respect to ground using the muscles crossing the hip joint (i.e., ILPSO,

GMAX, and HAMS). Excitations (u) are calculated over time (t) by the following equations:

$$u_C = K_C \tag{1}$$

$$u_{L+} = \max\left(0, K_{L+}\left[l(t - t_D) - l_o\right]\right) \tag{2}$$

$$u_{V+} = \max\left(0, K_{V+}\left[v(t - t_D)\right]\right) \tag{3}$$

$$u_{F\pm} = \pm K_{F\pm} F(t - t_D) \tag{4}$$

$$u_{PD} = K_p\left[\theta(t - t_D) - \theta_o\right] + K_v\left[\dot{\theta}(t - t_D)\right] \tag{5}$$

States calculated from the model were muscle length (l), muscle velocity (v), muscle force (F),

and pelvis tilt orientation (θ) and velocity ($\dot{\theta}$). In total, there were 70 free parameters of the

optimization, including the controller gains (K_C, K_{L+}, K_{V+}, $K_{F\pm}$, K_p, and K_v), and offsets for muscle

length feedback (l_o) and proportional feedback of θ (θ_o). For all positive feedback and PD control

laws, the parameter for time delay, t_D, was set for each muscle depending on the most proximal

joint over which the muscle crosses: t_D was 5 ms for the hip, 10 ms for the knee, and 20 ms for the

ankle [20]. For the F- law from the soleus to the tibialis anterior, t_D was 40 ms. When considered

with a first-order activation time constant of 10 ms, this delay better represented the short-latency

TA suppression (56 to 74 ms) observed in experiments [69]. Overall, however, the delays used

were still shorter than those measured experimentally [70].

Fig 10. The gait controller used a combination of a state machine and low-level control laws

to determine excitations. The state machine had three states in stance (i.e., early stance (ES), mid-

stance (MS), and pre-swing (PS)) and two states in swing (i.e., swing (S) and landing preparation

(LP)), and it determined when low-level control laws were active. Low-level control laws included

constant signals (C) and feedback terms based on muscle length (L), muscle velocity (V), muscle

force (F), and proportional-derivative control (PD) based on the pelvis tilt angle. Positive and

negative feedback are denoted by (+) and (-), respectively. All feedback laws based on muscle

states acted upon the same muscle, except for a negative force feedback from the soleus to the

tibialis anterior.

Simulation and optimization framework

OpenSim was used to form the model's equations of motion and integrate them [56,57].

The state of the model was initialized by first setting the pelvis's horizontal position to 0 m. Then,

we set initial angular positions and velocities, and the speed of the pelvis according to free

parameters in the optimization. Although optimizing velocities may add extra energy into the

system, we do not think this affected our results as we found step-to-step variability to be small,

and we only analyzed kinematics and kinetics after the first two steps taken. Finally, the vertical

position of the pelvis was set such that the GRF was equal to half of the model's body weight.

Integration continued until the desired simulation time was reached or until the model fell, which

we defined as the model's center of mass (COM) height falling below 0.8 times the height of the

COM at the start of the simulation.

To evaluate a set of design parameters, we defined an objective function, J, which we

sought to minimize. The objective, J, quantified high-level tasks of walking:

$$J = J_{cot} + w_{spd}J_{spd} + w_{inj}J_{inj} + w_{head}J_{head} \qquad (6)$$

The goal was to minimize the gross cost of transport (J_{cot}) while maintaining a prescribed step-to-

step speed within a range, avoiding falling (J_{spd}), avoiding ligament injury (J_{inj}), and stabilizing

the head (J_{head}). To balance competing objectives, the weights were manually adjusted to the

following: $w_{spd} = 10{,}000$, $w_{inj} = 0.1$, and $w_{head} = 0.25$. These weights emphasized finding a gait that

did not fall over one that minimized cost of transport, injury, and accelerations at the head. For

optimized solutions, the contribution from the J_{head} term was larger than those from the J_{spd} and

J_{inj} terms, and thus our results were most sensitive to our choice of w_{head}. Because of this, our

choice for w_{head} represents a trade-off between head stability, especially during early stance, and

cost of transport.

J_{cot} penalized simulations in which the model walked in a less metabolically efficient

manner [51] and was calculated by summing the basal and per-muscle metabolic rates [71,72] to

estimate the gross metabolic rate ($\dot{E}$) over the duration of the simulation (t_{end}), and normalizing by

the mass of the model (m) and distance travelled (d):

$$J_{cot} = \frac{\int_0^{t_{end}} \dot{E}dt}{md} \tag{7}$$

554   J_{spd} penalized simulations in which the model fell at a time (t_{fall}) before the desired

555   simulation end time (t_{des}) and had step speeds outside of a prescribed range. J_{spd} was calculated

556   using the following equations:

$$J_{spd} = 1 - \sum_s \frac{t_s v_{s,pen}}{v_{min}} \tag{8}$$

$$v_{s,pen} = \begin{cases} 0, & t_{end} = t_{fall} \\ \max\left[0, 1 - \phi(v_s, v_{min}, v_{max})\right], & t_{end} = t_{des} \end{cases} \tag{9}$$

$$\phi(\varphi, \varphi_{min}, \varphi_{max}) = \begin{cases} 0, & \varphi_{min} < \varphi < \varphi_{max} \\ \varphi_{min} - \varphi, & \varphi < \varphi_{min} \\ \varphi - \varphi_{max}, & \varphi > \varphi_{max} \end{cases} \tag{10}$$

where s denotes a step, t_s is the duration of a step, $v_{s,pen}$ is a penalty for a given step (ranging

between 0 and 1, where 1 is no penalty), v_s is the step speed, v_{min} and v_{max} define the minimum and

maximum speeds allowed without a penalty, and Φ applies a linear penalty if v_s is not between

v_{min} and v_{max}. If all steps were within the speed range, J_{spd} was 0, and if the model immediately

fell, J_{spd} was 1. For any given step in which the speed was outside of the range, the step was linearly

penalized more heavily (i.e., $v_{s,pen}$ was closer to 0) the further the speed was out of the range, and

its final contribution to J_{spd} was scaled by t_s. For simulations with a prescribed speed, the speed

range was ±0.05 m/s of the prescribed speed. For example, to prescribe a speed of 1.25 m/s, v_{min} =

1.20 m/s and v_{max} = 1.30 m/s. For simulations without a prescribed speed, v_{min} was 0.75 m/s, and

v_{max} was arbitrarily large.

J_{inj} penalized simulations in which the model engaged the ligaments and was calculated by

integrating the sum of joint torques (T_j) squared generated by the model's ligaments:

572
$$J_{\text{inj}} = \int_0^{t_{\text{end}}} \sum_j T_j^2 dt \tag{11}$$

573   Since head stability is commonly regarded as an important task during gait [73], J_{head}

574   penalized simulations in which the model had excessively large accelerations and was calculated

575   by the following equation and Eq 10:

576
$$J_{\text{head}} = \int_0^{t_{\text{end}}} \left[\phi(a_x, a_{x,\min}, a_{x,\max})^2 + \phi(a_y, a_{y,\min}, a_{y,\max})^2\right] dt \tag{12}$$

where a_x and a_y are the horizontal and vertical accelerations at a point located near the center of

the head, and $a_{x,\min}$, $a_{x,\max}$, $a_{y,\min}$, and $a_{y,\max}$ determine the bounds between which no penalty is

applied if a_x and a_y are within the bounds. We set our acceleration bounds as $a_{x,\min} = -0.25g$, $a_{x,\max}$

$= 0.25g$, $a_{y,\min} = -0.50g$, and $a_{y,\max} = 0.50g$, which we approximated from experimental data of

human walking [52].

To solve our optimization problem, we used an evolutionary strategy called Covariance

Matrix Adaptation Evolutionary Strategy (CMA-ES) with a rank-μ update [74]. In total, there were

90 design variables: 74 for the gains, offsets, and transition parameters in the gait controller, and

16 for the parameters that set the model's initial state. We set the CMA-ES parameters such that

the population size of each generation (λ) was 16 and the update step for each generation used only

the 8 best solutions (μ). Because these types of problems tend to have noisy, non-convex search

spaces, and CMA-ES is a stochastic algorithm, we ran a set of multiple parallel optimizations with

the same initial guess, and used the best solution from the set to seed the next set of optimizations

as has been done previously [23,53]. Refer to S2 Table for the initial standard deviations used for

each parameter. We chose to stop when the best solution from a set had an objective function value

that was no better than 5% of the one from the previous set, as we observed optimized solutions

that were within 5% of each other were qualitatively similar.

Generating simulations of gait

To generate simulations of gait with a prescribed speed, we used 10 second simulations

(t_{des} = 10 s) and a set of 20 parallel optimizations, each with a maximum of 3000 generations for

CMA-ES. There were 7 simulations that were generated at prescribed speeds between 0.50 m/s

and 2.00 m/s at an interval of 0.25 m/s. We first trained the model at the middle speed, 1.25 m/s,

and then used those results to seed the neighboring speeds, continuing until we reached the lowest

and highest speeds. For example, the solution for our 1.25 m/s case was used to seed the 1.00 m/s

case, which was then used to seed the 0.75 m/s case.

To generate simulations of gait with a self-selected speed, we used longer, 30 second

simulations (t_{des} = 30 s), as we found that solutions with a shorter desired simulation time would

systematically fall forward at the end without a tight bound on step speed. We used a set of 10

parallel optimizations, each with a maximum of 1500 generations for CMA-ES. We generated 3

simulations of unimpaired gait at a self-selected speed, seeding from solutions of the slowest (0.50

m/s), middle (1.25 m/s), and fastest (2.00 m/s) cases from above. We also generated 18 simulations

of impaired gait, starting with the unimpaired solution and seeding the next more severe case. For

example, the solution from the unimpaired case was used to seed the mild SOL weakness case,

which was used to seed the moderate case, which was finally used to seed the severe case.

Acknowledgments

We would like to thank Apoorva Rajagopal and Jennifer Yong for their insightful feedback on this manuscript, and Christopher Dembia, Shrinidhi Lakshmikanth, Ajay Seth, and Thomas Uchida for helpful technical discussions. We thank the Stanford Research Computing Center for providing computational resources and support that contributed to these research results.

References

1. Neptune RR, Kautz SA, Zajac FE. Contributions of the individual ankle plantar flexors to support, forward progression and swing initiation during walking. J Biomech. 2001;34: 1387–1398. doi:10.1016/S0021-9290(01)00105-1

2. Liu MQ, Anderson FC, Schwartz MH, Delp SL. Muscle contributions to support and progression over a range of walking speeds. J Biomech. 2008;41: 3243–3252. doi:10.1016/j.jbiomech.2008.07.031

3. Handsfield GG, Meyer CH, Abel MF, Blemker SS. Heterogeneity of muscle sizes in the lower limbs of children with cerebral palsy. Muscle Nerve. 2016;53: 933–945. doi:10.1002/mus.24972

4. Noble JJ, Fry NR, Lewis AP, Keevil SF, Gough M, Shortland AP. Lower limb muscle volumes in bilateral spastic cerebral palsy. Brain Dev. 2014;36: 294–300. doi:10.1016/J.BRAINDEV.2013.05.008

5. Barber L, Hastings-Ison T, Baker R, Barrett R, Lichtwark G. Medial gastrocnemius muscle volume and fascicle length in children aged 2 to 5 years with cerebral palsy. Dev Med Child Neurol. 2011;53: 543–548. doi:10.1111/j.1469-8749.2011.03913.x

6. Kim CM, Eng JJ. The Relationship of Lower-Extremity Muscle Torque to Locomotor

Performance in People With Stroke. Phys Ther. 2003;83: 49–57. doi:10.1093/ptj/83.1.49

7.   Hoogerwaard EM, Bakker E, Ippel PF, Oosterwijk JC, Majoor-Krakauer DF, Leschot NJ,

et al. Signs and symptoms of Duchenne muscular dystrophy and Becker muscular

dystrophy among carriers in The Netherlands: a cohort study. Lancet. 1999;353: 2116—

2119. doi:10.1016/s0140-6736(98)10028-4

8.   Bushby K, Finkel R, Birnkrant DJ, Case LE, Clemens PR, Cripe L, et al. Diagnosis and

management of Duchenne muscular dystrophy, part 1: diagnosis, and pharmacological and

psychosocial management. Lancet Neurol. 2010;9: 77–93. doi:10.1016/S1474-

4422(09)70271-6

9.   Sabir M, Lyttle D. Pathogenesis of pes cavus in Charcot-Marie-Tooth disease. Clin

Orthop Relat Res. 1983; 173—178.

10.   Barber L, Barrett R, Lichtwark G. Passive muscle mechanical properties of the medial

gastrocnemius in young adults with spastic cerebral palsy. J Biomech. 2011;44: 2496–

2500. doi:10.1016/j.jbiomech.2011.06.008

11.   Mathewson MA, Ward SR, Chambers HG, Lieber RL. High resolution muscle

measurements provide insights into equinus contractures in patients with cerebral palsy. J

Orthop Res. 2015;33: 33–39. doi:10.1002/jor.22728

12.   Lin P-Y, Yang Y-R, Cheng S-J, Wang R-Y. The Relation Between Ankle Impairments

and Gait Velocity and Symmetry in People With Stroke. Arch Phys Med Rehabil.

2006;87: 562–568. doi:10.1016/J.APMR.2005.12.042

13.   Gao F, Grant TH, Roth EJ, Zhang L-Q. Changes in Passive Mechanical Properties of the

Gastrocnemius Muscle at the Muscle Fascicle and Joint Levels in Stroke Survivors. Arch

Phys Med Rehabil. 2009;90: 819–826. doi:10.1016/J.APMR.2008.11.004

14.   Mercuri E, Bushby K, Ricci E, Birchall D, Pane M, Kinali M, et al. Muscle MRI findings

in patients with limb girdle muscular dystrophy with calpain 3 deficiency (LGMD2A) and

early contractures. Neuromuscul Disord. 2005;15: 164–171.

doi:10.1016/J.NMD.2004.10.008

15.   Bushby K, Finkel R, Birnkrant DJ, Case LE, Clemens PR, Cripe L, et al. Diagnosis and

management of Duchenne muscular dystrophy, part 2: implementation of

multidisciplinary care. Lancet Neurol. 2010;9: 177–189.

doi:https://doi.org/10.1016/S1474-4422(09)70272-8

16.   Sutherland DH, Cooper L, Daniel D. The role of the ankle plantar flexors in normal

walking. JBJS. 1980;62.

17.   Steele KM, Seth A, Hicks JL, Schwartz MS, Delp SL. Muscle contributions to support

and progression during single-limb stance in crouch gait. J Biomech. 2010;43: 2099–2105.

doi:10.1016/J.JBIOMECH.2010.04.003

18.   Steele KM, van der Krogt MM, Schwartz MH, Delp SL. How much muscle strength is

required to walk in a crouch gait? J Biomech. 2012;45: 2564–2569.

doi:10.1016/J.JBIOMECH.2012.07.028

19.   van der Krogt MM, Bar-On L, Kindt T, Desloovere K, Harlaar J. Neuro-musculoskeletal

simulation of instrumented contracture and spasticity assessment in children with cerebral

palsy. J Neuroeng Rehabil. 2016;13: 64. doi:10.1186/s12984-016-0170-5

20.   Geyer H, Herr H. A Muscle-Reflex Model That Encodes Principles of Legged Mechanics

Produces Human Walking Dynamics and Muscle Activities. IEEE Trans Neural Syst

Rehabil Eng. 2010;18: 263–273. doi:10.1109/TNSRE.2010.2047592

21.   Wang JM, Hamner SR, Delp SL, Koltun V. Optimizing Locomotion Controllers Using

682   Biologically-Based Actuators and Objectives. ACM Trans Graph. 2012;31: 25.

683   doi:10.1145/2185520.2185521

684   22.   Dorn TW, Wang JM, Hicks JL, Delp SL. Predictive Simulation Generates Human

685   Adaptations during Loaded and Inclined Walking. PLoS One. 2015;10: e0121407.

686   doi:10.1371/journal.pone.0121407

687   23.   Song S, Geyer H. A neural circuitry that emphasizes spinal feedback generates diverse

688   behaviours of human locomotion. J Physiol. 2015;593: 3493–3511. doi:10.1113/JP270228

689   24.   Song S, Geyer H. Predictive neuromechanical simulations indicate why walking

690   performance declines with ageing. J Physiol. 2018;596: 1199–1210.

691   doi:10.1113/JP275166

692   25.   Schwartz MH, Rozumalski A, Trost JP. The effect of walking speed on the gait of

693   typically developing children. J Biomech. 2008;41: 1639–1650.

694   doi:10.1016/j.jbiomech.2008.03.015

695   26.   Ralston HJ. Energy-speed relation and optimal speed during level walking. Int Zeitschrift

696   für Angew Physiol einschließlich Arbeitsphysiologie. 1958;17: 277–283.

697   doi:10.1007/BF00698754

698   27.   Martin PE, Rothstein DE, Larish DD. Effects of age and physical activity status on the

699   speed-aerobic demand relationship of walking. J Appl Physiol. 1992;73: 200–206.

700   doi:10.1152/jappl.1992.73.1.200

28.   Browning RC, Baker EA, Herron JA, Kram R. Effects of obesity and sex on the energetic

cost and preferred speed of walking. J Appl Physiol. 2006;100: 390–398.

doi:10.1152/japplphysiol.00767.2005

29.   Wren TAL, Do KP, Rethlefsen SA, Healy B. Cross-correlation as a method for comparing

dynamic electromyography signals during gait. J Biomech. 2006;39: 2714–2718.

doi:10.1016/J.JBIOMECH.2005.09.006

30.   Ziv I, Blackburn N, Rang M, Koreska J. Muscle growth in normal and spastic mice. Dev

Med Child Neurol. 1984;26: 94–99. doi:10.1111/j.1469-8749.1984.tb04412.x

31.   Delp SL, Statler K, Carroll NC. Preserving plantar flexion strength after surgical treatment

for contracture of the triceps surae: A computer simulation study. J Orthop Res. 1995;13:

96–104. doi:10.1002/jor.1100130115

32.   Abel MF, Damiano DL. Strategies for increasing walking speed in diplegic cerebral palsy.

J Pediatr Orthop. 1996;16: 753—758. doi:10.1097/00004694-199611000-00010

33.   Turnbull GI, Charteris J, Wall JC. A comparison of the range of walking speeds between

normal and hemiplegic subjects. Scand J Rehabil Med. 1995;27: 175—182.

34.   Khodadadeh S, McClelland MR, Patrick JH. Variations of Gait Parameters in Duchenne

Muscular Dystrophy. Proc Inst Mech Eng Part H J Eng Med. 1990;204: 241–243.

doi:10.1243/PIME_PROC_1990_204_262_02

35.   Newman CJ, Walsh M, O'Sullivan R, Jenkinson A, Bennett D, Lynch B, et al. The

characteristics of gait in Charcot-Marie-Tooth disease types I and II. Gait Posture.

2007;26: 120–127. doi:10.1016/J.GAITPOST.2006.08.006

36.   Don R, Serrao M, Vinci P, Ranavolo A, Cacchio A, Ioppolo F, et al. Foot drop and plantar

flexion failure determine different gait strategies in Charcot-Marie-Tooth patients. Clin

724  Biomech. 2007;22: 905–916. doi:10.1016/J.CLINBIOMECH.2007.06.002

725 37. Dreher T, Buccoliero T, Wolf SI, Heitzmann D, Gantz S, Braatz F, et al. Long-term

726  results after gastrocnemius-soleus intramuscular aponeurotic recession as a part of

727  multilevel surgery in spastic diplegic cerebral palsy. J Bone Joint Surg Am. 2012;94:

728  627—637. doi:10.2106/jbjs.k.00096

729 38. Dietz FR, Albright JC, Dolan L. Medium-term follow-up of Achilles tendon lengthening

730  in the treatment of ankle equinus in cerebral palsy. Iowa Orthop J. 2006;26: 27–32.

731 39. Segal LS, Thomas SE, Mazur JM, Mauterer M. Calcaneal gait in spastic diplegia after

732  heel cord lengthening: a study with gait analysis. J Pediatr Orthop. 1989;9: 697—701.

733 40. Garbarino JL, Clancy M. A geometric method of calculating tendo Achillis lengthening. J

734  Pediatr Orthop. 1985;5: 573—576.

735 41. Gage JR. Surgical treatment of knee dysfunction in cerebral palsy. Clin Orthop Relat Res.

736  1990; 45—54.

737 42. Fulford GE. Surgical management of ankle and foot deformities in cerebral palsy. Clin

738  Orthop Relat Res. 1990; 55—61.

739 43. Wren TAL, Rethlefsen S, Kay RM. Prevalence of specific gait abnormalities in children

740  with cerebral palsy: influence of cerebral palsy subtype, age, and previous surgery. J

741  Pediatr Orthop. 2005;25: 79—83. doi:10.1097/00004694-200501000-00018

742 44. Wren TAL, Do KP, Kay RM. Gastrocnemius and soleus lengths in cerebral palsy equinus

743  gait—differences between children with and without static contracture and effects of

744  gastrocnemius recession. J Biomech. 2004;37: 1321–1327.

745  doi:10.1016/J.JBIOMECH.2003.12.035

45.   Perry J, Burnfield JM. Gait Analysis: Normal and pathological function. 2010. SLACK

Inc. : 85–534.

46.   Gage JR, Novacheck TF. An update on the treatment of gait problems in cerebral palsy. J

Pediatr Orthop B. 2001;10: 265—274. doi:10.1097/00009957-200110000-00001

47.   Kim CM, Eng JJ. Magnitude and pattern of 3D kinematic and kinetic gait profiles in

persons with stroke: relationship to walking speed. Gait Posture. 2004;20: 140–146.

doi:10.1016/J.GAITPOST.2003.07.002

48.   Newham DJ, Hsiao SF. Knee muscle isometric strength, voluntary activation and

antagonist co-contraction in the first six months after stroke. Disabil Rehabil. 2001;23:

379—386. doi:10.1080/0963828001006656

49.   Herbert RD, Crosbie J. Rest length and compliance of non-immobilised and immobilised

rabbit soleus muscle and tendon. Eur J Appl Physiol Occup Physiol. 1997;76: 472–479.

doi:10.1007/s004210050277

50.   Smith LR, Lee KS, Ward SR, Chambers HG, Lieber RL. Hamstring contractures in

children with spastic cerebral palsy result from a stiffer extracellular matrix and increased

in vivo sarcomere length. J Physiol. 2011;589: 2625–2639.

doi:10.1113/jphysiol.2010.203364

51.   Selinger JC, O'Connor SM, Wong JD, Donelan JM. Humans Can Continuously Optimize

Energetic Cost during Walking. Curr Biol. 2015;25: 2452–2456.

doi:10.1016/J.CUB.2015.08.016

52.   Menz HB, Lord SR, Fitzpatrick RC. Acceleration patterns of the head and pelvis when

walking on level and irregular surfaces. Gait Posture. 2003;18: 35–46. doi:10.1016/S0966-

768 6362(02)00159-5

769 53. Ong CF, Hicks JL, Delp SL. Simulation-Based Design for Wearable Robotic Systems: An

770   Optimization Framework for Enhancing a Standing Long Jump. IEEE Trans Biomed Eng.

771   2016;63: 894–903. doi:10.1109/TBME.2015.2463077

772 54. Heess N, TB D, Sriram S, Lemmon J, Merel J, Wayne G, et al. Emergence of Locomotion

773   Behaviours in Rich Environments. 2017; Available: arxiv:1707.02286

774 55. Kidziński Ł, Mohanty SP, Ong CF, Hicks JL, Carroll SF, Levine S, et al. Learning to Run

775   Challenge: Synthesizing Physiologically Accurate Motion Using Deep Reinforcement

776   Learning. In: Escalera S, Weimer M, editors. The NIPS '17 Competition: Building

777   Intelligent Systems. Cham: Springer International Publishing; 2018. pp. 101–120.

778 56. Delp SL, Anderson FC, Arnold AS, Loan P, Habib A, John CT, et al. OpenSim: open-

779   source software to create and analyze dynamic simulations of movement. IEEE Trans

780   Biomed Eng. 2007;54: 1940—1950. doi:10.1109/tbme.2007.901024

781 57. Seth A, Hicks JL, Uchida TK, Habib A, Dembia CL, Dunne JJ, et al. OpenSim:

782   Simulating musculoskeletal dynamics and neuromuscular control to study human and

783   animal movement. PLOS Comput Biol. 2018;14: 1–20. doi:10.1371/journal.pcbi.1006223

784 58. Delp SL, Loan JP, Hoy MG, Zajac FE, Topp EL, Rosen JM. An interactive graphics-

785   based model of the lower extremity to study orthopaedic surgical procedures. IEEE Trans

786   Biomed Eng. 1990;37: 757–767. doi:10.1109/10.102791

787 59. Thelen DG. Adjustment of Muscle Mechanics Model Parameters to Simulate Dynamic

788   Contractions in Older Adults. J Biomech Eng. 2003;125: 70–77. doi:10.1115/1.1531112

789 60. Rajagopal A, Dembia CL, DeMers MS, Delp DD, Hicks JL, Delp SL. Full-Body

Musculoskeletal Model for Muscle-Driven Simulation of Human Gait. IEEE Trans

Biomed Eng. 2016;63: 2068–2079. doi:10.1109/TBME.2016.2586891

61.   Millard M, Uchida T, Seth A, Delp SL. Flexing Computational Muscle: Modeling and

Simulation of Musculotendon Dynamics. J Biomech Eng. 2013;135: 21005–21011.

doi:10.1115/1.4023390

62.   Arnold EM, Hamner SR, Seth A, Millard M, Delp SL. How muscle fiber lengths and

velocities affect muscle force generation as humans walk and run at different speeds. J

Exp Biol. 2013; doi:10.1242/jeb.075697

63.   Whittington B, Silder A, Heiderscheit B, Thelen DG. The contribution of passive-elastic

mechanisms to lower extremity joint kinetics during human walking. Gait Posture.

2008;27: 628–634. doi:10.1016/J.GAITPOST.2007.08.005

64.   Ward SR, Eng CM, Smallwood LH, Lieber RL. Are Current Measurements of Lower

Extremity Muscle Architecture Accurate? Clin Orthop Relat Res. 2009;467: 1074–1082.

doi:10.1007/s11999-008-0594-8

65.   Arnold EM, Ward SR, Lieber RL, Delp SL. A Model of the Lower Limb for Analysis of

Human Movement. Ann Biomed Eng. 2010;38: 269–279. doi:10.1007/s10439-009-9852-5

66.   Thelen DG, Chumanov ES, Best TM, Swanson SC, Heiderscheit BC. Simulation of biceps

femoris musculotendon mechanics during the swing phase of sprinting. Med Sci Sports

Exerc. 2005;37: 1931—1938. doi:10.1249/01.mss.0000176674.42929.de

67.   Hunt KH, Crossley FRE. Coefficient of Restitution Interpreted as Damping in

Vibroimpact. J Appl Mech. 1975;42: 440–445. doi:10.1115/1.3423596

68.   Sherman MA, Seth A, Delp SL. Simbody: multibody dynamics for biomedical research.

Procedia IUTAM. 2011;2: 241–261. doi:10.1016/J.PIUTAM.2011.04.023

69.   Duysens J, Trippel M, Horstmann GA, Dietz V. Gating and reversal of reflexes in ankle

814   muscles during human walking. Exp Brain Res. 1990;82: 351–358.

815   doi:10.1007/BF00231254

816   70.   Frijns CJM, Laman DM, van Duijn MAJ, van Duijn H. Normal values of patellar and

817   ankle tendon reflex latencies. Clin Neurol Neurosurg. 1997;99: 31–36.

818   doi:10.1016/S0303-8467(96)00593-8

819   71.   Umberger BR, Gerritsen KGM, Martin PE. A model of human muscle energy

820   expenditure. Comput Methods Biomech Biomed Engin. 2003;6: 99—111.

821   doi:10.1080/1025584031000091678

822   72.   Uchida TK, Hicks JL, Dembia CL, Delp SL. Stretching Your Energetic Budget: How

823   Tendon Compliance Affects the Metabolic Cost of Running. PLoS One. 2016;11: 1–19.

824   doi:10.1371/journal.pone.0150378

825   73.   Bril B, Ledebt A. Head Coordination as a Means to Assist Sensory Integration in Learning

826   to Walk. Neurosci Biobehav Rev. 1998;22: 555–563. doi:10.1016/S0149-7634(97)00044-

827   4

828   74.   Hansen N, Kern S. Evaluating the CMA Evolution Strategy on Multimodal Test

829   Functions. In: Yao X, Burke EK, Lozano JA, Smith J, Merelo-Guervós JJ, Bullinaria JA,

830   et al., editors. Parallel Problem Solving from Nature - PPSN VIII. Berlin, Heidelberg:

831   Springer Berlin Heidelberg; 2004. pp. 282–291.

Supporting information

S1A Fig. Simulated walking at 0.55 m/s vs experimental data at 0.50 ± 0.20 m/s. To facilitate comparison, normalized speed from experimental data was converted using the model's leg length (0.856 m).

S1B Fig. Simulated walking at 0.79 m/s vs experimental data at 0.84 ± 0.16 m/s. To facilitate comparison, normalized speed from experimental data was converted using the model's leg length (0.856 m).

S1C Fig. Simulated walking at 1.04 m/s vs experimental data at 0.84 ± 0.16 m/s. To facilitate comparison, normalized speed from experimental data was converted using the model's leg length (0.856 m).

S1D Fig. Simulated walking at 1.22 m/s vs experimental data at 1.24 ± 0.15 m/s. To facilitate comparison, normalized speed from experimental data was converted using the model's leg length (0.856 m).

S1E Fig. Simulated walking at 1.46 m/s vs experimental data at 1.62 ± 0.18 m/s. To facilitate comparison, normalized speed from experimental data was converted using the model's leg length (0.856 m).

S1F Fig. Simulated walking at 1.71 m/s vs experimental data at 1.62 ± 0.18 m/s. To facilitate comparison, normalized speed from experimental data was converted using the model's leg length (0.856 m).

S1G Fig. Simulated walking at 1.96 m/s vs experimental data at 2.01 ± 0.27 m/s. To facilitate comparison, normalized speed from experimental data was converted using the model's leg length (0.856 m).

S2 Fig. Key kinematic and kinetic parameters for all deficits. This plot accompanies Fig 5 and

shows key parameters for walking with simulated weakness (blue) and contracture (orange) of the

SOL, GAS, or both (PF) along with simulated healthy gait (dotted line). Increasing color intensity

indicates increasing deficit severity.

S3A Fig. Simulated walking with mild, moderate, and severe soleus (SOL) weakness.

S3B Fig. Simulated walking with mild, moderate, and severe gastrocnemius (GAS) weakness.

S3C Fig. Simulated walking with mild, moderate, and severe soleus and gastrocnemius (PF)

weakness.

S3D Fig. Simulated walking with mild, moderate, and severe soleus (SOL) contracture.

S3E Fig. Simulated walking with mild, moderate, and severe gastrocnemius (GAS)

contracture.

S3F Fig. Simulated walking with mild, moderate, and severe soleus and gastrocnemius (PF)

contracture.

S1 Table. Muscle parameters for the unimpaired musculoskeletal model.

S2 Table. Initial Covariance Matrix Adaptation Evolutionary Strategy (CMA-ES) standard

deviation for each free parameter.

S1 Video. Highlights.

S2 Video. Compilation of all simulations.

Figure 2

Figure 3

Figure 4

Figure 6

Figure 7

Figure 8

Figure 10

Figure 1

Figure 9

Figure 5