Malaria vaccine candidates displayed on novel virus-like particles are immunogenic

and induce transmission-blocking activity

Authors/Affiliations:

Jo-Anne Chan[1,2], David Wetzel[3,4], Linda Reiling[1], Kazutoyo Miura[5], Damien Drew[1], Paul R

Gilson[1], David A Anderson[1], Jack S Richards[1,6], Carole A Long[5], Manfred Suckow[3], Volker

Jenzelewski[3], Takafumi Tsuboi[7], Michelle J Boyle[8], Michael Piontek[3] and James G

Beeson[1,2,6]

[1]Burnet Institute, Melbourne, VIC, Australia

[2]Department of Immunology, Central Clinical School, Monash University, VIC, Australia

[3]ARTES Biotechnology GmbH, Langenfeld, Germany

[4]Technical University of Dortmund, Laboratory of Plant and Process Design, Dortmund,

Germany

[5]Laboratory of Malaria and Vector Research, National Institute of Allergy and Infectious

Disease, National Institutes of Health, Rockville, Maryland, USA

[6]Department of Medicine, University of Melbourne, VIC, Australia

[7]Proteo-Science Centre, Ehime University, Matsuyama, Ehime, Japan

[8]QIMR-Berghofer Medical Research Institute, Herston, QLD, Australia

Corresponding Author:

James Beeson

Burnet Institute

85 Commercial Road, Melbourne, VIC 3001, Australia

beeson@burnet.edu.au

Ph: 61-3-9282-2442

Fax: 61-3-9282-2100

ABSTRACT

The development of effective malaria vaccines remains a global health priority. Currently, the most advanced vaccine, known as RTS,S, has only shown modest efficacy in clinical trials. Thus, the development of more efficacious vaccines by improving the formulation of RTS,S for increased efficacy or to interrupt malaria transmission are urgently needed. The RTS,S vaccine is based on the presentation of a fragment of the sporozoite antigen on the surface of virus-like particles (VLPs) based on human hepatitis B virus (HBV). In this study, we have developed and evaluated a novel VLP platform based on duck HBV (known as Metavax) for malaria vaccine development. This platform can incorporate large and complex proteins into VLPs and is produced in a *Hansenula* cell line compatible with cGMP vaccine production. Here, we have established the expression of leading *P. falciparum* malaria vaccine candidates as VLPs. This includes Pfs230 and Pfs25, which are candidate transmission-blocking vaccine antigens. We demonstrated that the VLPs effectively induce antibodies to malaria vaccine candidates with minimal induction of antibodies to the duck-HBV scaffold antigen. Antibodies to Pfs230 also recognised native protein on the surface of gametocytes, and antibodies to both Pfs230 and Pfs25 demonstrated transmission-reducing activity in standard membrane feeding assays. These results establish the potential utility of this VLP platform for malaria vaccines, which may be suitable for the development of multi-component vaccines that achieve high vaccine efficacy and transmission-blocking immunity.

INTRODUCTION

Mortality caused by *Plasmodium falciparum* malaria is estimated at 216 million cases annually, with approximately 500,000 deaths occurring worldwide [1]. Despite on-going efforts, malaria control has stalled with little reduction of malaria cases observed in the past few years [1]. The spread of anti-malarial drug resistance together with insecticide resistance in parasite vectors has further escalated the need for an effective malaria

vaccine. Malaria vaccine strategies can be broadly classified into three approaches; pre-

erythrocytic vaccines that target sporozoites and/or infected hepatocytes, blood-stage

vaccines that generally target merozoites and antigens on the surface of infected red blood

cells, and transmission-blocking vaccines that target the sexual stages of malaria or

mosquito-stage antigens [2]. While vaccines targeting pre-erythrocytic stages and blood

stages aim to directly prevent infection and disease, there is a growing focus on vaccines

that can interrupt or reduce malaria transmission, highlighted by key global organisations

including the World Health Organisation (WHO), Bill and Melinda Gates Foundation and

PATH Malaria Vaccine Initiative [3]. The most advanced *P. falciparum* vaccine RTS,S

(Mosquirix^{TM}) is based on the pre-erythrocytic stage of the parasite life cycle and is the only

malaria vaccine to have completed phase III clinical trials [4] and is currently undergoing

implementation trials in several African countries [1]. However, vaccine efficacy was low in

young children [5] and antibodies induced by vaccination waned quickly in the year after

immunisation [6]. The WHO and their partners have set an objective of developing a malaria

vaccine with 75% efficacy [3]. To achieve this goal, future vaccine development may be

dependent on novel strategies that induce sufficiently high levels of functional antibodies[2].

Transmission-blocking vaccines will need to induce a potent antibody response within the

host to inhibit the downstream development of parasites in the mosquito vector after a blood

meal [7]. This will prevent or reduce the subsequent spread of malaria parasites throughout

an endemic population. However, the advancement of such vaccines remains hampered by

the lack of knowledge and tools required to study the sexual, transmissible stages of *P.*

falciparum. The progress of sexual-stage antigens as vaccine candidates is further limited, in

part, by the difficulty to express high yields of properly folded recombinant antigens, and the

requirement for vaccine approaches that generate high and sustained levels of antibodies

for effective transmission-blocking activity. Leading vaccine candidates that are expressed

during the *P. falciparum* sexual-stage include Pfs230 and Pfs25. Pfs230 is expressed on the

surface of gametocytes that reside within the human host, while Pfs25 is expressed on the

surface of female gametes, zygotes and ookinetes in the mosquito vector [8]. Numerous

studies have shown that individuals naturally exposed to malaria acquire antibodies that

target Pfs230 (reviewed in [9]). Therefore, immunity afforded by vaccines based on Pfs230,

and other major antigens expressed on the gametocyte surface, have the added benefit of

potential for antibody boosting from natural malaria exposure. Pfs25 represents the only

sexual-stage antigen that has completed human clinical trials [10-12], and Pfs230 is

currently undergoing clinical trials. Antibodies generated through vaccination with Pfs25

have been reported to inhibit the development of parasites within the mosquito midgut [11-

13], demonstrating the potential of vaccines to interrupt the transmission of malaria

throughout a human population.

The development of potential vaccines requires the induction of strong and sustained

immune responses in humans, which has been challenging to achieve. The classic

approach of a subunit vaccine includes the delivery of a recombinant antigen formulated with

the appropriate adjuvant (reviewed in [14]). Expression of antigens in virus-like particles

(VLPs) have been successfully developed for the hepatitis B virus and human papilloma

virus (reviewed in [15]). Key characteristic features of viruses such as their repetitive surface

geometry and activation of specific immunological receptors are maintained by VLPs.

However, they do not have the ability to undergo gene replication as they lack a viral

genome [16]. Therefore, they have been established as a safe and effective vaccine delivery

platform for use in humans (reviewed in [15]). Furthermore, the particle size of VLPs

(ranging from 20–200nm) is optimal for immunogenicity and uptake by dendritic cells [17,18].

VLP-based vaccines elicit strong humoral and cellular responses, supported by their

capacity to cross-link B cell receptors for successful B cell activation [19]. The use of VLPs

to display and present antigens to the immune system appeared to generate a stronger

cellular and humoral response compared to the soluble antigen alone [20,21]. The only VLP-

based transmission-blocking vaccine designed for use in human clinical trials involves a

chimeric, non-enveloped VLP expressing Pfs25 fused to the Alfalfa mosaic virus coat protein

[22]. However, although the vaccine induced Pfs25-specific antibodies in a clinical trial, there

was limited inhibition of parasite transmission to mosquitoes [22], suggesting the need for

improved vaccine formulations.

Here, we describe a novel platform that incorporates sexual-stage malaria vaccine

candidates, Pfs230 and Pfs25, into a VLP scaffold that is based on the duck hepatitis B virus

(DHBV) small surface protein (dS). In contrast to the human HBV platform, the DHBV

delivery platform allows the expression of large and complex proteins [16], which are typical

of malaria vaccine candidates. In this platform, resulting VLPs contain a lipid envelope that

favours the optimal presentation of transmembrane and GPI-anchored proteins. *P.*

falciparum sexual-stage antigens, Pfs25 and the domains Pfs230c [23] and Pfs230D1M [24]

derived from the full-length Pfs230 protein, were genetically fused to the dS and the resulting

fusion proteins were co-expressed with wild-type dS in methylotrophic yeast *Hansenula*

polymorpha [16]. The successful display of antigens on the chimeric VLPs (referred to as

Pfs25-dS/dS, Pfs230c-dS/dS and Pfs230D1M-dS/dS) was confirmed using Western blot

analyses and visualized by electron microscopy and super resolution microscopy (presented

in the accompanying manuscript by Wetzel D et al 2019, submitted) [25]. In this manuscript,

we report the immunogenicity of these sexual-stage chimeric VLPs through animal

immunisations and further characterized their functional antibody response through

mosquito feeding assays.

METHODS

Production and purification of chimeric VLPs

The production and purification of the three different kinds of chimeric VLPs (Pfs230c-dS/dS,

Pfs230D1M-dS/dS and Pfs25-dS/dS) was conducted as previously described [16,25]. Genes

encoding the three fusion proteins Pfs230c-dS, Pfs230D1M-dS and Pfs25-dS were designed

and codon-optimized for expression in *Hansenula polymorpha* (Genbank ref.: MH142260,

MH142261, MH142262). Pfs230c-dS consists of amino acids (aa) 443-1132 (630 aa

fragment) from full-length Pfs230 [23] and the shorter variant Pfs230D1M-dS consists of aa

542-736 (194 aa fragment) from full-length Pfs230 [24]. The fusion protein Pfs25-dS is

comprised of aa 23-193 (170 aa fragment) of the cysteine-rich Pfs25 protein fused to the dS.

Each of the fusion proteins encoding genes were co-expressed with a gene encoding the

wildtype dS (Genbank ref: MF510122) in a recombinant *Hansenula*-derived yeast cell line.

The purification of the chimeric VLPs was based on a downstream process approved for

human VLP vaccine production from yeast [26] including adaptations for the purification of

VLPs containing the *P. falciparum* antigens [25]. The relative incorporation of the Pfs230-

dS/dS or Pfs25-dS/dS fusion proteins into VLPs, compared to total protein, was estimated

by densitometry analysis of Coomassie stained gels of VLPs, as described in the

accompanying manuscript [25]. The relative incorporation rates were ~30% for Pfs230c-

dS/dS VLPs, ~24% for Pfs230D1M-dS/dS VLPs, and ~3% for Pfs25-dS/dS VLPs.

Expression of monomeric recombinant proteins

The expression of monomeric recombinant Pfs230D1M was performed in HEK293F cells as

previously described [27]. Briefly, a truncated form of Pfs230 containing the first 6-cys

domain of Pfs230 has previously been expressed as a monomeric recombinant protein in *P.*

pastoris termed Pfs230D1H [24]. To assess vaccine-induced antibodies to Pfs230, we

expressed a modified form of monomeric recombinant Pfs230D1H in the mammalian

HEK293 cell expression system, which we termed Pfs230D1M. Monomeric recombinant

Pfs25 was expressed in a wheatgerm cell-free expression system as previously described

[28].

Measuring antibodies to chimeric VLPs by ELISA

Chimeric VLPs or monomeric recombinant proteins were coated onto Maxisorp microtiter

plates (Nunc) at 1µg/ml in PBS and incubated overnight at 4°C. Plates were blocked with 1%

casein in PBS (Sigma-Aldrich) for 2h at 37°C before primary antibodies were added

(polyclonal mouse anti-Pfs230 or anti-Pfs25 antibodies; or rabbit antibodies against Pfs230

or Pfs25). Secondary HRP-conjugated antibodies (polyclonal goat anti-mouse IgG at 1/1000

or anti-rabbit IgG at 1/2500 from Millipore) were used to detect antibody binding. Colour

detection was developed using ABTS or TMB liquid substrate (Sigma-Aldrich), which was

subsequently stopped using 1% SDS (for ABTS) or 1M sulphuric acid (for TMB). PBS was

used as a negative control and plates were washed thrice using PBS with 0.05% Tween in

between antibody incubation steps. The level of antibody binding was measured as optical

density at 405nm (for ABTS) or 450nm (for TMB).

Measuring antibody affinity by ELISA

Antibody affinity was assessed using standard ELISA, with the addition of an antibody

dissociation step using increasing concentrations of ammonium thiocynate (in PBS). This

was incubated for 20 min at room temperature, following incubation with antibody samples

(R1917 and 1918 were tested at 1/800, R 1919 and R1920 were tested at 1/100). Colour

detection was developed using ABTS (measured at optical density at 405nm).

Measuring antibodies to the gametocyte surface by flow cytometry

Mature stage V gametocyte-infected erythrocytes were generated as previously described

[27,29] and treated with saponin to permeabilize the infected erythrocyte membrane.

Gametocytes were subsequently incubated with whole serum from rabbits immunized with

Pfs230c-dS/dS VLPs, followed by an AlexaFluor 488-conjugated donkey anti-rabbit IgG

(1/500) with ethidium bromide to distinguish between infected and uninfected erythrocytes

(1/1000), with washing between each step. Data was acquired by flow cytometry (FACS

Verse, BD Biosciences) and analysed using FlowJo software. Antibody levels are expressed

as the geometric mean fluorescence intensity (MFI; arbitrary units).

Immunofluorescence Microscopy

Thin blood smears of stage V 3D7 gametocyte-infected erythrocytes were fixed in 90% acetone and 10% methanol for 5 min at -20°C as previously described [30]. Briefly, slides were rehydrated in PBS and blocked with 6% BSA for 30 minutes before incubation with primary antibodies (rabbit antibodies were tested at 1/100) for 2h. Slides were subsequently incubated with the secondary AlexaFluor 488-conjugated IgG (1/1000; Thermo Scientific) for 1h. Slides were washed thrice in PBS between antibody incubation steps. Slides were mounted in medium containing DAPI to label the parasite nucleus. Images were collected using a Plan-Apochromat (100×/1.40) oil immersion phase-contrast lens (Carl Zeiss) on an AxioVert 200M microscope (Carl Zeiss) equipped with an AxioCam Mrm camera (Carl Zeiss). Images were processed using Photoshop CS6 (Adobe).

Rabbit immunisations

Pfs230c-dS/dS and Pfs230D1M-dS/dS VLPs were formulated with equal volumes of sterile Alhydrogel aluminium hydroxide vaccine adjuvant (Brenntag, Denmark) and incubated on the shaker for 5 min prior to immunisations. New Zealand White rabbits were immunised 2 weeks apart with 3 doses of chimeric VLPs. The total amount of protein and adjuvant used in each immunisation is presented in Supplementary Tables S1-S3 for Pfs230c-dS/dS, Pfs230D1M-dS/dS and Pfs25-dS/dS VLPs, respectively. Briefly, rabbits (R1864-R1871) were immunised with two vaccine doses of Pfs230c-dS/dS VLPs formulated with and without Alhydrogel. As standard doses, we used 20 μg or 100 μg to total VLP protein for immunizations; the respective amount of Pfs230c protein is reported in Supplementary Tables S1-S3. Rabbits were immunised with Pfs230D1M-dS/dS VLPs formulated with Freund's adjuvant (R1917, R1918) or Alhydrogel (R1919, R1920) at a single vaccine dose (of 100 μg to total VLP protein). Rabbits were immunised with Pfs25-dS/dS VLPs formulated with Freund's adjuvant (R1825, R1826) at a single vaccine dose (of 100 μg to total VLP protein). Animal immunisations and the formulation of Pfs25dS/dS VLPs with Freund's Adjuvant were conducted by the Antibody Facility at the Walter and Eliza Hall Institute. Rabbits were euthanised with a terminal dose of sodium pentobarbitone administered

intravenously. Animal immunisations were approved by the Animal Ethics Committee of the

Walter and Eliza Hall Institute, Australia (#2017.018).

Measuring transmission-blocking activity by standard membrane feeding assays

IgG purification from individual rabbit serum samples was performed using Protein G

columns (GE Healthcare) according to the manufacturer instructions and adjusted to a final

concentration of 20 mg/ml in PBS. The standardized methodology for performing the

standard membrane feeding assays (SMFA) was described previously [31]. Briefly, 16-18

days old gametocyte cultures of the *P. falciparum* NF54 line were mixed with purified test

IgG at 7.5 mg/ml, and the final mixture, which contained human complement, was

immediately fed to ~50 female *Anopheles stephensi* mosquitoes through a membrane-

feeding apparatus. Mosquitoes were kept for 8 days and dissected to enumerate the oocysts

in the midgut (n=40 per group for the negative control group, and n=20 for test groups). As

the negative controls, total IgG purified from pre-immune rabbit sera were utilised. Only

midguts from mosquitoes with any eggs in their ovaries at the time of dissection were

analyzed. The human serum and red blood cells used for the SMFA were purchased from

Interstate Blood Bank (Memphis, TN).

Statistical analysis

The best estimate of % inhibition in oocyst density (% transmission-reducing activity,

%TRA), the 95% confidence intervals (95%CI), and *p*-values from single or multiple feeds

were calculated using a zero-inflated negative binomial random effects model (ZINB model)

described previously [32]. All statistical tests were performed in R (version 3.5.1), and *p*-

values <0.05 were considered significant.

RESULTS

Sexual-stage vaccine candidates are displayed on chimeric VLPs

Chimeric VLPs expressing sexual-stage *P. falciparum* antigens, Pfs230 and Pfs25, were

produced and purified as described [25]. Antigens were incorporated into the VLP scaffold

through genetic fusion of Pfs230 and Pfs25 with the dS protein. For the expression of the

chimeric Pfs230 VLPs, we used 2 different constructs. Due to the large size and complexity

of Pfs230, we did not attempt expression of full length Pfs230 in VLPs; instead we

expressed two truncated forms based on published constructs that have demonstrated

induction of transmission-blocking antibodies when used as recombinant protein vaccines in

experimental animals. The first was a truncated construct of Pfs230, termed Pfs230c, which

includes the first two 6-cysteine domains of Pfs230 [23], was used to generate fusion

proteins for VLP formation. Pfs230c has been previously shown to elicit transmission-

blocking antibodies [23]. The expression and purification of Pfs230c-dS VLPs was

challenging [25]. Therefore, a shorter construct of Pfs230, termed Pfs230D1M (based on the

construct described by [24]) which includes only the first 6-cysteine domain of Pfs230, was

designed and fused to the dS antigen. Further, Pfs230D1M is a leading transmission-

blocking vaccine candidate currently in clinical trials. Here, we evaluated the display of these

sexual-stage antigens on the surface of native VLPs. We used specific antibodies generated

against Pfs230 and Pfs25 to detect the expression of these antigens on the chimeric VLPs

by ELISA (Fig 1). We found that a Pfs230-specific polyclonal antibody recognised the

surface of Pfs230c-dS/dS (Fig 1A) and Pfs230D1M-dS/dS VLPs (Fig 1B). Similarly, a Pfs25-

specific monoclonal antibody recognised the surface of Pfs25-dS VLPs (Fig 1C). These

results confirm the expression of sexual-stage antigens on the surface of chimeric VLPs.

There was no recognition of a plain dS VLP without a Pfs230 or Pfs25 fusion protein (Fig

1D).

VLPs expressing sexual-stage antigens induced antibodies in rabbits

To understand whether these chimeric VLPs displaying sexual-stage antigens were capable

of eliciting an immune response, the VLP constructs were used to immunise rabbits (n=8 for

Pfs230c-dS/dS, n=4 for Pfs230D1M-dS/dS, n=2 for Pfs25-dS/dS). Serum from immunised

rabbits had significant antibody recognition of monomeric recombinant Pfs230D1M (Fig 2A-

D) or Pfs25 (Fig 2E, 2F) in a concentration-dependent manner by ELISA. For immunisations

with Pfs230c-dS/dS VLP, we investigated two vaccine doses (20 or 100 μg of total VLP

protein) formulated with or without Alhydrogel, which is an adjuvant suitable for clinical use.

The higher dose appeared to induce a higher antibody response in rabbits, and formulation

with the Alhydrogel adjuvant also contributed to higher antibody levels (Fig 2A, B). Given the

findings with Pfs230c-dS/dS VLPs, immunisations with Pfs230D1M-dS/dS VLP were only

performed at a single dose (using the higher dose of 100μg of total VLP protein), formulated

with Alhydrogel or with Freund's adjuvant for comparison. Both formulations induced

substantial antibodies to monomeric recombinant Pfs230D1M (Fig 2C, D); antibodies were

higher for formulations with Freund's, which is a more potent adjuvant, but cannot be used

clinically. For Pfs25-dS/dS VLPs, immunizations were only performed with Freund's adjuvant

as a proof-of-principle, since Pfs25 has already been evaluated with different adjuvants

using different platforms in published studies and has advanced into clinical trials [11,13].

Rabbit antibodies induced by chimeric VLPs recognized native Pfs230 expressed on

gametocytes

We evaluated whether rabbit antibodies generated against Pfs230c-dS/dS VLPs were

capable of recognizing native antigens expressed on the surface of mature stage V

gametocytes by flow cytometry (Fig 3A). The majority of the rabbit sera had substantial

antibody reactivity to the surface of mature gametocytes, suggesting that vaccine-induced

antibodies recognize native Pfs230 (Fig 3A). This finding was confirmed using

immunofluorescence microscopy, which showed that rabbit antibodies generated against

Pfs230c-dS/dS VLPs labelled native Pfs230 expressed on the surface of fixed mature

gametocytes (Fig 3B).

Rabbit antibodies induced by chimeric VLPs have transmission-reducing activity

To address the functional significance of rabbit antibodies generated against sexual-stage

chimeric VLPs, we examined the ability of these antibodies to inhibit mosquito infection

through standard membrane feeding assays (SMFA) in the presence of human complement

[31]. Functional transmission-blocking activity is defined as the reduction in oocyst count

compared to a negative control group. When antibodies against Pfs230c-dS/dS VLPs were

tested at 7.5mg/ml, none of them showed significant inhibition (Table 1). In contrast, 2 out of

4 rabbits immunized with Pfs230D1M-dS/dS VLPs demonstrated significant inhibition in

SMFA (Table 2). Antibodies generated against Pfs25-dS/dS VLPs from one of the two

rabbits also successfully blocked the development of oocysts within the mosquito midgut

(Table 3; 97.4% inhibition). It was interesting that the second rabbit did not significantly

inhibit transmission despite substantial induction of antibodies detected by ELISA,

suggesting that antibody titre and specificity may be important for functional activity.

Interestingly, there was no clear relationship between IgG levels and activity in SMFA by

rabbit antibodies generated to Pfs230D1M-dS/dS. While VLPs formulated with Freund's

adjuvant generated higher IgG reactivity to Pfs230, IgG generated using VLPs formulated

with alum tended to have stronger inhibitory activity in SMFA (Table 2). To understand these

differences further, we estimated the avidity of IgG binding to Pfs230D1M for different rabbit

antibodies (Supplementary Figure S1). This indicated no major difference in IgG avidity that

would explain difference in activity in SMFA, suggesting differences in functional activity may

be explained by differences in epitope targeting of antibodies.

DISCUSSION

Our work describes the immunogenicity of transmission-blocking malaria vaccine candidates

presented on the surface of VLPs using a novel platform based on the duck hepatitis B virus.

Leading sexual-stage antigens, Pfs230 and Pfs25 were engineered into chimeric VLPs and

their surface expression characterised using specific antibodies. Data is presented in the

accompanying manuscript [25] on the production and purification of these chimeric VLPs.

These showed that chimeric VLPs expressed proteins of expected size corresponding to

Pfs25 and the respective Pfs230 domains, including the small surface protein dS. Imaging

by electron microscopy identified homogenous, particulate structures corresponding to the

chimeric VLPs and super resolution microscopy further confirmed the recognition of Pfs230-

derived domains and Pfs25 on the chimeric VLP surface, and their colocalisation with the dS

antigen, using specific polyclonal antibodies. In this study, we showed that the VLPs were

reactive by ELISA with antibodies to Pfs230 or Pfs25, further demonstrating the display of

malaria vaccine candidates on the VLP surface. In rabbit immunisation studies, we showed

that the VLPs were immunogenic and were capable of inducing substantial antibody

reactivity to monomeric recombinant Pfs230D1M or Pfs25 protein. Further, rabbit antibodies

generated against Pfs230c-dS/dS and Pfs230D1M-dS/dS VLPs recognised the surface of

native gametocytes by flow cytometry and this was also visualised by immunofluorescence

microscopy. Importantly, rabbit antibodies against Pfs230D1M-dS/dS and Pfs25-dS/dS

VLPs were capable of blocking mosquito infection measured through standard membrane

feeding assays.

A conformational-dependent monoclonal antibody, 4B7, was used to characterise the

display of antigen expressed by the Pfs25-dS/dS VLP. There were high levels of antibody

recognition measured by ELISA to the Pfs25-dS/dS VLP, which correlated with the coating

concentration of monomeric recombinant Pfs25. Similarly, using a Pfs230-specific polyclonal

antibody, the display of antigen expressed by the Pfs230c-dS/dS and Pfs230D1M-dS/dS

VLP was confirmed. This suggests that the Pfs25 and Pfs230 constructs had successfully

been incorporated as fusion proteins into the chimeric VLPs. The incorporation rate of

Pfs25-dS into VLPs was relatively low and optimization may be needed to increase

incorporation rates prior to further evaluation of Pfs25 VLPs using this platform.

Our immunogenicity studies showed that the chimeric VLPs could elicit an immune response

in rabbits. For immunisations with Pfs230c-dS/dS VLP, two dosing regimens were used in

the presence and absence of Alhydrogel. As the higher dose appeared to induce a better

antibody response in rabbits, and formulation with Alhydrogel was needed for high antibody

levels, immunisations with Pfs230D1M-dS/dS VLP were done at a single dose (using the

higher dose) and formulated with Alhydrogel. Pfs230D1M-dS/dS VLPs were also formulated

with Freund's adjuvant for comparison. VLPs formulated with Alhydrogel induced substantial

antibody responses, but responses were higher with Freund's adjuvant. This was expected

because Freund's is a more potent adjuvant than Alhydrogel. Our results suggest that

investigating other clinically-acceptable adjuvants that may induce higher antibody

responses than Alhydrogel would be valuable. Further, rabbits were immunised with Pfs25-

dS/dS VLP using only Freund's adjuvant as a proof-of-principle to evaluate immunogenicity

and transmission-blocking activity. Antibodies generated against the Pfs25-dS/dS VLP

recognised monomeric recombinant Pfs25 by ELISA. Together, these findings suggest that

immunisation with chimeric VLPs induced a good immune response in rabbits, further

supporting the potential for the use of this VLP platform to present malaria antigens for

vaccination. However, further detailed studies are needed to evaluate the immunogenicity

and functional activity of antibodies generated by antigens presented as VLPs compared to

monomeric recombinant proteins. In addition, we measured the ability of rabbit antibodies to

recognise the surface of native gametocytes by flow cytometry. We found that rabbit

antibodies generated against Pfs230c-dS/dS VLPs were capable of recognising the surface

of mature, stage V gametocytes. This was also confirmed by immunofluorescence

microscopy using fixed smears of mature gametocytes.

Functional antibody responses were evaluated using standard membrane feeding assays,

which are widely used to assess transmission-reducing activity [32]. Antibodies from rabbits

immunised with Pfs25-dS/dS VLPs and Pfs230D1M-dS/dS could strongly inhibit the

development of oocysts within the mosquito midgut, indicating the successful inhibition of

parasite transmission. However, inhibitory activity was variable between immunized rabbits.

While further studies with larger numbers of animals, and including mice or rats, will be

required to better understand variability in vaccine responses, the findings here are sufficient

to establish a proof-of-concept for generating transmission-blocking activity using VLPs. Our

findings support previous work that showed that rabbit antibodies against Pfs230D1M had

transmission-reducing activity by SMFA [24]. Recent work has also reported that only

constructs containing domain 1 of Pfs230 were capable of inducing transmission-reducing

activity by SMFA, compared to constructs lacking that particular domain [33].

Further, the Pfs230D1M construct is currently undergoing phase I clinical trials. Together,

these findings support the importance of Pfs230D1M in the induction of functional

transmission-reducing antibodies. Of note, transmission-reducing activity was not observed

for antibodies generated against Pfs230c-dS/dS VLPs. This could have been due to the sub-

optimal folding of Pfs230c, such that key functional epitopes were not displayed, or

important antibody-binding epitopes were masked. These factors potentially resulted in low

purity and reduced yield of chimeric VLP and thus led to the lack of functional antibodies

generated. Interestingly, there was no clear relationship between IgG reactivity levels

quantified by ELISA and transmission-blocking activity in SMFA. This was particularly

evident for Pfs230D1M-dS/dS; immunization with VLPs formulated with Freund's adjuvant

generated higher IgG reactivity than using alum as the adjuvant, but IgG generated using

alum tended to have greater activity in SMFA. There was no substantial difference in avidity

between IgG induced using different adjuvants, which suggests the differences in

transmission-blocking activity are most likely explained by differences in epitope-specificity

of IgG. Future studies with a larger number of animals and immunization regimens may help

further understand this.

In conclusion, we have demonstrated the successful display of sexual-stage antigens

Pfs230 and Pfs25 on the surface of chimeric VLPs. We have established a proof-of-concept

which showed that these VLPs generated significant immune responses that recognised

homologous recombinant protein and native sexual-stage antigens expressed on

gametocytes. Further, these antibodies had the ability to block the transmission of parasites

to mosquitoes through membrane feeding assays. Together, our results support the further

evaluation of chimeric VLPs as a novel delivery platform for leading malaria vaccine

candidates. Future studies to optimise antigen incorporation and presentation in VLPs and to

evaluate different adjuvants and dosing regimens for immunization will further inform the

potential utility of this strategy for malaria vaccine development.

Author contributions statement

Conceptualization, JC, DW, LR, PG, JR, DA, MS, MP, and JB; Methodology, JC, DW, LR,

DD, VJ, KM, TT and MP; Investigation, JC, DW, LR, DD and KM; Administration, VJ, JR,

DA, MP and JB; Manuscript preparation, JC, DW and JB (and critically reviewed by all

authors); Funding Acquisition, JB and MP. Supervision, JB, MP, CL, MB. All authors

approved the final manuscript.

Acknowledgements

The authors gratefully acknowledge Colleen Woods for ongoing advice and project

feedback, Heribert Helgers, Renske Klassen, Christine Langer, Ashley Lisboa-Pinto and

Thomas Rohr for technical and academic assistance.

Funding

Funding was provided by PATH Malaria Vaccine Initiative and the National Health and

Medical Research Council (NHMRC) of Australia (Senior Research Fellowship and Program

Grant to JB, Career Development Fellowship to MB). Burnet Institute is supported by funding

from the NHMRC Independent Research Institutes Infrastructure Support Scheme and a

Victorian State Government Operational Infrastructure grant. The SMFA work performed

here was supported was supported in part by the intramural program of the National Institute

of Allergy and Infectious Diseases/NIH. The funders had no role in study design, data

collection and analysis, decision to publish or preparation of the manuscript. ARTES

Biotechnology GmbH provided support in the form of salaries for authors (DW, MS, VJ and

MP) and generation of VLPs, but did not have any additional role in the study design, data

collection and analysis, decision to publish or preparation of the manuscript. The specific

roles of these authors are articulated in the 'Author Contributions' section.

Competing Interests

The authors DW, MS, VJ and MP are associated with ARTES Biotechnology GmbH which

owns the license for the VLP technology [34,35]. This does not alter our adherence to PLOS

ONE policies on sharing data and materials.

REFERENCES

457   1.   World Health Organization. World malaria report 2018. 2018: 1–210.

458   2.   Beeson JG, Kurtovic L, Dobaño C, Opi DH, Chan J-A, Feng G, et al. Challenges and
459   strategies for developing efficacious and long-lasting malaria vaccines. Sci Transl
460   Med. American Association for the Advancement of Science; 2019;11: eaau1458.
461   doi:10.1126/scitranslmed.aau1458

462   3.   Malaria Vaccine Funders Group. Malaria Vaccine Technology Roadmap 2013. In:
463   httpswww.malariavaccine.orgmalaria-and-vaccinesmalaria-vaccine- roadmap.

464   4.   RTS,S Clinical Trials Partnership. Efficacy and safety of RTS,S/AS01 malaria vaccine
465   with or without a booster dose in infants and children in Africa: final results of a phase
466   3, individually randomised, controlled trial. Lancet. 2015;386: 31–45.
467   doi:10.1016/S0140-6736(15)60721-8

468   5.   Agnandji ST, Lell B, Soulanoudjingar SS, Fernandes JF, Abossolo BP, Conzelmann
469   C, et al. First results of phase 3 trial of RTS,S/AS01 malaria vaccine in African
470   children. N Engl J Med. 2011;365: 1863–1875. doi:10.1056/NEJMoa1102287

471   6.   White MT, Verity R, Griffin JT, Asante KP, Owusu-Agyei S, Greenwood B, et al.
472   Immunogenicity of the RTS,S/AS01 malaria vaccine and implications for duration of
473   vaccine efficacy: secondary analysis of data from a phase 3 randomised controlled
474   trial. Lancet Infect Dis. 2015;15: 1450–1458. doi:10.1016/S1473-3099(15)00239-X

475   7.   Sauerwein RW, Bousema T. Transmission blocking malaria vaccines: Assays and
476   candidates in clinical development. Vaccine. 2015;33: 7476–7482.
477   doi:10.1016/j.vaccine.2015.08.073

478   8.   Schneider P, Reece SE, van Schaijk BCL, Bousema T, Lanke KHW, Meaden CSJ, et
479   al. Quantification of female and male Plasmodium falciparum gametocytes by reverse

transcriptase quantitative PCR. Mol Biochem Parasitol. 2015;199: 29–33.
doi:10.1016/j.molbiopara.2015.03.006

9. Stone WJR, Dantzler KW, Nilsson SK, Drakeley CJ, Marti M, Bousema T, et al.
Naturally acquired immunity to sexual stage P. falciparum parasites. Parasitology.
2016;143: 187–198. doi:10.1017/S0031182015001341

10. Malkin EM, Durbin AP, Diemert DJ, Sattabongkot J, Wu Y, Miura K, et al. Phase 1
vaccine trial of Pvs25H: a transmission blocking vaccine for Plasmodium vivax
malaria. Vaccine. 2005;23: 3131–3138. doi:10.1016/j.vaccine.2004.12.019

11. Sagara I, Healy SA, Assadou MH, Gabriel EE, Kone M, Sissoko K, et al. Safety and
immunogenicity of Pfs25H-EPA/Alhydrogel, a transmission-blocking vaccine against
Plasmodium falciparum: a randomised, double-blind, comparator-controlled, dose-
escalation study in healthy Malian adults. Lancet Infect Dis. 2018;18: 969–982.
doi:10.1016/S1473-3099(18)30344-X

12. Wu Y, Ellis RD, Shaffer D, Fontes E, Malkin EM, Mahanty S, et al. Phase 1 trial of
malaria transmission blocking vaccine candidates Pfs25 and Pvs25 formulated with
montanide ISA 51. PLoS ONE. 2008;3: e2636. doi:10.1371/journal.pone.0002636

13. Talaat KR, Ellis RD, Hurd J, Hentrich A, Gabriel E, Hynes NA, et al. Safety and
Immunogenicity of Pfs25-EPA/Alhydrogel®, a Transmission Blocking Vaccine against
Plasmodium falciparum: An Open Label Study in Malaria Naïve Adults. PLoS ONE.
2016;11: e0163144. doi:10.1371/journal.pone.0163144

14. Draper SJ, Angov E, Horii T, Miller LH, Srinivasan P, Theisen M, et al. Recent
advances in recombinant protein-based malaria vaccines. Vaccine. 2015;33: 7433–
7443. doi:10.1016/j.vaccine.2015.09.093

15.   Mohsen MO, Zha L, Cabral-Miranda G, Bachmann MF. Major findings and recent

504   advances in virus-like particle (VLP)-based vaccines. Semin Immunol. 2017.

505   doi:10.1016/j.smim.2017.08.014

506   16.   Wetzel D, Rolf T, Suckow M, Kranz A, Barbian A, Chan J-A, et al. Establishment of a

507   yeast-based VLP platform for antigen presentation. Microb Cell Fact. 2018;17: 17.

508   doi:10.1186/s12934-018-0868-0

509   17.   Grgacic EVL, Anderson DA. Virus-like particles: passport to immune recognition.

510   Methods. 2006;40: 60–65. doi:10.1016/j.ymeth.2006.07.018

511   18.   Bachmann MF, Jennings GT. Vaccine delivery: a matter of size, geometry, kinetics

512   and molecular patterns. Nat Rev Immunol. Nature Publishing Group; 2010;10: 787–

513   796. doi:10.1038/nri2868

514   19.   Kündig TM, Senti G, Schnetzler G, Wolf C, Prinz Vavricka BM, Fulurija A, et al. Der p

515   1 peptide on virus-like particles is safe and highly immunogenic in healthy adults. J

516   Allergy Clin Immunol. 2006;117: 1470–1476. doi:10.1016/j.jaci.2006.01.040

517   20.   Brune KD, Leneghan DB, Brian IJ, Ishizuka AS, Bachmann MF, Draper SJ, et al.

518   Plug-and-Display: decoration of Virus-Like Particles via isopeptide bonds for modular

519   immunization. Sci Rep. Nature Publishing Group; 2016;6: 19234.

520   doi:10.1038/srep19234

521   21.   Keller SA, Bauer M, Manolova V, Muntwiler S, Saudan P, Bachmann MF. Cutting

522   edge: limited specialization of dendritic cell subsets for MHC class II-associated

523   presentation of viral particles. J Immunol. 2010;184: 26–29.

524   doi:10.4049/jimmunol.0901540

525   22.   Chichester JA, Green BJ, Jones RM, Shoji Y, Miura K, Long CA, et al. Safety and

526   immunogenicity of a plant-produced Pfs25 virus-like particle as a transmission

blocking vaccine against malaria: A Phase 1 dose-escalation study in healthy adults.

Vaccine. 2018;36: 5865–5871. doi:10.1016/j.vaccine.2018.08.033

23.　Williamson KC, KEISTER DB, Muratova O, Kaslow DC. Recombinant Pfs230, a

Plasmodium falciparum gametocyte protein, induces antisera that reduce the

infectivity of Plasmodium falciparum to mosquitoes. Mol Biochem Parasitol. 1995;75:

33–42.

24.　Macdonald NJ, Nguyen V, Shimp R, Reiter K, Herrera R, Burkhardt M, et al.

Structural and Immunological Characterization of Recombinant 6-Cysteine Domains

of the Plasmodium falciparum Sexual Stage Protein Pfs230. J Biol Chem. 2016;291:

19913–19922. doi:10.1074/jbc.M116.732305

25.　Wetzel D, Chan J-A, Suckow M, Barbian A, Weniger M, Jenzelewski V, et al. Display

of malaria transmission-blocking antigens on chimeric duck hepatitis B virus-derived

virus-like particles produced in *Hansenula polymorpha*. bioRxiv doi:10.1101/595538

26.　Schaefer S, Piontek M, Ahn S, Papendieck A, Janowicz Z, Timmermans I, et al.

Recombinant hepatitis B vaccines—disease characterization and vaccine production.

In: G G, editor. Hansenula polymorpha—biology and applications. 2002. pp. 175–210.

27.　Chan J-A, Drew DR, Reiling L, Lisboa-Pinto A, Dinko B, Sutherland CJ, et al. Low

Levels of Human Antibodies to Gametocyte-Infected Erythrocytes Contrasts the

PfEMP1-Dominant Response to Asexual Stages in P. falciparum Malaria. Front

Immunol. 2018;9: 3126. doi:10.3389/fimmu.2018.03126

28.　Tsuboi T, Takeo S, Iriko H, Jin L, Tsuchimochi M, Matsuda S, et al. Wheat germ cell-

free system-based production of malaria proteins for discovery of novel vaccine

candidates. Infection and Immunity. American Society for Microbiology Journals;

2008;76: 1702–1708. doi:10.1128/IAI.01539-07

29. Dinko B, King E, Targett GAT, Sutherland CJ. Antibody responses to surface antigens

of Plasmodium falciparum gametocyte-infected erythrocytes and their relation to

gametocytaemia. Parasite Immunol. John Wiley & Sons, Ltd; 2016;38: 352–364.

doi:10.1111/pim.12323

30. Chan J-A, Howell KB, Reiling L, Ataíde R, Mackintosh CL, Fowkes FJI, et al. Targets

of antibodies against Plasmodium falciparum-infected erythrocytes in malaria

immunity. J Clin Invest. 2012. doi:10.1172/JCI62182

31. Miura K, Deng B, Tullo G, Diouf A, Moretz SE, Locke E, et al. Qualification of

standard membrane-feeding assay with Plasmodium falciparum malaria and potential

improvements for future assays. PLoS ONE. Public Library of Science; 2013;8:

e57909. doi:10.1371/journal.pone.0057909

32. Miura K, Swihart BJ, Deng B, Zhou L, Pham TP, Diouf A, et al. Transmission-blocking

activity is determined by transmission-reducing activity and number of control oocysts

in Plasmodium falciparum standard membrane-feeding assay. Vaccine. 2016;34:

4145–4151. doi:10.1016/j.vaccine.2016.06.066

33. Tachibana M, Miura K, Takashima E, Morita M, Nagaoka H, Zhou L, et al.

Identification of domains within Pfs230 that elicit transmission blocking antibody

responses. Vaccine. 2019;37: 1799–1806. doi:10.1016/j.vaccine.2019.02.021

34. Anderson D, Grgacic E. Viral vectors expressing fusion of viral large envelope protein

and protein of interest. 2004.

35. Grgacic E, Anderson D, Loke P, Anders R. Recombinant proteins and virus-like

particles comprising L and S polypeptides of avian hepadnaviridae and methods,

nucleic acid constructs, vectors and host cells for producing same. 2006.

Table 1 Assessment of transmission-blocking activity of rabbit antibodies against

Pfs230c-dS/dS VLPs by SMFA

Rabbit ID (+/- Adjuvant)	Protein content[1]	Antibody levels[2]	% inhibition[3]	95% confidence intervals	*p* value
R1864 (-Alhydrogel)	20µg VLP ~6µg Pfs230c-dS	0.29	-20.9	-162.2 – 43.5	0.62
R1865 (-Alhydrogel)	20µg VLP ~6µg Pfs230c-dS	0.30	6	-100.0 – 57.2	0.85
R1866 (+Alhydrogel)	20µg VLP ~6µg Pfs230c-dS	1.17	-30.8	-187.4 – 43.5	0.52
R1867 (+Alhydrogel)	20µg VLP ~6µg Pfs230c-dS	1.40	-34.8	-193.4 – 37.7	0.46
R1868 (-Alhydrogel)	100µg VLP ~30µg Pfs230c-dS	0.92	-13.7	-143.7 – 48.3	0.72
R1869 (-Alhydrogel)	100µg VLP ~30µg Pfs230c-dS	2.16	-28.9	-182.5 – 41.8	0.53
R1870 (+Alhydrogel)	100µg VLP ~30µg Pfs230c-dS	0.87	-8.1	-124.4 – 50.5	0.90
R1871 (+Alhydrogel)	100µg VLP ~30µg Pfs230c-dS	1.88	-16.6	-156.6 – 48.0	0.70

[1]Refers to total VLP protein content and estimated incorporation rate of Pfs230c-dS

[2]Mean antibody levels by ELISA at 1/100 serum dilution (OD 405nm)

[3]% inhibition is the best estimate from 1 feed

All rabbit antibodies were tested at an IgG concentration of 7.5 mg/ml.

Table 2 Assessment of transmission-blocking activity of rabbit antibodies against

Pfs230D1M-dS/dS VLPs by SMFA

Rabbit ID (Adjuvant)	Antibody levels[1]	% inhibition[2]	95% confidence intervals	*p* value
R1917 (Freund's)	3.84	43.7	4.1 – 68.0	**0.03**
R1918 (Freund's)	3.74	9.2	-51.3 – 47.5	0.74
R1919 (Alhydrogel)	1.72	40.7	-0.2 – 65.9	0.05
R1920 (Alhydrogel)	2.51	91.7	85.7 – 95.4	**0.001**

[1]Mean antibody levels by ELISA at 1/100 serum dilution (OD 405nm)

[2]% inhibition is the best estimate from 2 feeds

Protein content was 100µg to total VLP protein and 24µg Pfs230D1M-dS; all rabbit

antibodies were tested at an IgG concentration of 7.5 mg/ml; *p* values that achieved

statistical significance are presented in bold.

Table 3 Assessment of transmission-blocking activity of rabbit antibodies against

Pfs25-dS/dS VLPs by SMFA

Rabbit ID (Adjuvant)	Antibody levels[1]	% inhibition[2]	95% confidence intervals	*p* value
R1825 (Freund's)	3.42	97.4	95.4 – 98.8	**0.001**
R1826 (Freund's)	2.91	-1.4	-69.4 – 40.5	0.96

592   [1]Mean antibody levels by ELISA at 1/100 serum dilution (OD 405nm)

593   [2]% inhibition is the best estimate from 2 feeds

594   Protein content was 100µg to total VLP protein and 3µg Pfs25-dS; all rabbit antibodies were

595   tested at an IgG concentration of 7.5 mg/ml; *p* value that achieved statistical significance is

596   presented in bold.

FIGURES

Fig 1 Characterising the expression of sexual-stage antigens on chimeric VLPs

Total antibody binding to (**A**) Pfs230c-dS/dS, (**B**) Pfs230D1M-dS/dS, (**C**) Pfs25-dS/dS and

(**D**) plain dS VLPs were measured by ELISA. VLPs were coated at varying concentrations

(µg/ml) and probed with either a Pfs25 or Pfs230-specific polyclonal antibody (1µg/ml). In

(**D**), plain dS VLPs were probed with anti-Pfs230 antibody. The level of antibody binding is

expressed as optical density (OD) measured at 405nm; symbols represent the mean and

error bars represent the range between samples tested in duplicate.

Fig 2 Immunogenicity of sexual-stage chimeric VLPs in rabbits

The level of antibody binding to a titration of monomeric recombinant (**A**, **C**) Pfs230D1M and

(**E**) Pfs25 was measured in whole rabbit serum (1/100). Rabbits were immunised with

chimeric Pfs230c-dS/dS (**A**, **B**), Pfs230D1M-dS/dS (**C**, **D**) or Pfs25-dS/dS (**E**, **F**) VLPs.

Monomeric recombinant proteins were serially diluted from 4µg/ml.

The level of antibody binding to monomeric recombinant (**B**, **D**) Pfs230D1M and (**F**) Pfs25

(coated at 1µg/ml) was measured by titrating whole rabbit serum. Rabbits were immunised

with chimeric Pfs230c-dS/dS (**A**, **B**), Pfs230D1M-dS/dS (**C**, **D**) or Pfs25-dS/dS (**E**, **F**) VLPs.

Whole rabbit serum was serially diluted from 1/100.

For all graphs, antibody binding is expressed as optical density (OD) measured at 405nm;

symbols represent the mean and error bars represent the range between samples tested in

duplicate (n=8 for Pfs230c-dS/dS, n=4 for Pfs230D1M-dS/dS, n=2 for Pfs25-dS/dS). Rabbits

R1864-R1871 received Pfs230c-dS/dS VLPs formulated with and without Alhydrogel (**A**, **B**;

Alh). Rabbits R1917 and R1918 received Pfs230D1M-dS/dS VLPs formulated with Freund's

adjuvant (Fre), while R1919 and R1920 received Pfs230D1M-dS/dS VLPs formulated with

Alhydrogel (**C**, **D**). Rabbits R1825 and R1826 received Pfs25-dS/dS VLPs formulated with

Freund's adjuvant (**E**, **F**). Adjuvants and total VLP protein content used for immunisations

are also presented in Supplementary Tables S1-S3.

Fig 3 Characterising the antibody function of rabbits immunised with chimeric VLPs

A. Total antibody binding to the surface of native *P. falciparum* gametocytes (stage V gametocyte-infected erythrocytes were permeabilised with saponin) measured by flow cytometry. Whole serum from rabbits immunised with Pfs230c-dS/dS VLPs were used (n=8). Antibody levels are expressed as geometric mean fluorescence intensity (MFI); bars represent mean and range of samples tested in duplicate.

B. Immunofluorescence microscopy demonstrates the recognition of the native gametocyte surface by serum antibodies from rabbits immunised with Pfs230c-dS/dS VLPs (R1870; green). Cells were fixed with 90% acetone and 10% methanol, and DAPI was used to stain nuclear DNA (blue). Representative images taken of gametocytes labelled with R1870 are shown.

644 **SUPPLEMENTARY MATERIALS**

645 **Supplementary Table S1 Vaccine groups for Pfs230c-dS/dS VLP rabbit immunisations**

Rabbit ID	Total VLP protein (µg)	Total Pfs230c-dS protein (µg)	Alhydrogel (with or without)
R1864, R1865	20	6	-
R1866, R1867	20	6	+
R1868, R1869	100	30	-
R1870, R1871	100	30	+

646 Note: Estimated incorporation of Pfs230c-dS into VLPs was ~30%, determined by
647 Coomassie stained gels.

649 **Supplementary Table S2 Vaccine groups for Pfs230D1M-dS/dS VLP rabbit**
650 **immunisations**

Rabbit ID	Total VLP protein (µg)	Total Pfs230D1M-dS protein (µg)	Adjuvant
R1917, R1918	100	24	Freund's
R1919, R1920	100	24	Alhydrogel

651 Note: Estimated incorporation of Pfs230D1M-dS into VLPs was ~24%, determined by
652 Coomassie stained gels.

654 **Supplementary Table S3 Vaccine groups for Pfs25-dS/dS VLP rabbit immunisations**

Rabbit ID	Total VLP protein (µg)	Total Pfs25-dS protein (µg)	Adjuvant
R1825, R1826	100	3	Freund's

Note: Estimated incorporation of Pfs25-dS into VLPs was ~3%, determined by Coomassie
stained gels.

Supplementary Fig S1

Fig S1 Avidity of rabbit antibodies to bind monomeric recombinant Pfs230D1M

The avidity of antibody binding to monomeric recombinant Pfs230D1M was assessed in

serum from rabbits immunised with Pfs230D1M-dS/dS VLPs. The affinity of rabbit antibodies

to bind monomeric recombinant Pfs230D1M was similar, irrespective of adjuvant formulation

used for rabbit immunisations. Rabbits R1917 and R1918 received Pfs230D1M-dS/dS VLPs

formulated with Freund's adjuvant (Fre), while R1919 and R1920 received Pfs230D1M-

dS/dS VLPs formulated with Alhydrogel (Alh). Antibody affinity was assessed by the

dissociation of antibodies using increasing concentrations of ammonium thiocynate.

Antibody binding is defined as the OD of ammonium thiocynate-treated samples/OD of

untreated samples x100.