Vegetative Cell and Spore Proteomes of *Peptoclostridium difficile* show finite differences

and reveal potential biomarkers.

Wishwas R. Abhyankar[1,2*], Linli Zheng[1,2], Stanley Brul[1+], Chris G. de Koster[2+], Leo J. de

Koning[2]

[1]Department of Molecular Biology and Microbial Food Safety, University of Amsterdam,

Amsterdam, the Netherlands;

[2]Department of Mass Spectrometry of Bio-Macromolecules, University of Amsterdam,

Amsterdam, the Netherlands.

* Corresponding author E-mail: w.r.abhyankar@uva.nl

[+]These authors contributed equally to this work.

Abstract

Clostridium difficile-associated infection (CDI) is a health-care-associated infection mainly transmitted via highly resistant endospores from one person to the other. *In vivo*, the spores need to germinate in to cells prior to establishing an infection. Bile acids and glycine, both available in sufficient amounts inside the human host intestinal tract, serve as efficient germinants for the spores. It is therefore, for better understanding of *Peptoclostridium difficile* virulence, crucial to study both the cell and spore states with respect to their genetic, metabolic and proteomic composition. In the present study, mass spectrometric relative protein quantification, based on the ^{14}N/^{15}N peptide isotopic ratios, has led to quantification of over 700 proteins from combined spore and cell samples. The analysis has revealed that the proteome turnover between a vegetative cell and a spore for this organism is moderate. Additionally, specific cell and spore surface proteins, vegetative cell proteins CD1228, CD3301 and spore proteins CD2487, CD2434 and CD0684 are identified as potential biomarkers for *P. difficile* infection.

Introduction

Peptoclostridium (previously *Clostridium*) *difficile,* an anaerobic, gram-positive pathogen, is the causative agent of *Clostridium difficile* infection (CDI) characterized by pseudomembranous colitis and nosocomial diarrhoea. An over-extensive use of antibiotics is implicated for the spread of CDI and high rates of asymptomatic colonization by *P. difficile* make its diagnosis challenging[1]. This necessitates designing strategies and algorithms to optimize the diagnostic tools[2]. Pathogenesis of CDI is manifested via its Rho-glycosylating toxins TcdA and TcdB[3]. Moreover, in response to adverse conditions, *P. difficile* can form endospores - multi-layered, highly resistant cellular entities - which are the main transmissible forms in *P. difficile* infections.

The spores can germinate in the intestinal environment upon interaction with bile acid mediated

by the CspC protease[4], subsequently manifesting the infection.

To date, substantial research has been done on *P. difficile* vegetative cells and spores to

understand the survival mechanisms of this pathogen. However, research performed on spores

has gained more importance, owing to their crucial role in survival of *P. difficile*. The number of

spore-related genes identified in *Clostridia* is significantly smaller than that in *Bacilli*[5]. Less than

25% of the spore coat proteins of *B. subtilis* have homologues in *P. difficile* and unlike in *B.*

subtilis, in *P. difficile*, neither does the activation of σ^G rely on σ^E nor is it required for the

production and σ^K activation[6]. Furthermore, in *P. difficile* spores, cortex hydrolysis occurs before

the release of Ca^{2+}-dipicolinic acid complex[7], whereas in *B. subtilis* spores, the release of this

complex precedes cortex hydrolysis.

In the past decade, a few transcriptomic studies[8,9], quantitative proteomic studies[10-16] and a

lipoproteomic study[17] have been done on *P. difficile* vegetative cells. Lawley and colleagues

described a protocol to purify spores and performed an extensive proteomic characterization of *P.*

difficile 630 spores[18]. Shortly after, we used a gel-free proteomic method and a one-pot sample

processing method that focused on the spore surface layers of *B. cereus* and *P. difficile*[19].

Moreover, previous studies have described exosporium removal methods for *P. difficile* spores,

and examined the exosporium protein components[20] as well as spore surface glycoproteins[21]. Yet,

none of these studies focusses on a comparative analysis that functionally links the spore and

vegetative cell proteome. In the present study we have quantitatively characterized the *P. difficile*

vegetative cell proteome relative to that of spores. To this end spores are mixed with ^{15}N-

metabolically labelled vegetative cells and the mixture is processed with our recently developed

one-pot method for mass spectrometric analyses, where the $^{14}N/^{15}N$ isotopic protein ratios

represent the relative spore over vegetative cell protein abundances. We aim to deduce putative

spore- and vegetative cell-predominant protein biomarkers for *P. difficile.*

Results

Metabolic labelling of *P. difficile* vegetative cells using ^{15}N-yeastolate. As illustrated in **Fig. 1**

our culturing methods successfully yielded ^{15}N labelled vegetative cells and ^{14}N spores. For a

number of identified ^{15}N labelled vegetative cell peptides, the ^{15}N label content has been

calculated on the basis of their mass spectrometric isotope patterns using the NIST isotope

calculator[22]. This shows that the present metabolic labelling method achieves a ^{15}N label content

of ≥ of 99.5% which is amply sufficient for accurate protein quantification.

Identification and quantification of cell and spore proteins. A total of 1095 proteins has been

identified from *P. difficile* spores and vegetative cells of which 796 have been relatively and

reproducibly quantified between spores and vegetative cells (**Supplementary Table S1**). **Fig. 2**

represents a distribution of quantified proteins, where the abundance of the combined spore and

vegetative cell proteins indicated by the $\log_{10}$ values of their MASCOT scores are plotted against

the corresponding relative protein levels in the two morphotypes indicated by the $\log_2$ values of

the light/heavy ratios. Eighty seven proteins are considered to be predominant present in spores

with a light/heavy ratio > 20, while 81 proteins are considered to be predominant present in

vegetative cells with light/heavy ratios < 0.05. From the remaining 628 commonly shared

proteins 18% are enriched in spores with light/heavy ratio between 1 and 20 while 82% are

enriched in cells with a light/ heavy ratio between 1 and 0.05. In total, 167 proteins have been

classified as membrane proteins by the TMHMM analysis (**Supplementary Table S2**). The

cellular overview based on pathway analysis of the quantified proteins is represented in

Supplementary Fig S1 indicating the pathways to which these commonly shared proteins

belong. As seen

Spore-predominant proteins. These include proteins from the spore coat and exosporium

layers, classified under UniProt Keywords virion and capsid proteins, along with some rotamase

proteins and metalloproteases (**Table 1**). SspA is the most abundant protein in this category,

whereas an uncharacterized protein CD2657 (with 13 times higher levels in spores than in

vegetative cells) is the least abundant (**Fig 2**). The TMHMM analysis classified 26 proteins from

this category as membrane proteins (**Supplementary Table S2**). Most of these are

uncharacterized membrane proteins but some are known proteins such as SpoVD, SpoVAC,

SpoVFB, FtsH, and DacF.

Cell-predominant proteins. These include the cytoplasmic proteins such as aminotransferases,

arginine biosynthesis, elongation factors, cell shape and peptidoglycan synthesis proteins (**Table**

1). The surface layer protein SlpA is the most abundant and unique protein in this category,

whereas an uncharacterized protein CD0594 (with levels 27 times higher in vegetative cells than

in spores) is the least abundant (**Fig 2**). Eighteen membrane proteins are predominant in

vegetative cells, as predicted by TMHMM (**Supplementary Table S2**).

Proteins shared between spores and vegetative cells. The proteins shared between spores and

cells are mostly ribosomal proteins, cell cycle-regulating and/or associated proteins, and

cytosolic proteins involved in pathways required for anabolism and catabolic pathways of energy

metabolism distributed over 46 categories by DAVID (**Table 1, Supplementary Figure S1**).

These also include products of 25 essential genes[23] such as peptidoglycan synthesis protein

MurG (CD2651) and formate-tetrahydrofolate ligase CD0718 (2 and 2.4 times higher levels in

spores than in vegetative cells, respectively), and S-adenosylmethionine synthase MetK as well

as a rubrerythrin CD0825 (~4 and ~9 times higher levels in vegetative cells than in spores,

respectively) (**Fig 2**). The TMHMM analysis of the shared proteins identified 123 membrane

proteins (**Supplementary Table S2**), such as the phosphotransferase system (PTS) of sugar

transporters, ABC-type transporters, and V-type ATPases. These also included most proteins

involved in the Wood-Ljungdahl pathway (**Fig 3**). From **Table 1** it can be deduced that proteins

from this category that are present in spores are essentially those that are required for

hibernation, the initiation of growth and the resumption of metabolism upon germination and

initiation of outgrowth.

Discussion

To our knowledge, the proteomes of *P. difficile* vegetative cells and spores have been explored

for the first time in a single experimental set-up to understand the fundamental differences

between these two morphological forms of the bacterium. To this end, spores and ^{15}N-labelled

vegetative cells have been mixed for relative quantification. Metabolic labelling using ^{15}N

isotopes is a highly accurate means of proteome quantification[24]. However, a method for

labelling *P. difficile* was unavailable until recently[16]. Here a method for metabolic labelling using

^{15}N-labelled yeastolate medium has been developed which provides a simple, economical, and

rapid means to perform quantitative proteomics of a variety of pathogenic and non-pathogenic

microbes. Our analyses show that 80% of the quantified proteins are common for both cells and

spores, indicating that pathogenic *P. difficile* employs a relatively modest proteomic changeover

to enable long-term survival as a dormant spore. The corresponding pathways are shared

between vegetative cells and spores (**Supplementary Figure S1**) however the relative quantities

of these proteins vary between cells and spores. A discussion of quantified and functionally key

proteins of *P. difficile* is presented below.

Peptoclostridium difficile expresses an array of cell surface proteins, including the S-layer

protein SlpA and its paralogues from the cell wall protein (CWP) family. Eight Cwp family

proteins are quantified: Cwp18 and 22 are identified in both morphotypes, with higher levels in

spores, whereas Cwp2, 5, 6, 19, 84, and CwpV are identified in vegetative cells. Cwp22, a

functional homologue of LD-transpeptidase (Ldt_{cd2}), is an important spore protein that plays a

role in peptidoglycan remodelling and confers resistance to β-lactam antibiotics[25,26]. CwpV

promotes *P. difficile* aggregation and its strain-dependent structural variations may assist in

evading the host antibody response[27] or to launch an anti-phage strategy[28]. *Peptoclostridium*

difficile spores frequently have an interspace region between the spore coat and the fragile,

heterogeneous exosporium[29,30]. Although most known and putative exosporium proteins

described previously[20] have been identified in this study, the BclA family of proteins have not

(except BclA1 (CD0332), identified in one replicate and thus not quantified). An absence of hair-

like structures in the *P. difficile* 630 exosporium[30] may underlie this finding. Other identified

proteins such as CD1474, CD2845, and CD1524 - all rubrerythrins -likely present in the

exosporium, may play a role in fighting reactive oxygen species and oxidative stress[31].

Peptoclostridium difficile relies heavily on the phosphoenolpyruvate-dependent phospho-

transferase system (PTS) for uptake and regulation of various sugars and sugar derivatives[32]. The

PTS is clearly advantageous for germinating spores, since various sugars are readily available in

the human gut and may thus be used to facilitate outgrowth and establish infection[33,34]. We have

identified 22 PTS proteins, of which 15 could be quantified (**Supplementary Table S1**). The

quantified PTS proteins, except CD3027, are shared between vegetative cells and spores. Along

with the central PTS proteins HPr (PtsH/CD2756) and Enzyme I (PtsI/CD2755), these proteins

function in the transport of fructose (FruABC/CD2269, CD2486-88), glucose (PtsG/CD2667,

CD3027, CD3089), mannitol (MtlF/CD2332 and MtlA/CD2334)), mannose (CD3013-14),

xyloside (XynB/CD3068), and β-glucoside (BglF5/CD3137). Although shared, CD3013-14,

CD2486-88, CD3089, PtsH, and MtlA-F showed relatively higher levels in spores, whereas

PtsG, PtsI, BglF5, FruABC, CD3027, and XynB show lower levels in spores. Interestingly, a

previous study has shown that in germinating spores, *bglF5* and *ptsG* transcripts are

downregulated whereas those of *fruABC* and *cd2486-87* are upregulated[35]. In mouse infections,

P. difficile CD2487 is upregulated 14 h post-infection whereas proteins XynB, CD3027, and PtsI

are downregulated 38 h post-infection[8]. In pig infections of *P. difficile* PtsI, BglF5 (4-12 h post-

infection) and MtlA (only 12 h post-infection) are upregulated[36] and XynB and CD3013-14 are

downregulated. Furthermore, MtlA and MtlF can repress *tcdA* and *tcdB* toxin expression in *P.*

difficile[37]. Put together, these studies indicate that PtsI, BglF5, CD2487, and MtlA potentially

play a role in the pathogenesis of *P. difficile* infections; CD2487 and MtlA are predominantly

present in spores, making them important targets in understanding spore persistence.

Non-PTS transport systems are also involved in carbohydrate uptake in clostridia[32]. Of all

ATPases and related proteins quantified, only 6 and 4 proteins belong to the cell-predominant

and spore-predominant categories, respectively. Four V-type ATPases have been quantified;

however, only AtpC is spore-predominant whereas the other three are also expressed in cells.

AtpC is associated with proton transport, possesses a hydrolase activity, and contains a CodY-

binding region[38] potentially repressing toxin expression in *P. difficile*[39]. It also regulates

synthesis and circulation of pyruvate and 2-oxoglutarate in the cell[40], providing proteomic

flexibility during spore revival. The spore-predominant ATPases include a cation (Ca^{2+})-

transporting ATPase (CD2503), which shares 42% identity with *B. subtilis* YloB ATPase, likely

responsible for accumulating intracellular calcium and reinforcing thermal resistance[41]. Two

ABC-type transporters - lipoproteins CD2365 and SsuA - are expressed in both *P. difficile*

vegetative cells and spores[17] but are present at higher levels in spores. These are alkanesulfonate

and taurine binding proteins, respectively and SsuA is involved in sulfur metabolism[42].

Interestingly, the taurine side chain of taurocholate selectively binds its potential receptor site[43]

and taurine itself is an alkane sulfonate, thus higher levels of SsuA and CD2365 in spores could

indicate potential taurine interaction during germination, similar to CD3298[44]. Identified protein

CD0114 shares 25% identity with CD3298, making it worth studying for a possible role in spore

germination. Additionally, CD3669 - an orthologue of GerM -might be involved in sporulation[45].

The 'stay-green' family protein CD3613 is a putative exosporium protein. Usually, these proteins

perform chlorophyll degradation[46] but the upregulation of CD3613 during sporulation in a mouse

model[8] suggests a potential role in pathogenesis. CD2434 has an UBA_NAC_like bacterial

protein domain commonly found in proteins involved in ubiquitin-dependent proteolysis[47].

Although not a direct evidence, this observation suggests involvement of CD2434 in

pathogenesis; a previous study has shown that the *E. coli* toxin CNF1 utilizes the ubiquitin-

proteasome assembly of host cells to partially inactivate their Rho GTPases[48], a mechanism

similar to that of TcdA and TcdB toxins[49]. The spore envelope protein CD2635 may be involved

in germination[9]. This protein, similar to CD2636, contains a characteristic YIEGIA domain and

both could play significant roles in spore assembly as well as disintegration.

Amino acids play a crucial role in spore germination and functioning of the Stickland pathways

in *P. difficile*. CD3458 and CD1555 are putative amino acid permeases identified to be slightly

more abundant in spores than in vegetative cells. CD3458 contains a putative amino acid

permease domain and an SLC5-6-like_sbd superfamily domain, thus qualifying as an amino acid

permease and sodium/glucose co-transporter. Another amino acid permease CD2612, identified

in vegetative cells, is upregulated in the presence of cysteine[42], implying a role in sulfur

metabolism. CD2344 contains an Asp-Al_Ex domain found in aspartate-alanine antiporters and

might be capable of developing a membrane potential enough to carry ATP synthesis via FoF1

ATPase[50].

Peptidases and proteases are crucial for vegetative cell and spore survival. In this study, 29

peptidases and 12 proteases are identified and quantified. These include proteins involved in

germination, such as Gpr (CD2470) and CspBA. In the present study, CspC protease has been

detected in only one replicate and thus not quantified. Other proteins involved in cellular

regulatory processes, such as ATP-dependent Clp proteases, zinc metalloproteases, serine

proteases, Lon proteases, have also been quantified in the present study. These peptidases belong

to various families such as aminopeptidase (M1), metalloendopeptidase (M16), aminopeptidase

(M18), membrane dipeptidase (M19), glutamate carboxypeptidase (M20), glycoprotease (M22),

methionyl aminopeptidase (M24), prolyl oligopeptidase (S9), and collagenase (U32). It is

speculated that proteins belonging to the M22 and U32 family (CD0150 and CD1228,

respectively) function in spore germination. In fact, BA0261, a CD0150 orthologue in *B.*

anthracis, is suggested to play a role in spore germination[51] and a collagenase is implicated in

virulence of *B. cereus* endophthalmitis[52] indicating a similar potential for CD1228. A previous

study suggested that a Lon protease in *Brucella* sp. is involved in BALB/c mice infections[53].

Thus, the Lon protease CD3301, present in vegetative cells and spores, also likely plays a role in

infection. CD0684 has been previously reported to be present in *P. difficile* spores under σ^G

regulation and suggested to be involved in stress resistance[9]. Notably, in the present study, none

of the identified peptidases or proteases are found to be predominant in vegetative cells.

From the quantified dataset 198 proteins are classified to the metabolic pathways category by

DAVID analysis. Although the spores are metabolically dormant, the proteins belonging to the

amino acid biosynthesis, purine metabolism, glycolysis, fatty acid metabolism, and nitrogen

metabolism are present and form the core protein set in spores. Moreover, in spores, arginine

biosynthesis pathway proteins are present at ~20% of the levels detected in vegetative cells. This

indicates that germinating spores require *de novo* synthesis of these proteins post-germination to

assist the outgrowing cells. Ribosomal proteins - except the 50S ribosomal protein L30

(CD0881) - are also present in low amounts in *P. difficile* spores. CD0881 has a ferredoxin-like

fold, resembling the structure of yeast L7 proteins, and is likely involved in processing

precursors of large rRNAs[54], a function that could well aid the outgrowing spores. The phosphate

butyryltransferases (CD0715 and CD0112/Ptb) involved in the butanoate metabolism pathway

are present not only in vegetative cells but also in spores, thus conferring on the spores a

metabolic flexibility.

Peptoclostridium difficile may deploy several sulfur and nitrogen metabolism proteins while

surviving in anaerobic conditions. Of these, only CD2431, a nitrite/sulfite reductase, is abundant

in spores. This protein also contains a 4Fe-4S domain and can catalyse the reduction of sulfite to

sulfide and nitrite to ammonia[55]. *Peptoclostridium difficile* and other acetogenic clostridia have

acquired such metabolic flexibility that they can directly utilize the CO_2 and H_2 from air and

yield a variety of products including acetate and methane[56]. The Wood-Ljungdahl pathway of

acetogenesis is believed to be the first biochemical pathway to have emerged on earth[57] and all

proteins involved in this pathway are identified in *P. difficile* 630, which reinforces the

acetogenic nature of *P. difficile* growth. Of these, CD3405, CD3407 and CD0730 have been

detected only in single replicates and thus are not quantified. The other Wood-Ljungdahl

pathway proteins have all been quantified, with only three proteins - MetF (CD0722), CD0728,

and CD3258 - being highly abundant in spores. In contrast, only a single protein - CD0893 - is

predominant in vegetative cells.

The acetogenic mode of life of *P. difficile* requires specific enzymes, such as acetyl-CoA

synthases/CO dehydrogenases (CD0174, CD0176, and CD0727), formate dehydrogenases

(CD2179), and iron-only hydrogenases (CD0893, CD3258, and CD3406). Enzymes CD0174 and

CD0176 synthesize the key metabolite acetyl-CoA from CO, methyl corrinoid, and CoASH. The

formate dehydrogenases can be seleno (CD3317) or non-seleno (CD0769 and CD2179)

enzymes. Protein CD2179, an anaerobic dehydrogenase, reduces CO_2 to formate which is further

metabolized to acetyl-CoA through enzymatic reactions, one of them involving another acetyl-

CoA synthase/CO dehydrogenase with a methyltransferase subunit. In acetogens lacking

cytochromes, the Rnf complex (encoded by CD1137-42) is the putative coupling site for energy

conservation[56]. In the present study, all components of the Rnf complex, except CD1140-41,

have been identified. The Rnf complex proteins, together with electron transport flavoproteins

etfA2/B2 (CD1055-56), are employed in butyrate formation[58]. However, the present study has

identified only etfA1/B1 and etfA3/B3 proteins. These proteins are predominant in vegetative

cells, indicating that they likely function exclusively during the vegetative life cycle of *P.*

difficile.

The iron-only hydrogenases, 10 times more efficient in hydrogen production than [NiFe]

hydrogenases[59], are abundant in clostridia. *Peptoclostridium difficile* encodes two trimeric and

three monomeric hydrogenases[60]. Moreover, proteins CD3405-3407 function as electron-

bifurcating hydrogenases whereby physiological electron carriers such as ferredoxin are used for

H_2 production[61]. In the present study, CD3258 is seen predominantly in spores whereas CD0893

occurs mostly in vegetative cells. Both proteins are monomeric, ferredoxin dependent[61], and

contain a H-cluster i.e. a centre for hydrogen production[62]. However, CD3258 has a sequence of

eight cysteines for stabilizing two [4Fe4S] clusters transferring electrons from the surface to the

protein's active site[63] whereas CD0893 has a single FeS domain with a $(Cx_{1-4}Cx_{5-9}Cx_3C)$

arrangement at its N-terminus and the H-cluster has an additional cysteine residue

(TSCCCPxW[60]). The predominant expression of CD3258 and CD0893 in spores and vegetative

cells, respectively, indicates the distinct roles of these proteins in *P. difficile* physiology.

In addition, a few noteworthy and yet uncharacterized proteins are detected in our study.

CD1470, a sulfotransferase, may be involved in cyanide detoxification. PdaA1 (CD1430) and

PdaA2 (CD2719) have recently been shown to be important for cortex muramic acid-δ-lactam

synthesis; spores lacking it are heat sensitive, deficient in germination, and exhibit late

virulence[64]. CD2719 is not identified in the present study; however, CD1556, an orthologue of

PdaA2 of *B. cereus* var. *anthracis* is identified. Thus, CD1556 may be important for spore

structure and germination. CD1319, an orthologue of YlxY, of *B. subtilis*[65] may be important for

sporulation.

Concluding remarks

The one-pot sample processing method along with ^{15}N metabolic labelling has enabled a

reproducible, combined cell and spore quantitative proteome analysis of the anaerobic pathogen

P. difficile 630. The analysis outlines a relatively modest proteomic adaptation of this

evolutionarily and clinically important anaerobic pathogen, when as a survival strategy, it

completes spore formation. In addition to specific cell and spore surface proteins, the study has

qualified vegetative cell proteins CD1228, CD3301 and spore proteins CD2487, CD2434 and

CD0684 as potential biomarkers for *P. difficile* infections.

Materials and Methods

Bacterial strains, cell culture, and sporulation

Peptoclostridium difficile strain 630 (ATCC® BAA1382™), acquired from the Leibniz Institute

of Microorganisms and Cell Cultures, Germany, was used to derive vegetative cells and spores.

All cultivations were performed at 37°C in an anaerobic chamber (Whitley DG250) supplied

with a gas mixture comprising 10% hydrogen, 10% carbon dioxide, and 80% nitrogen. The cells

were first grown overnight in Schaedler anaerobic broth (Oxoid, CM0497) and further passaged

thrice through the newly developed ^{15}N-yeastolate medium (described below) to obtain ^{15}N-

labelled vegetative cells. After the third passage, the cells were grown overnight until $OD_{600} \approx$

1.7 and harvested by centrifugation. These cells were then aliquoted and stored at -20°C until

further use. To obtain spores, the vegetative cells were pre-cultured overnight in Columbia broth

and inoculated in Clospore medium [66]. Typically, bottles containing 500 ml of Clospore medium,

kept in the anaerobic chamber overnight, were inoculated with the pre-cultures. Spores were

harvested after 2 weeks of incubation and intensively purified using a combination of ultra-

sonication, enzyme treatment (lysozyme, trypsin, and proteinase K), and washing with sterile

milli-Q water[18,66]. The spore crops were subjected to density gradient centrifugation by layering

spores suspended in 20% Histodenz (Sigma-Aldrich, USA) on top of 50% Histodenz in 2 ml

Eppendorf tubes, and centrifuging for 25 min at 15000 $\times g$.

Preparation of ^{15}N-yeastolate medium

Saccharomyces cerevisiae CEN. PK1137D was grown at 37°C in a defined CBS medium[67]

modified with ^{15}NH$_4$Cl (replacing (NH$_4$)$_2$SO$_4$) as the sole nitrogen source. Yeast cells were

harvested by centrifugation (5000 $\times g$, 30 min) and washed with water. The protocol to generate

yeastolate was adapted from previous studies[68,69]. The yeast cells were made into a 30% (w/v)

slurry, ultrasonicated by a tip ultrasonicator for 30 min. The pH of the slurry was adjusted to 7.5

using NaOH before incubating under continuous shaking for 5 days at 50°C. Thereafter, the

slurry was ultrasonicated again and centrifuged at 20000 $\times g$ for 30 min to collect the supernatant.

The pellet was washed twice and the supernatants were combined and lyophilized, to generate

powdered yeastolate. The final yeastolate medium contained 2% ^{15}N-yeastolate, 2% glucose, and

0.2% NaCl.

One-pot sample processing

The one-pot protocol has been previously described in detail[14]. Typically, 0.5 mg (dry weight) of

spores/cells was suspended in lysis buffer and disrupted for seven cycles with 0.1 mm zirconium

beads (BioSpec Products, Bartlesville, OK, USA) using a Precellys 24 homogenizer (Bertin

Technologies, Aix en Provence, France). The tubes were incubated for 1 h at 56°C and alkylated

using 15 mM iodoacetamide for 45 min at room temperature in the dark. The reaction was

quenched with 20 mM thiourea and digestion with Lys-C (at 1:200 protease/protein ratio) was

carried out for 3 h at 37°C. Samples were diluted with 50 mM ammonium bicarbonate buffer and

digested with trypsin (at 1:100 protease/protein ratio) was carried out at 37°C for 18 h. The

tryptic digest was freeze-dried. Before use, the freeze-dried samples were re-dissolved in 0.1%

TFA and desalted using Omix µC18 pipette tips (80 µg capacity, Varian, Palo Alto, CA, USA)

according to the manufacturer's instructions.

Fractionation of peptides

ZIC-HILIC chromatography was used to fractionate the freeze-dried peptide samples. Dried

digests were dissolved in 500 µl of Buffer A (85% acetonitrile, 5 mM ammonium acetate, 0.4%

acetic acid, pH 5.8), centrifuged to remove any undissolved components, and injected into the

chromatography system. An isocratic flow with 100% Buffer A for 10 min was followed by a

gradient of 0-30% Buffer B (30% acetonitrile, 5 mM ammonium acetate, 0.5% acetic acid, pH

3.8) in the first phase and 30-100% of Buffer B in the second phase (flow rate 400 µl/min). The

peptides were eluted and collected in 10 fractions, freeze-dried, and stored at -80°C until further

use.

LC-FT-ICR MS/MS analysis

ZIC-HILIC fractions were re-dissolved in 0.1% TFA, peptide concentrations were determined by

measuring absorbance at 205nm and 300ng tryptic peptide mixtures were injected for analyses.

LC-MS/MS data of each ZIC-HILIC fraction were acquired with an Apex Ultra Fourier

transform ion cyclotron resonance mass spectrometer (Bruker Daltonics, Bremen, Germany)

equipped with a 7 T magnet and a Nano electrospray Apollo II Dual Source coupled to an

Ultimate 3000 (Dionex, Sunnyvale, CA, USA) HPLC system. LC conditions and acquisition

parameters were as described previously[14].

Data analysis and bioinformatics

Each raw FT-MS/MS data set was mass calibrated better than 1.5 ppm on the peptide fragments

from the co-injected GluFib calibrant. The 10 ZIC-HILIC fractions were jointly processed as a

multifile with the MASCOT DISTILLER program (version 2.4.3.1, 64 bits), MDRO 2.4.3.0

(MATRIX science, London, UK), including the Search toolbox and the Quantification toolbox.

Peak-picking for both MS and MS/MS spectra was optimized for the mass resolution of up to

60000. Peaks were fitted to a simulated isotope distribution with a correlation threshold of 0.7,

with minimum signal to noise ratio of 2. The processed data were searched in a MudPIT

approach with the MASCOT server program 2.3.02 (MATRIX science, London, UK) against the

P. difficile 630 ORF translation database. The MASCOT search parameters were as follows:

enzyme - trypsin, allowance of two missed cleavages, fixed modification - carboamidomethylat-

ion of cysteine, variable modifications - oxidation of methionine and deamidation of asparagine

and glutamine, quantification method - metabolic ^{15}N labelling, peptide mass tolerance and

peptide fragment mass tolerance - 50 ppm. MASCOT MudPIT peptide identification threshold

score of 20 and FDR of 2% were set to export the reports.

Using the quantification toolbox, the quantification of the light spore peptides relative to the

corresponding heavy cell peptides was determined as light/heavy ratio using Simpson's

integration of the peptide MS chromatographic profiles for all detected charge states. The

quantification parameters were: Correlation threshold for isotopic distribution fit - 0.98, ^{15}N label

content - 99.6%, XIC threshold - 0.1, all charge states on, max XIC width -120 seconds, elution

time shift for heavy and light peptides - 20 seconds. All isotope ratios were manually validated

by inspecting the MS spectral data. The protein isotopic ratios were then calculated as the

average over the corresponding peptide ratios. For each of the three replicas, the identification

and quantification reports were imported into a custom made program to facilitate data

combination and statistical analysis. Protein identification was validated with identifications in at

least two replicas. For these identified proteins the relative quantification was calculated as the

geometric mean of at least two validated light/ heavy ratios. All identification and quantification

protein data are listed in **Table S1**. The mass spectrometry proteomics data have been deposited

as a partial submission to the ProteomeXchange Consortium via the PRIDE[70] partner repository

with the dataset identifier PXD012030.

Transmembrane proteins were predicted using the default parameters on the TMHMM Server

(version 2.0 http://www.cbs.dtu.dk/services/TMHMM/). DAVID Bioinformatics Resources tool

(version 6.8) was used[71] to retrieve the functional annotation data of UniProt key word and

KEGG pathway classifications. The BioCyc pathway analysis tool[72] was used to generate a

cellular overview of the quantified proteins.

Acknowledgements

W.R.A is supported by the grant NWO ALWOP.260. L.Z. acknowledges the Erasmus Mundus

program (EMEA3) and TNO (Healthy Living) for funding this research.

Author Contributions

W.R.A analysed the data, prepared the figures and tables and wrote the main manuscript text.

L.Z. performed the experiments. L. dK conceptualized and designed the experiments as well as

curated and processed the proteomics data. S.B. and C. dK supervised and mentored the research.

All authors reviewed the manuscript.

Competing interests

The authors declare no competing interests.

References

Shin, J. H., Chaves-Olarte, E. & Warren, C. A. *Clostridium difficile* Infection. *Microbiol Spectr* **4**, doi:10.1128/microbiolspec.EI10-0007-2015 (2016).

Bagdasarian, N., Rao, K. & Malani, P. N. Diagnosis and treatment of *Clostridium difficile* in adults: a systematic review. *JAMA* **313**, 398-408, doi:10.1001/jama.2014.17103 (2015).

Aktories, K., Schwan, C. & Jank, T. *Clostridium difficile* Toxin Biology. *Annu Rev Microbiol* **71**, 281-307, doi:10.1146/annurev-micro-090816-093458 (2017).

Francis, M. B., Allen, C. A., Shrestha, R. & Sorg, J. A. Bile acid recognition by the *Clostridium difficile* germinant receptor, CspC, is important for establishing infection. *PLoS Pathog* **9**, e1003356, doi:10.1371/journal.ppat.1003356 (2013).

Galperin, M. Y. *et al.* Genomic determinants of sporulation in Bacilli and Clostridia: towards the minimal set of sporulation-specific genes. *Environ Microbiol* **14**, 2870-2890, doi:10.1111/j.1462-2920.2012.02841.x (2012).

Paredes-Sabja, D., Shen, A. & Sorg, J. A. *Clostridium difficile* spore biology: sporulation, germination, and spore structural proteins. *Trends Microbiol* **22**, 406-416, doi:10.1016/j.tim.2014.04.003 (2014).

Francis, M. B., Allen, C. A. & Sorg, J. A. Spore Cortex Hydrolysis Precedes Dipicolinic Acid Release during *Clostridium difficile* Spore Germination. *J Bacteriol* **197**, 2276-2283, doi:10.1128/JB.02575-14 (2015).

8   Janoir, C. *et al.* Adaptive strategies and pathogenesis of *Clostridium difficile* from in vivo

transcriptomics. *Infect Immun* **81**, 3757-3769, doi:10.1128/IAI.00515-13 (2013).

9   Saujet, L. *et al.* Genome-wide analysis of cell type-specific gene transcription during

spore formation in *Clostridium difficile*. *PLoS Genet* **9**, e1003756,

doi:10.1371/journal.pgen.1003756 (2013).

10   Chen, J. W. *et al.* Proteomic comparison of historic and recently emerged hypervirulent

Clostridium difficile strains. *J Proteome Res* **12**, 1151-1161, doi:10.1021/pr3007528

(2013).

11   Chilton, C. H. *et al.* Comparative proteomic analysis of *Clostridium difficile* isolates of

varying virulence. *J Med Microbiol* **63**, 489-503, doi:10.1099/jmm.0.070409-0 (2014).

12   Jain, S., Graham, C., Graham, R. L., McMullan, G. & Ternan, N. G. Quantitative

proteomic analysis of the heat stress response in *Clostridium difficile* strain 630. *J*

Proteome Res **10**, 3880-3890, doi:10.1021/pr200327t (2011).

13   Otto, A. *et al.* The protein inventory of *Clostridium difficile* grown in complex and

minimal medium. *Proteomics Clin Appl* **10**, 1068-1072, doi:10.1002/prca.201600069

(2016).

14   Swarge, B. N. *et al.* "One-Pot" Sample Processing Method for Proteome-Wide Analysis

of Microbial Cells and Spores. *Proteomics Clin Appl* **12**, e1700169,

doi:10.1002/prca.201700169 (2018).

15   Ternan, N. G., Jain, S., Srivastava, M. & McMullan, G. Comparative transcriptional

analysis of clinically relevant heat stress response in *Clostridium difficile* strain 630.

PLoS One **7**, e42410, doi:10.1371/journal.pone.0042410 (2012).

16   Trautwein-Schult, A., Maass, S., Plate, K., Otto, A. & Becher, D. A Metabolic Labeling

Strategy for Relative Protein Quantification in *Clostridioides difficile*. *Front Microbiol* **9**,

2371, doi:10.3389/fmicb.2018.02371 (2018).

17   Charlton, T. M., Kovacs-Simon, A., Michell, S. L., Fairweather, N. F. & Tate, E. W.

Quantitative Lipoproteomics in *Clostridium difficile* Reveals a Role for Lipoproteins in

Sporulation. *Chem Biol* **22**, 1562-1573, doi:10.1016/j.chembiol.2015.10.006 (2015).

18   Lawley, T. D. *et al.* Proteomic and genomic characterization of highly infectious

Clostridium difficile 630 spores. *J Bacteriol* **191**, 5377-5386, doi:10.1128/JB.00597-09

(2009).

19   Abhyankar, W. R. *et al.* In pursuit of protein targets: proteomic characterization of

bacterial spore outer layers. *J Proteome Res* **12**, 4507-4521, doi:10.1021/pr4005629

(2013).

20   Díaz-González, F. *et al.* Protein composition of the outermost exosporium-like layer of

Clostridium difficile 630 spores. *J Proteomics* **123**, 1-13, doi:10.1016/j.jprot.2015.03.035

(2015).

21   Strong, P. C. *et al.* Identification and characterization of glycoproteins on the spore

surface of *Clostridium difficile*. *J Bacteriol* **196**, 2627-2637, doi:10.1128/JB.01469-14

(2014).

22   Kilpatrick, E. L., Liao, W. L., Camara, J. E., Turko, I. V. & Bunk, D. M. Expression and

characterization of [15]N-labeled human C-reactive protein in *Escherichia coli* and *Pichia*

pastoris for use in isotope-dilution mass spectrometry. *Protein Expr Purif* **85**, 94-99,

doi:10.1016/j.pep.2012.06.019 (2012).

23   Larocque, M., Chenard, T. & Najmanovich, R. A curated *C. difficile* strain 630 metabolic

network: prediction of essential targets and inhibitors. *BMC Syst Biol* **8**, 117,

doi:10.1186/s12918-014-0117-z (2014).

24   Bantscheff, M., Schirle, M., Sweetman, G., Rick, J. & Kuster, B. Quantitative mass

spectrometry in proteomics: a critical review. *Anal Bioanal Chem* **389**, 1017-1031,

doi:10.1007/s00216-007-1486-6 (2007).

25   Peltier, J. *et al. Clostridium difficile* has an original peptidoglycan structure with a high

level of N-acetylglucosamine deacetylation and mainly 3-3 cross-links. *J Biol Chem* **286**,

29053-29062, doi:10.1074/jbc.M111.259150 (2011).

26   Ternan, N. G., Jain, S., Graham, R. L. & McMullan, G. Semiquantitative analysis of

clinical heat stress in *Clostridium difficile* strain 630 using a GeLC/MS workflow with

emPAI quantitation. *PLoS One* **9**, e88960, doi:10.1371/journal.pone.0088960 (2014).

27   Reynolds, C. B., Emerson, J. E., de la Riva, L., Fagan, R. P. & Fairweather, N. F. The

Clostridium difficile cell wall protein CwpV is antigenically variable between strains, but

exhibits conserved aggregation-promoting function. *PLoS Pathog* **7**, e1002024,

doi:10.1371/journal.ppat.1002024 (2011).

28   Sekulovic, O., Ospina Bedoya, M., Fivian-Hughes, A. S., Fairweather, N. F. & Fortier, L.

C. The *Clostridium difficile* cell wall protein CwpV confers phase-variable phage

resistance. *Mol Microbiol* **98**, 329-342, doi:10.1111/mmi.13121 (2015).

29   Permpoonpattana, P. *et al.* Surface layers of *Clostridium difficile* endospores. *J Bacteriol*

193, 6461-6470, doi:10.1128/JB.05182-11 (2011).

30   Pizarro-Guajardo, M., Calderon-Romero, P., Castro-Cordova, P., Mora-Uribe, P. &

Paredes-Sabja, D. Ultrastructural Variability of the Exosporium Layer of *Clostridium*

difficile Spores. *Appl Environ Microbiol* **82**, 2202-2209, doi:10.1128/AEM.03410-15

(2016).

31 Sztukowska, M., Bugno, M., Potempa, J., Travis, J. & Kurtz, D. M., Jr. Role of

rubrerythrin in the oxidative stress response of *Porphyromonas gingivalis*. *Mol Microbiol*

44, 479-488 (2002).

32 Mitchell, W. J. & Tangney, M. in *Handbook on Clostridia* (ed P. Dürre) Ch. 8, 196-225

(CRC Press, 2005).

33 Chassard, C. & Lacroix, C. Carbohydrates and the human gut microbiota. *Curr Opin Clin*

Nutr Metab Care **16**, 453-460, doi:10.1097/MCO.0b013e3283619e63 (2013).

34 Flint, H. J., Scott, K. P., Louis, P. & Duncan, S. H. The role of the gut microbiota in

nutrition and health. *Nat Rev Gastroenterol Hepatol* **9**, 577-589,

doi:10.1038/nrgastro.2012.156 (2012).

35 Dembek, M., Stabler, R. A., Witney, A. A., Wren, B. W. & Fairweather, N. F.

Transcriptional analysis of temporal gene expression in germinating *Clostridium difficile*

630 endospores. *PLoS One* **8**, e64011, doi:10.1371/journal.pone.0064011 (2013).

36 Scaria, J. *et al. Clostridium difficile* transcriptome analysis using pig ligated loop model

reveals modulation of pathways not modulated in vitro. *J Infect Dis* **203**, 1613-1620,

doi:10.1093/infdis/jir112 (2011).

37 Dupuy, B. & Sonenshein, A. L. Regulated transcription of *Clostridium difficile* toxin

genes. *Mol Microbiol* **27**, 107-120 (1998).

38 Dineen, S. S., McBride, S. M. & Sonenshein, A. L. Integration of metabolism and

virulence by *Clostridium difficile* CodY. *J Bacteriol* **192**, 5350-5362,

doi:10.1128/JB.00341-10 (2010).

39   Dineen, S. S., Villapakkam, A. C., Nordman, J. T. & Sonenshein, A. L. Repression of

Clostridium difficile toxin gene expression by CodY. *Mol Microbiol* **66**, 206-219,

doi:10.1111/j.1365-2958.2007.05906.x (2007).

40   Sonenshein, A. L. Control of key metabolic intersections in *Bacillus subtilis*. *Nat Rev*

Microbiol **5**, 917-927, doi:10.1038/nrmicro1772 (2007).

41   Raeymaekers, L., Wuytack, E., Willems, I., Michiels, C. W. & Wuytack, F. Expression

of a P-type Ca($^{2+}$)-transport ATPase in *Bacillus subtilis* during sporulation. *Cell Calcium*

32, 93 (2002).

42   Dubois, T. *et al.* Control of *Clostridium difficile* Physiopathology in Response to

Cysteine Availability. *Infect Immun* **84**, 2389-2405, doi:10.1128/IAI.00121-16 (2016).

43   Howerton, A., Ramirez, N. & Abel-Santos, E. Mapping interactions between germinants

and *Clostridium difficile* spores. *J Bacteriol* **193**, 274-282, doi:10.1128/JB.00980-10

(2011).

44   Kochan, T. J. *et al.* Intestinal calcium and bile salts facilitate germination of *Clostridium*

difficile spores. *PLoS Pathog* **13**, e1006443, doi:10.1371/journal.ppat.1006443 (2017).

45   Rodrigues, C. D., Ramirez-Guadiana, F. H., Meeske, A. J., Wang, X. & Rudner, D. Z.

GerM is required to assemble the basal platform of the SpoIIIA-SpoIIQ transenvelope

complex during sporulation in *Bacillus subtilis*. *Mol Microbiol* **102**, 260-273,

doi:10.1111/mmi.13457 (2016).

46   Park, S. Y. *et al.* The senescence-induced staygreen protein regulates chlorophyll

degradation. *Plant Cell* **19**, 1649-1664, doi:10.1105/tpc.106.044891 (2007).

47   Hofmann, K. & Bucher, P. The UBA domain: a sequence motif present in multiple

enzyme classes of the ubiquitination pathway. *Trends Biochem Sci* **21**, 172-173 (1996).

48   Doye, A. *et al.* CNF1 exploits the ubiquitin-proteasome machinery to restrict Rho

539   GTPase activation for bacterial host cell invasion. *Cell* **111**, 553-564 (2002).

540   49   Voth, D. E. & Ballard, J. D. *Clostridium difficile* toxins: mechanism of action and role in

541   disease. *Clin Microbiol Rev* **18**, 247-263, doi:10.1128/CMR.18.2.247-263.2005 (2005).

542   50   Abe, K. *et al.* Plasmid-encoded asp operon confers a proton motive metabolic cycle

543   catalyzed by an aspartate-alanine exchange reaction. *J Bacteriol* **184**, 2906-2913 (2002).

544   51   Liu, H. *et al.* Formation and composition of the *Bacillus anthracis* endospore. *J Bacteriol*

545   **186**, 164-178 (2004).

546   52   Beecher, D. J., Olsen, T. W., Somers, E. B. & Wong, A. C. Evidence for contribution of

547   tripartite hemolysin BL, phosphatidylcholine-preferring phospholipase C, and

548   collagenase to virulence of *Bacillus cereus* endophthalmitis. *Infect Immun* **68**, 5269-5276

549   (2000).

550   53   Robertson, G. T. *et al.* The *Brucella abortus* CcrM DNA methyltransferase is essential

551   for viability, and its overexpression attenuates intracellular replication in murine

552   macrophages. *J Bacteriol* **182**, 3482-3489 (2000).

553   54   Dunbar, D. A., Dragon, F., Lee, S. J. & Baserga, S. J. A nucleolar protein related to

554   ribosomal protein L7 is required for an early step in large ribosomal subunit biogenesis.

555   *Proc Natl Acad Sci U S A* **97**, 13027-13032, doi:10.1073/pnas.97.24.13027 (2000).

556   55   Zeghouf, M., Fontecave, M. & Coves, J. A simplifed functional version of the

557   *Escherichia coli* sulfite reductase. *J Biol Chem* **275**, 37651-37656,

558   doi:10.1074/jbc.M005619200 (2000).

559   56   Köpke, M., Straub, M. & Dürre, P. *Clostridium difficile* is an autotrophic bacterial

560   pathogen. *PLoS One* **8**, e62157, doi:10.1371/journal.pone.0062157 (2013).

57   Russell, M. J. & Martin, W. The rocky roots of the acetyl-CoA pathway. *Trends Biochem*

Sci **29**, 358-363, doi:10.1016/j.tibs.2004.05.007 (2004).

58   Aboulnaga el, H. *et al.* Effect of an oxygen-tolerant bifurcating butyryl coenzyme A

dehydrogenase/electron-transferring flavoprotein complex from *Clostridium difficile* on

butyrate production in *Escherichia coli*. *J Bacteriol* **195**, 3704-3713,

doi:10.1128/JB.00321-13 (2013).

59   Buckel, W. in *Handbook on Clostridia* (ed P. Dürre) Ch. 9, 226-283 (CRC Press, 2005).

60   Calusinska, M., Happe, T., Joris, B. & Wilmotte, A. The surprising diversity of clostridial

hydrogenases: a comparative genomic perspective. *Microbiology* **156**, 1575-1588,

doi:10.1099/mic.0.032771-0 (2010).

61   Schut, G. J. & Adams, M. W. The iron-hydrogenase of *Thermotoga maritima* utilizes

ferredoxin and NADH synergistically: a new perspective on anaerobic hydrogen

production. *J Bacteriol* **191**, 4451-4457, doi:10.1128/JB.01582-08 (2009).

62   Cammack, R. Hydrogenase sophistication. *Nature* **397**, 214-215, doi:10.1038/16601

(1999).

63   Florin, L., Tsokoglou, A. & Happe, T. A novel type of iron hydrogenase in the green alga

Scenedesmus obliquus is linked to the photosynthetic electron transport chain. *J Biol*

Chem **276**, 6125-6132, doi:10.1074/jbc.M008470200 (2001).

64   Coullon, H. *et al.* N-Deacetylases required for muramic-delta-lactam production are

involved in *Clostridium difficile* sporulation, germination, and heat resistance. *J Biol*

Chem **293**, 18040-18054, doi:10.1074/jbc.RA118.004273 (2018).

65 Traag, B. A., Pugliese, A., Eisen, J. A. & Losick, R. Gene conservation among

endospore-forming bacteria reveals additional sporulation genes in *Bacillus subtilis*. *J*

Bacteriol **195**, 253-260, doi:10.1128/JB.01778-12 (2013).

66 Perez, J., Springthorpe, V. S. & Sattar, S. A. Clospore: a liquid medium for producing

high titers of semi-purified spores of *Clostridium difficile*. *J AOAC Int* **94**, 618-626

(2011).

67 Verduyn, C., Postma, E., Scheffers, W. A. & Van Dijken, J. P. Effect of benzoic acid on

metabolic fluxes in yeasts: a continuous-culture study on the regulation of respiration and

alcoholic fermentation. *Yeast* **8**, 501-517, doi:10.1002/yea.320080703 (1992).

68 Egorova-Zachernyuk, T. A., Bosman, G. J., Pistorius, A. M. & DeGrip, W. J. Production

of yeastolates for uniform stable isotope labelling in eukaryotic cell culture. *Appl*

Microbiol Biotechnol **84**, 575-581, doi:10.1007/s00253-009-2063-z (2009).

69 Opitz, C., Isogai, S. & Grzesiek, S. An economic approach to efficient isotope labeling in

insect cells using homemade ^{15}N-, ^{13}C- and ^{2}H-labeled yeast extracts. *J Biomol NMR* **62**,

373-385, doi:10.1007/s10858-015-9954-3 (2015).

70 Vizcaino, J. A. *et al.* ProteomeXchange provides globally coordinated proteomics data

submission and dissemination. *Nat Biotechnol* **32**, 223-226, doi:10.1038/nbt.2839 (2014).

71 Huang da, W., Sherman, B. T. & Lempicki, R. A. Systematic and integrative analysis of

large gene lists using DAVID bioinformatics resources. *Nat Protoc* **4**, 44-57,

doi:10.1038/nprot.2008.211 (2009).

72 Paley, S. M. & Karp, P. D. The Pathway Tools cellular overview diagram and Omics

Viewer. *Nucleic Acids Res* **34**, 3771-3778, doi:10.1093/nar/gkl334 (2006).

Figures and Tables

Fig 1. Preparation workflow of (A) ^{15}N-labelled vegetative cells and (B) ^{14}N spores of

Peptoclostridium difficile **630.** See the Materials and Methods section for more details.

Fig 2. **Distribution of proteins in *Peptoclostridium difficile* 630 spores and vegetative cells.**

MASCOT score indicates the combined spore and cell abundance of a protein versus its
light/heavy protein isotopic ratio, which represents the relative level of the protein in spores over
vegetative cells. The orange dots indicate spore-predominant proteins (light/heavy ratios > 20),
blue dots indicate vegetative cell-predominant proteins (light/heavy ratios < 0.05), and white dots
indicate proteins common between spores and vegetative cells (20> light/heavy ratios >0.05).
Black arrow - SspA; green arrow- CD2657; red arrow - SlpA; purple arrow- CD0594; brown
arrow - CD0825; and yellow arrow - CD0718. (See the text and **Table S1** for more details).

Fig 3. Classification of proteins associated with the Wood-Ljungdahl pathway identified in

***Peptoclostridium difficile* 630.** Proteins presented in red, green, and blue fall under the

categories of vegetative cell-predominant, spore-predominant, and shared proteins, respectively.

Table 1. Uniprot keywords annotation enrichment of quantified *Peptoclostridium difficile*

630 spore- and vegetative cell proteins based on DAVID functional annotation analysis.

UniProt Keyword[a]	No. of proteins		
	Spore proteome	Vegetative cell proteome	Commonly shared
Aminotransferase		10	
Arginine biosynthesis		7	
Cell shape		8	
Elongation factor		5	
Peptidoglycan synthesis		7	
Cytoplasm	130	136	127
Transferase	115	121	110
Nucleotide-binding	107	114	105
Hydrolase	107	104	94
ATP-binding	89	95	87
Metal-binding	82	80	78
Oxidoreductase	61	65	54
Ribonucleoprotein	48	48	48
RNA-binding	49	48	48
Ligase	46	48	46
Ribosomal protein	47	47	47
Protein biosynthesis	34	36	34
Lyase	31	36	28

Magnesium	33	33	33
rRNA-binding	32	32	32
Zinc	30	28	28
Amino-acid biosynthesis	22	27	22
Isomerase	25	26	24
Aminoacyl-tRNA synthetase	23	24	23
Protease	25	20	20
GTP-binding	15	16	15
Pyridoxal phosphate	13	14	12
Glycosyltransferase	14	13	13
Cell cycle	11	13	11
Cell division	11	13	11
Flavoprotein	14	12	11
FAD	11	10	9
NADP	11	12	11
tRNA-binding	11	11	11
NAD	12	11	10
Ion transport	10	11	10
Pyruvate	11	11	11
Chaperone	10	10	10
Manganese	10	9	9
ATP synthesis	9	9	9
Aminopeptidase	9	9	9

Lysine biosynthesis	8	8	8
Hydrogen ion transport	8	8	8
Glycolysis	8	7	7
Stress response	7	7	7
Diaminopimelate biosynthesis	6	6	6
Pyrimidine biosynthesis	6	6	6
CF(1)	5	5	5
One-carbon metabolism	5	5	5
DNA-directed RNA polymerase	4	4	4
Methionine biosynthesis	4	4	4
Virion	10		
Capsid protein	9		
Rotamase	5		
Metalloprotease	6		

632   [a] EASE score i.e. p-value threshold for the keyword annotation enrichment was set to 0.05

Supplemental Material

Supplementary Figure S1. Cellular overview of quantified proteins from spores and vegetative cells of *Peptoclostridium difficile* 630. The quantified proteins that are spore-predominant (red), commonly shared but still higher in spores (purple), commonly shared but higher in cells (orange) and cell-predominant proteins (green) are represented with the pathways to which they belong. Refer to **Supplementary Table S1** for the details.

Supplementary Table S1. Proteins identified and quantified from *Peptoclostridium difficile* 630 vegetative cells and spores. Score (AA): Arithmetic average of Mascot scores from three replicates; SEM (AA): Standard Error of Means of arithmetic averages; L/H (GM): Geometric means of Light/Heavy ratios from three replicates; SEM (AA): Standard error of means in arithmetic averages of Light/Heavy ratios; SD (GM): Geometric standard deviation in Light/Heavy ratios; Proteins identified only in a single replicate are shown in red; Proteins encoded by essential genes are shown in blue.

Supplementary Table S2. Predicted membrane proteins from *Peptoclostridium difficile* 630 vegetative cells and spores. All the default parameters were used for TMHMM predictions.