Rapid, ultra-local adaptation facilitated by phenotypic plasticity

Syuan-Jyun Sun[1*], Andrew M. Catherall[1], Sonia Pascoal[1], Benjamin J. M. Jarrett[1,2], Sara E. Miller[3], Michael J. Sheehan[3], Rebecca M. Kilner[1]

[1] Department of Zoology, University of Cambridge, Downing Street, Cambridge, CB2 3EJ, UK

[2] Department of Entomology, Michigan State University, 288 Farm Lane Room 243, East Lansing, MI 48824, USA

[3] Department of Neurobiology and Behavior, Cornell University, 215 Tower Road, Ithaca, NY 14853, USA

*Corresponding author: Syuan-Jyun Sun,

Phone: 01223 (3)34466

e-mail: sjs243@cam.ac.uk

Keywords: interspecific competition, phenotypic plasticity, plasticity-led evolution, burying beetles, *Nicrophorus vespilloides,* niche expansion, local adaptation.

Abstract

Models of 'plasticity-first' evolution are attractive because they explain the rapid

evolution of new complex adaptations. Nevertheless, it is unclear whether plasticity can

still facilitate rapid evolution when diverging populations are connected by gene flow.

Here we show how plasticity has generated adaptive divergence in fecundity in wild

populations of burying beetles *Nicrophorus vespilloides*, which are still connected by

gene flow, which occupy distinct Cambridgeshire woodlands that are just 2.5km apart

and which diverged from a common ancestral population c. 1000-4000 years ago. We

show that adaptive divergence is due to the coupling of an evolved increase in the

elevation of the reaction norm linking clutch size to carrion size (i.e. genetic

accommodation) with plastic secondary elimination of surplus offspring. Working in

combination, these two processes have facilitated rapid adaptation to fine-scale

environmental differences, despite ongoing gene flow.

Introduction

Understanding the scale and pace of local adaptation is a long-standing problem in

evolutionary biology that has acquired new significance for predicting how natural

populations will respond to man-made environmental change[1–6]. In theory, phenotypic

plasticity can potentially both accelerate the pace of evolution, and fine-tune the scale of local

adaptation, through two distinct processes. The extent of phenotypic plasticity in any trait is

described by a reaction norm, which relates environmental variation to the phenotype it

induces. 'Plasticity-first' evolution can speed up the pace of evolutionary change in complex

traits because the shape, slope and elevation of a reaction norm each have genetic

components, upon which selection can act[1–6]. Recent experimental work on natural

populations has shown that selection in a new environment can greatly reduce the slope of the

reaction norm, causing the constitutive expression of novel, canalised traits[7,8]. New

adaptations can therefore evolve through the 'genetic assimilation' (*sensu*[9]) of traits that were

once induced environmentally.

Genetic assimilation yields rapid change but is predicted to occur only when there is

little or no gene flow between populations that inhabit different local environments[10–12].

While gene flow still persists, theory predicts that selection will instead act to maintain

plasticity in adaptive traits[11,12]. Plasticity is adaptive under these conditions because it

'forgives' costly mistakes incurred through the expression of locally maladaptive traits.

Nevertheless, local adaptation through genetic change is still possible, if selection fine-tunes

the slope or elevation of the reaction norm within each population to match local conditions

through genetic accommodation[13,14]. However, whether this mechanism is at work in natural

animal populations is unclear (but see[15] for evidence from a plant species).

We tested whether plasticity has facilitated recent, ultra-local adaptation in wild

populations of burying beetles *Nicrophorus vespilloides*. We focused on populations

occupying Gamlingay Wood and Waresley Wood in Cambridgeshire, UK, which are c. 2.5

km apart. We have previously shown that the two beetle populations cannot be differentiated

at neutral genetic markers, indicating there is ongoing gene flow[16]. Gamlingay Wood and

Waresley Wood have existed as distinct woodlands since at least 1086 because they are both

recorded in the Domesday Book (a land survey commissioned by William the Conqueror).

However, until 3000-4000 years they were almost certainly connected as part of the 'Wild

Wood', the ancient forest that once covered England but which was deforested from the

Bronze Age onwards[17].

Burying beetles (*Nicrophorus* spp.) breed on small dead vertebrates, such as rodents,

and it is common for several *Nicrophorus* species to co-exist within the same woodland.

Competition for carrion partitions the carrion niche so that larger burying beetle species tend

to breed on larger carrion, while the smallest species (*N. vespilloides* in the UK) typically

uses the smallest carrion[18–21]. Burying beetles match the number of eggs they lay to the size

of the carcass they obtain[22], and facultatively cull any surplus larvae through partial filial

cannibalism[23]. The adaptive clutch size and brood size therefore depend on the size of the

carrion that is routinely available for reproduction.

Here we show that the burying beetle guilds differ between the Gamlingay and Waresley

Woods, and that this changes the range in carrion size available for *N. vespilloides* to breed

upon. We demonstrate corresponding adaptive divergence in levels of fecundity shown by *N.*

vespilloides from each population, and show how plasticity has facilitated such rapid adaptive

divergence on such an ultra-local scale.

Results

The Nicrophorus guild differs between Gamlingay and Waresley Woods

In general, we found that the two woodlands harboured a similar number of *Nicrophorus*

beetles: we caught a total of 1873 *Nicrophorus* individuals in Gamlingay Wood over the

four-year sampling period compared with 1806 *Nicrophorus* individuals in Waresley Wood.

However, whereas Gamlingay Wood routinely supports four *Nicrophorus* species, Waresley

Wood is routinely inhabited by only two species (Fig. 1a). The average abundance per trap of

each *Nicrophorus* species differed between the two woods (species x woodland interaction: χ^2

= 142.68, d.f. = 4, $p < 0.001$). *N. vespilloides* was by far the most abundant in each woodland,

comprising 80.6% (1510 individuals) of all *Nicrophorus* beetles trapped in Gamlingay Wood

and 93.9% (1695 individuals) of those trapped in Waresley Wood. Both sites also contained

stable populations of the largest species, *N. humator* (Fig. 1b; Supplementary Tables 1 and

Supplementary Tables 2). Only Gamlingay Wood contained populations of intermediate-

sized *N. interruptus* and *N. investigator* in all four years of the study (Fig. 1b; Supplementary

Tables 1 and Supplementary Tables 2), and in significantly greater abundance in Waresley

Woods (Tukey HSD, $z = 7.90$, $p < 0.001$ and $z = 5.80$, $p < 0.001$, respectively). The

Nicrophorus guild is therefore significantly different between the two woodlands

(PERMANOVA (permutational multivariate analysis of variance) test $F = 0.024$, $p < 0.001$;

Supplementary Fig. 1b).

No evidence for divergence in the small mammal population between Gamlingay and

Waresley Woods

We sampled the small mammal population in Gamlingay and Waresley Woods to estimate

the abundance and type of carrion that might be available to the burying beetles to breed

upon (see Methods). In Gamlingay, 32 animals were caught across 50 trap sessions (23 new

catches and 9 recaptures); in Waresley 41 animals were caught across 50 trap sessions (30

new catches and 11 recaptures). Across both woods, bank voles (*Myodes glareolus*; range:

15-40 g) and wood mice (*Apodemus sylvaticus*; range: 13-27 g) were the dominant species,

constituting 53% and 43% of all trapped mammals respectively. There was no difference in

the mean body mass of small mammals sampled between the two sites ($\chi^2 = 0.19$, d.f. = 1, $p =$

0.661; Supplementary Fig. 2).

The Gamlingay guild is 'ancestral', whereas the Waresley guild is derived

We conclude from these data that there are approximately similar numbers of *Nicrophorus*

beetles within each woodland competing for an approximately similar size, abundance and

type of rodent carrion to breed upon. The key difference lies in the number of burying beetle

species in each wood.

We further infer from the ecological data that the more speciose burying beetle guild in

Gamlingay Wood more closely approximates the ancestral burying beetle guild that was

present in the 'Wild Wood', whereas the guild in Waresley Wood represents a more recent,

derived condition. This is because burying beetle guilds in pristine ancient forests in North

America are more speciose than those present in degraded woodlands[24], in keeping with the

general observation that there is a positive relationship between habitat size and species

richness[25]. Furthermore, smaller-bodied generalist carrion beetles are more likely to survive

in fragmented forests[24].

Niche expansion by N. vespilloides beetles in Waresley Wood

The ecological data additionally suggest that in Gamlingay Wood, *N. vespilloides* is more

likely to be confined to breeding only on smaller carrion. In general, larger *Nicrophorus*

species appear to be under selection to breed on larger carcasses[26,27], forcing smaller beetles

to breed on smaller carrion. Accordingly, *N. vespilloides* is more than twice as likely to be

found on small carcasses than on large carcasses in continental forests, which are rich in

Nicrophorus species[21]. In Waresley Wood, by contrast, *N. vespilloides* is more likely to breed

on larger carrion as well because it more rarely faces competition from *N. interruptus* and *N.*

investigator for larger carcasses (woodland effect: $\chi^2 = 27.75$, d.f. = 1, $p < 0.001$; Fig. 1c; see

also ref. [18]).

By measuring reproductive performance in the laboratory, we determined the carrion

niche occupied by the following species, in decreasing order of size (Fig. 1b): *N. investigator*

(from Gamlingay Wood), *N. interruptus* (from Gamlingay Wood), and *N. vespilloides* beetles

(from both Gamlingay and Waresley Woods) (see Methods). For each species, we

experimentally varied the size of carrion available for reproduction, presenting pairs with

either a small mouse carcass (range: 12-20 g; mean ± S.E.M: 16.80 ± 0.55 g) or a large

mouse carcass (range: 25-31 g; mean ± S.E.M: 28.27 ± 0.53 g; natural carcass range = 8.5-41

g). To quantify reproductive performance on each carcass size, we measured 'carcass use

efficiency', calculated by dividing the total brood mass at the end of larval development

(measured when larvae had dispersed from the carcass) by the size of the carcass the brood

was reared on. We predicted that if there is local adaptation, *N. investigator* and *N.*

interruptus should each exhibit greatest efficiency when breeding on a large carcass, while *N.*

vespilloides from Gamlingay Wood should exhibit greatest efficiency when breeding on a

small carcass. *N. vespilloides* from Waresley Wood should be more efficient at breeding on a

large carcass than *N. vespilloides* from Gamlingay Wood.

We found that the efficiency of converting the carcass into larvae varied with carcass

size, but in a different way for each *Nicrophorus* species (carcass size x *Nicrophorus* species

interaction term: $\chi^2 = 28.85$, d.f. = 3, $p < 0.001$; Fig. 2 and Supplementary Table 3). *N.*

investigator exhibited greater efficiency when breeding on large carcasses rather than on

small carcasses ($t = 3.51$, $p < 0.001$). *N. interruptus* was similarly efficient when breeding on

both large and small carcasses ($t = 0.99$, $p = 0.325$). These species are therefore adapted to

breed on larger carrion. In contrast, *N. vespilloides* from Gamlingay Wood exhibited greatest

reproductive efficiency on small carcasses ($t = -4.16$, $p < 0.001$). Waresley beetles were

similarly efficient to Gamlingay beetles when breeding on a small carcass ($t = 0.58$, $p =$

0.938) but were significantly more efficient at breeding on a large carcass than *N.*

vespilloides from Gamlingay Wood ($t = -3.11$, $p = 0.011$). We conclude that the population of

N. vespilloides in Waresley Wood has adaptively expanded its carrion niche to breed on

larger carcasses, in response to the loss of competition for larger carrion from *N. interruptus*

and *N. investigator*.

Niche expansion by Waresley N. vespilloides is due to divergent reaction norms

To test whether niche expansion by Waresley *N. vespilloides* was facilitated by genetic

accommodation, we generated reaction norms relating carcass size to clutch size for *N.*

vespilloides from the two woodland populations (see Methods). From the results in Fig. 2, we

predicted that the reaction norm for Waresley *N. vespilloides* would be significantly steeper

than the reaction norm for Gamlingay *N. vespilloides*. In fact, we found that the reaction

norms differed in their intercept rather than in their slope (Fig. 3a). Although female *N.*

vespilloides from both woodlands laid more eggs when given a larger carcass to breed upon

(carcass size effect: $\chi^2 = 10.97$, d.f. $= 1$, $p < 0.001$; Supplementary Table 4), Waresley *N.*

vespilloides consistently laid more eggs than Gamlingay *N. vespilloides* (woodland effect: χ^2

$= 21.07$, d.f. $= 1$, $p < 0.001$; Supplementary Table 4), irrespective of carcass size (carcass size

x woodland interaction: $\chi^2 = 1.44$, d.f. $= 1$, $p = 0.231$) or female size (woodland x female size

interaction: $\chi^2 = 1.30$, d.f. $= 1$, $p = 0.254$).

Likewise, although female *N. vespilloides* from both woodlands laid smaller eggs when

given a larger carcass to breed upon (carcass size effect: $\chi^2 = 5.90$, d.f. $= 1$, $p = 0.015$; Fig.

3b), Waresley *N. vespilloides* consistently laid smaller eggs than Gamlingay *N. vespilloides*

(woodland effect: $\chi^2 = 4.59$, d.f. $= 1$, $p = 0.032$), irrespective of carcass size (carcass size x

woodland interaction: $\chi^2 = 0.33$, d.f. $= 1$, $p = 0.565$). Mean egg volume per clutch did not

predict mean larval mass per brood at dispersal ($\chi^2 = 0.28$, d.f. $= 1$, $p = 0.595$), regardless of

the population of origin. This suggests that any under-provisioning of eggs is compensated by

the over-abundance of resources available on the carcass after hatching[28,29].

Genetic accommodation is facilitated by plastic secondary culling of surplus offspring

When we related brood size (measured at the end of reproduction) to carcass size, we found

the predicted difference between Gamlingay and Waresley *N. vespilloides* in the slope of this

reaction norm (woodland x carcass size interaction: $\chi^2 = 4.80$, d.f. = 1, $p = 0.029$; Fig. 4;

Supplementary Table 4). On a large carcass, Waresley *N. vespilloides* produced more larvae

than Gamlingay *N. vespilloides* ($z = -1.97$, $p = 0.049$), whereas on a small carcass the number

of larvae produced did not differ between the two populations ($z = 0.76$, $p = 0.447$).

Therefore Waresley *N. vespilloides* lays more eggs on a small carcass than Gamlingay *N.*

vespilloides, but there is plastic culling of surplus offspring after egg-laying, probably due to

partial filial cannibalism[23].

In short, Waresley *N. vespilloides* has evolved the ability to lay extra eggs whatever the

size of the carcass it locates for reproduction. This brings fitness gains if the beetle locates a

large carcass and correspondingly increases the efficiency with it converts carrion into

offspring. Yet it brings fitness costs if the beetle instead finds a small carcass to breed upon,

because there is a pronounced trade-off between larval number and larval size[30] and small

larvae have markedly lower fitness[31]. However, with a secondary mechanism for dispensing

with excess young on smaller carcasses (probably filial cannibalism), these costs are

eliminated. We suggest that genetic accommodation of the reaction norm relating clutch size

to carrion has evolved and is adaptive only because there is an accompanying plastic

mechanism for correcting the over-production of offspring on a small carcass. The same

mechanism can also prevent costly excess fecundity if there is gene flow from Waresley to

Gamlingay.

Divergence at loci associated with oogenesis in N. vespilloides from Gamlingay v. Waresley

Woods

To be certain that differences in the elevation of the reaction norm were due to evolutionary

change, we sought evidence of associated genetic divergence. We generated low-coverage

whole genome sequences for 40 individuals collected from each wood ($n = 80$ chromosomes

per population). We found minimal divergence across the genome with no instances of

extreme outliers (Supplementary Fig. 4), probably because the divergent traits are controlled

by many loci. This is typical for behavioural and life history traits and consistent with the

predicted quantitative genetic basis of genetic accommodation[6]. The highest *Fst*-windows in

the genome showed only modest absolute values of divergence. Nevertheless, they were

extreme outliers due to the otherwise consistently low pattern of *Fst* between populations (see

Methods).

By comparing differentiation between both woodlands and a third population from

300km away in Swansea, Wales, UK, we polarised the divergence between populations. We

calculated all pairwise *Fst* and assigned the relative divergence of each population using

population branch statistics (PBS) to understand which population was driving divergence

across the genome[32,33]. Not surprisingly, the distant Welsh population showed the highest

genome-wide PBS. We found that population from Waresley Wood showed higher

differentiation compared to the population from Gamlingay Wood (mean PBS: Waresley =

0.0074, Gamlingay = 0.0056, Wales = 0.0109). These analyses therefore support the

ecological data in indicating that the Gamlingay *N. vespilloides* represents the ancestral

condition, whereas the Waresley *N. vespilloides* are the derived population (Supplementary

Fig. 5).

To visualise the relative divergence between populations across the genome, we

generated a scatterplot of the PBS values for 2kb non-overlapping windows for each

population (Fig. 5a). The analysis highlighted multiple potential candidate genes associated

with the differences in egg-laying behaviour, again consistent with this trait being controlled

by many loci of small effect. For example, homologs of three of the highly differentiated

genes in the Waresley Wood population – *obg-like ATPase*, *transmembrane protein 214*, and

liprin-alpha – show elevated expression in the ovaries of fruit flies[34] and other arthropods[35],

suggesting a plausible role in regulating egg production. Another gene, *kekkon1*, is a

transmembrane protein known to regulate the activity of the epidermal growth factor receptor

(EGFR) during oogenesis in *Drosophila*[36,37].

Finally, we asked whether genes involved in oogenesis generally showed elevated levels

of differentiation in each population, in comparison with the rest of the genome. For each

population, we ranked genes by the highest PBS score in 500bp windows overlapping with

the gene-body and conducted a gene set enrichment analysis for each population. *N.*

vespilloides from both Waresley and Gamlingay Woods showed enrichment in multiple GO

terms associated with ovaries and oogenesis (Table 1).

Discussion

Our analyses support the three principle criteria required to show evidence for plasticity-first

evolution[6]. We show that the ancestral reaction norm linking carrion size to clutch size has

evolved to have a greater intercept in *N. vespilloides* from Waresley Wood. This change has

enabled Waresley N. *vespilloides* beetles to expand their carrion niche in the absence of rival

congenerics. It is adaptive because Waresley *N. vespilloides* are now more effective than

Gamlingay *N. vespilloides* at converting carrion that ranges widely in size into offspring.

Whereas previous analyses of plasticity-first evolution in wild animal populations have

emphasised how new adaptations can evolve from the genetic assimilation of plastic traits[6–

8,38], our study is different in showing how adaptive traits in nature can also result from

genetic accommodation.

Importantly, plasticity not only precedes the evolution of new local adaptations but also

provides a secondary correcting mechanism that ameliorates the costs of expressing these

new adaptations in the wrong local environment. Indeed, the presence of a plastic secondary

correcting mechanism is key to explaining how there has been genetic accommodation of the

reaction norm in *N. vespilloides* from Waresley Woods, despite gene flow with *N.*

vespilloides from Gamlingay Woods. We have demonstrated that when evolution by genetic

accommodation in one trait is coupled with a second plastically expressed correcting

mechanism, then new adaptations can emerge rapidly and on an ultra-fine scale. We suggest

that whenever these two mechanisms operate together, in two related traits, there is immense

potential for rapid evolutionary change that is finely tuned to local environmental conditions.

Methods

The history of Gamlingay and Waresley Woods

We focused on two woodlands: Gamlingay Wood (Latitude: 52.15555°; Longitude:

−0.19286°), and Waresley Wood (Latitude: 52.17487°; Longitude: −0.17354°). They are

woodland islands of approximately the same size (c 50ha) in a landscape dominated by arable

farming (Supplementary Fig. 1a). In common with other woodlands recorded in the

Domesday book, these woods have stayed approximately the same size since 1086[39].

Gamlingay Wood was acquired and managed by Merton College, Oxford for c. 800 years[40].

Its ecology was described in detail in 1912[40]. The modern history of Waresley Wood is less

well-known[39,41]. Both sites are now designated as 'ancient woodland' and are managed by the

Bedfordshire, Cambridgeshire and Northamptonshire Wildlife Trusts.

Competition for carrion in Gamlingay and Waresley Woods

Burying beetle trapping

Each year from 2014-2017 we set five beetle traps per woodland, at exactly the same five

locations within each wood (Supplementary Fig. 1a), and checked the contents every 2-3

weeks from June until October, each time rebaiting the trap with fresh compost and a dead

mouse. Carrion-baited soil-filled traps were suspended at each site, with traps set at least 150

m apart from each other. We collected and identifed all the *Nicrophorus* spp. within each trap

and measured the pronotum width (to the nearest 0.01 mm) as an index of body size (see

below).

We caught five species in total (in increasing order of size): *N. vespilloides*, *N.*

interruptus, *N. vespillo*, *N. investigator* and *N. humator*. However, *N. vespillo* were only

caught 5 and 3 times within the four years in Gamlingay and Waresley respectively,

indicating that there is no stable population of *N. vespillo* in either wood. A clear separation

of the guild structure between the woods can be seen for each sampling time across the four

years in NMDS (nonmetric multidimensional scaling) ordination (Supplementary Fig. 1b).

Evidence from other populations suggests that our measurements reflect long-term

differences in guild structure because abundance measures of *Nicrophorus* are robust over

time, even when there are marked change in woodland management[40].

Burying beetle husbandry in the lab

After removing any phoretic mites, beetles were retained and kept individually in plastic

boxes (12cm x 8cm x 2cm), which were filled with moist soil in a laboratory kept at 20°C

and on a 16:8 light to dark cycle. Beetles were fed twice a week with minced beef. We kept

all field-caught individuals for at least two weeks before breeding to ensure that they were

sexually mature and to reduce any variation in nutritional status. We then maintained stock

populations of both Gamlingay and Waresley Woods by breeding pairs of unrelated

individuals on 8-16 g mice carcasses.

Size distributions of the Nicrophorus spp.

Body size was measured for *Nicrophorus* spp. collected from 2014 to 2017 (except in 2016).

In total, 838 *N. vespilloides*, 41 *N. humator*, 127 *N. interruptus*, 54 *N. investigator*, and 5 *N.*

vespillo were measured for Gamlingay Wood, whereas 824 *N. vespilloides*, 51 *N. humator*,

25 *N. interruptus*, 4 *N. investigator*, and 6 *N. vespillo* were measured for Waresley Wood.

Mean body size of *Nicrophorus* spp. significantly varied among species (GLMM: χ^2 =

2069.44, d.f. = 4, *p* < 0.001; Supplementary Table 2). Post-hoc comparisons revealed that *N.*

vespilloides was smaller than the other *Nicrophorus* spp. (Supplementary Table 1 and

Supplementary Table 2). A Kolmogorov-Smirnov test comparing pairwise differences in

Nicrophorus spp. body size frequency distribution revealed similar patterns found in

differences of mean body size (Fig. 1 and Supplementary Table 5).

Mark-recapture experiment

In 2014, we investigated the interconnectivity of populations of burying beetles between

Gamlingay and Waresley Woods using a mark-recapture survey. In total, 98 *N. vespilloides*,

9 *N. humator*, 17 *N. interruptus*, 9 *N. investigator,* and 2 *N. vespillo* were marked for

Gamlingay Wood, whereas 113 *N. vespilloides*, 5 *N. humator*, 1 *N. interruptus*, 1 *N.*

investigator were marked for Waresley Wood. Beetles were marked with a numbered plastic

bee tag on either the right or left elytra for those found in Gamlingay or Waresley Woods

respectively. To identify any previously caught beetles that lost their tags, we permanently

marked them by cutting a small portion of the elytra (less than 2%). All marked beetles from

each wood were released from a designated location at the geographic midpoint with the

minimum total distance to all trapping sites for Gamlingay (Latitude: 52.16294°; Longitude:

−0.18984°) and Waresley Woods (Latitude: 52.176508°; Longitude: −0.156776°). We found

no evidence of migration between the two sites. We recaught 8 of 98 marked *N. vespilloides*

from Gamlingay Wood in Gamlingay Wood and 8 out of 113 marked *N. vespilloides* from

Waresley Wood in Waresley Wood. None of the other marked *Nicrophorus* spp. was

recaptured in either wood.

Small mammal trapping

To assess the rodent carrion available for *Nicrophorus* spp. reproduction, we sampled the

small mammal communities in the two woodlands. In general, rodent populations peak in the

autumn, because breeding for the year has just ceased and there has yet to be any winter-

induced mortality[42]. Sampling at this time is therefore ideal for detecting which species are

present and for determining their relative abundance. We placed Longworth traps in both

woodlands in November 2016. Traps were baited with oats and blowfly maggots (with hay

provided as bedding) and set in pairs within 20 m of each original beetle trapping site

(Supplementary Fig. 1a), with 10 traps set per wood. We continuously trapped rodents for

three days, generating 50 trap sessions per woodland. Traps were checked daily at

approximately 0830 and 1500 (generating a total of 30 trap sessions overnight and 20 trap

sessions in daylight hours). Trapped mammals were identified, weighed, sexed, marked by a

fur clip on either the right or left rear flank, and released *in situ*. Any recaptured mammal was

recorded in subsequent censuses. All traps were reset and rebaited immediately after

checking. In addition to the results reported in the main text, one yellow-necked mouse

(*Apodemus flavicollis*; range: 14-45 g) and one common shrew (*Sorex araneus*; range: 5-14

g) were caught in Waresley. We have no reason to think that the mortality of these rodent

species should differ between woodlands that are in such close proximity and that are

subjected to similar levels of ecological management.

Division of the carrion niche by Nicrophorus beetles in Gamlingay and Waresley Woods

During the field seasons in 2017 and 2018, pairs of wild-caught *N. vespilloides*, *N.*

interruptus, and *N. investigator* were bred on either small (12-20 g; 16.80 ± 0.55 g) or large

carcasses (25-31 g; 28.27 ± 0.53 g) within a breeding box (17 x 12 x 6 cm) filled with 2 cm

of moist soil. All field-caught beetles were kept for two weeks in the laboratory and fed twice

a week prior to breeding. In total, we established eight treatments: large (n = 47) and small (n

= 48) carcasses for Gamlingay *N. vespilloides*; large (n = 42) and small (n = 33) carcasses for

Waresley *N. vespilloides*; large (n = 25) and small (n = 18) carcasses for *N. interruptus*; large

(n = 9) and small (n = 13) carcasses for *N. investigator*. *N. interruptus* and *N. investigator*

were both drawn from Gamlingay Wood as the populations of these species in Waresley were

too small to be used experimentally.

Approximately eight days after parents are given a carcass to breed on, larvae switch

from aggregating on the carcass to dispersing away into the soil to pupate. When one or more

larvae from each brood switched their behaviour in this way, we scored the whole brood as

having reached the dispersal stage. At this point, all larvae were counted and total brood mass

was weighed to the nearest 0.001 g. We also calculated average larval mass for each brood by

dividing total brood mass by number of larvae.

In addition to the results reported in the main text, we found that Gamlingay *N.*

vespilloides was less efficient on larger carcasses than both *N. interruptus* (t = -3.92, p <

0.001) and *N. investigator* (t = -2.69, p = 0.038).

Niche expansion in N. vespilloides by genetic accommodation

This experiment was conducted in the laboratory, over two blocks in 2017, using the second

and third descendant generations of field-caught beetles from Gamlingay and Waresley

Woods. By rearing beetles from both woodlands in the lab in a common garden environment

for at least one generation prior to testing, we minimized any residual environmental effects

when quantifying the reaction norm for each population. To pair beetles for reproduction, we

began by haphazardly casting broods into dyads when larvae had matured into adults. Within

each dyad, we haphazardly chose four males from one brood, and paired them with four

haphazardly chosen females from the second brood. Two of these pairs were then given a

small mouse to breed upon (12-17 g; 15.03 ± 0.67 g), while the remaining two pairs were

given a large mouse to breed upon (26-31 g; 28.86 ± 0.67 g). By using sibships to generate

pairs in this way, we were able to compare how very similar genotypes responded to the

opportunity to breed on either a small or large mouse. Each pair, and their mouse, was housed

in a clear plastic box (17 x 12 x 6 cm), with 2 cm depth of Miracle-Gro compost. The box

was placed in a dark cupboard for eight days after pairing the beetles, until larvae started to

disperse away from the carcass. In the second block of the experiment, we measured clutch

size. Fifty-six hours after we introduced the beetles to the carcass, we photographed the base

of each transparent breeding box. Using Digimizer ver. 5.1.0, we then counted the number of

visible eggs, and also measured the length (L) and width (w) of all eggs that were able to be

measured accurately (i.e. those that were fully visible and lying flat on the base of the box).

All counting and measuring of eggs was performed blind to the carcass size treatment and the

population from whence the breeding beetles came. Egg volume was then calculated using

the formula $V = 1/6 \cdot \pi \cdot w^2 \cdot L$, which assumes eggs to be a prolate spheroid (following ref. [43]).

In both blocks, we also measured the number of larvae present at dispersal and weighed the

whole brood.

Clutch size

In total, 2518 eggs were counted across 132 breeding boxes, of which 1633 could be

measured. No eggs could be seen in four of these boxes, and these were excluded from

further analysis (hence $n = 128$). The total number of eggs counted (observed clutch size),

and the number of eggs which could accurately be measured, each correlated positively with

brood size (number of eggs counted: $\chi^2 = 27.18$, d.f. = 1, $p < 0.001$; number of eggs

measured: $\chi^2 = 24.26$, d.f. = 1, $p < 0.001$), indicating these are accurate estimates of true

clutch size. The total volume of all eggs in a clutch did not differ between populations ($\chi^2 =$

2.16, d.f. = 1, $p = 0.141$; Supplementary Table 4) or carcass size treatments ($\chi^2 = 2.38$, d.f. =

1, $p = 0.123$; Supplementary Table 4).

Brood size

We found that when given a larger carcass for reproduction, *N. vespilloides* from Gamlingay

Wood produced only a marginally larger brood than when breeding on a small carcass ($z =$

1.82, $p = 0.068$). By contrast, Waresley *N. vespilloides* produced significantly larger broods

on larger carcasses than on smaller carcasses ($z = 5.53$, $p < 0.001$). Whether they bred on

small or large carcasses, the variance was similar for Gamlingay and Waresley *N.*

vespilloides in both brood size (Bartlett's test, $p = 0.615$) and carcass use efficiency

(Bartlett's test, $p = 0.943$).

Offspring size

Females produced heavier larvae at dispersal when breeding on a large carcass than on a

small carcass ($\chi^2 = 139.05$, d.f. = 1, $p < 0.001$; Supplementary Table 4), irrespective of their

woodland of origin ($\chi^2 = 1.06$, d.f. = 1, $p = 0.304$; Supplementary Table 4). We could find no

difference in the size of adult *N. vespilloides* trapped in Gamlingay and Waresley Woods, (χ^2

$= 2.02$, d.f. = 1, $p = 0.156$; Supplementary Fig. 3a), even after controlling for sex ($\chi^2 = 1.41$,

d.f. = 1, $p = 0.235$). We could not detect any differences in body size distribution between

Gamlingay and Waresley *N. vespilloides* ($D = 0.044$, $p = 0.410$; Supplementary Fig. 3b).

Thus, given an abundance of resources for reproduction, *N. vespilloides* allocates them to

producing more offspring rather than larger offspring.

Divergence at loci associated with oogenesis in N. vespilloides from Gamlingay v. Waresley

Woods

We generated low-coverage whole genome sequences for three populations of *N.*

vespilloides: from Waresley Wood, Gamlingay Wood and Swansea, Wales, UK. Twenty-two

N. vespilloides from three ancient woodlands near Swansea in Wales were trapped by Dr

Chris Cunningham in 2017. The three sites in Wales (Park Wood (Latitude: 51.57258°;

Longitude: −4.03096°); Clyne Valley Wood (Latitude: 51.61262°; Longitude: −4.02293°);

and Caswell Bay Wood (Latitude: 51.57258°; Longitude: −4.03096°)) are approximately 300

km away from our two study sites in Cambridgeshire.

DNA was individually extracted from beetle heads using the DNeasy Blood and Tissue

kit (Qiagen) and subsequently quantified and quality checked using NanoDrop and Qubit.

DNA was then shipped to Cornell University, where libraries were prepared using partial

reactions of a Nexterra kit by the Cornell Genomics Core. Libraries were subsequently

sequenced by Novogene (Davis, CA, USA) at an average depth of approximately 5x

coverage. Paired-end 550 bp insert libraries were prepared for each sample with the Nextera

library preparation kit. Libraries were sequenced using the Illumina HiSeq (Novogene, Davis,

CA) at an average coverage of 3.4X. Trimmomatic (v0.36) was used to removed adaptors and

poor-quality sequence. Trimmed reads were mapped to the *N. vespilloides* reference genome

using the Burrows-Wheeler Aligner (v0.7.13)[44]. SNPs were identified using Picard (v2.8.2)

and GATK (v3.6) HaplotypeCaller following best practice recommendations[45]. After

alignment, SNPs were hard filtered using the parameters: QD < 2.0 || SOR > 3.0 || FS > 200.

We used VCFtools to calculate population genetic statistics for each population. To examine

population structure, we generated a thinned VCF file with one SNP per 5kb and used Tassel

(version 5)[46] to calculate generate an MDS plot. For calculation of *Fst* and PBS values,

aligned bam files were analysed in ANGSD (v 0.911)[47], which is specifically designed for

analysis of low-coverage genome sequencing data. Data for this project is available at the

NCBI Sequence Read Archive under Bioproject PRJNA530213.

Genes were assigned gene ontology (GO) terms using the BLAST2GO workflow

(v5.1.1). In brief, gene identity was determined based upon BLAST searches to the

Arthropod or *Drosophila* non-redundant protein databases and protein domains were

identified based on matches to Interpro database. GO terms were assigned to each gene

model based upon mapping results. GO terms were filtered with the "Filter Annotation by

GO Taxa" option to remove GO terms that are incompatible for Arthropods.

Population genetic summary statistics were similar for both populations (mean of 2 kb

windows – Gamlingay Pi = 0.0055 ± 1.02e-5, Tajima's D = -0.70 ± 0.002; Waresley Pi =

0.0056 ± 1.0e-5, Tajima's D = -0.67 ± 0.002). Consistent with previous microsatellite

analyses[16], we found little to no genetic differentiation between populations from Gamlingay

and Waresley Woods (unweighted *Fst* = 0.0069; weighted *Fst* = 0.013), strongly suggesting

there is ongoing gene flow between the two populations.

Among the few loci that diverged between the populations, the top window of

divergence between the two populations fell in *transmembrane protein 214* (*Fst* = 0.11, *zFst*

= 19.2, *p* = 7.2e-82), which is highly expressed in the ovaries in *Drosophila melanogaster*,

and is therefore a potential candidate gene influencing differences in *N. vespilloides* egg

laying behaviour between Gamlingay and Waresley Woods.

PBS analysis identified further candidate genes. We found that a different portion of

kek1 is moderately differentiated in *N. vespilloides* from Gamlingay Wood (Fig. 5a),

suggesting that differentiation related to oogenesis is not limited to *N. vespilloides* from

Waresley Wood. One of the stronger signals of differentiation in Waresley Wood that may

contribute to the phenotypic differences observed between populations is found in a serotonin

receptor (Fig. 5b). Serotonin has been linked to reproduction via effects on the production of

ecdysteroids such as juvenile hormone in multiple insects[48–50]. Serotonin has also been

related to the intensity of aggressive behaviour in contests across diverse insect groups[51–54].

Consistent with the outlier analyses, oogenesis-related GO-terms tended to have higher

enrichment scores in *N. vespilloides* from Waresley Wood compared to Gamlingay Wood

(Table 1). The gene enrichment analysis also revealed local divergence between beetles from

Waresley and Gamlingay Woods at genes associated with other traits, including learning and

memory and sensory systems (Supplementary Data 1).

Statistical analyses

All statistical analyses were performed in R version 3.4.3 (R Development Core Team), with

NMDS in the package *vegan*, GLM and GLMM in the package *lme4*[55], and Tukey's HSD post

hoc comparisons in the package *lsmeans*[56].

Competition for carrion in Gamlingay and Waresley Woods: field data

To test for a difference in the *Nicrophorus* guild between Gamlingay and Waresley Woods, a

frequency table of beetle communities was analysed using permutational multivariate

analysis of variance (PERMANOVA) on two-dimensional nonmetric multidimensional

scaling (NMDS) based on Bray-Curtis distances[57], with the 'adonis' function (vegan

package). We tested the effect of study site on the composition of the beetle community,

using sampling year as strata. The analysis was based on 10000 permutations of the data. We

visualized the difference of beetle community between Gamlingay and Waresley population

in two dimensions of a NMDS plot. NMDS two-dimensional stress values (a measure of

goodness of fit) were below 0.1 (0.085), indicating the ordination provides a good fit to the

data[58].

We used general linear mixed models (GLMM) with a Poisson error structure to test for

differences across beetle species and sites on average abundance for each species per trap.

Beetle species and population (Gamlingay/Waresley) were included as fixed effects, whereas

trap ID and sampling year were included as random factors. We also tested for differences in

the probability of co-occurrence (co-occurring = 1, non-co-occurring = 0) between *N.*

vespilloides and at least one other *Nicrophorus* spp. across sites using a binomial GLMM.

Population was included as a fixed effect and year as a random effect.

We compare the distributions of body size between Gamlingay and Waresley *N.*

vespilloides, and among *Nicrophorus* spp., using the Kolmogorov-Smirnov (K-S) two-sample

test, which tests whether the cumulative distributions of two data sets are derived from the

same distribution. We also tested for significant differences in mean body size between *N.*

vespilloides in Gamlingay and Waresley, using a GLMM that included population

(Gamlingay/Waresley) and sex (male/female) as fixed effects, and sampling year as a random

factor. Differences of mean body size among *Nicrophorus* spp. was assessed in a GLMM that

included species and sex as fixed effects, and sampling year as a random factor.

We analysed the body mass of rodents in Gamlingay and Waresley using a GLM by

including rodent species and population (Gamlingay/Waresley) as fixed effects.

Division of the carrion niche by Nicrophorus beetles in Gamlingay and Waresley Woods

To test for differences in reproductive performance between species, we conducted a GLMM

regression to analyse differences in efficiency (total brood mass divided by carcass mass),

which was logit transformed prior to analysis. Beetle species, carcass size (small/large), and

their interaction were included as explanatory variables. Sampling year was included as a

random factor. In this analysis, we included beetle species as *N. interruptus*, *N. investigator*,

Gamlingay *N. vespilloides* and Waresley *N. vespilloides* to fully compare differences not only

between *Nicrophorus* beetle species, but also *N. vespilloides* between populations.

Niche expansion in N. vespilloides by genetic accommodation

For the reaction norm experiment, we used GLMMs to test the interacting effect of

population and carcass size on brood size and average larval mass, with dyad identity nested

within block included as a random factor. In all models, brood size and average larval mass

were analysed with a Poisson and Gaussian error distribution, respectively. A similar

statistical approach was used for analyses of clutch traits to test for the significant differences

on clutch size and average egg volume in GLMMs with a Poisson and Gaussian error

distribution, respectively. The effect of population of origin, carcass size, and their

interaction were included as fixed effects, whereas dyad identity was included as a random

factor. If a significant interaction was found, a Tukey's post-hoc test was performed to detect

significant effects using multiple pairwise comparisons.

Acknowledgements

We thank P. Pilbeam for carrying out small mammal trapping in the field. We thank A.

Attisano, C. Swannack, and R. Mashoodh for helping with beetle trapping. S.-J.S. was

supported by the Taiwan Cambridge Scholarship from the Cambridge Commonwealth,

European & International Trust. A.M.C. was supported by a studentship from the National

Environmental Research Council's Earth System Sciences Doctoral Training Partnership at

Cambridge University and a Victoria Brahm Schild Internship Grant from Homerton College.

R.M.K. was supported by a European Research Council Consolidator's grant 301785

BALDWINIAN_BEETLES and a Wolfson Merit Award from the Royal Society.

Author Contributions

R.M.K. conceived the idea. S.-J.S., A.M.C. S.P. and R.M.K designed the study. S.-J.S.,

A.M.C. and B.J.M.J. acquired the phenotypic data, which was analysed by S.-J.S. and

A.M.C. M.J.S. and S.P. designed the genetic's component, which was collected by S.P. and

analysed by M.J.S and S.E.M. All authors contributed to discussion of the results and writing

the manuscript.

Competing Interests Statement

The authors declare no competing interests.

References

1.   Baldwin, M. A new factor in evolution. *Am. Nat.* **30,** 441–451 (1896).

2.   Morgan, C. L. On modification and variation. *Science* **4,** 733–740 (1896).

3.   Robinson, B. W. & Dukas, R. The influence of phenotypic modifications on evolution:
the Baldwin effect and modern perspectives. *Oikos* **85,** 582 (1999).

4.   West-Eberhard, M. J. *Developmental plasticity and evolution*. (Oxford University
Press, 2003).

5.   Pfennig, D. W. *et al.* Phenotypic plasticity's impacts on diversification and speciation.
Trends Ecol. Evol. **25,** 459–467 (2010).

6.   Levis, N. A. & Pfennig, D. W. Evaluating 'plasticity-first' evolution in nature: key
criteria and empirical approaches. *Trends Ecol. Evol.* **31,** 563–574 (2016).

7.   Levis, N. A., Isdaner, A. J. & Pfennig, D. W. Morphological novelty emerges from
pre-existing phenotypic plasticity. *Nat. Ecol. Evol.* **2,** 1289–1297 (2018).

8.   Corl, A. *et al.* The genetic basis of adaptation following plastic changes in coloration
in a novel environment. *Curr. Biol.* **28,** 2970–2977.e7 (2018).

9.   Waddington, C. H. Genetic assimilation of an acquired character. *Evolution.* **7,** 118–

126 (1953).

10.   Lenormand, T. Gene flow and the limits to natural selection. *Trends Ecol. Evol.* **17,** 183–189 (2002).

11.   Sultan, S. E. & Spencer, H. G. Metapopulation structure favors plasticity over local adaptation. *Am. Nat.* **160,** 271–83 (2002).

12.   Crispo, E. Modifying effects of phenotypic plasticity on interactions among natural selection, adaptation and gene flow. *J. Evol. Biol.* **21,** 1460–1469 (2008).

13.   West-Eberhard, M. J. Developmental plasticity and the origin of species differences. *Proc. Natl. Acad. Sci. U. S. A.* **102 Suppl,** 6543–9 (2005).

14.   Thompson, J. N. *Relentless evolution*. (University of Chicago Press, 2013).

15.   Bock, D. G., Kantar, M. B., Caseys, C., Matthey-Doret, R. & Rieseberg, L. H. Evolution of invasiveness by genetic accommodation. *Nat. Ecol. Evol.* **2,** 991–999 (2018).

16.   Pascoal, S. & Kilner, R. M. Development and application of 14 microsatellite markers in the burying beetle *Nicrophorus vespilloides* reveals population genetic differentiation at local spatial scales. *PeerJ* **5,** e3278 (2017).

17.   Rackham, O. *The history of the countryside : the classic history of Britain's landscape, flora and fauna*. (Phoenix Press, 2000).

18.   Scott, M. P. The ecology and behavior of burying beetles. *Annu. Rev. Entomol* **43,** 595–618 (1998).

19.   Otronen, M. The effects of body size on the outcome of Fig.hts in burying beetles (*Nicrophorus*). *Ann. Zool. Fennici* **25,** 191–201 (1988).

20.   Safryn, S. A. & Scott, M. P. Sizing up the competition: do burying beetles weigh or measure their opponents? *J. Insect Behav.* **13,** 291–297 (2000).

21.   Urbański, A. & Baraniak, E. Differences in early seasonal activity of three burying beetle species (Coleoptera: Silphidae: *Nicrophorus* F.) in Poland. *Coleopt. Bull.* **69,** 283–292 (2015).

22. Müller, J. K., Eggert, A.-K. & Furlkröger, E. Clutch size regulation in the burying
beetle *Necrophorus vespilloides* Herbst (Coleoptera: Silphidae). *J. Insect Behav.* **3,**
265–270 (1990).

23. Bartlett, J. Filial cannibalism in burying beetles. *Behav. Ecol. Sociobiol.* **21,** 179–183
(1987).

24. Gibbs, J. P. & Stanton, E. J. Habitat fragmentation and arthropod community change:
carrion Beetles, phoretic mites, and flies. *Ecol. Appl.* **11,** 79 (2001).

25. Lomolino, M. V. Ecology's most general, yet protean 1 pattern: the species-area
relationship. *J. Biogeogr.* **27,** 17–26 (2000).

26. Smith, R. J. & Heese, B. Carcass selection in a high altitude population of the burying
beetle, *Nicrophorus investigator* (Silphidae). *Southwest. Nat.* **40,** 50–55 (1995).

27. Hopwood, P. E., Moore, A. J., Tregenza, T. & Royle, N. J. Niche variation and the
maintenance of variation in body size in a burying beetle. *Ecol. Entomol.* **41,** 96–104
(2016).

28. Russell, A. F., Langmore, N. E., Cockburn, A., Astheimer, L. B. & Kilner, R. M.
Reduced egg investment can conceal helper effects in cooperatively breeding birds.
Science **317,** 941–944 (2007).

29. Smiseth, P. T., Andrews, C. P., Mattey, S. N. & Mooney, R. Phenotypic variation in
resource acquisition influences trade-off between number and mass of offspring in a
burying beetle. *J. Zool.* **293,** 80–83 (2014).

30. Schrader, M., Jarrett, B. J. M. & Kilner, R. M. Parental care masks a density-
dependent shift from cooperation to competition among burying beetle larvae.
Evolution. **69,** 1077–1084 (2015).

31. Lock, J. E., Smiseth, P. T. & Moore, A. J. Selection, inheritance, and the evolution of
parent-offspring interactions. *Am. Nat.* **164,** 13–24 (2004).

32. Yi, X. *et al.* Sequencing of 50 human exomes reveals adaptation to high altitude.
Science **329,** 75–8 (2010).

33.   Fumagalli, M. *et al.* Greenlandic Inuit show genetic signatures of diet and climate
626   adaptation. *Science* **349,** 1343–1347 (2015).

627   34.   Chintapalli, V. R., Al Bratty, M., Korzekwa, D., Watson, D. G. & Dow, J. A. T.
628   Mapping an atlas of tissue-specific *Drosophila melanogaster* metabolomes by high
629   resolution mass spectrometry. *PLoS One* **8,** e78066 (2013).

630   35.   Cao, X. & Jiang, H. An analysis of 67 RNA-seq datasets from various tissues at
631   different stages of a model insect, *Manduca sexta*. *BMC Genomics* **18,** 796 (2017).

632   36.   Ghiglione, C. *et al.* The transmembrane molecule kekkon 1 acts in a feedback loop to
633   negatively regulate the activity of the *Drosophila* EGF receptor during oogenesis. *Cell*
634   **96,** 847–56 (1999).

635   37.   Wittes, J. & Schüpbach, T. A gene expression screen in *Drosophila melanogaster*
636   identifies novel JAK/STAT and EGFR targets during oogenesis. *G3 Genes, Genomes,*
637   *Genet.* **9,** 47–60 (2019).

638   38.   Scoville, A. G. & Pfrender, M. E. Phenotypic plasticity facilitates recurrent rapid
639   adaptation to introduced predators. *Proc. Natl. Acad. Sci. U. S. A.* **107,** 4260–3 (2010).

640   39.   Rackham, O. *Trees and woodland in the British landscape*. (Phoenix Giant, 1996).

641   40.   Adamson, R. S. An ecological study of a Cambridgeshire woodland. *J. Linn. Soc.*
642   *London, Bot.* **40,** 339–387 (1912).

643   41.   Darby, H. C. Domesday woodland. *Econ. Hist. Rev.* **3,** 21 (1950).

644   42.   Jackson, D. M., Trayhurn, P. & Speakman, J. R. Associations between energetics and
645   over-winter survival in the short-tailed field vole *Microtus agrestis*. *J. Anim. Ecol.* **70,**
646   633–640 (2001).

647   43.   Berrigan, D. The allometry of egg size and number in insects. *Oikos* **60,** 313 (1991).

648   44.   Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler
649   transform. *Bioinformatics* **25,** 1754–1760 (2009).

650   45.   Van der Auwera, G. A. *et al.* in *Current Protocols in Bioinformatics* **43,** 11.10.1-
651   11.10.33 (John Wiley & Sons, Inc., 2013).

652   46.   Bradbury, P. J. *et al.* TASSEL: software for association mapping of complex traits in
653   diverse samples. *Bioinformatics* **23,** 2633–2635 (2007).

654   47.   Korneliussen, T. S., Albrechtsen, A. & Nielsen, R. ANGSD: Analysis of Next
655   Generation Sequencing Data. *BMC Bioinformatics* **15,** 356 (2014).

656   48.   Woodring, J. & Hoffmann, K. H. The effects of octopamine, dopamine and serotonin
657   on juvenile hormone synthesis, In vitro, in the cricket, *Gryllus bimaculatus*. *J. Insect*
658   *Physiol.* **40,** 797–802 (1994).

659   49.   Shimada-Niwa, Y. & Niwa, R. Serotonergic neurons respond to nutrients and regulate
660   the timing of steroid hormone biosynthesis in *Drosophila*. *Nat. Commun.* **5,** 5778
661   (2014).

662   50.   Cook, C. N., Lawson, S. P., Brent, C. S. & Rehan, S. M. Biogenic amines shift during
663   the pre-reproductive to reproductive transition in the small carpenter bee, *Ceratina*
664   *calcarata*. *Apidologie* **50,** 90–99 (2019).

665   51.   Dierick, H. A. & Greenspan, R. J. Serotonin and neuropeptide F have opposite
666   modulatory effects on fly aggression. *Nat. Genet.* **39,** 678–682 (2007).

667   52.   Aleksandrowicz, O. & Komosiński, K. in *Protection of Coleoptera in the Baltic Sea*
668   *Region* (ed. J. Skłodowski, S. Huruk, A. Barševskis, S. T.) 147–153 (Warsaw
669   Agricultural University Press, 2005).

670   53.   Bubak, A. N., Renner, K. J. & Swallow, J. G. Heightened serotonin influences contest
671   outcome and enhances expression of high-intensity aggressive behaviors. *Behav. Brain*
672   *Res.* **259,** 137–142 (2014).

673   54.   Stevenson, P. A. & Rillich, J. Controlling the decision to Fig.ht or flee: the roles of
674   biogenic amines and nitric oxide in the cricket. *Curr. Zool.* **62,** 265–275 (2016).

675   55.   Bates, D., Maechler, M., Bolker, B. & Walker, S. Fitting linear mixed-effects models
676   using lme4. *R package version* 1.0-6 (2015).

677   56.   Lenth, R. V. Least-squares means: the R package lsmeans. *J. Stat. Softw.* **69,** 1–33
678   (2016).

679   57.   Clarke, K. R. Non-parametric multivariate analyses of changes in community

680   structure. *Aust. J. Ecol.* **18,** 117–143 (2016).

681   58.   Ramette, A. Multivariate analyses in microbial ecology. *FEMS Microbiol. Ecol.* **62,**

682   142–160 (2007).

683
684

685
686

Figure 1. **a**, Number of *Nicrophorus* beetles caught per trap from 2014-2017. **b**, Body size and frequency distribution of field-caught *Nicrophorus* spp., illustrated withbox-and-whisker plots (left) and kernel density estimation (right). Points indicate outliers. **c**, The proportion of traps set at which *N. vespilloides* was trapped with another *Nicrophorus* species, in the two woodlands. Error bars indicate mean ± S.E.M.

694
695

696
697 **Figure 2.** Efficiency (%) of carcass use (total brood mass divided by carcass mass) of *N.*
698 *interruptus*, *N. investigator*, and *N. vespilloides* from Gamlingay Wood and *N. vespilloides*
699 from Waresley Wood. Values represent the mean ± S.E.M.
700
701

702
703
704 **Figure 3.** The effect of carcass size on (**a**) clutch size and (**b**) average egg volume produced
705 by *N. vespilloides*. *n* = 27 Gamlingay *N. vespilloides* per carcass size treatment, and *n* = 37
706 Waresley *N. vespilloides* per carcass size treatment. Values represent the mean ± S.E.M.
707

708
709

710
711
712 **Figure 4.** The effect of carcass size on (**a**) brood size and (**b**) average larval mass for *N.*
713 *vespilloides* in the reaction norm experiment (*n* = 46 Gamlingay *N. vespilloides* per carcass
714 size treatment, and *n* = 62 Waresley *N. vespilloides* per carcass size treatment). Values
715 represent the mean ± S.E.M.
716
717

a

b

Figure 5. Differentiation at putative oogenesis genes. a, A scatterplot of PBS values for Waresley and Gamlingay in 2kb windows genome-wide. Loci in the lower right hand of the figure show high differentiation in Waresley but not in Gamlingay. Loci with PBS scores greater than 0.05 are highlighted – Waresley = blue, Gamlingay = red, Both = purple. Notable genes are annotated. **b**, Sliding window analysis (window = 500bp, slide = 100bp) of PBS values at the *5HT receptor 2a-like* receptor. The peak PBS in Waresley (blue) falls in the last intron of the gene.

Table 1. Population differences in enrichment scores in multiple GO terms associated with ovaries and oogenesis.

GO ID	GO Name	Size	Waresley		Gamling		Wales	
			NES	FDR q-val	NES	FDR q-val	NES	FDR q-val
GO:0030707	ovarian follicle cell development	345	3.36	0.00E+00	3.27	1.2E-05	2.52	1.35E-03
GO:0007297	ovarian follicle cell migration	129	3.32	0.00E+00	2.86	1.0E-04	3.09	4.10E-05
GO:1905879	regulation of oogenesis	66	2.22	7.31E-03	2.07	1.4E-02		
GO:0048599	oocyte development	165	2.16	9.70E-03	1.83	4.1E-02		
GO:0007304	chorion-containing eggshell formation	78	2.12	1.20E-02	2.41	2.0E-03		
GO:0007308	oocyte construction	157	2.07	1.55E-02	1.83	4.2E-02		
GO:0030703	eggshell formation	79	2.04	1.80E-02	2.50	1.3E-03		
GO:1905881	positive regulation of oogenesis	37	2.04	1.83E-02				
GO:0009994	oocyte differentiation	194	2.01	2.05E-02	2.25	5.1E-03		
GO:0007309	oocyte axis specification	145	1.90	3.44E-02	1.85	3.9E-02		
GO:0030728	ovulation	11	1.83	4.58E-02				
GO:0007306	eggshell chorion assembly	66			2.04	1.6E-02		
GO:0060281	regulation of oocyte development	29			1.79	4.9E-02		

Supplementary Information

a

b

Supplementary Fig. 1. Study sites and population differences in community structure. a, Location of traps in Gamlingay Wood (red points) and Waresley Wood (blue points). **b**, NMDS ordination of Gamlingay ($n = 141$) and Waresley ($n = 134$) community structure. Each point represents the sum of the beetle community collected in each trap per sampling time.

Supplementary Fig. 2. Mean body weight of small mammals. Gamlingay ($n = 21$, red) and Waresley Woods ($n = 30$, blue). One bank vole and one wood mouse, each caught in in Gamlingay Wood, escaped before they could be measured.

Supplementary Fig. 3. Differences in (a) body size and (b) frequency distribution of field-caught *N. vespilloides* from Gamlingay and Waresley Woods. Measurements depict pronotum width, the standard index for measuring beetle body size. Median values, inter-quartile range, maximum, and minimum are as illustrated in box-and-whisker plots. Outliers are depicted as points. Number of *N. vespilloides* sampled: Gamlingay (*n* = 839) and Waresley Woods (*n* = 824).

Supplementary Fig. 4. Density plot of *Fst* value between Gamlingay and Waresley Woods. The plot shows the narrow distribution of low *Fst* values between the two woodland populations. Though there is a long tail to the right, the extreme values are modest in absolute terms.

Supplementary Fig. 5. MDS plot of three burying beetle populations. The plot shows the first two dimensions of a multidimensional scaling analysis of genetic diversity among burying beetles in three populations. The populations do not separate out, indicating little to no genetic structure.

Supplementary Table 1. *Nicrophorus* **spp. body size (given by pronotum size, in mm).**

Species	*N*	Mean	SD	Minimum	Q1	Median	Q3	Maximum
N. vespilloides	1662	4.76	0.46	3.25	4.43	4.76	5.08	6.03
N. interruptus	152	5.59	0.64	4.28	5.10	5.60	6.10	7.07
N. investigator	58	6.19	0.61	4.89	5.80	6.18	6.58	7.59
N. humator	92	6.86	0.89	4.85	6.30	6.75	7.31	9.25
N. vespillo	11	5.74	0.59	4.86	5.44	5.76	5.98	6.93

(*N*): Number of *Nicrophorus* spp.; (SD): Standard Deviation; 25% (Q1): Quantile; (Q3): 75% Quantile.

Supplementary Table 2. Post-hoc Tukey HSD comparing mean body size between
Nicrophorus **spp.**

Species	N. vespilloides		N. interruptus		N. investigator		N. humator	
	t	*p*	*t*	*p*	*t*	*p*	*t*	*p*
N. interruptus	18.69	**<0.001**	-	-	-	-	-	-
N. investigator	20.62	**<0.001**	-7.61	**<0.001**	-	-	-	-
N. humator	38.15	**<0.001**	18.54	**<0.001**	7.69	**<0.001**	-	-
N. vespillo	6.31	**<0.001**	-0.93	0.268	2.68	0.058	6.85	**<0.001**

p values <0.05 are highlighted in bold. Dashes indicate comparisons that were not made.

Supplementary Table 3. Results of the ANOVAs for division of carrion niche by *Nicrophorus* spp.

Dependent variable	Explanatory variables	χ^2	d.f.	*p* value
Efficiency (%)	Carcass size	17.26	1	**<0.001**
	Beetle species	20.42	3	**<0.001**
	Carcass size x Beetle species	28.85	3	**<0.001**

p values <0.05 are highlighted in bold.

Supplementary Table 4. Results of the ANOVAs for reaction norm experiment.

Dependent variable	Explanatory variables	χ^2	d.f.	*p* value
Brood size	Carcass size	3.33	1	0.068
	Population	3.89	1	**0.048**
	Carcass size x Population	4.80	1	**0.029**
Average larval mass	Carcass size	139.05	1	**<0.001**
	Population	1.06	1	0.304
Clutch size	Carcass size	11.33	1	**<0.001**
	Population	21.07	1	**<0.001**
	Female size	4.26	1	**0.039**
Average egg volume	Carcass size	5.90	1	**0.015**
	Population	4.59	1	**0.032**
Total egg volume	Carcass size	2.38	1	0.123
	Population	2.16	1	0.141

p values <0.05 are highlighted in bold.

Supplementary Table 5. Differences in body size frequency distribution between *Nicrophorus* **spp.**

Species	*N. vespilloides*		*N. interruptus*		*N. investigator*		*N. humator*	
	D	p	D	p	D	p	D	p
N. interruptus	0.54	**<0.001**	-	-	-	-	-	-
N. investigator	0.81	**<0.001**	0.41	**<0.001**	-	-	-	-
N. humator	0.91	**<0.001**	0.62	**<0.001**	0.38	**<0.001**	-	-
N. vespillo	0.70	**<0.001**	0.31	0.268	0.43	0.061	0.69	**<0.001**

D statistic and corresponding *p*-values from Kolmogorov-Smirnov tests. *p* values <0.05 are highlighted in bold. Dashes indicate comparisons that were not made.

Supplementary Data 1. Spreadsheet file of gene set enrichment analysis results of the multiple GO terms for Gamlingay and Waresley *N. vespilloides*.