Investigating Population-scale Allele Specific Expression in Wild Populations of *Oithona similis* (Cyclopoida, Claus 1866)

Romuald Laso-Jadart[1,6*], Kevin Sugier[1], Emmanuelle Petit[2], Karine Labadie[2], Pierre Peterlongo[3], Christophe Ambroise[4], Patrick Wincker[1,6], Jean-Louis Jamet[5], Mohammed-Amin Madoui[1,6*]

[1]Génomique Métabolique, Genoscope, Institut François Jacob, CEA, CNRS, Univ Evry, Université Paris-Saclay, Evry, France.

[2]CEA, Genoscope, Institut de Biologie François Jacob, Université Paris-Saclay, Evry, 91057, France.

[3]Univ Rennes, CNRS, Inria, IRISA - UMR 6074, F-35000 Rennes.

[4]LaMME, CNRS, Univ Evry, Université Paris-Saclay, Evry, France

[5]Université de Toulon, Mediterranean Institute of Oceanology (MIO), AMU-UTLN UM110, CNRS UMR7294, IRD UMR235, Equipe Ecologie Marine et Biodiversité (EMBIO), Bat. R-121., CS 60584, 83041 Toulon Cedex 9, France

[6]Research Federation for the study of Global Ocean Systems Ecology and Evolution, FR2022/Tara Oceans GO-SEE, 3 rue Michel-Ange, 75016 Paris, France

[*]Corresponding authors. Emails : rlasojad@genoscope.cns.fr & amadoui@genoscope.cns.fr

Abstract

Allele-specific expression (ASE) is a widely studied molecular mechanism at cell, tissue and organism levels. Here, we extrapolated the concept of ASE to the population-scale (psASE), aggregating ASEs detected at smaller scales. We developed a novel approach to detect psASE based on metagenomic and metatranscriptomic data of environmental samples containing communities of organisms. This approach which measures the deviation between the frequency and the relative expression of biallelic loci, was applied on samples collected during the *Tara* Oceans expedition (2009-2013), in combination to new *Oithona similis* transcriptomes, a widespread marine copepod. Among a total of 25,768 single nucleotide variants (SNVs) of *O. similis*, 587 (2.3%) were targeted by psASE in at least one population. The distribution of SNVs targeted by psASE in different populations is significantly shaped by population genomic differentiation (p-value = 9.3×10^{-9}), supporting a partial genetic control of psASE. To investigate the link between evolution and psASE, loci under selection were compared to loci under psASE. A significant amount of SNVs (0.6%) were targeted by both selection and psASE (p-values $< 9.89 \times 10^{-3}$), supporting the hypothesis that natural selection and ASE may lead to the same phenotype. Population-scale ASE offers new insights into the gene regulation control in populations and its link with natural selection.

Introduction

Allele-specific expression (ASE), or allelic imbalance, refers to the difference of expression between two alleles of a locus in a heterozygous genotype. Recently, several studies led to a better understanding of ASE thanks to the development of advanced tools (Skelly et al. 2011; Mayba et al. 2014; Castel et al. 2015; Harvey et al. 2015; Lu et al. 2015; Rivas et al. 2015; Miao et al. 2018) allowing its detection at cellular (Milani et al. 2009; Ginart et al. 2016; Huang et al. 2017; Dong & Jiang 2019) tissue (McKean et al. 2016; Zhuo et al. 2017; Liu et al. 2018) and/or inter-individual levels (Josephs et al. 2015; Tung et al. 2015; Lonsdale et al. 2017; Wang et al. 2017). Moreover, studies began to question the relative contribution of genetics and environment on gene expression using ASE in human (Cheung et al. 2008; Buil et al. 2014; Cheung et al. 2008; Moyerbrailean et al. 2016; Knowles et al. 2017) and fruit flies (Leon-Novelo et al. 2017). Because the availability of reference genomes, numerous individual RNA-seq and whole-genome genotyping data still constitutes an important challenge for many species, there are few studies investigated ASE in natural populations (Tung et al. 2015; Wang et al. 2017).

ASEs are mainly due to epigenetic or genetic processes in *cis*-regulatory variations. Through DNA methylation, chromatin state or histone modifications, epigenetics could repress a disadvantageous or a specific parental allele, leading in some cases to monoallelic expression, as demonstrated in a variety of organisms including *Arabidopsis thaliana*, mouse, maize or bumblebee (Szabo & Mann 1995; Zhang & Borevitz 2009; Wei & Wang 2013; Ginart et al. 2016; Lonsdale et al. 2017). ASE may also have a genetic origin through, for example, mutations in transcription factor binding sites (Bailey et al. 2015; Cavalli et al. 2016), or post-transcriptional mechanisms like non-sense mediated decay (Castel et al. 2015; Rivas et al. 2015; Pirinen et al. 2015).

However, ASE at population-scale (psASE) has never been investigated. At this level, psASE may aggregate classical ASE at cell, tissue or individual levels (Figure 1a), but also differential expression between two homozygous genotypes. However, measurement of psASE would require the sequencing of

several individuals at genomic and transcriptomic levels separately. To measure psASE on small

organisms, an alternative approach could be to take advantage of metagenomic and metatranscriptomic

data that already contains sequences from pooled individuals. In this context, intra-species variants of a

single species have to be extracted, allowing to evaluate if the population-scale relative expression of an

allele deviates from its genomic frequency (Figure 1b).

Copepods, and particularly species belonging to the *Oithona* genus, are small crustaceans forming the

most abundant metazoan on Earth (Humes 1994; Gallienne 2001; Kiørboe 2011). This abundance,

reflecting strong adaptive capacities to environmental fluctuations, together with large hypothetic effective

population size (Peijnenburg & Goetze 2013; Riginos et al. 2016; Madoui et al. 2017; Arif et al. 2018)

make this species suitable for population genomics analyses. In addition, they play an ecological key role

in biogeochemical cycles and in the marine trophic food chain (Wassmann et al. 2006). In this study, we

propose to test our model to detect psASE by focusing on the widespread epipelagic copepod, *Oithona*

similis (Cyclopoida, Claus 1866). We used environmental samples collected by the *Tara* Oceans

expedition (Karsenti et al. 2011; Pesant et al. 2015) during its Arctic phase (2013), an area where

O. similis is known to be highly abundant (Blachowiak-Samolyk et al. 2008; Dvoretsky 2012; Zamora-

Terol et al. 2013; Castellani 2016) and for which both metagenomic and metatranscriptomic data are

available. Variants of *O. similis* were extracted and tested for psASE. In second time, we tried to decipher

the potential link between psASE, genomic differentiation and natural selection.

Material and Methods

Variant calling using *Tara* Oceans metagenomic and metatranscriptomic data

We used metagenomic and metatranscriptomic reads generated from samples of the size fraction 20–180

µm collected in seven *Tara* Oceans stations (TARA_155, 158, 178, 206, 208, 209 and 210)(Figure 2a)

according to protocols described by Alberti et *al.* (Supplementary Table S2). Because of the lack of a

reference genome, the reference-free variant caller *DiscoSNP++* (Uricaru et al. 2014; Peterlongo et al.

2017) was used to extract SNVs simultaneously from raw metagenomic and metatranscriptomic reads, and

was ran using parameter –b 1. Only SNVs corresponding to biallelic loci with a minimum of 4x of depth

of coverage in all stations were initially selected. Then, to capture loci belonging to *Oithona similis*, SNVs

were clustered based on their loci co-abundance across samples using density-based clustering algorithm

implemented in the R package dbscan (Ester et al. 1996; Ram et al. 2010) and ran with parameters epsilon

= 10 and minPts = 10. This generated three SNVs clusters, the largest of which contained 102,258 SNVs.

To ensure only the presence of *O. similis* SNVs, we observed the fitting of the depth of coverage to the

expected negative binomial distribution in each population (Supplementary Figure S2). As the variant

calling step is reference-free, the alternative allele (here, B-allele) is arbitrary set by *DiscoSNP++*. For

each variant in each population, the B-allele frequency (BAF) and the population-level B-allele relative

expression (BARE) were computed as follow, $BAF = \frac{G_B}{G_B+G_A}$ and $BARE = \frac{T_B}{T_B+T_A}$, with G_A, G_B the

metagenomic read counts supporting the reference and alternative alleles respectively and T_A, T_B the

metatranscriptomic read counts supporting the reference and alternative alleles respectively.

Biallelic loci were then filtered based on their metagenomic coverage. For each sample, the median and

standard deviation σ of the distribution of metagenomic coverage of all biallelic loci were estimated.

Biallelic loci must be characterized by a metagenomic coverage between a limit of median ± 2σ, with a

minimum and maximum of 5x and 150x coverage in each sample to avoid low covered and multicopy

genomic regions. To keep out rare alleles and potential calling errors, only variants characterized by a

BAF comprised between 0.9 and 0.1, and a BARE between 0.95 and 0.05 in at least one population were

chosen for the final dataset resulting in 25,768 biallelic loci.

To ensure that these loci belong to *O. similis*, the global *F*-statistics (or Wright's fixation index (Wright

1951; B. S. Weir and C. Clark Cockerham 1984)) over the seven populations was computed as follow, F_{ST}

$= \frac{\sigma^2}{\bar{p}(1-\bar{p})}$, with $\bar{p}$ being the mean allele frequency across all the samples , and its distribution was tested for

unimodality via a Hartigans' dip test (Hartigan & Hartigan 1985). Moreover, LK statistics (R.C. & J.

1973) was computed as follow, LK $= \frac{n-1}{\overline{F_{ST}}} F_{ST}$, and compared to the expected chi-squared distribution with

df = n-1, with n being the number of populations and $\overline{F_{ST}}$ being the mean F_{ST} across all loci.

Population-scale ASE detection using metagenomic and metatranscriptomic data

In each population, we first selected variants for BAF $\neq$ {0,1}. We tested the correlation between BAF and

BARE and then we computed $D = BAF\text{-}BARE$, as the deviation between the BAF and the BARE. In the

seven samples, D followed a Gaussian distribution centered on 0, allowing the estimation of the Gaussian

distribution parameters and test the probability of a variant to belong to this distribution (this is later

referred here as "D-test" or "deviation test"). Given the large number of tests, we applied the Benjamini

and Hochberg approach (Benjamini & Hochberg 1995) to control the False Discovery Rate (FDR). We

also computed a "low expression bias" test by comparing the read counts T_A and T_B to the observed

metagenomic proportion 1-BAF and BAF respectively with a chi-squared test and applied the Benjamini

and Hochberg correction for multiple testing.

The deviation and the low expression bias tests were applied to all loci in the seven populations

separately. Loci with a deviation test q-value < 0.1 and a low expression bias test q-value < 0.1 were

considered under psASE. This generates seven sets of candidate loci under psASE, one set for each

population.

Estimation of genomic differentiation and detection of variants under selection

Pairwise-F_{ST} was computed for each locus between each pair of population and the median pairwise-F_{ST}

was retained to measure the genomic differentiation between each population. A Mantel test was

performed to test for isolation-by-distance between median pairwise-F_{ST} and geographic Euclidean

distances using *vegan* v2.5-2 (Oksanen et al. 2018) and *geosphere* v1.5-7 (Hijmans 2017) R packages. The

pcadapt R package v4.0.2 (Luu et al. 2017) was used to detect selection among populations from the B-

allele frequency matrix. The computation was run on "*Pool-seq*" mode, with a minimum allele frequency

of 0.05 across the populations, and variants with a corrected Benjamini and Hochberg p-value < 0.05 were

considered under selection.

Modelling psASE with genomic differentiation

The seven sets of candidate loci under psASE were crossed to identify variants under psASE in several

populations (named "shared psASEs") and all non-empty crossings between populations were represented

by an upset plot.

To test if populations characterized by weak genetic differentiation tend to share more loci under psASE

than genetically distant populations, we modelled the number of shared psASEs between populations by

genomic differentiation using a non-linear model: $E(Y) \sim a\ exp(bX) + c$, with Y being the number of

shared ASE and X the genomic differentiation. The latter was estimated by computing the median-F_{ST} of

each non-empty crossing. To find the starting values, the model was linearized as follow, $log(E(Y)-c0) \approx$

$log(a)+bX$, with $c0 = min(Y) * 0.5$ and a and b parameters were estimated via the *lm* R function. The non-

linear model was then applied, and least squares estimates were used via the *nls* R function. Pearson

correlation between the fitted and empirical values was then computed via the *cor.test* R function.

psASE and link with natural selection

To identify alleles targeted by both psASE and natural selection, the set of variants under ASE in each

population was crossed with the set of loci detected under selection. The size of the intersection was tested

by a hypergeometric test, Hypergeometric(q,m,n,k), with q being number of alleles under ASE in the

population and under selection (size of intersection), m being the total number of alleles under selection, n

being the total number of variants under neutral evolution, and k being the total number of alleles under

psASE in the tested population. We considered that, in a given population, the number of alleles under

both psASE and selection was significantly higher than expected by chance for p-value < 0.05.

Material sampling, mRNA extraction and Mediterranean *O. similis* transcriptomes sequencing

To conduct a functional analysis, Mediterranean *O. similis* transcriptomes were produced. *Oithona similis* specimens were sampled at the North of the Large Bay of Toulon, France (Lat 43°06' 02.3" N and Long 05°56' 53.4" E). Sampling took place in November 2016. The samples were collected from the upper water layers (0-10m) using zooplankton nets with a mesh of 90µm and 200µm (0.5 m diameter and 2.5 m length). Samples were preserved in 70% ethanol and stored at -4°C. From the Large Bay of Toulon samples, *O. similis* individuals were isolated under the stereomicroscope (Rose 1933; Nishida 1985). We selected two different development stages: four copepodites (juveniles) and four adult males. Each individual was transferred separately and crushed, with a tissue grinder (Axygen) into a 1.5 mL tube (Eppendorf). Total mRNAs were extracted using the 'RNA isolation' protocol from NucleoSpin RNA XS kit (Macherey-Nagel) and quantified on a Qubit 2.0 with a RNA HS Assay kit (Invitrogen) and on a Bioanalyzer 2100 with a RNA 6000 Pico Assay kit (Agilent). cDNA were constructed using the SMARTer-Seq v4 Ultra low Input RNA kit (ClonTech). The libraries were built using the NEBNext Ultra II kit, and were sequenced with an Illumina.

Transcriptomes assembly and annotation

Each read set was assembled with Trinity v2.5.1 (Haas et al. 2013) using default parameters and transcripts were clustered using cd-hit v4.6.1 (Fu et al. 2012) (Supplementary Table S1). To ensure the classification of the sampled individuals, each ribosomal read set were detected with SortMeRNA (Kopylova et al. 2012) and mapped with bwa v0.7.15 using default parameters (Li & Durbin 2009) to 82 ribosomal 28S sequences of *Oithona* species used in Cornils et al., 2017 (Supplementary Figure S1). The transcriptome assemblies were annotated with Transdecoder v5.1.0 (Haas et al. 2013) to predict the open reading frames (ORFs) and protein sequences (Supplementary Table S1). In parallel, homology searches were also included as ORF retention criteria; the peptide sequences of the longest ORFs were aligned on *Oithona nana* proteome (Madoui et al. 2017) using DIAMOND v0.9.22 (Buchfink et al. 2014). Protein domain annotation was performed on the final ORF predictions with Interproscan v5.17.56.0 (Jones et al.

2014) and a threshold of e-value $<10^{-5}$ was applied for Pfam annotations. Finally, homology searches of

the predicted proteins were done against the nr NCBI database, restricted to Arthropoda (taxid: 6656),

with DIAMOND v0.9.22.

Variant annotation

The variant annotation was conducted in two steps. First, the variant sequences were relocated on the

previously annotated *O. similis* transcripts using the "VCF_creator.sh" program of *DiscoSNP++*.

Secondly, a variant annotation was carried out with SNPeff (Cingolani et al. 2012) to identify the location

of variants within transcripts (i.e., exon or UTR) and to estimate their effect on the proteins (missense,

synonymous or nonsense). The excess of candidate variant annotations was tested in the following classes:

missenses, synonymous, 5' and 3'UTR. A significant excess was considered for a hypergeometric test p-

value < 0.05.

Gene enrichment analysis

To identify specific biological function or processes associated to the variants, a domain-based analysis

was conducted. The Pfam annotation of the transcripts carrying variants targeted by ASE and selection

was used as entry for dcGO Enrichment (Fang & Gough 2013)**.** A maximum of the best 300 GO-terms

were chosen based on their z-score and FDR p-value ($<10^{-3}$) in each ontology category. To reduce

redundancy, these selected GO-terms were processed using REVIGO (Supek et al. 2011), with a similarity

parameter of 0.5 against the whole Uniprot catalogue under the SimRel algorithm. To complete the

domain-based analysis, the functional annotations obtained from the homology searches against the nr

were manually curated.

Results

Extracting polar *Oithona similis* variants from environmental samples

From metagenomic and metatranscriptomic raw data of seven sampling stations (Figure 2a), we identified 102,258 variants using a reference-free approach. Among them, 25,768 expressed *O. similis* variants were retrieved after filtering. To ensure that the variants belonged to *O. similis*, we performed three different analyses. First, in each sample, the distributions of variable loci depth of coverage were unimodal (Figure 2b) and fitted the expected negative binomial distributions (Supplementary Figure S2). Second, 97% of 25,768 variants were relocated on Mediterranean *O. similis* transcriptomes (Figure 2a). Third, the global distribution of F_{ST} of the seven populations was unimodal (Hartigans' dip test, D=0.0012, p-value=0.19) with a low median F_{ST} at 0.1 (Figure 2c), confirmed by the pairwise-F_{ST} distributions (Supplementary Figure S3d). Finally, the LK distribution over all the loci followed the expected chi-squared distribution (Figure 2d), showing that most of the loci follow the neutral evolution model, as expected in a single species.

***Oithona similis* genomic differentiation in Arctic Seas**

The seven populations were globally characterized by a weak to moderate differentiation, with a maximum median pairwise-F_{ST} of 0.12 between populations from TARA_210 and 155/178 (Figure 3b, Supplementary Figure S3). Populations from stations TARA_158 (Norway Current), 206 and 208 (Baffin Bay) were genetically closely related, with the lowest median pairwise-F_{ST} (0.02), despite TARA_158 did not co-geolocalize with the two other stations. The four other populations (TARA_155, 178, 209, and 210) were equally distant from each other (0.1-0.12). Finally, TARA_158, 206 and 208 on one side, and TARA_155, 178, 210 and 209 on the other side showed the same pattern of differentiation (0.05-0.07). A Mantel test was performed and revealed no correlation between F_{ST} and geographic distances (r = 0.34, p-value = 0.13) (Supplementary Figure S5).

Detection of population-scale ASE

The number of SNVs tested for psASE varied between 13,454 and 22,578 for TARA_210 and 206 respectively. As expected, most of the loci presented a strong correlation between B-allele frequency and B-allele relative expression (Figure 3a, Table 1). Thus, we measured the D deviation, representing the deviation between BAF and BARE, which followed a Gaussian distribution in each population (Figure 3a). Variants under psASE (i.e. having a D significantly higher or lower than expected and passing the low expression bias test) were found in every population, ranging from 26 to 162 variants for TARA_178 and 206 respectively (Table 1). Overall, we found 587 variants under psASE, including 535 population-specific ASEs, and 52 ASEs shared by several populations (Figure 3c). Remarkably, 30 ASEs out of the 52 were present in the populations from TARA_158, 206 and 208 that correspond to the genetically closest populations. By comparing the number of shared psASEs in the different sets of populations to their genomic differentiation, we found a negative trend between the two (with a strong negative b estimate), illustrated by a significant correlation between non-linear fitted and empiric values (0.89, p-value 9.3×10^{-9}, Figure 3d). This modelling shows that genetically close populations tend to share the same variants under psASE.

Loci targeted by population-level ASE and selection in Arctic populations

The set of variants was tested for selection using *pcadapt*. The PCA decomposed the genomic variability in six components (Supplementary Figure S3a,b,c); the first two components discriminated TARA_155 and 178 from the others (32% and 28.1% variance explained respectively), and the third component differentiated TARA_210 and 209 (19.5%). The fourth principal component separated TARA_209 and 210 from 158/206/208 (11.3 %), with the last two concerning TARA_158/206/208. Globally, these results dovetailed with the F_{ST} analysis, with details discussed later. Finally, we detected 674 variants under selection, representing 2.6% of the dataset (corrected p-value < 0.05).

The seven sets of variants under psASE were crossed with the set of variants under selection (Figure 4a). The size of the intersections ranged from 6 to 42 variants (TARA_155 and 210/206) and was significantly

248 higher than expected by chance for all the populations (Figure 4b, hypergeometric test p-value < 0.05). It

249 represented a total of 152 unique variants under selection and ASE in at least one population,

250 corresponding to 23% and 26% of variants under psASE and under selection respectively.

Functional analysis of genes targeted by population-scale ASE and selection

Among the 152 loci targeted by psASE and selection, 145 were relocated on *O. similis* transcripts

(Supplementary Table S4). Amid these transcripts, 137 (90%) had a predicted ORF, 97 (64%) were linked

to at least one Pfam domain and 90 (59%) to a functional annotation. Fifteen SNVs were missense

variations, 59 synonymous, 31 and 29 were in 5' and 3' UTR, without any significant excess

(Supplementary Table S4 and S5). Based on homology searches (Supplementary Table S4), eight genes

were linked to nervous system (Table 2). Among them, two genes were involved in glutamate metabolism

(omega-amidase NIT2 and 5-oxoprolinase), three were predicted to be glycine, γ-amino-butyric acid

(GABA) and histamine neuroreceptors. Finally, four were also implicated in arthropods photoreceptors.

The domain-based analysis confirmed these results, with an enrichment in GO-terms biological process

also linked to nervous system (Supplementary Figure S6).

Discussion

Genomic and transcriptomic variation data belong to a single *O. similis* lineage

Because genomes of small animals like copepods are difficult to reconstruct, we used *DiscoSNP++*, a

reference-free variant caller to extract variants from metagenomic data, that already showed its good

accuracy on *Tara* Oceans metagenomic data (Arif et al. 2018).

Global populations of *O. similis* are known to be composed of cryptic lineages across oceanic basins

(Cornils et al. 2017). It is also known that this species in highly abundant among other copepods in Arctic

Seas (Blachowiak-Samolyk et al. 2008; Dvoretsky 2012; Zamora-Terol et al. 2013; Castellani 2016).

Thus, the assessment that the extracted variants from the seven samples used in our study belong to the

same *O. similis* cryptic lineage was a prerequisite for further analyses. Three different analyses support

this assumption. First, the distribution of depth of coverage in each of the seven samples followed the

expected negative binomial distribution (Supplementary Figure S2). Indeed, the possibility to observe

these patterns in the presence of different species would require them to be equally co-abundant, which is

unlikely. Thus, this covariation of the depth of coverage of these variants supports the single species

genome origin. Secondly, the high proportion of variants (97%) mapped on the Mediterranean *O. similis*

transcriptomes, another cryptic lineage (Cornils et al. 2017), showed that the variant clustering method

was efficient to regroup loci of *O. similis*. Finally, the unimodal distribution of F_{ST} showed that these

populations of *O. similis* belong to the same polar cryptic species, and that all loci are under neutral

evolution. Altogether, these results show that we were able to retrieve polymorphic data of a single

species, *O. similis*, on which population genomics and psASE detection can be undertaken.

***O. similis* populations are weakly structured within the Arctic Seas**

We observed that the seven populations examined showed low genomic differentiation, despite the large

distances separating them, which was illustrated by a non-significant Mantel test for isolation-by-distance

(Figure 2a). F_{ST} and *pcadapt* analyses both showed the same patterns of genomic differentiation. Going

into details, three different cases can be described. First, the differentiation of populations from

TARA_155 and 178 is relatively high compared to the others. Secondly, the geographically close

populations from TARA_210 and 209 present a relatively high differentiation (median pairwise-F_{ST} of

0.11, PC3). This could be explained by the West Greenland current acting as a physical barrier between

the populations, which could lead to reduced gene flow (Myers et al. 2008). At last, the strong link

between TARA_158 from Northern Atlantic current and TARA_206/208 from the Baffin Bay is the most

intriguing. Despite the large distances that separate the first one from the others, these three populations

are well connected. Based on this weak structure and that most of loci follows a neutral evolution (Figure

2d), outliers detected by *pcadapt* probably are truly under selection and not due to specific population

differentiation.

Metagenomic data enable to draw the silhouette of the population genetics but lacks resolution when

dealing with intra-population structure. However, our findings are concordant with previous studies

underpinning the large-scale dispersal, interconnectivity of marine zooplankton populations in other

oceans, at diverse degrees (Goetze 2005; Peijnenburg & Goetze 2013; Blanco-Bercial & Bucklin 2016;

Höring et al. 2017). Weak genetic structure in the polar region was highlighted for other major Arctic

copepods like *Calanus glacialis* (Weydmann et al. 2016), and *Pseudocalanus* species (Aarbakke et al.

2014). The absence of structure was explained by ancient diminutions of effective population size due to

past glaciations (Bucklin & Wiebe 1998; Edmands 2001; Aarbakke et al. 2014), or high dispersal and

connectivity between the present-day populations due to marine currents (Weydmann et al. 2016). Using

Lagrangian time travel or dispersal probabilities could help to estimate how much marine currents explain

this observed genomic differentiation.

Population-scale ASE in *O. similis* populations

Usually, ASE analyses are achieved by measuring the difference in RNA-seq read counts of a

heterozygous site at cellular, tissue or individual scales. Obtaining a comprehensive and exhaustive view

of ASE at all these scales is merely possible due to sampling challenges for small organisms. Particularly,

gathering a large number of individuals, sampling the tissues or organs of interest remain a technical

barrier especially for uncultured small animals, when the amount of DNA retrieved from a single

individual is not sufficient for high-throughput sequencing. Here, the detection of psASE was possible by

comparing the observed allele frequencies based on metagenomic and metatranscriptomic data, which

bypasses the obstacles previously described. The null hypothesis, supposing a strong correlation between

the expression of an allele and its genomic abundance at the population-scale was confirmed in the seven

populations, allowing us to continue and detect outliers targeted by psASE with our approach. In the case

of psASE, we expect to detect strong patterns of unbalanced allelic expression present in a large fraction

of the individuals of a population. Thus, a variant can be detected under psASE because of intra- or inter-

individual ASEs. Intra-individual ASE detected at population-level should be present in the majority of

individuals at cell or organ levels which constitutes the classical type of ASE mechanism. Inter-individual

ASEs encompass ASEs due to physiological states, developmental stages, or sex differentiation. In

O. similis, the proportion of each developmental stage and sex is known to vary within and between

populations and across seasons (Lischka & Hagen 2005; Dvoretsky 2012).

In our study, the amount of detected psASE in each population was always lower than 1% of tested

heterozygous variants, which altogether correspond to 2% of the total set of variants. In humans (Zhang et

al. 2014), baboons (Tung et al. 2015) and flycatchers (Wang et al. 2017), 17%, 23% of genes and 7.5% of

transcripts were affected by ASE respectively, meaning that psASE possibly tends to dilute individual

ASE.

Genomic drivers of population-scale ASE

The role of genomic background on psASE was deciphered by explaining the amount of shared psASEs

by the genomic differentiation. From the strong correlation and the estimates of the model, we conclude

that highly connected populations tend to share more loci under psASE than less connected populations,

characterized by weak differentiation. However, even if only a small proportion of variants were targeted

by psASE in different populations, our results show that genomic background partially drives psASE. To

have a better understanding of the factors explaining psASE, environmental or population dynamics

parameters (sex-ratio bias, proportions of larvae for example) could also be used as explanatory variables.

Toward the link between population-scale ASE and natural selection

As genomic background among the seven populations partially explains psASE, we estimated the number

of loci under psASE that were also targeted by natural selection. A significant amount of SNVs (152)

were subject to selection among the seven populations and to psASE in at least one population, meaning

that the two mechanisms can target the same genomic regions more than expected randomly.

Three main features of ASE can be under selection. First, the observed variant can be in linkage with

another variation in upstream *cis*-regulatory elements like transcription factors fixation sites, or epialleles

(Castel et al. 2015). Secondly, our annotation of candidate variants with SNPeff revealed a majority of

variants located in 5' and 3'UTRs, which are variations known to both affect transcription efficiency

through mRNA secondary structures, stability and location (Mignone et al. 2002; Matoulkova et al. 2012;

Dvir et al. 2013). For variants located in exons, a majority were identified as synonymous mutations,

growingly described as potential target of selection by codon usage bias, codon context, mRNA secondary

structure or transcription and translation dynamics (Ingvarsson 2010; Shabalina et al. 2013). Also, fifteen

missense mutations were spotted, but with moderate predicted impact on protein amino acid composition.

Nevertheless, we did not find premature nonsense mutation, even if variants under ASE has been

described to trigger or escape potential nonsense-mediated decay, but the possibility that the causal

variation is located in introns cannot be ruled out. However, due to the large possible origins of ASE and

the lack of a proper reconstructed *O. similis* genome and genotypes, no major cause was identified from

this study.

The process of adaptation through gene expression was studied in human populations and investigated

thanks to the large amount of accessible data. In a first study, a link has been established between gene

expression and selection, affecting particular genes and phenotypes, looking at *cis*-acting SNPs (Fraser

2013). In a second study, the team was able to detect loci under ASE and selection at the same time in

different human populations (Tian et al. 2018). Finally, approaches in a plant model, *Capsella grandiflora*,

a species characterized by weak population structure and large effective population size, emphasized the

relative impact of purifying selection and positive selection on *cis*-regulatory variation in populations

(Josephs et al. 2015; Steige et al. 2017). In our results, we identified a large fraction of candidate loci

(Supplementary Figure S5) being under psASE in one population with a very low genomic frequency and

tending to fixation in other populations (as exampled in Figure 4c). This could be linked to an adaptation

of a population to environmental changing conditions during constant oceanic migration, leading to a

purifying selection on the considered allele or a positive selection on the alternative allele. Our

observations thus complete previous analyses, as they quantify the link between psASE and selection in

natural marine populations composed of many individuals and reveal the evolutive potency of ASE, for

the first time at the population-scale. Together, these results show that the two mechanisms lead to a

similar molecular phenotype in allelic expression. This raises questions on whether ASE and selection are

independent or sequential mechanisms.

Nervous system and visual perception are important targets of the natural selection and population-

scale ASE in *Oithona similis*

This evolutive link between psASE and selection is supported by the biological functions associated to the

targeted genes, which are involved notably in the copepods nervous system in two ways. The first result is

the presence of genes implicated in glutamate metabolism and glycine and/or GABA receptors. Glutamate

and GABA are respectively excitatory and inhibitory neurotransmitters in arthropods motor neurons

(Smarandache-Wellmann 2016). Glycine and GABA receptors have already been described as a target of

selection in *Oithona nana* in Mediterranean Sea (Madoui et al. 2017; Arif et al. 2018). Secondly, the

functional analysis revealed also the importance of the eye and visual perception in the *O. similis*

evolution.

Together, this shows that psASE has an evolutive impact on functions linked to the nervous system of

copepods. The latter constitutes a key trait for its reproduction and survival, and based on our data and

previous work, a prime target for evolution, allowing higher capacity of perceiving and fast reacting

leading to more efficient predator escape, prey catching and mating. This can explain the great evolutive

success of these animals (Svensen 2000; Kiørboe et al. 2010; Kiørboe 2011).

Conclusion

Gene expression variation is thought to play a crucial role in evolutive and adaptive history of natural

populations. Herein, we integrated metagenomic and metatranscriptomic data to detect ASE at the

population level for the first time. Then, we demonstrated the link between psASE and genomic

background of the populations on one hand and with natural selection on the other hand, by providing a

quantitative observation of this phenomenon and its impact on specific biological features of copepods. In

the future, we will try to expand these observations to other organisms and question the nature of the link

between psASE and natural selection.

Acknowledgments

This work was supported by sponsors who participated in the *Tara* Oceans Expedition 2009–2013: Centre

National de la Recherche Scientifique, European Molecular Biology Laboratory,

Genoscope/Commissariat à l'Energie Atomique, the French Government "Investissements d'Avenir"

programmes OCEANOMICS (ANR-11- BTBR-0008), FRANCE GENOMIQUE (ANR-10-INBS-09-08),

Agnes b., the Veolia Environment Foundation, Region Bretagne, World Courier, Illumina, Cap L'Orient,

the Electricite de France (EDF) Foundation EDF Diversiterre, Fondation pour la Recherche sur la

Biodiversite, the Prince Albert II de Monaco Foundation, Etienne Bourgois and the *Tara* schooner and its

captain and crew. *Tara* Oceans would not exist without continuous support from 23 institutes (oceans.tara-

expeditions.org). We acknowledge the support of Vincent Segura and Leila Tirichine for fruitful scientific

discussions and support on the analyses and manuscript. This is contribution number XX from *Tara*

Oceans.

Author's contributions

410 Individuals for transcriptome production were sampled by J-LJ and KS. KS extracted RNA, EP and KL

411 prepared the libraries and sequencing, MAM assembled the reads and RLJ annotated transcriptomes. PP

412 and CA gave expertise support on *DiscoSNP++* and statistical framework respectively. RLJ and MAM

413 performed the analyses and wrote the manuscript. MAM designed and supervised the study. J-LJ and PW

414 offered scientific support.

Competing interests

The authors declare no competing interests.

References

Aarbakke ONS, Bucklin A, Halsband C, Norrbin F. 2014. Comparative phylogeography and demographic history of five sibling species of Pseudocalanus (Copepoda: Calanoida) in the North Atlantic Ocean. J. Exp. Mar. Bio. Ecol. 461:479–488. doi: 10.1016/j.jembe.2014.10.006.

Alberti A et al. 2017. Viral to metazoan marine plankton nucleotide sequences from the Tara Oceans expedition. Sci. Data. 4:170093. doi: 10.1038/sdata.2017.93.

Arif M et al. 2018. Discovering Millions of Plankton Genomic Markers from the Atlantic Ocean and the Mediterranean Sea. Mol. Ecol. Resour. 0–3. doi: 10.1111/1755-0998.12985.

B. S. Weir and C. Clark Cockerham. 1984. Estimating F-Statistics for the Analysis of Population Structure. Evolution (N. Y). 38:1358–1370. doi: 10.2307/2408641.

Bailey SD, Virtanen C, Haibe-Kains B, Lupien M. 2015. ABC: A tool to identify SNVs causing allele-specific transcription factor binding from ChIP-Seq experiments. Bioinformatics. 31:3057–3059. doi: 10.1093/bioinformatics/btv321.

Benjamini Y, Hochberg Y. 1995. Controlling the False Discovery Rate : A Practical and Powerful Approach to Multiple Testing. J. R. Stat. Soc. Ser. B. 57:289–300. doi: 10.2307/2346101.

Blachowiak-Samolyk K, Kwasniewski S, Hop H, Falk-Petersen S. 2008. Magnitude of mesozooplankton variability: A case study from the Marginal Ice Zone of the Barents Sea in spring. J. Plankton Res. 30:311–323. doi: 10.1093/plankt/fbn002.

Blanco-Bercial L, Bucklin A. 2016. New view of population genetics of zooplankton: RAD-seq analysis reveals population structure of the North Atlantic planktonic copepod Centropages typicus. Mol. Ecol. 25:1566–1580. doi: 10.1111/mec.13581.

Buchfink B, Xie C, Huson DH. 2014. Fast and sensitive protein alignment using DIAMOND. Nat. Methods. 12:59–60. doi: 10.1038/nmeth.3176.

Bucklin A, Wiebe PH. 1998. Low mitochondrial diversity and small effective population sizes of the copepods Calanus finmarchicus and Nannocalanus minor: Possible impact of climatic variation during recent glaciation. J. Hered. 89:383–392. doi: 10.1093/jhered/89.5.383.

Buil A et al. 2014. Gene-gene and gene-environment interactions detected by transcriptome sequence analysis in twins. Nat. Genet. 47:88–91. doi: 10.1038/ng.3162.

Castel SE, Levy-Moonshine A, Mohammadi P, Banks E, Lappalainen T. 2015. Tools and best practices for data processing in allelic expression analysis. Genome Biol. 16:195. doi: 10.1186/s13059-015-0762-6.

Castellani. 2016. Contribution to the Themed Section : ' The Role of Zooplankton in Marine Biogeochemical Cycles : From Fine Scale to Global Marine zooplankton and the Metabolic Theory of Ecology : is it a predictive tool ? J. Plankton Res. 38:762–770.

Cavalli M et al. 2016. Allele-specific transcription factor binding to common and rare variants associated with disease and gene expression. Hum. Genet. 135:485–497. doi: 10.1007/s00439-016-1654-x.

Cheung VG et al. 2008. Monozygotic Twins Reveal Germline Contribution to Allelic Expression Differences. Am. J. Hum. Genet. 82:1357–1360. doi: 10.1016/j.ajhg.2008.05.003.

Cingolani P et al. 2012. A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of Drosophila melanogaster strain w1118; iso-2; iso-3. Fly (Austin). doi: 10.4161/fly.19695.

Claus C. 1866. Die Copepoden-Fauna von Nizza. Ein Beitrag zur Charakteristik der Formen und deren Abänderungen 'im Sinna Darwin's'. Elwebt'sche Univ. Marbg. Leipzig. 1:1:34.

Cornils A, Wend-Heckmann B, Held C. 2017. Global phylogeography of Oithona similis s.l. (Crustacea, Copepoda, Oithonidae) – A cosmopolitan plankton species or a complex of cryptic lineages? Mol. Phylogenet. Evol. 107:473–485. doi: 10.1016/j.ympev.2016.12.019.

Denno ME, Privman E, Borman R, Wolin D, Venton BJ. 2016. Quantification of histamine and carcinine in Drosophila melanogaster tissues. ACS Chem Neurosci. 7:407–414. doi: 10.1021/acschemneuro.5b00326.

Dong M, Jiang Y. 2019. Single-Cell Allele-Specific Gene Expression Analysis. Comput. Methods Single-Cell Data Anal. 1935.

Dvir S et al. 2013. Deciphering the rules by which 5'-UTR sequences affect protein expression in yeast. Proc. Natl. Acad. Sci. 110:E2792–E2801. doi: 10.1073/pnas.1222534110.

Dvoretsky VG. 2012. Seasonal mortality rates of Oithona similis (Cyclopoida) in a large Arctic fjord. Polar Sci. 6:263–269. doi: 10.1016/j.polar.2012.09.001.

Edmands S. 2001. Phylogeography of the intertidal copepod Tigriopus californicus reveals substantially reduced population differentiation at northern latitudes. Mol. Ecol. 10:1743–1750. doi: 10.1046/j.0962-1083.2001.01306.x.

Ester M, Kriegel H-P, Sander J, Xu X. 1996. *A Density-Based Algorithm for Discovering Clusters in Large Spatial Databases with Noise*. www.aaai.org (Accessed January 8, 2019).

Fang H, Gough J. 2013. DcGO: Database of domain-centric ontologies on functions, phenotypes, diseases and more. Nucleic Acids Res. 41. doi: 10.1093/nar/gks1080.

Fraser HB. 2013. Gene expression drives local adaptation in humans. Genome Res. 1089–1096. doi: 10.1101/gr.152710.112.

Fu L, Niu B, Zhu Z, Wu S, Li W. 2012. CD-HIT: Accelerated for clustering the next-generation sequencing data. Bioinformatics. 28:3150–3152. doi: 10.1093/bioinformatics/bts565.

Gallienne CP. 2001. Is Oithona the most important copepod in the world's oceans? J. Plankton Res. 23:1421–1432. doi: 10.1093/plankt/23.12.1421.

Ginart P et al. 2016. Visualizing allele-specific expression in single cells reveals epigenetic mosaicism in an H19loss-of-imprinting mutant. Genes Dev. 30:567–578. doi: 10.1101/gad.275958.115.

Goetze E. 2005. Global Population Genetic Structure and Biogeography of the Oceanic Copepods Eucalanus Hyalinus and E. Spinifer. Evolution (N. Y). 59:2378. doi: 10.1554/05-077.1.

Gurudev N, Yuan M, Knust E. 2014. chaoptin, prominin, eyes shut and crumbs form a genetic network
controlling the apical compartment of Drosophila photoreceptor cells. Biol. Open. 3:332–341. doi:
10.1242/bio.20147310.

Haas BJ et al. 2013. De novo transcript sequence reconstruction from RNA-seq using the Trinity platform
for reference generation and analysis. Nat. Protoc. 8:1494–1512. doi: 10.1038/nprot.2013.084.

Hartigan JA, Hartigan PM. 1985. The dip test of unimodality. Ann. Stat. 13:70–84.

Harvey CT et al. 2015. QuASAR: Quantitative allele-specific analysis of reads. Bioinformatics. 31:1235–
1242. doi: 10.1093/bioinformatics/btu802.

Hijmans RJ. 2017. geosphere: Spherical Trigonometry. https://cran.r-project.org/package=geosphere.

Höring F, Cornils A, Auel H, Bode M, Held C. 2017. Population genetic structure of Calanoides natalis
(Copepoda, Calanoida) in the eastern Atlantic Ocean and Benguela upwelling system. J. Plankton Res.
39:618–630. doi: 10.1093/plankt/fbx035.

Huang WC et al. 2017. Diverse Non-genetic, Allele-Specific Expression Effects Shape Genetic
Architecture at the Cellular Level in the Mammalian Brain. Neuron. 93:1094–1109.e7. doi:
10.1016/j.neuron.2017.01.033.

Humes AG. 1994. How Many Copepods? Hydrobiologia. 293:1–7. doi: 10.1007/BF00229916.

Ingvarsson PK. 2010. Natural Selection on Synonymous and Nonsynonymous Mutations Shapes Patterns
of Polymorphism in Populus tremula. Mol. Biol. Evol. 27:650–660. doi: 10.1093/molbev/msp255.

Jones P et al. 2014. InterProScan 5: genome-scale protein function classification. Bioinformatics.
30:1236–1240. doi: 10.1093/bioinformatics/btu031.

Josephs EB, Lee YW, Stinchcombe JR, Wright SI. 2015. Association mapping reveals the role of
purifying selection in the maintenance of genomic variation in gene expression. Proc. Natl. Acad. Sci.
112:15390–15395. doi: 10.1073/pnas.1503027112.

Karsenti E et al. 2011. A Holistic Approach to Marine Eco-Systems Biology. PLoS Biol. 9:e1001177. doi:
10.1371/journal.pbio.1001177.

Kiørboe T. 2011. What makes pelagic copepods so successful? J. Plankton Res. 33:677–685. doi:
10.1093/plankt/fbq159.

Kiørboe T, Andersen A, Langlois VJ, Jakobsen HH. 2010. Unsteady motion: Escape jumps in planktonic
copepods, their kinematics and energetics. J. R. Soc. Interface. 7:1591–1602. doi: 10.1098/rsif.2010.0176.

Knowles DA et al. 2017. Allele-specific expression reveals interactions between genetic variation and
environment. Nat. Methods. 14:699–702. doi: 10.1038/nmeth.4298.

Kopylova E, Noé L, Touzet H. 2012. SortMeRNA: Fast and accurate filtering of ribosomal RNAs in
metatranscriptomic data. Bioinformatics. 28:3211–3217. doi: 10.1093/bioinformatics/bts611.

Krantz DE, Zipursky SL. 1990. Drosophila chaoptin, a member of the leucine-rich repeat family, is a
photoreceptor cell-specific adhesion molecule. EMBO J. 9:1969–77.
http://www.ncbi.nlm.nih.gov/pubmed/2189727 (Accessed January 30, 2019).

Leon-Novelo L, Gerken AR, Graze RM, McIntyre LM, Marroni F. 2017. Direct Testing for Allele-

Specific Expression Differences Between Conditions. G3 GENES, GENOMES, Genet. 8:g3.300139.2017.
doi: 10.1534/g3.117.300139.

Leung V et al. 2016. The planar cell polarity protein Vangl2 is required for retinal axon guidance. Dev.
Neurobiol. 76:150–165. doi: 10.1002/dneu.22305.

Li H, Durbin R. 2009. Fast and accurate long-read alignment with Burrows-Wheeler transform.
Bioinformatics. 26:589–595. doi: 10.1093/bioinformatics/btp698.

Lischka S, Hagen W. 2005. Life histories of the copepods Pseudocalanus minutus, P. acuspes (Calanoida)
and Oithona similis (Cyclopoida) in the Arctic Kongsfjorden (Svalbard). Polar Biol. 28:910–921. doi:
10.1007/s00300-005-0017-1.

Liu Z, Dong X, Li Y. 2018. A Genome-Wide Study of Allele-Specific Expression in Colorectal Cancer.
Front. Genet. 9:1–11. doi: 10.3389/fgene.2018.00570.

Lonsdale Z et al. 2017. Allele specific expression and methylation in the bumblebee, *Bombus terrestris*.
PeerJ. 5:e3798. doi: 10.7717/peerj.3798.

Lu R et al. 2015. Analyzing allele specific RNA expression using mixture models. BMC Genomics.
16:566. doi: 10.1186/s12864-015-1749-0.

Luu K, Bazin E, Blum MGB. 2017. pcadapt: an R package to perform genome scans for selection based
on principal component analysis. Mol. Ecol. Resour. 17:67–77. doi: 10.1111/1755-0998.12592.

Madoui MA et al. 2017. New insights into global biogeography, population structure and natural selection
from the genome of the epipelagic copepod Oithona. Mol. Ecol. 26:4467–4482. doi: 10.1111/mec.14214.

Masai I, Okazakit A, Hosoyat T, Hottatt Y. 1993. Drosophila retinal degeneration A gene encodes an eye-
specific diacylglycerol kinase with cysteine-rich zinc-finger motifs and ankyrin repeats (signal
transduction/phosphatidylinositol   metabolism).   Neurobiology.   90:11157–11161.   doi:
10.1073/pnas.90.23.11157.

Matoulkova E, Michalova E, Vojtesek B, Hrstka R. 2012. The role of the 3′ untranslated region in post-
transcriptional regulation of protein expression in mammalian cells. RNA Biol. 9:563–576. doi:
10.4161/rna.20231.

Mayba O et al. 2014. MBASED: allele-specific expression detection in cancer tissues and cell lines.
Genome Biol. 15:405. doi: 10.1186/s13059-014-0405-3.

McKean DM et al. 2016. Loss of RNA expression and allele-specific expression associated with
congenital heart disease. Nat. Commun. 7:1–9. doi: 10.1038/ncomms12824.

Miao Z, Alvarez M, Pajukanta P, Ko A. 2018. ASElux: An ultra-fast and accurate allelic reads counter.
Bioinformatics. 34:1313–1320. doi: 10.1093/bioinformatics/btx762.

Mignone F, Gissi C, Liuni S, Pesole G, others. 2002. Untranslated regions of mRNAs. Genome Biol.
3:0004–1. doi: 10.1186/gb-2002-3-3-reviews0004.

Milani L et al. 2009. Allele-specific gene expression patterns in primary leukemic cells reveal regulation
of gene expression by CpG site methylation Allele-specific gene expression patterns in primary leukemic
cells reveal regulation of gene expression by CpG site methylation. Genome Res. 1–11. doi:
10.1101/gr.083931.108.

Monastirioti M. 1999. Biogenic amine systems in the fruit fly Drosophila melanogaster. Microsc. Res.
Tech. 45:106–121. http://onlinelibrary.wiley.com/doi/10.1002/(SICI)1097-
0029(19990415)45:2%3C106::AID-JEMT5%3E3.0.CO%5Cn2-3/pdf%5Cnpapers://5b5dd7df-cb43-4c9c-
aba9-19017de79ffc/Paper/p13846.

Moyerbrailean GA et al. 2016. High-throughput allele-specific expression across 250 environmental
conditions. Genome Res. 26:1627–1638. doi: 10.1101/gr.209759.116.

Myers PG, Donnelly C, Ribergaard MH. 2008. Structure and variability of the West Greenland Current in
Summer derived from 6 repeat standard sections. Prog. Oceanogr. doi: 10.1016/j.pocean.2008.12.003.

Nishida S. 1985. Taxonomy and distribution of the family Oithonidae (Copepoda, Cyclopoida) in the
Pacific and Indian Oceans. Bull. Ocean Res. Institute, Univ. Tokyo. 20:1–167.

Oksanen J et al. 2018. vegan: Community Ecology Package. https://cran.r-project.org/package=vegan.

Peijnenburg KTCA, Goetze E. 2013. High evolutionary potential of marine zooplankton. Ecol. Evol.
3:2765–2781. doi: 10.1002/ece3.644.

Pesant S et al. 2015. Open science resources for the discovery and analysis of Tara Oceans data. Sci. Data.
2:1–16. doi: 10.1038/sdata.2015.23.

Peterlongo P, Riou C, Drezen E, Lemaitre C. 2017. DiscoSnp++: de novo detection of small variants from
raw unassembled read set(s). bioRxiv. 209965. doi: 10.1101/209965.

Pirinen M et al. 2015. Assessing allele-specific expression across multiple tissues from RNA-seq read
data. Bioinformatics. 31:2497–2504. doi: 10.1093/bioinformatics/btv074.

R.C. L, J. K. 1973. Distribution of gene frequency as a test of the theory of the selective neutrality of
polymorphisms. Genetics. 74:175–195. doi: 10.1186/1475-925X-13-94.

Ram A, Jalal S, Jalal AS, Kumar M. 2010. A Density Based Algorithm for Discovering Density Varied
Clusters in Large Spatial Databases. Int. J. Comput. Appl. 3:1–4. doi: 10.5120/739-1038.

Rawls AS. 2003. Strabismus requires Flamingo and Prickle function to regulate tissue polarity in the
Drosophila eye. Development. 130:1877–1887. doi: 10.1242/dev.00411.

Riginos C, Crandall ED, Liggins L, Bongaerts P, Treml EA. 2016. Navigating the currents of seascape
genomics: How spatial analyses can augment population genomic studies. Curr. Zool. 62:581–601. doi:
10.1093/cz/zow067.

Rivas MA et al. 2015. Effect of predicted protein-truncating genetic variants on the human transcriptome.
Science (80-.). 348:666–669. doi: 10.1126/science.1261877.

Rose. 1933. Faune de France - Copépodes pélagiques. Lechevalier. Paris. www.faunedefrance.org.
132:767–767. doi: 10.1038/132767a0.

Shabalina SA, Spiridonov NA, Kashina A. 2013. Sounds of silence: Synonymous nucleotides as a key to
biological regulation and complexity. Nucleic Acids Res. 41:2073–2094. doi: 10.1093/nar/gks1205.

Skelly DA, Johansson M, Madeoy J, Wakefield J, Akey JM. 2011. A powerful and flexible statistical
framework for testing hypotheses of allele-specific gene expression from RNA-seq data. Genome Res.
21:1728–1737. doi: 10.1101/gr.119784.110.

Smarandache-Wellmann CR. 2016. Arthropod neurons and nervous system. Curr. Biol. 26:R960–R965.
doi: 10.1016/j.cub.2016.07.063.

Steige KA, Laenen B, Reimegård J, Scofield DG, Slotte T. 2017. Genomic analysis reveals major
determinants of cis-regulatory variation in Capsella grandiflora. Proc. Natl. Acad. Sci. U. S. A. 114:1087–
1092. doi: 10.1073/pnas.1612561114.

Stuart AE. 1999. From fruit flies to barnacles, histamine is the neurotransmitter of arthropod
photoreceptors. Neuron. 22:431–433. doi: 10.1016/S0896-6273(00)80699-6.

Supek F, Bošnjak M, Škunca N, Šmuc T. 2011. Revigo summarizes and visualizes long lists of gene
ontology terms. PLoS One. 6. doi: 10.1371/journal.pone.0021800.

Svensen C. 2000. Remote prey detection in Oithona similis: hydromechanical versus chemical cues. J.
Plankton Res. 22:1155–1166. doi: 10.1093/plankt/22.6.1155.

Szabo PE, Mann JR. 1995. Allele-specific expression and total expression levels of imprinted genes
during early mouse development: implications for imprinting mechanism. Genes Dev. 9:3097–3108. doi:
10.1101/gad.9.24.3097.

Tian L et al. 2018. Genome-wide comparison of allele-specific gene expression between African and
European populations. Hum. Mol. Genet. 27:1067–1077. doi: 10.1093/hmg/ddy027.

Tung J et al. 2015. Allele-specific gene expression in a wild nonhuman primate population. Mol. Ecol.
2:147–185. doi: 10.1111/j.1365-294X.2010.04970.x.

Uricaru R et al. 2014. Reference-free detection of isolated SNPs. Nucleic Acids Res. doi:
10.1093/nar/gku1187.

Wang M, Uebbing S, Ellegren H. 2017. Bayesian inference of allele-specific gene expression indicates
abundant Cis-regulatory variation in natural flycatcher populations. Genome Biol. Evol. 9:1266–1279.
doi: 10.1093/gbe/evx080.

Wang T, Montell C. 2007. Phototransduction and retinal degeneration in Drosophila. Pflugers Arch. Eur.
J. Physiol. 454:821–847. doi: 10.1007/s00424-007-0251-1.

Wassmann P et al. 2006. Food webs and carbon flux in the Barents Sea. Prog. Oceanogr. 71:232–287. doi:
10.1016/j.pocean.2006.10.003.

Wei X, Wang X. 2013. A computational workflow to identify allele-specific expression and epigenetic
modification in maize. Genomics, Proteomics Bioinforma. 11:247–252. doi: 10.1016/j.gpb.2013.05.006.

Weydmann A, Coelho NC, Serrão EA, Burzyński A, Pearson GA. 2016. Pan-Arctic population of the
keystone copepod Calanus glacialis. Polar Biol. 39:2311–2318. doi: 10.1007/s00300-016-1898-x.

Wright S. 1951. the Genetical Structure of Populations. Ann. Eugen. 15:323–354. doi: 10.1111/j.1469-
1809.1949.tb02451.x.

Zamora-Terol S, Nielsen TG, Saiz E. 2013. Plankton community structure and role of Oithona similis on
the western coast of Greenland during the winter-spring transition. Mar. Ecol. Prog. Ser. 483:85–102. doi:
10.3354/meps10288.

Zhang S et al. 2014. Genome-wide identification of allele-specific effects on gene expression for single
and multiple individuals. Gene. 533:366–373. doi: 10.1016/j.gene.2013.09.029.

Zhang X, Borevitz JO. 2009. Global analysis of allele-specific expression in Arabidopsis thaliana.
Genetics. 182:943–954. doi: 10.1534/genetics.109.103499.

Zhuo Z, Lamont SJ, Abasht B. 2017. RNA-Seq Analyses Identify Frequent Allele Specific Expression and
No Evidence of Genomic Imprinting in Specific Embryonic Tissues of Chicken. Sci. Rep. 7:1–10. doi:
10.1038/s41598-017-12179-9.

Tables

Table 1: Allele-specific expression detection and link with selection by population

Table 2: Functional annotations of variants targeted by ASE and selection implicated in nervous system

Supplementary Table S1: *Oithona similis* Mediterranean transcriptomes summary

Supplementary Table S2: *Tara* Oceans and *Oithona similis* Mediterranean transcriptomes samples
accession numbers

Supplementary Table S3: Variants dataset

Supplementary Table S4: Variants targeted by ASE and selection functional annotations

Supplementary Table S5: Variant annotation by SNPeff

Figures

Figure 1: Allele-specific expression at population-scale. a, Different scales of allele-specific expression
detection for a heterozygous gene, from population to cellular levels. For a heterozygous genotype, ASE is
understood as the difference in expression between to alleles of a single gene, opposed to strict biallelic
expression. For clarity, the example of ASE presented here is monoallelic expression. **b,** Null hypotheses
to detect psASE. On the upper panel, BAF and BARE are plotted for each variant (points). In grey, the
majority of variants, with allele relative expression following genomic frequency. In red (blue), variants
under potential psASE for allele A (B). The black dotted line stands for the theoretical regression curve
reflecting proportionality between BAF and BARE. On the lower panel, the deviation between BAF and
BARE is computed and should theoretically follow Gaussian distribution centered on 0. In the grey areas,
variants with no apparent deviation between BAF and BARE; in red (blue) area, variants in potential
psASE for allele A (B), with a significant difference between BAF and BARE.

Figure 2: Genomic variation data of *O. similis*. a, Scheme representing the study, from samplings to
analyses **b,** Genomic depth of coverage distributions of the set of 25,768 variants by sample **c,** F_{ST}
distribution across the seven samples **d,** LK distribution. The red line represents χ^2 theoretical distribution
(df = 6).

Figure 3: Population-scale allele-specific expression and its link with genomic differentiation. a, Each column corresponds to a population. Upper panel represents the relation between BAF and BARE, each hexagone corresponds to an area containing the number of variants indicated by the color scale. Black lines are the linear regression curves. Lower panel represents the deviation D distribution. The red lines correspond to the Gaussian distribution estimated from the data. **b,** Pairwise-F_{ST} matrix. The median (mean) of each pairwise-F_{ST} distribution computed on allele frequencies is indicated **c,** Upset plot of psASE detection in the seven populations. Each bar of the upper plot corresponds to the number of variants under psASE in the combination of population(s) indicated by black dots in the lower plot. **d,** Genomic differentiation and shared psASE. Each dot is a combination of population as presented in the lower panel of the upset plot. The blue line represents the non-linear regression curve estimated from the data and 95% confidence interval in grey.

Figure 4: Crossing psASE and selection. a, Scheme representing the method; in yellow the total dataset of SNVs, in green the tested SNVs in one population, in blue SNVs under psASE in this population, in red the SNVs targeted by selection. **b,** Crossing the candidate variants under psASE (blue) and those under selection (red) for each population. The hypergeometric p-value corresponds to the significance of the amount of variants both under psASE in the considered population and under selection (purple). **c,** Metagenomic and metatranscriptomic profiles of the SNV 1219525, under psASE in TARA_210 (*) and reaching near fixation in the other populations. BAF and BARE for each population in golden and blue respectively.

Supplementary Fig S1: Validation of taxonomic assignation

Supplementary Fig S2: *O. similis* depth of coverage of biallelic loci in seven *Tara* Oceans samples

Supplementary Fig S3: Population genomic differentiation

Supplementary Fig S4: Genomic differentiation and geographic distance

Supplementary Fig S5: Metagenomic and metatranscriptomic profiles of candidate loci

Supplementary Fig S6: Functional analysis of *O. similis* transcripts targeted by ASE and selection

696 **Table 1: Allele-specific expression detection and link with selection by population**. Adjusted-r² corresponds to linear regression performed
697 between BAF and BARE of all tested loci in the respective populations.

Population	Genomic median depth of coverage	Number of tested variants	Adjusted-r²	Number of variants under ASE	Number of variants under ASE and selection	Hypergeometric test p-value
TARA_155	25	18,812	0.77	91	6	9.89×10^{-3}*
TARA_158	35	21,476	0.91	131	29	5.06×10^{-20}*
TARA_178	24	18,145	0.67	26	9	5×10^{-11}*
TARA_206	55	22,578	0.95	162	42	4.82×10^{-31}*
TARA_208	48	21,469	0.93	133	14	2.2×10^{-6}*
TARA_209	12	13,956	0.73	62	24	1.05×10^{-23}*
TARA_210	14	13,454	0.71	69	42	8.89×10^{-51}*
Overall	-	25,768	-	587 (2.3%)	152 (0.59%)	-

701   **Table 2: Functional annotations of variants targeted by ASE and selection implicated in nervous system.**

VarID	Ref	Alt	Homology search	Pfam	SnpEff Localization	SnpEff Impact	References
722267	A	G	histamine H1 receptor	PF00001	3 ' UTR	MODIFIER	(Monastirioti 1999; Stuart 1999; Denno et al. 2016)
9665345	T	G	chaoptin	PF13306 | PF13855	synonymous variant	LOW	(Krantz & Zipursky 1990; Gurudev et al. 2014)
15623788	G	A	eye-specific diacylglycerol kinase	PF13637	synonymous variant	LOW	(Masai et al. 1993; Wang & Montell 2007)
23795359	A	T	vang-like protein 2-B	PF06638	synonymous variant	LOW	(Rawls 2003; Leung et al. 2016)
1276227	C	T	glycine receptor subunit alpha-2 / gamma-aminobutyric acid receptor subunit alpha-6	PF02932 | PF2931	3 ' UTR	MODIFIER	-
1404415	G	C	omega-amidase NIT2	PF00795	3 ' UTR	MODIFIER	-
11174785	A	G	5-oxoprolinase	PF02538 | PF05378 | PF01968	5 ' UTR	MODIFIER	-
11690229	A	T	glycine receptor subunit alpha-2	PF02931	synonymous variant	LOW	-

b

Genomic depth of coverage

TARA_155 TARA_158 TARA_178 TARA_206 TARA_208 TARA_209 TARA_210

Populations

c

count

4000

3000

2000

1000

Hartigan's Dip-Test
D=0.0012, p-value=0.19

0.00   0.25   0.50   0.75   1.00

F_{ST}

d

density

0.15

0.10

0.05

0.00

χ2 expected distribution
(df=6)

0   10   20   30   40

LK

a

5°55'0"E   5°0'0"E

Toulon harbour

● Sampling site

Little bay   Large bay

43°5'0"N

Mediterranean Sea

2 km

Sampling
Taxonomic sorting

Males   Juveniles

mRNA extraction
Sequencing
Assembly
Annotation

TARA_178
TARA_206
TARA_208   TARA_155
TARA_209
TARA_210

☐ Northern Atlantic
☐ Baffin Bay
☐ Kara Sea
☐ Labrador Sea

TARA Ocean sampling

Metatranscriptomic reads   Metagenomic reads

Variant calling
Clustering
Filtering

25,768
Variant sequences

Mapping
(97% variant sequences)

Transcriptomes

Populations   Populations

Variants

B-allele
relative
expression

B-allele
frequency

Population-scale
allele-specific expression

Selection   Genomic differentiation   Functional annotation

a

Heterozygous variants
in population TARA_X

All loci

psASE

selection

b

TARA_155
91   674
6*
*p-value = 9.89 x 10⁻³

TARA_158
131   674
29*
*p-value = 5.06 x 10⁻²⁰

TARA_178
26   674
9*
*p-value = 5.00 x 10⁻¹¹

TARA_206
162   674
42*
*p-value = 4.82 x 10⁻³¹

TARA_208
133   674
14*
*p-value = 2.2 x 10⁻⁶

TARA_209
62   674
24*
*p-value = 1.05 x 10⁻²³

TARA_210
69   674
42*
*p-value = 8.89 x 10⁻⁵¹

c

*

Frequency

BAF
BARE

TARA_155
TARA_158
TARA_178
TARA_206
TARA_208
TARA_209
TARA_210

Populations