A decrease in transcription capacity limits growth rate upon translation inhibition

Qing Zhang[1,#a], Elisa Brambilla[1], Rui Li[2], Hualin Shi[2,3], Marco Cosentino Lagomarsino[4,#b] and Bianca Sclavi[1,#c*].

1. LBPA, UMR 8113 CNRS, ENS Paris-Saclay, 94235 Cachan, France

2. CAS Key Laboratory of Theoretical Physics, Institute of Theoretical Physics, Chinese Academy of Sciences, Beijing 100190, China

3. School of Physical Sciences, University of Chinese Academy of Sciences, Beijing, China

4. LCQB, UMR 7238 CNRS, Sorbonne Université, 75005 Paris, France

Current affiliations:

[#a] Current Address: CAS Key Laboratory of Theoretical Physics, Institute of Theoretical Physics, Chinese Academy of Sciences, Beijing 100190, China

[#b] Current Address: Dipartimento di Fisica, Università degli Studi di Milano, 20133 Milano and INFN, Italy, and IFOM, FIRC Institute for Molecular Oncology, Milan, Italy

[#c] Current Address : LCQB, UMR 7238 CNRS, Sorbonne Université, Paris, France

* Corresponding author

E-mail : bianca.sclavi@sorbonne-universite.fr

Abstract

In bacterial cells, inhibition of ribosomes by sublethal concentrations of antibiotics leads to a decrease in growth rate despite an increase in ribosome content. The limitation of ribosomal activity results in an increase in the level of expression from ribosomal promoters; this can deplete the pool of RNA polymerase (RNAP) that is available for the expression of non-ribosomal genes. However, the magnitude of this effect remains to be quantified. Here, we use the change in the activity of constitutive promoters with different affinities for RNAP to quantify the change in the concentration of free RNAP. The data are consistent with a significant decrease in the amount of RNAP available for transcription of both ribosomal and non ribosomal genes. Results obtained with different reporter genes reveal an mRNA length dependence on the amount of full-length translated protein, consistent with the decrease in ribosome processivity affecting more strongly the translation of longer genes. The genes coding for the β and β' subunits of RNAP are amongst the longest genes in the *E. coli* genome, while the genes coding for ribosomal proteins are among the shortest genes. This can explain the observed decrease in transcription capacity that favors the expression of genes whose promoters have a high affinity for RNAP, such as ribosomal promoters.

Importance

Exposure of bacteria to sublethal concentrations of antibiotics can lead to bacterial adaptation and survival at higher doses of inhibitors, which in turn can lead to the emergence of antibiotic resistance. The presence of sublethal concentrations of antibiotics targeting translation results in an increase in the amount of ribosomes per cell and a decrease in the cells' growth rate. In this work, we have found that inhibition of ribosome activity can result in a decrease in the amount of free RNA polymerase available for transcription, thus limiting the protein expression

46 rate via a different pathway than what was expected. This result can be explained by our

47 observation that long genes, such as those coding for RNA polymerase subunits, have a higher

48 probability of premature translation termination in the presence of ribosome inhibitors, while

49 expression of short ribosomal genes is affected less, consistent with their increased

50 concentration.

Introduction

Bacteria often encounter sub-lethal levels of antibiotics produced by other microorganisms in

their environment. A decrease in growth rate in these conditions can allow a strain to survive

long enough until the inhibitor is no longer present or, in some cases, until the bacteria becomes

resistant to the antibiotic via the selection of pre-existing mutations or an increase in mutation

rates (1–4). However, the mechanistic details of these response pathways often remain to be

described. The cellular response to the limitation of translation activity is thought to be related

to the pathway involved in the stringent response, the regulatory mechanism that decreases

ribosome production in response to a decrease in amino acid availability. This is mediated by

the change in concentration of the secondary messenger molecule, (p)ppGpp, that is produced

by the RelA enzyme when the pool of amino acids decreases and ribosomes are not loaded with

charged tRNAs (5, 6). ppGpp can directly inhibit ribosome assembly (7, 8) and the activity of

RNA polymerase (RNAP) at ribosomal promoters, while increasing the activity of the promoters

of genes for amino acid biosynthesis (9). The transcription of ribosomal operons can use a large

fraction of the free RNA polymerase pool in the cell because of the high affinity of the ribosomal

promoters for the enzyme and a high frequency of transcription initiation. Therefore, the

regulation of ribosomal promoter activity can be a means by which the pool of free RNA

polymerase can be repartitioned between ribosomal operon transcription and non-ribosomal

mRNA synthesis. This has been referred to as the "passive control" of transcription regulation

(10–14).

The ppGpp dependent feedback loop also plays a role in the regulation of ribosome content as

a function of growth rate (15–17). Growth rate dependent regulation of gene expression

determines the allocation of cellular resources between the production of ribosomes and that

of other proteins and results in a linear increase in ribosome content with increasing growth

rate (15, 18–20). In richer growth media, when the amount of amino acids is higher, ppGpp

levels are lower, favoring ribosome production and a higher fraction of active ribosomes (16).

In poorer media it is the inverse, accumulation of ppGpp slows down the production of new

ribosomes and a smaller fraction of the ribosome pool is in an active form (19).

When ribosome activity is inhibited by sublethal concentrations of antibiotics, amino acids are

used more slowly and their concentration increases, which can result in a decrease in the

intracellular ppGpp pool (5). The cellular response, as predicted by the ppGpp feedback loop, is

to produce a higher amount of ribosomes and an increased translation rate, however, despite

this increase, the cell's growth rate is reduced (18, 19). This has been proposed to result from a

decrease in the resources available for the production of non-ribosomal proteins that become

limiting for cellular metabolism (18). More recent results point to a decrease in the fraction of

active ribosomes to explain the decrease in the total protein production rate (19).

To measure the effect that the inhibition of ribosome activity can have on gene expression

resulting from a possible repartition of RNAP, we have compared the activity of a ribosomal

promoter to that of constitutive promoters with different affinities for RNAP. This approach

stems from a well-established protocol developed by Hans Bremer and coworkers of using

quantitative measurements of changes in constitutive and ribosomal promoter activity as

reporters of changes in the amount of free RNAP and of ppGpp (21–24).

In parallel, we analyzed transcriptomics and proteomics data from the literature on the direct

and indirect effects of changing ppGpp concentration and translation limitation on gene

expression (25, 26). The results from this analysis are consistent with a linear decrease with

decreasing growth rate in the concentration of free RNAP available for promoter binding and

transcription. We propose a model that can explain this decrease in transcriptional activity

based on the observation that gene length is an important parameter on the change in protein

expression in the presence of sublethal levels of chloramphenicol and that RNAP contains two

of the longest gene products in the *E. coli* genome.

Results

Transcription regulation by ppGpp does not suffice to explain gene expression changes with

increasing translation inhibition by chloramphenicol. In order to measure the effect of

increasing ribosome inhibition on RNAP repartition between ribosomal and non-ribosomal

promoters we have chosen three reporter cassettes. The first contains a shortened version of

the well-characterized ribosomal RNA operon promoter *rrnB*P1, here called P1, that terminates

at -69 from the transcription start site (27). The binding sites for Fis and the higher affinity H-NS

binding site are thus omitted from this construct. This promoter has a GC-rich discriminator

region at the transcription initiation site that makes the open complex sensitive to changes in

negative supercoiling and to inhibition by ppGpp (Table S1) (23, 28, 29). The second promoter

used here is a constitutive promoter, P5, that has consensus -10 and -35 sequences and no

discriminator region. The third is PLtet, also a strong constitutive promoter with no

discriminator region but with a lower affinity for RNA polymerase due to a non-consensus -10

sequence (30). Bremer and coworkers have shown that the activity of the *rrnB*P1 promoter is

proportional to the concentration of ppGpp *in vivo* (23), and that the activity of constitutive

promoters can be used to estimate the amount of free RNA polymerase in the cell (21, 23). Each

of these promoters was placed upstream of the *gfpmut2* gene and this cassette was inserted in

the chromosome together with a kanamycin resistance gene expressed divergently from the

chosen promoter (Fig. S1). Growth of these strains in a 96-well plate allowed us to measure the

changes in growth rate, the GFP concentration and the resulting GFP production rate (Gpr) as a

function of chloramphenicol concentration (Figs. 1 and S2). We compared four different growth

media, M9 with glucose, M9 with glycerol and these two media supplemented with casamino

acids (cAA). This results in four different growth rates. Furthermore, it has been already shown

that cells growing in a growth medium containing amino acids have a lower concentration of

ppGpp (16, 23), allowing us to compare the effects due to changing concentrations of this key

metabolite without the use of mutant strains that can result in secondary effects on cell

metabolism due to the multiple targets of ppGpp (8, 31).

Figure 1. Promoters with different affinities for RNAP and different regulation by ppGpp react differently to translation limitation. A. Change in GFP concentration (RFU GFP/OD$_{600}$) as a function of growth rate. Four growth media were used, from the slowest to the fastest: M9-glycerol, M9-glucose, M9-glycerol+casaminoacids, M9-glucose+casaminoacids. P5 and PLtet are both constitutive promoters with different affinities for RNAP while P1 is a shortened version of the rrnBP1 ribosomal RNA promoter with a RNAP affinity similar to P5 but regulated by ppGpp. **B.** Change in GFP concentration as a function of increasing concentration of chloramphenicol in the four growth media. The growth media with casamino acids are in red the ones without casamino acids are in green. The

In the absence of translation inhibition, the change in promoter activity measured as a function

of growth rate is consistent with previous measurements on constitutive promoters and *rrnB*P1

derived promoters (11, 27, 32) (Fig. 1A). The concentration of GFP from the constitutive

promoters tends to decrease at the faster growth rates due to their lack of specific growth rate

dependent regulation and the increased dilution rate (33), while the concentration of GFP

expressed from the rrnBP1 promoter increases with growth rate until the last point, where Fis

activation, absent in this construct, has been shown to be required for continued increased

expression (27). The PLtet promoter has a lower affinity for RNAP than P5 does (see below),

however when RNAP binds at the PLtet promoter it initiates transcription with a higher

frequency than at P5 (30), resulting in a higher promoter activity (Fig. S2) and consequently a

higher GFP concentration (Fig. 1A).

Previous work has shown that as the concentration of chloramphenicol is increased, the total

RNA content relative to the total protein mass increases -reflecting the increase in ribosomal

RNA- and the concentration of a reporter protein expressed from a constitutive promoter

decreases (18). Therefore, the expected result here is that the GFP concentration from a

ribosomal RNA promoter (P1) should increase while the concentration from a constitutive

promoter (P5 or PLtet) should decrease, with the lower affinity constitutive promoter (PLtet)

decreasing at a faster rate if increasing amounts of RNAP are being used for transcription of

ribosomal operons. The comparison of P1 and PLtet agrees with this prediction. Unexpectedly,

however, the patterns of the change in GFP production rate and GFP concentration for the P1

and P5 promoters are very similar (Figs 1 and S2). The pattern of the change of expression for

these two promoter constructs depends strongly on whether the growth medium contains

casamino acids, independently of the carbon source. In the absence of cAA the increase in GFP

concentration as a function of Cm is significantly greater than in their presence (Fig. 1 and Fig

S2). Increased expression from the P1 promoter in the growth media lacking cAA would be

expected from a decrease in ppGpp as a result of increased amino acid pools due to ribosome

inhibition; the P5 promoter on the other hand does not contain the GC-rich discriminator region

and is not expected to show increased activity upon a decrease in ppGpp. The similar increase

in GFP concentration of these two different promoters thus points to a stronger effect of ppGpp

on GFP expression at the post-transcriptional level.

Decrease in free RNAP concentration with increasing translation inhibition by

chloramphenicol. Since the translation rate of GFP is shared by the three promoter constructs,

it is possible to obtain an estimate of the magnitude of the promoter-specific effect of ppGpp,

and of changes in free RNAP, on the transcription rate by measuring the ratios of GFP production

rates. This operation "cancels out" the translation component of gene expression and isolates

the transcription-specific effect (see SI text and legend to Fig. 2). Fig. 2A shows the change in

the ratios of GFP production rates as a function of growth rate in the absence of

chloramphenicol. The ratio of P1 to P5 rates increases rapidly between M9-glu and M9-cAA-gly,

consistent with a lower level of ppGpp in the cells growing in the presence of cAA (23) increasing

the probability of transcription initiation specifically from P1.

In the presence of Cm, the fold increase of gene expression from P1 is greater than the one of

P5 in the cells that are grown without cAA, consistent with a decrease in ppGpp levels by the

addition of the antibiotic (Fig. 2C). As the Cm concentration is increased further, the difference

between the two promoters decreases again to the initial level. On the other hand, in the

growth media with cAA, and thus lower levels of ppGpp, the P1 to P5 ratio decreases, indicating

that the change in GFP production rate from P1 is lower than that of P5 as a function of

increasing Cm. A similar result is also observed for the PLtet to P5 ratio, independently of the

growth medium (Fig. 2B).

The comparison of two promoters with differing affinities for RNAP can be used to estimate the

change in the amount of free RNAP that is available for transcription (23, 34). In the absence of

Cm, the ratio of PLtet to P5 GFP production rate increases linearly with increasing growth rate

(Fig. 2A), in agreement with previous estimates of the change in free RNAP as a function of

doubling time (22, 33).

Figure 2. Ratios of GFP production rate for the different promoters can be used to estimate the changes in the
concentration of free RNAP. A. Ratio of PLtet to P5 and P1 to P5 as a function of growth rate. **B.** Ratio of PLtet to
P5 with increasing chloramphenicol concentration. **C.** Ratio of P1 to P5 with increasing chloramphenicol
concentration. **D.** Estimated decrease in the concentration of free RNAP from the change in the ratios of promoter

binding as a function of chloramphenicol concentration. To obtain this estimate, the relative activity of the promoters as a function of free RNAP concentration (c_f) was obtained from the respective RNAP binding constants, $K_i \in \{K_1, K_5, K_{Ltet}\}$ using $c_f/(K_i + c_f)$. The constant K_5 is known, K_1 and K_{Ltet} were derived by the formula $K_5 \exp(\Delta E)$, where ΔE is the difference between the binding energies of RNAP with P1 (or PLtet) and P5 (see Supporting Information). The black vertical bars indicate the concentrations of RNAP that give the measured GFP production rate ratios shown in (B) for different Cm concentrations, shown above the plot. The ratio of PLtet and P5 activities for example is given by: $\mathrm{Gpr(PLtet)/Gpr(P5)} = a(K_5 + c_f)/(K_{Ltet} + c_f)$, where a is a scaling factor. K_5 and a were fixed by fitting the data in absence of Cm from our experiments and the literature. Data from the cells growing in cAA-glu was used so that ppGpp-dependent regulation of P1 is small and can be ignored. Fig. S3 in Supplementary Materials shows the estimation of RNAP concentration for the four growth media used. Fig.S4 shows an estimation of the change in ppGpp as a function of chloramphenicol concentration. See details of the analysis in the Supporting Information.

From the values of the RNAP affinities for these two promoters, PLtet and P5, and the relative changes in GFP production rate it is possible to estimate the change in the amount of free RNAP in the cells as a function of increasing translation limitation (see SI text). Fig. 2D and Fig. S3 illustrate this point. The RNAP binding affinity for each promoter was estimated based on a statistical-mechanical selection model developed by Berg and von Hippel (35) (see SI text). Fig. 2D shows the estimated binding curves for RNAP to each promoter; the vertical bars indicate the ratio of promoter activity corresponding to changes in the ratio measured experimentally, which were used to estimate the RNAP concentration. The values of the Cm concentration for each of these ratios is shown on the top axis. The decrease in free RNAP with increasing chloramphenicol has a stronger effect on PLtet first, then on P1 and finally on P5. This simple model of RNAP-dependent capacity can explain the data in Fig. 2B on the change in the PLtet to P5 ratio, while the P1 to P5 ratio is also influenced by changes in ppGpp concentration. Thus, using the P1 to P5 GFP production rate ratio it is possible to estimate the change in ppGpp as a function of growth rate and as a function of Cm (Fig. S4), in a similar fashion to the approach validated by Bremer and colleagues (23).

In summary, the transcriptional changes measured by the decrease in the PLtet to P5 and P1 to

P5 ratios, which are in line with a decrease in free RNAP concentration, must be independent

of a ppGpp-mediated repartition between ribosomal and non-ribosomal promoters, since they

can also be observed in the growth media with cAA, where ppGpp levels have been previously

shown to be very low (16).

The decrease in ribosome processivity by chloramphenicol reduces the expression of longer

genes more than shorter ones. In addition to the effect of RNAP availability, we hypothesized

that the results observed might be dependent on the reporter gene used -*gfpmut2*, 714 bp long,

or *lacZ*, 3072 bp long, coding for the β-galactosidase enzyme - since translation of longer genes

would have a higher probability to end prematurely in the presence of ribosome inhibitors such

as chloramphenicol acting during elongation.

Figure 3. Gene length can influence gene expression under translation limitation. A. Change in GFP (238 aa) and
β-galactosidase (1024 aa) expressed from the P5 promoter as a function of increasing chloramphenicol
concentrations. The bacteria were grown in M9 glu+cAA with chloramphenicol (Cm) at concentrations of 0, 2 μM,
4 μM , or 6 μM in flasks. The concentrations were normalized by dividing the data by the point without Cm. **B.** Fit
of the model to the ratio of GFP to β-galactosidase concentrations from the data in (A). The growth rate at each
chloramphenicol concentration is the average over the two strains. The error bars correspond to the SEM from
three independent experiments.

In order to produce an expectation, we reasoned as follows. The processivity of translation has

been shown to decrease exponentially with increasing gene length (36). If, in addition, ribosome

processivity were decreased by an inhibitor, then the probability to finish the translation of a

long gene would be lower compared to a shorter gene, decreasing the rate of expression of the

longer gene to a greater extent. To test this hypothesis, we compared the changes in gene

expression from the same constitutive promoter, P5, of two different genes, *gfpmut2*, and *lacZ*,

in the presence of increasing chloramphenicol concentrations (Fig. 3A). The results show that

while the concentration of GFP increases as a function of Cm concentration, β-galactosidase

concentration decreases (Fig. 3A). Fig. 3B shows the change in the ratio of the shorter to the

longer protein as a function of Cm concentration. The black line shows the fit obtained to a

model of ribosome processivity. In the model, we derived the expression ratio of a short gene

to a long gene based on the scenario where chloramphenicol hits a translating ribosome,

causing the stalling of this and the following ribosomes and leading to the nonsymmetric

degradation of mRNA (as depicted by Dai et al. (19), see details in SI text). It is possible to

estimate the probability that a ribosome will be inhibited by the antibiotic before it reaches the

end of the mRNA, P_{hit} (19). The equation can be extended to describe the dependence of P_{hit}

on protein length, i.e.

$$P_{hit} = 1 - \exp(-k_{on}[Cm]L/v) \quad (1)$$

where k_{on} denotes the binding constant of Cm with ribosome ($k_{on} = 0.034(\mu M \cdot min)^{-1}$

(37)), L indicates protein length, and v is the translation elongation rate dependent on the

RNA/protein mass ratio (see SI text). Using equation 1, at 8 μM Cm (the highest concentration

used here) P_{hit} is 23% for LacZ (1024 aa), while for GFP it is 6% (238 aa). These results therefore

indicate that a gene's length, in addition to its promoter's affinity for RNAP, can influence how

its expression levels change in the presence of sub-lethal concentrations of ribosome inhibitors.

A possible cause for the decrease in the pool of free RNAP independently of the changes in

ppGpp is a decrease in the total amount of RNAP per cell. The RNA polymerase holoenzyme is

composed of 5 subunits: β, β', ω, 2 subunits of α, and a σ factor. β, β' are among the longest

genes in *E. coli* with a length of 4029 bp and 4224 bp respectively (the average gene length in *E.*

coli is about 900 bp) and they could be subject to the length effect we found in our reporters.

A decrease in translation processivity can result in decreased expression of late operon genes.

Transcription and translation are coupled via a physical interaction between RNA polymerase

and the first ribosome translating the mRNA, which can be mediated by NusG or RfaH (38–41).

Therefore, an additional factor that could decrease the expression of longer genes, and of late

genes within an operon, in the presence of translation inhibitors is the loss of this RNAP-

ribosome interaction exposing the mRNA for degradation and/or decreasing RNAP speed and

processivity (42).

Previous work has shown that inhibiting ribosome activity with a higher concentration of

antibiotics than was used here can result in decoupling of transcription and translation (42). To

test whether these sublethal concentrations of Cm could have a similar effect on the RNAP-

ribosome interaction, constructs were made where two genes of equal length coding for a red

and a green fluorescent protein are placed one after the other within the same operon (Fig. 4A).

In this case, there is an equal probability that a ribosome will stall during translation of either

gene. However, if the first ribosome translating the upstream gene is inhibited, the one in

contact with RNAP, it will also decrease the probability that the downstream gene will be

transcribed by affecting the stability of the mRNA and of the transcription complex. Indeed, in

these constructs we observe a decrease in the GFP to RFP ratio with increasing Cm. This could

be either due to a decrease in transcription processivity from the loss of the RNAP-ribosome

interaction or to an increase in the probability that the operon mRNA is degraded before RNAP

finishes GFP transcription due to early termination of RFP translation, or a combination of both.

The formation of a terminator hairpin in the transcribed mRNA can be used to detect the

presence of a ribosome-RNAP interaction (43). If a ribosome is bound on the mRNA as it is being

extruded from RNAP, the terminator hairpin structure cannot fold. If inhibition of ribosome

activity decouples translation from the ongoing transcription, then the RNA will be allowed to

fold and transcription will stop. Transcription terminators of different strength have been

inserted between the two genes (43). The efficiency of these terminators is determined by the

distance between the stop codon of the upstream gene and the hairpin loop forming sequence.

Termination will be less efficient when this distance is shorter. If transcription and translation

are decoupled, the RNAP to ribosome distance will increase and we expect that the efficiency

of the weaker terminators will increase with increasing Cm. However, we observe the same

fractional decrease in GFP/RFP in all the constructs, independently of the distance from the stop

codon, or of the presence of the terminator (Fig. 4 and S8). Therefore, it seems that at these

low Cm concentrations, the probability of decoupling of transcription and translation is not

significant enough to allow for the folding of the hairpin loop. The stalling of a ribosome

however, independently of its interaction with RNAP, can result in an increased probability of

mRNA degradation. In this case, the stalling of non-leading ribosomes, which are in greater

number than the one interacting with RNAP and thus a more probable target, can decrease the

lifetime of the operon's mRNA, decreasing the probability that translation will be completed at

the *gfp* gene. These results can be reproduced by a model using the parameters values for the

probability of translation termination obtained from the comparison in GFP vs LacZ translation

(Fig. 3) and the terminator strengths measured in the previous study (43) (Fig. 4B) (see SI for the

details of the model).

Figure 4. Operon position can influence gene expression under translation limitation. (**A**) Sketch for GFP and RFP
fused in an operon with an intergenic terminator of efficiency y (see the details of the construction in (43)) and the
formula of the expression ratio of GFP to RFP. When the efficiency y is zero, it corresponds to the case without
intergenic terminator (see no ter in (B)). (**B**) The expression ratio between downstream GFP and upstream RFP (in
the units of RFU(GFP)/RFU(RFP)), as a function of growth rate decreases with increasing Cm concentration. The
experimental ratios (symbols) are fit with the model (lines) on Rb-stalling-induced mRNA degradation (See
Supplementary Materials). `No ter' and `ter 1-4' correspond to terminator sequences of increasing efficiency (`I21'
(no terminator), `R9', `R17', `W13' and `R32' respectively in (43)). The cells were grown in M9 minimal medium
containing glucose. Cm was added to a final concentration of 0, 2, or 4 µM.

Finally, in light of these results we have analyzed data from previously published transcriptomics

and proteomics studies as a function of the presence of ppGpp and at increasing sublethal

chloramphenicol concentration respectively (25, 31) and we have found that the effect of

promoter affinity, gene length and operon position observed here on this set of promoters and

reporter proteins can help explain the global changes in gene expression in the presence of

sublethal levels of chloramphenicol (SI text and SI Figs S10 - S14).

Discussion

A linear decrease in transcription capacity with increased translation limitation. While the amount of free RNAP decreases with decreasing growth rate as a function of nutrient content (Fig. 2A), the decrease in the presence of increasing translation limitation, within the same growth medium, has a steeper, linear, slope (Fig. 2B and Fig. S3). The evidence provided here points to a possible cellular adaptation mechanism leading to a reduction in transcription capacity when ribosome activity is compromised. This adaptation decreases the cost of transcription of untranslated mRNAs (44) and allows for more resources to be available for the synthesis of increased amounts of ribosomes to respond to the presence of translation inhibitors. Importantly, we can quantify the contribution of the change in free RNAP to changes in transcription capacity: the comparison of two different constitutive promoters with differing RNAP affinity, PLtet and P5, shows a striking difference in the change in GFP production rate and the resulting GFP concentration with increasing ribosome inhibition (Fig. 1). The decrease in the amount of free RNAP estimated by measuring the ratio of GFP production rates from these two promoters (PLtet/P5) is about 10-fold, independently of the presence of amino acids in the growth medium (Fig. 2 and Fig. S3) and therefore of the change in ppGpp concentration (see below).

We found that, as the sublethal concentration of ribosome inhibitor is increased, the growth rate decreases linearly with the decrease in the concentration of free RNA polymerase, pointing to a possible growth-limiting role for this enzyme in these conditions (Fig. 2B). The decrease in free RNAP could be due to different factors affecting the nonspecific interactions of the enzyme with the genome (22), however, the results obtained here from the comparison of the expression of two proteins of different lengths, β-gal and GFP (Fig. 3) suggest that a decrease in

the amounts of the full length protein may have a significant contribution to this effect. The

RNA polymerase core contains two of the longest proteins in *E. coli*, the β and β' subunits (Fig.

S12), increasing the probability that a ribosome will stall before reaching the end of the mRNA.

This interpretation is further supported by the proteomics analysis of Hui *et al*. (25). They

measured the change in protein fraction of over 1000 proteins in the presence of increasing

concentrations of Cm by quantitative mass spectrometry. Their results show that the β and β'

subunits of RNAP remain a constant fraction of the proteome with increasing Cm and decreasing

growth rate. Since decreasing growth rate is associated with decreased protein production rate,

and in these conditions ribosomes are in excess, these results are in line with RNAP playing a

limiting role in determining the total rate of protein production and the cell's growth rate.

Moreover, these results can shed light on a recent study by Dai et al (19) where it was proposed

that in the presence of sublethal concentrations of Cm, despite an increase in the translation

elongation rate due to a higher concentration of ternary complexes, the reduction in the total

protein production rate results from a decrease in the active ribosome fraction, or the fraction

of ribosomes that can reach the end of a mRNA in the presence of the inhibitor (19). Here we

identify RNAP as one of the genes that is likely to be most affected by the decrease in ribosome

processivity due to the length of its β and β' subunits, while shorter genes are affected to a

lesser extent.

Is the extreme length of RNA polymerase genes a feature conserved for the coupling of

translation and transcription rates? The extreme length of the β and β' subunits of RNAP is

conserved throughout bacteria (45). In the case of *Helicobacteraceae* and *Wolbachia* the two

genes are even fused together (45, 46). The gene length of β and β' subunits can vary in different

strains since they are composed of independent structural modules separated by spacers of

differing length (45). In *E. coli* the spacer sequences, that account for more than 25% of the total

sequence, can be deleted without causing a significant decrease in transcription activity. In

archea and chloroplasts some of the conserved protein modules are found in separate genes

and in *E. coli* they can be split from each other to produce an active enzyme (47), suggesting

that the length of these genes is not imposed by functional constraints. The reason why the

RNAP and ribosomal proteins find themselves at opposite ends of the spectrum of gene lengths

in bacteria is likely linked to the assembly process, structural flexibility and stability of the final

multi-protein complex (48, 49); however, these results suggest that it could also play an

important role for the cell's survival in the presence of antibiotics that decrease translation

processivity.

The passive control model. Another factor that can decrease the pool of RNAP available for

transcription of non-ribosomal genes is the increase in the transcription from ribosomal

promoters resulting from the decrease in ppGpp, the inverse of what takes place during the

stringent response and what has been referred to as the "passive control" of transcription

regulation (10–14). However, the results presented here show a similar slope of the decrease

in the concentration of free RNAP (Fig. 2 and S3), independently of the initial amounts of ppGpp

found in the cells growing in the presence or absence of casamino acids (23). Therefore, while

changes in ppGpp do affect the activity of ribosomal promoters, the magnitude of this effect is

not strong enough to further decrease the activity of non-ribosomal promoters such as PLtet

and P5. The main factor that partitions the limited pool of RNAP between ribosomal and non-

ribosomal promoters is therefore the affinity of the specific RNAP-promoter complex formed

for transcription initiation. These results can also help explain a previous report showing that

the induction of persistence and of β-lactam tolerance by chloramphenicol is independent of

the presence of the RelA enzyme, that, together with SpoT, is responsible for ppGpp synthesis

(50).

Change in total translation rate in response to translation limiting inhibitors. The similarity in

the pattern of the change in GFP expression from the rrnBP1 and P5 promoters despite their

different regulation by ppGpp indicates that upon translation limitation the main effect of

ppGpp on gene expression may be via a change in the translation rate rather than the

transcription rate (Fig. 1 and Fig. S15). The translation of short proteins such as GFP is barely

affected by the presence of sublethal concentrations of Cm (8 % probability of drop off at the

highest concentration used here) and can therefore be used to estimate the changes in total

translation activity and to estimate how the translation capacity is affected by chloramphenicol

challenges (Fig. S15). As the concentration of ribosome inhibitor is increased, the translation

rate of GFP increases and mirrors the change of the effect of ppGpp on the P1 promoter (Fig.

S15 and S4A respectively). Several factors can contribute to this increase in translation rate in

the presence of Cm: the shorter than average length of ribosomal genes (Fig. S12), whose

translation is not affected as strongly by translation inhibition compared to longer genes; the

high affinity of RNAP for ribosomal promoters, insuring transcription despite a decrease in the

concentration of free RNAP (Fig. S11); the decrease in ppGpp in the growth media lacking

casamino acids (Fig. S4), resulting in an increase in both ribosomal promoter activity (Fig. 1) (5,

51) and in the fraction of active ribosomes due to the effect of ppGpp on ribosome assembly (7,

8, 19), in addition to the increase in translation elongation rate (19). The higher increase in

translation rate in the growth media without cAA is consistent with a larger fraction of inactive

ribosomes stored in the cells growing at these slower growth rates that can be quickly

reactivated by a decrease in ppGpp concentration, not only for rapid adaptation to changes in

local nutrient content, but also to respond to the presence of growth inhibitors (19, 52–55). On

the other hand, in rich media, when ppGpp levels are low, a decrease in transcription rate when

translation is compromised might play a crucial role for the cell's survival, as the potential of the

cell to increase its translation capacity to respond to the presence of the inhibitor is limited by

the smaller fraction of inactive ribosomes (19, 56). Decoupling of transcription and translation

can have several deleterious effects, including mRNA degradation and R-loop accumulation that

can interfere with DNA replication, causing genome instability and increased mutation rates

(57).

The decrease in growth rate due to limiting transcription is unexpected, as ribosome activity is

usually thought to always be rate-limiting for bacterial growth, however, depending on the

growth conditions, transcription has also been seen to become limiting in eukaryotic cells (68),

pointing to different strategies of cellular adaptation to changing growth conditions and

limitations. Understanding how bacteria modulate their growth rate and resource allocation in

response to inhibition of growth has paramount importance in biotechnological and health

applications (58–61). The results presented here provide a new cellular mechanism by which

bacterial cells can decrease their growth rate in response to antibiotic stress (1, 56, 62). In

summary, in the presence of sublethal concentrations of chloramphenicol, it is not translation

that becomes limiting for the cell's growth rate, or the ppGpp dependent repartition of RNAP

between ribosomal and non-ribosomal promoters, but it is the decrease in total transcription

capacity (Fig. 5). It remains to be established whether this is a common response to other

translation limiting factors, although a similar pattern of a decrease in growth rate despite a

proportional increase in ribosomal RNA content and translation rate has been observed in the

past with antibiotics such as tetracycline, erythromycin, and neomycin, and limiting expression

of initiator factors 2 and 3 (18, 19).

Figure 5. Summary. Orange arrows show translation effects and blue arrows show transcription effects.
A. In the absence of translation limitation ribosomes translate both ribosomal and non-ribosomal genes
with similar rates. The amount of RNAP available is regulated in part by the changes in ppGpp and the
ensuing transcription rate of ribosomal promoters. **B.** The decrease in ribosome processivity by inhibitors
such as chloramphenicol (Cm) results in translation of longer genes being prematurely terminated more
frequently than translation of shorter genes. RNAP subunits β and β' are among the longest genes in *E.
coli*, while ribosomal proteins are among the shortest. The decrease in free RNAP can be measured by a
decreasing ratio of high affinity to low affinity promoter genes expression rate. The decrease in ribosome
processivity increases the probability of mRNA degradation, thus penalizing the expression of genes at
the end of operons. In nutrient poor media, inhibition of ribosome activity by chloramphenicol increases
the pool of amino acids and decreases the levels of ppGpp, increasing both ribosome production and
ribosome activity. Ab: antibiotic.

Materials and Methods

Strains, promoters and reporters

GFP and β-galactosidase were used as the reporter proteins to measure the rate of gene

expression from a specific promoter. The GFP gene used is *gfpmut2* coding for a fast-folding

GFP (63). The β-galactosidase gene is a 5'-end-modified *lacZ* from the pCMVbeta plasmid (64).

Comparison with the wild type *lacZ* gene shows that the additional 23 amino acids do not

change the results obtained with this version of the reporter gene (data not shown). The

promoters include two constitutive promoters (P5, obtained from T5 phage and PLtet, i.e. P_{LtetO-}

$_1$ (30) and a shortened version of a rRNA promoter (*rrnB*P1 without the upstream Fis and H-NS

sites) (Table S1). The constructs of P1-*gfpmut2*, P5-*gfpmut2*, PLtet-*gfpmut2* and P5-*lacZ* with a

divergent kanamycin resistance gene were inserted in the chromosome of the BW25113

Escherichia coli strain. P1-*gfpmut2* and P5-*gfpmut2* (for Fig. 1) were inserted at position 258235

between the convergent *crl* and *phoE* genes, PLtet-*gfpmut2* was at position 356850 between

cynR and *codA*, and P5-*lacZ* and P5-*gfpmut2* (for Fig. 3) were at position 1395689 between *uspE*

and *ynaJ*. Genome position did not have an effect on the change in reporter gene expression as

a function of Cm. The double fluorescent protein system (RFP-GFP constructs) has been

described previously (43). The ribosome binding sites (RBS) i.e. Shine-Dalgarno sequences, used

in above constructs are all similar to the consensus UAAGGAGGU (65). The RBS for GFP

(*gfpmut2*) and β-gal (*lacZ*) is GAAGGAGAU, for RFP (*mCherry*) it is AGAGGAGAA.

Bacterial growth and fluorescence measurements.

Bacterial growth was carried out in M9 minimal growth medium supplemented with 0.5%

glycerol (gly), 0.5% glucose (glu), 0.5% glycerol+0.2% casamino acids (cAA+gly) and 0.5%

glucose+0.2% casamino acids (cAA+glu). The pre-culture was obtained from the inoculation of

one bacterial colony in LB growth medium. After overnight growth, the seed culture was washed

once with PBS and diluted 200 times with the corresponding growth medium containing a

specific concentration of chloramphenicol (Cm). This culture was diluted again 200 times once

it reached exponential phase. The cultures were grown in flasks, shaking at 37°C and 170 rpm,

and optical density and fluorescence were measured with a plate reader (Tecan, Infinite 200Pro)

every 30 - 50 min. Alternatively, the cultures were grown in a 96 well plate, with 150 µL of

bacterial culture per well covered by 70µL mineral oil. The culture plate was kept at 37°C in the

plate reader, shaking and measuring fluorescence and OD_{600} every 5 minutes. The auto-

fluorescence measured from the wild type strain Bw25113 was subtracted from the

fluorescence of the fluorescent strains at the same OD (dependent on the medium). The

experimental procedure of the β-galactosidase assay followed the protocol of Zhang et al.[66]

except that the bacterial strains were cultivated in flasks instead of 48-well plates[18, 67, 68].

The measurement of RFP-GFP constructs followed the protocol described previously [43].

Analysis of GFP reporter expression data.

Experimental data obtained from the plate reader were analyzed with Matlab to obtain growth

rate, protein concentration and protein expression rate. The pipeline is shown in Fig. S1. The

window in the growth curve corresponding to the exponential growth phase was defined as a

linear range between an upper and a lower threshold in the growth curve plot of $\log(OD_{600})$

versus time (the thresholds determined manually or from an automated method [69] gave

similar results). Growth rate was derived from the slope of $\log(OD_{600})$ versus time in exponential

phase (Fig. S2B and C). GFP concentration was derived as the slope of the plot of GFP versus

OD_{600} in the exponential growth phase (Fig. S2B). β-galactosidase concentration in Miller Units

was obtained by the following formula (Fig. S2C)

$$\beta - \text{gal activity} \triangleq 1000 \cdot A \cdot \frac{1}{0.01} \cdot 20 \cdot \frac{1}{OD_{600}} = 2 \times 10^6 \times \frac{A}{OD_{600}} \quad (2)$$

where A comes from the fit of OD_{600} as a function of time with the formula $A(1 - e^{-\gamma t})/\gamma$ and γ is a decay factor from taking into account that the reaction product o-nitrophenol is volatile(66). The rate of protein expression is defined as the product of protein concentration and growth rate (μ).

Acknowledgments

The authors would like to thank Luca Cialdrini and Gilles Fischer for useful comments on the manuscript. This work was supported by HFSP grant RGY0070/2014 (BS, MCL and QZ) and the Strategic Pilot Project of Chinese Academy of Sciences (XDA17010504) (HS and QZ).

References

References

1. Mathieu A, Fleurier S, Frénoy A, Dairou J, Bredeche M-F, Sanchez-Vizuete P, Song X, Matic I. 2016.

Discovery and Function of a General Core Hormetic Stress Response in E. coli Induced by

Sublethal Concentrations of Antibiotics. Cell Reports 17:46–57.

2. Kohanski MA, Dwyer DJ, Hayete B, Lawrence CA, Collins JJ. 2007. A Common Mechanism of

Cellular Death Induced by Bactericidal Antibiotics. Cell 130:797–810.

3. Levin-Reisman I, Ronin I, Gefen O, Braniss I, Shoresh N, Balaban NQ. 2017. Antibiotic tolerance

facilitates the evolution of resistance. Science (New York, NY) 355:826–830.

4. Andersson DI, Hughes D. 2014. Microbiological effects of sublethal levels of antibiotics. Nature

Reviews Microbiology 12:465–478.

5. Hauryliuk V, Atkinson GC, Murakami KS, Tenson T, Gerdes K. 2015. Recent functional insights into

the role of (p)ppGpp in bacterial physiology. Nat Rev Micro 13:298–309.

6. Murray HD, Schneider DA, Gourse RL. 2003. Control of rRNA Expression by Small Molecules Is

Dynamic and Nonredundant. Molecular Cell 12:125–134.

7. Nunes-Alves C. 2016. Bacterial physiology: (p)ppGpp target ribosome assembly. Nat Rev Micro

14:266–266.

8. Zhang Y, Zborníková E, Rejman D, Gerdes K. 2018. Novel (p)ppGpp Binding and Metabolizing

Proteins of Escherichia coli. MBio 9.

9. Haugen SP, Ross W, Gourse RL. 2008. Advances in bacterial promoter recognition and its control

by factors that do not bind DNA. Nat Rev Microbiol 6:507–519.

10. Gummesson B, Magnusson LU, Lovmar M, Kvint K, Persson Ö, Ballesteros M, Farewell A, Nyström

T. 2009. Increased RNA polymerase availability directs resources towards growth at the expense

of maintenance. The EMBO Journal 28:2209–2219.

11. Klumpp S, Zhang Z, Hwa T. 2009. Growth rate-dependent global effects on gene expression in

bacteria. Cell 139:1366–75.

12. Mauri M, Klumpp S. 2014. A model for sigma factor competition in bacterial cells. PLoS Comput

Biol 10:e1003845.

13. Nyström T. 2004. Growth versus maintenance: a trade-off dictated by RNA polymerase

availability and sigma factor competition? Mol Microbiol 54:855–862.

14. Stracy M, Lesterlin C, Garza de Leon F, Uphoff S, Zawadzki P, Kapanidis AN. 2015. Live-cell

superresolution microscopy reveals the organization of RNA polymerase in the bacterial nucleoid.

Proc Natl Acad Sci USA 112:E4390-4399.

15. Paul BJ, Ross W, Gaal T, Gourse RL. 2004. rRNA transcription in Escherichia coli. Annu Rev Genet

38:749–770.

16. Zhang X, Liang S-T, Bremer H. 2006. Feedback control of ribosome synthesis in Escherichia coli is

dependent on eight critical amino acids. Biochimie 88:1145–1155.

17. Potrykus K, Murphy H, Philippe N, Cashel M. 2011. ppGpp is the major source of growth rate

control in E. coli. Environmental Microbiology 13:563–575.

18. Scott M, Gunderson CW, Mateescu EM, Zhang Z, Hwa T. 2010. Interdependence of cell growth

and gene expression: origins and consequences. Science 330:1099–1102.

19. Dai X, Zhu M, Warren M, Balakrishnan R, Patsalo V, Okano H, Williamson JR, Fredrick K, Wang Y-P,

Hwa T. 2016. Reduction of translating ribosomes enables Escherichia coli to maintain elongation

rates during slow growth. Nature Microbiology 2:16231.

20. Bremer H, Dennis P. 1996. Modulation of Chemical Composition and Other Parameters of the Cell

by Growth Rate, p. 1553–1569. *In* Neidhardt, FC, Ingraham, JL, Low, KB, Magasanik, B,

Schaechter, M, Umbarger, HE (eds.), Escherichia coli and Salmonella typhimurium: cellular and

Molecular Biology, 2nd ed. American Society for Microbiology, Washington D.C.

21. Bremer H, Dennis P, Ehrenberg M. 2003. Free RNA polymerase and modeling global transcription

in Escherichia coli. Biochimie 85:597–609.

22. Patrick M, Dennis PP, Ehrenberg M, Bremer H. 2015. Free RNA polymerase in Escherichia coli.

Biochimie 119:80–91.

23. Zhang X, Dennis P, Ehrenberg M, Bremer H. 2002. Kinetic properties of rrn promoters in

Escherichia coli. Biochimie 84:981–996.

24. Liang S-T, Bipatnath M, Xu Y-C, Chen S-L, Dennis P, Ehrenberg M, Bremer H. 1999. Activities of

constitutive promoters in Escherichia coli. Journal of Molecular Biology 292:19–37.

25. Hui S, Silverman JM, Chen SS, Erickson DW, Basan M, Wang J, Hwa T, Williamson JR. 2015.

Quantitative proteomic analysis reveals a simple strategy of global resource allocation in

bacteria. Molecular systems biology 11:784.

26. Traxler MF, Summers SM, Nguyen H-T, Zacharia VM, Hightower GA, Smith JT, Conway T. 2008.

The global, ppGpp-mediated stringent response to amino acid starvation in Escherichia coli.

Molecular microbiology 68:1128–1148.

27.   Dennis PP, Ehrenberg M, Bremer H. 2004. Control of rRNA synthesis in Escherichia coli: a systems

586   biology approach. Microbiol Mol Biol Rev 68:639–668.

587   28.   Felth L, Molzahn C, Wang S, Henderson K, Ruff E, Poulos M, Chhabra M, Bieter L, Artsimovitch I,

588   Record MT. 2016. Systematic Studies of Effects of the Discriminator on Stabilization of the Open

589   Complex and Abortive and Productive Initiation. FASEB J 30:1055.6-1055.6.

590   29.   Pemberton IK, Muskhelishvili G, Travers AA, Buckle M. 2000. The G+C-rich discriminator region of

591   the tyrT promoter antagonises the formation of stable preinitiation complexes. J Mol Biol

592   299:859–864.

593   30.   Lutz R, Bujard H. 1997. Independent and tight regulation of transcriptional units in Escherichia

594   coli via the LacR/O, the TetR/O and AraC/I1-I2 regulatory elements. Nucleic acids research

595   25:1203–1210.

596   31.   Traxler MF, Zacharia VM, Marquardt S, Summers SM, Nguyen H-T, Stark SE, Conway T. 2011.

597   Discretely calibrated regulatory loops controlled by ppGpp partition gene induction across the

598   "feast to famine" gradient in Escherichia coli. Mol Microbiol 79:830–845.

599   32.   Ryals J, Little R, Bremer H. 1982. Control of rRNA and tRNA syntheses in Escherichia coli by

600   guanosine tetraphosphate. J Bacteriol 151:1261–8.

601   33.   Klumpp S, Hwa T. 2008. Growth-rate-dependent partitioning of RNA polymerases in bacteria.

602   Proc Natl Acad Sci U S A 105:20245–50.

603   34.   Sendy B, Lee DJ, Busby SJW, Bryant JA. 2016. RNA polymerase supply and flux through the lac

604   operon in Escherichia coli. Philos Trans R Soc Lond, B, Biol Sci 371.

605   35.   Berg OG, von Hippel PH. 1987. Selection of DNA binding sites by regulatory proteins. Statistical-

606   mechanical theory and application to operators and promoters. J Mol Biol 193:723–750.

607   36.   Tsung K, Inouye S, Inouye M. 1989. Factors affecting the efficiency of protein synthesis in

608   Escherichia coli. Production of a polypeptide of more than 6000 amino acid residues. Journal of

609   Biological Chemistry 264:4428–4433.

610   37.   Harvey RJ, Koch AL. 1980. How partially inhibitory concentrations of chloramphenicol affect the

611   growth of Escherichia coli. Antimicrob Agents Chemother 18:323–337.

612   38.   Artsimovitch I. 2018. Rebuilding the bridge between transcription and translation. Mol Microbiol

613   108:467–472.

614   39.   Demo G, Rasouly A, Vasilyev N, Svetlov V, Loveland AB, Diaz-Avalos R, Grigorieff N, Nudler E,

615   Korostelev AA. 2017. Structure of RNA polymerase bound to ribosomal 30S subunit. eLife

616   Sciences 6:e28560.

617   40.   Fan H, Conn AB, Williams PB, Diggs S, Hahm J, Gamper HB, Hou Y-M, O'Leary SE, Wang Y, Blaha

618   GM. 2017. Transcription–translation coupling: direct interactions of RNA polymerase with

619   ribosomes and ribosomal subunits. Nucleic Acids Res 45:11043–11055.

620   41.   Kohler R, Mooney RA, Mills DJ, Landick R, Cramer P. 2017. Architecture of a transcribing-

621   translating expressome. Science 356:194–197.

622   42.   Proshkin S, Rahmouni AR, Mironov A, Nudler E. 2010. Cooperation Between Translating

623   Ribosomes and RNA Polymerase in Transcription Elongation. Science 328:504–508.

624   43.   Li R, Zhang Q, Li J, Shi H. 2015. Effects of cooperation between translating ribosome and RNA

625   polymerase on termination efficiency of the Rho-independent terminator. Nucleic acids research

626   gkv1285.

627   44.   Hausser J, Mayo A, Keren L, Alon U. 2019. Central dogma rates and the trade-off between

628   precision and economy in gene expression. Nature Communications 10:68.

45. Lane WJ, Darst SA. 2010. Molecular Evolution of Multi-subunit RNA Polymerases: Sequence
Analysis. J Mol Biol 395:671.

46. Zakharova N, Paster BJ, Wesley I, Dewhirst FE, Berg DE, Severinov KV. 1999. Fused and
Overlapping rpoB and rpoC Genes in Helicobacters, Campylobacters, and Related Bacteria.
Journal of Bacteriology 181:3857.

47. Severinov K, Mustaev A, Kukarin A, Muzzin O, Bass I, Darst SA, Goldfarb A. 1996. Structural
modules of the large subunits of RNA polymerase. Introducing archaebacterial and chloroplast
split sites in the beta and beta' subunits of Escherichia coli RNA polymerase. Journal of Biological
Chemistry 271:27969–74.

48. Davis JH, Williamson JR. 2017. Structure and dynamics of bacterial ribosome biogenesis. Phil
Trans R Soc B 372:20160181.

49. Murakami KS, Darst SA. 2003. Bacterial RNA polymerases: the wholo story. Curr Opin Struct Biol
13:31–39.

50. Kudrin P, Varik V, Oliveira SRA, Beljantseva J, Santos TDP, Dzhygyr I, Rejman D, Cava F, Tenson T,
Hauryliuk V. 2017. Subinhibitory Concentrations of Bacteriostatic Antibiotics Induce relA-
Dependent and relA-Independent Tolerance to β-Lactams. Antimicrobial Agents and
Chemotherapy 61:e02173-16.

51. Milon P, Tischenko E, Tomšic J, Caserta E, Folkers G, La Teana A, Rodnina MV, Pon CL, Boelens R,
Gualerzi CO. 2006. The nucleotide-binding site of bacterial translation initiation factor 2 (IF2) as a
metabolic sensor. Proceedings of the National Academy of Sciences 103:13962–13967.

52. Bakshi S, Choi H, Weisshaar JC. 2015. The spatial biology of transcription and translation in
rapidly growing Escherichia coli. Front Microbiol 6:636.

53.   Gausing K. 1982. Regulation of ribosome synthesis in E. coli. Trends in Biochemical Sciences 7:65–

652   67.

653   54.   McGinnis JL, Liu Q, Lavender CA, Devaraj A, McClory SP, Fredrick K, Weeks KM. 2015. In-cell

654   SHAPE reveals that free 30S ribosome subunits are in the inactive state. PNAS 112:2425–2430.

655   55.   Mori M, Schink S, Erickson DW, Gerland U, Hwa T. 2017. Quantifying the benefit of a proteome

656   reserve in fluctuating environments. Nat Commun 8:1225.

657   56.   Greulich P, Scott M, Evans MR, Allen RJ. 2015. Growth-dependent bacterial susceptibility to

658   ribosome-targeting antibiotics. Mol Syst Biol 11:n/a-n/a.

659   57.   Gowrishankar J, Harinarayanan R. 2004. Why is transcription coupled to translation in bacteria?

660   Molecular Microbiology 54:598–603.

661   58.   Scott M, Hwa T. 2011. Bacterial growth laws and their applications. Curr Opin Biotechnol 22:559–

662   565.

663   59.   Davies J, Spiegelman GB, Yim G. 2006. The world of subinhibitory antibiotic concentrations.

664   Current Opinion in Microbiology 9:445–453.

665   60.   Deris JB, Kim M, Zhang Z, Okano H, Hermsen R, Groisman A, Hwa T. 2013. The Innate Growth

666   Bistability and Fitness Landscapes of Antibiotic-Resistant Bacteria. Science 342:1237435.

667   61.   de Jong H, Geiselmann J, Ropers D. 2017. Resource Reallocation in Bacteria by Reengineering the

668   Gene Expression Machinery. Trends Microbiol 25:480–493.

669   62.   Balaban N. 2011. Persistence: mechanisms for triggering and enhancing phenotypic variability.

670   Current opinion in genetics & development 21:768–75.

671   63.   Cormack BP, Valdivia RH, Falkow S. 1996. FACS-optimized mutants of the green fluorescent

672   protein (GFP). Gene 173:33–38.

673   64.   MacGregor GR, Caskey CT. 1989. Construction of plasmids that express E. coli beta-galactosidase

674   in mammalian cells. Nucleic Acids Res 17:2365.

675   65.   Shine J, Dalgarno L. 1974. The 3'-Terminal Sequence of Escherichia coli 16S Ribosomal RNA:

676   Complementarity to Nonsense Triplets and Ribosome Binding Sites. Proc Natl Acad Sci U S A

677   71:1342–1346.

678   66.   Zhang Q, Li R, Li J, Shi H. 2018. Optimal Allocation of Bacterial Protein Resources under Nonlethal

679   Protein Maturation Stress. Biophys J 115:896–910.

680   67.   Zhang X, Bremer H. 1995. Control of the Escherichia coli rrnB P1 promoter strength by ppGpp.

681   Journal of Biological Chemistry 270:11181–11189.

682   68.   Shetty RP. 2008. Applying engineering principles to the design and construction of transcriptional

683   devices. Massachusetts Institute of Technology.

684   69.   Rudge TJ, Brown JR, Federici F, Dalchau N, Phillips A, Ajioka JW, Haseloff J. 2016. Characterization

685   of Intrinsic Properties of Promoters. ACS Synth Biol 5:89–98.

Supporting information

688   The supplementary information file includes:

689   Supplementary text

690   Figs. S1 to S16

691   Table S1

692   References for SI reference citations