A decrease in transcription capacity limits growth rate upon translation inhibition

Qing Zhang[1], Elisa Brambilla[1], Rui Li[2], Hualin Shi[2,3], Marco Cosentino Lagomarsino[4] and Bianca Sclavi[1].

1. LBPA, UMR 8113 CNRS, ENS Paris-Saclay, 94235 Cachan, France

2. CAS Key Laboratory of Theoretical Physics, Institute of Theoretical Physics, Chinese Academy of Sciences, Beijing 100190, China

3. School of Physical Sciences, University of Chinese Academy of Sciences, Beijing, China

4. LCQB, UMR 7238 CNRS, Sorbonne Université, 75005 Paris, France

Current affiliations:

Qing Zhang

    CAS Key Laboratory of Theoretical Physics, Institute of Theoretical Physics, Chinese Academy of Sciences, Beijing 100190, China

Marco Cosentino Lagomarsino

    Dipartimento di Fisica, Università degli Studi di Milano, 20133 Milano, Italy

    IFOM, FIRC Institute for Molecular Oncology, Milan, Italy

Bianca Sclavi

    LCQB, UMR 7238 CNRS, Sorbonne Université, Paris, France

Abstract

In bacterial cells, inhibition of ribosomes by sublethal concentrations of antibiotics leads to a decrease in growth rate despite an increase in ribosome content. This has been proposed to result from a decrease in the resources available for the production of non-ribosomal proteins that become limiting for cellular metabolism, mediated by changes in ppGpp concentration. Here, by the comparison of the activity of promoters that are differentially regulated by ppGpp and with different affinities for RNAP, we show that increased ribosome inhibition instead results in a decrease in total RNAP activity, independently of the changes in ppGpp. This is due to the decrease in ribosome processivity which affects more strongly the translation of longer genes, such as those coding for the β and β' subunits of RNAP, amongst the longest genes in the *E. coli* genome, while the expression of the very short ribosomal proteins is not limited to the same extent. Furthermore, the decrease in the total amount of free RNAP favors the expression of genes with promoters with a high RNAP affinity, such as ribosomal promoters, over the expression of other, growth-limiting, proteins. The decrease in transcription capacity could thus lead to increased survival whenever ribosome activity is compromised.

Bacteria often encounter sub-lethal levels of antibiotics produced by other microorganisms in their environment that can lead to a decrease in growth rate, allowing a strain to survive long enough until the inhibitor is no longer present or, in some cases, until the bacteria becomes resistant to the antibiotic via the selection of pre-existing mutations or an increase in mutation rates [1–4]. However, the mechanistic details of these response pathways often remain to be described. The cellular response to the limitation of translation activity is thought to be related to the pathway involved in the stringent response, the regulatory mechanism that decreases

ribosome production in response to a decrease in amino acid availability. This is mediated by

the change in concentration of the secondary messenger molecule, (p)ppGpp, that is produced

by the RelA enzyme when the pool of amino acids decreases and ribosomes are not loaded with

charged tRNAs[5,6]. ppGpp can directly inhibit ribosome assembly[7,8] and the activity of RNA

polymerase (RNAP) at ribosomal promoters, while increasing the activity of the promoters of

genes for amino acid biosynthesis[9]. The transcription of ribosomal operons can use a large

fraction of the free RNA polymerase pool in the cell because of the high affinity of the ribosomal

promoters for the enzyme and a high frequency of transcription initiation. Therefore, the

regulation of ribosomal promoter activity can be a means by which the pool of free RNA

polymerase can be repartitioned between ribosomal operon transcription and non-ribosomal

mRNA synthesis (Fig. 1A). This has been referred to as the passive control of transcription

regulation [10–14].

The ppGpp dependent feedback loop also plays a role in the regulation of ribosome content as

a function of growth rate[15–17]. Growth rate dependent regulation of gene expression determines

the allocation of cellular resources between the production of ribosomes and that of other

proteins and results in a linear increase in ribosome content with increasing growth rate[15,18–20]

. In richer growth media, when the amount of amino acids is higher, ppGpp levels are lower,

favoring ribosome production and a higher fraction of active ribosomes (Figs. 1 and S1A). In

poorer media it is the inverse, accumulation of ppGpp slows down the production of new

ribosomes and a smaller fraction of the ribosome pool is in an active form[19] (Fig. S1B). The

activity of ribosomal promoters is also regulated as a function of the growth rate by changes in

the activity of the Fis and H-NS proteins and by changes in the levels of negative supercoiling of

the DNA [21–24], which can explain the increase in rRNA with increasing growth rate in the mutant

strains devoid of ppGpp [25–27].

When ribosome activity is inhibited by sublethal concentrations of antibiotics, amino acids are

used more slowly and their concentration increases, thus decreasing the intracellular ppGpp

pool. The cellular response, as predicted by the ppGpp feedback loop, is to produce a higher

amount of ribosomes and an increased translation rate, however, despite this increase, the

cell's growth rate is reduced[18,19]. This has been proposed to result from a decrease in the

resources available for the production of non-ribosomal proteins that become limiting for

cellular metabolism [18,28]. More recent results point to a decrease in the fraction of active

ribosomes to explain the decrease in the total protein production rate[1] .

To measure the effect that ppGpp can play on gene expression in translation-limiting conditions,

and its possible effect on the repartition of RNAP, we have compared the activity of promoters

that are differentially regulated by ppGpp and that have different affinities for RNAP (Figs 1A

and S1). In parallel, we analyzed data from the literature on the direct and indirect effects of

changing ppGpp concentration on gene expression and compared them to the genes that

change expression upon translation inhibition[29,30]. The results from this analysis show that the

direct effect of ppGpp at the transcriptional level can only partially explain the change in protein

expression levels under translation limitation and that instead there is a decrease in RNA

polymerase production that increases the competition between high and low affinity

promoters, favoring the transcription of ribosomal operons. We propose that it is this decrease

in the total transcriptional capacity that results in a decrease the cell's growth rate even when

the translational capacity is increased.

Results

Decrease in total RNAP activity with increasing translation inhibition independently of

changes in ppGpp. In order to measure the effect of increasing ribosome inhibition on RNAP

activity via changes in ppGpp we chose three promoters with different RNAP affinities and that

are differently regulated by ppGpp. The first is a shortened version of the well characterized

ribosomal *rrnB* operon promoter *rrnB*P1, here called P1, that terminates at -69 from the

transcription start site[23]. This promoter has a GC-rich discriminator region at the transcription

initiation site that makes the open complex sensitive to ppGpp inhibition (Table S1). The second

is a constitutive promoter, P5, that has consensus -10 and -35 sequences and no discriminator

region. The third is PLtet, also a strong constitutive promoter with no discriminator region but

with a lower affinity for RNA polymerase due to a non-consensus -10 sequence[31]. Each of these

promoters was placed upstream of the *gfpmut2* gene and this cassette was inserted in the

chromosome together with a kanamycin resistance gene expressed divergently from the chosen

promoter (Fig. S2). Growth of these strains in a 96-well plate allowed us to measure the changes

in growth rate, the GFP production rate (Gpr), and the resulting GFP concentration as a function

of chloramphenicol concentration (Figs. 1B-D and S2). We compared four different growth

media, M9 with glucose, M9 with glycerol and these two media supplemented with casamino

acids (CAA). This results in four different growth rates. In addition, it has been shown that cells

growing in a growth medium containing amino acids have lower levels of ppGpp[16], allowing us

to compare the effects due to changing concentrations of this key metabolite without the use

of mutant strains that can result in secondary effects on cell metabolism[32].

In the absence of translation inhibition, the change in promoter activity measured as a function

of growth rate is consistent with previous measurements on constitutive promoters and *rrnB*P1

promoters[11,11] (Fig. 1B-D dashed lines). Previous work had shown that as the concentration of

chloramphenicol is increased, the ribosome content increases and the concentration of a

reporter protein expressed from a constitutive promoter decreases[18]. Therefore, the expected

result here is that the GFP concentration from a ribosomal promoter (P1) should increase while

the concentration from a constitutive promoter (P5 or PLtet) should decrease, with the lower

affinity constitutive promoter (PLtet) decreasing at a faster rate if increasing amounts of RNAP

are being used for transcription of ribosomal operons. Unexpectedly, however, the pattern of

the change in Gpr and GFP concentration for the P1 and P5 promoters is very similar. The

pattern of the change of expression for the three promoter constructs depends strongly on the

growth medium and whether or not it contains casamino acids. In the absence of CAA, in the

cells starting with a higher amount of ppGpp, the increase in Gpr at low Cm is significantly

greater than in their presence (blue lines compared to the red lines in Fig. 1). This shows that a

decrease in ppGpp increases not only ribosomal promoter activity but also the translation rate

of GFP, independently of promoter sequence, as expected from recent studies on the effect of

ppGpp on ribosome assembly and activity[7,8]. The similarity in the pattern of the change in GFP

expression from the *rrnB*P1 and P5 promoters indicates in fact that upon translation limitation

the main effect of ppGpp on gene expression is via the translation rate rather than the

transcription rate (Fig. 1 and Fig. S16). Nevertheless, for each promoter the change in Gpr results

in a distinct pattern in the change of GFP concentration as a function of the presence of CAA in

the growth medium. In the two growth media with casamino acids [GFP] increases at the lower

Cm concentrations for P1 and P5, to then decrease again, while for PLtet it decreases with

increasing Cm.

Figure 1. Effect of translation limitation on GFP expression from three different promoters. A. The negative feedback loop mediated by ppGpp maintains ribosome production and activity proportional to the amount of available amino acids[5,6]. During the stringent response, upon a decrease in nutrients, an increase in ppGpp reduces the activity of ribosomal promoters and more RNAP is available for the transcription of non-ribosomal genes. In the presence of translation inhibitors, on the other hand, the pool of available amino acids increases, decreasing the concentration of ppGpp (Fig. S1). The three promoters used are: P1, a shortened version of the rRNA *rrnB*P1 promoter, P5, a constitutive promoter with a high affinity for RNA polymerase and PLtet, a constitutive promoter with a lower affinity for RNAP. These promoters can be used as reporters of the changes in ppGpp, by comparing P1 and P5, and of the changes in the activity of RNA polymerase, by comparing PLtet and P5. **B-D** Change in GFP concentration and GFP production rate in the three strains as a function of growth rate at increasing Cm concentrations in growth media with and without casamino acids (red and blue lines respectively). In the growth media with casamino acids the cells have a lower concentration of ppGpp [16]. The dashed black lines connect the points obtained in the absence of Cm. The growth rate decreases with increasing chloramphenicol concentration (black arrow). The four points correspond to 0, 2, 4 and 8 µM final chloramphenicol concentration. The error bars represent the SEM from 3 independent experiments. The error bars smaller than the size of the symbols are not shown.

One of the predictions from previous studies is that the induction of the ribosomal operons

promoters by decreasing ppGpp will result in a decrease in the amount of RNAP that is available

to transcribe mRNA[10–14]. In order to obtain an estimate of the magnitude of the effect of ppGpp

on promoter activity we measured the ratios of Gpr. Since the translation rate of GFP is shared

similarly by the three promoter constructs, this operation will leave the ratio of the

transcription-specific effect (Fig. 2). The comparison of two promoters with differing affinities

for RNAP can be used to estimate the change in the amount of RNAP that is available for

transcription[33]. In the absence of Cm, the ratio of PLtet to P5 Gpr increases with increasing

growth rate, in agreement with previous estimates of the change in free RNAP as a function of

doubling time[34]. The ratio of P1 to P5 instead increases rapidly between M9-glu and M9-CAA-

gly, consistent with a lower level of ppGpp in the cells growing in the presence of CAA increasing

the probability of transcription initiation specifically from P1. The PLtet/P5 and P1/P5 ratios are

greater than 1 because while PLtet and P1 have a lower affinity for RNAP than P5, when RNAP

binds at the PLtet or at the P1 promoter it initiates transcription with a higher frequency than

at P5 [31].

When Cm is increased in the cells with a higher amount of ppGpp (Fig. 2, blue lines), the fold

induction of P1 is greater than the one of P5, consistent with a decrease in ppGpp levels. As the

Cm concentration is increased further, the difference between the two promoters decreases

again to the initial level, suggesting that the ppGpp levels could increase again with the increase

in the cellular ribosome content (Fig. S4). On the other hand, in the growth media with casamino

acids, and thus lower levels of ppGpp, the P1 to P5 ratio decreases, indicating that the change

in Gpr from P1 is smaller than that of P5 as a function of increasing Cm. Here the decrease in

ppGpp is not large enough to increase the transcription initiation rate from P1 to the same

extent as in the poorer growth media, therefore the repartition of RNAP between ribosomal

and non-ribosomal promoters is expected to have a smaller influence on transcription

regulation. However, this still does not explain why the P1/P5 ratio should decrease with

increasing translation inhibition. One possibility is that there is an increase in ppGpp, however

a similar result is also observed for the PLtet to P5 ratio, independently of the growth medium.

Instead, this is consistent with a decrease in the amount of free RNA polymerase in the cell with

increasing Cm, independently of ppGpp regulation of ribosomal promoters.

The values of the RNAP affinities for these promoters and the relative changes in Gpr can be

used to estimate the change in the amount of free RNAP taking place in the cells. Fig. 2C and

Fig. S3 illustrate this point. The RNAP binding affinity for each promoter was estimated based

on a statistical-mechanical selection model developed by Berg and von Hippel[35] (see details in

the SI). The binding curves for RNAP to each promoter are shown. The vertical bars indicate the

ratio of promoter activity at a given RNAP concentration. The values of the Cm concentration

for each of these ratios is shown on the top. The decrease in free RNAP with increasing

chloramphenicol will have a stronger effect on PLtet first, then on P1 and finally on P5. Similarly,

using the P1 to P5 Gpr ratio it is possible to estimate the change in ppGpp as a function of growth

rate and as a function of Cm (Fig. S4). In summary, the decrease in free RNAP, as measured by

the decrease in the PLtet to P5 and P1 to P5 ratios, appears to be independent of a ppGpp-

mediated repartition between ribosomal and non-ribosomal promoters, since it can also be

observed in the growth media where ppGpp levels are very low.

Figure 2. Ratios of GFP production rate for the different promoters reveal promoter sequence-specific effects of translation limitation and ppGpp concentration. A. Ratio of PLtet to P5. The activity of the lower affinity constitutive promoter, PLtet, decreases more than that of P5 upon increasing Cm. **B.** Ratio of P1 to P5. Changes in ppGpp concentration in the growth media without cAA (blue lines) have a specific effect on the P1 promoter's activity compared to the constitutive promoter P5. The dashed lines connect the points obtained in the absence of Cm. **C.** The change in the ratios of promoter binding as a function of chloramphenicol concentration can be used to estimate the decrease in the amount of free RNAP. The relative activity of the promoters as a function of free RNAP concentration (c_f) was estimated from the respective RNAP binding constants, $K_i \in \{K_1, K_5, K_{Ltet}\}$ using $c_f/(K_i + c_f)$. K_5 is known, K_1 and K_{Ltet} can be derived by the formula $K_5 \exp(\Delta E)$, where ΔE is the difference between the binding energies of RNAP with P1 (or PLtet) and P5 (see Supporting Information). The black vertical bars indicate the concentrations of RNAP that give the measured GFP production rate ratios shown in (A) for different Cm concentrations, shown above the plot. The ratio of PLtet and P5 activities for example is given by: $Gpr(PLtet)/Gpr(P5) = a(K_5 + c_f)/(K_{Ltet} + c_f)$, where a is a scaling factor. K_5 and a can be fixed by fitting the data in absence of Cm from our experiments and the literature. Data from the cells growing in cAA-glu was used so that ppGpp-dependent regulation of P1 is small and can be ignored. Fig. S3 in Supplementary Materials shows the estimation of RNAP concentration for the four growth media used. See details of the analysis in the Supporting Information.

The decrease in ribosome processivity by chloramphenicol decreases the expression of longer

genes more than shorter ones. A possible cause for the decrease in the pool of free RNAP

independently of the changes in ppGpp is a decrease in the amount of RNAP per cell. The RNA

polymerase holoenzyme is composed of 5 subunits: β, β', ω, 2 subunits of α, and a σ factor. β,

β' are among the longest genes in *E. coli* with a length of 4029 bp and 4224 bp respectively (the

average gene length in *E. coli* is about 900 bp). We hypothesized that translation of such long

genes would have a higher probability to end prematurely in the presence of ribosome

inhibitors such as chloramphenicol.

Figure 3. Gene length can influence gene expression under translation limitation. A. Change in GFP
(238 aa) and β-galactosidase (1024 aa) expressed from the P5 promoter as a function of increasing
chloramphenicol concentrations. The bacteria were grown in M9 glu+cAA with chloramphenicol (Cm) at
concentrations of 0, 2 μM , 4 μM , or 6 μM in flasks. The concentrations were normalized by dividing
the data by the point without Cm. **B.** Fit of the model to the ratio of GFP to β-galactosidase
concentrations from the data in (A). The growth rate at each chloramphenicol concentration is the
average over the two strains. The error bars correspond to the SEM from three independent
experiments.

The processivity of translation has been shown to decrease exponentially with increasing gene

length[36]. If, in addition, ribosome processivity were decreased by an inhibitor, then the

probability to finish the translation of a long gene would be lower compared to a shorter gene,

decreasing the rate of expression of the longer gene to a greater extent. To test this hypothesis,

we compared the changes in gene expression from the same constitutive promoter, P5, of two

different genes, *gfpmut2*, 714 bp long, and *lacZ*, 3072 bp long and coding for the enzyme β-

galactosidase, in the presence of increasing chloramphenicol concentrations (Fig. 3A). The

results show that while the concentration of GFP increases as a function of Cm concentration,

β-galactosidase concentration decreases (Fig. 3A). Fig. 3B shows the change in the ratio of the

shorter to the longer protein as a function of Cm concentration. The black line shows the fit

obtained to a model of ribosome processivity. Briefly, it is possible to estimate the probability

that a ribosome will be inhibited by the antibiotic before it reaches the end of the mRNA, P_{hit},

as described by Dai et al[19]. Their equation can be extended to describe the dependence of P_{hit}

on protein length, i.e.

$$P_{hit} = 1 - \exp(-k_{on}[Cm]L/v) \quad (1)$$

where k_{on} denotes the binding constant of Cm with ribosome ($k_{on} = 0.034(\mu M \cdot min)^{-1}$ [37]),

L indicates protein length, and v is the translation elongation rate dependent on the

RNA/protein mass ratio (see the Materials and Methods; ref.[19]). Using equation 1, at 8 µM Cm

(the highest concentration used here) P_{hit} is 23% for LacZ (1024 aa), while for GFP it is 6% (238

aa). These results therefore indicate that a gene's length, in addition to its promoter's affinity

for RNAP, can influence how its expression levels change in the presence of sub-lethal

concentrations of ribosome inhibitors.

We have extended this analysis and this model to measure the difference in the expression of

early and late genes within the same operon upon decreased translation processivity and we

have found that indeed late operon genes are less expressed than early ones due to mRNA

degradation induced by the presence of a stalled ribosome (SI Fig. S8). However, at these low

Cm concentrations transcription-translation coupling is not significantly perturbed (SI Fig. S8

and S9). Finally, we have analyzed data from previously published transcriptomics and

proteomics studies[29,32] and we have found that the effect of promoter affinity, gene length and

operon position can help explain the global changes in gene expression in the presence of

sublethal levels of chloramphenicol (SI Fig.s S12,S13,S15).

Discussion

Understanding how bacteria modulate their growth rate and resource allocation in response to

inhibition of growth has paramount importance in biotechnological and health applications[38–

41]. The evidence provided here points to a possible cellular adaptation mechanism leading to a

reduction of transcription capacity when ribosome activity is compromised. This decreases the

cost of transcription of untranslated mRNAs[42] and allows for more resources to be available for

the synthesis of increased amounts of ribosomes to respond to the presence of translation

inhibitors. Here we found that as the sublethal concentration of ribosome inhibitor is increased

the growth rate decreases linearly with the decrease in the amount of available RNA

polymerase, pointing to a growth-limiting role for this enzyme in these conditions (Fig. 2). These

results are coherent with the observation in a previous study on the quantitation of over 1000

proteins in the presence of increasing concentrations of Cm[29] that the β and β' subunits of RNAP

remain a constant fraction of the proteome with increasing Cm and decreasing growth rate. This

is what would be expected if RNAP were to play an important role in determining the total rate

of protein production and the cell's growth rate.

The comparison of two different constitutive promoters with differing RNAP affinity, PLtet and

P5, shows a striking difference in the change in Gpr and the resulting GFP concentration with

increasing ribosome inhibition (Fig. 1). The decrease in the amount of free RNAP with increasing

sublethal concentrations of chloramphenicol estimated by measuring the ratio of gene

expression from these two promoters (PLtet/P5) is about 10-fold (Fig. 2 and Fig. S3) and is

independent of the change in ppGpp in the different growth media (Fig. 2). This can be used to

predict the change in transcription rate for the P5 promoter, for example, and therefore the

GFP translation rate that results in the measured Gpr (Fig. S16). The translation of short proteins

such as GFP is barely affected by the presence of sublethal concentrations of Cm (8 % probability

of drop off at the highest concentration used here) and can therefore be used to estimate the

changes in total translation activity. The translation rate increases and mirrors the ratio of P1 to

P5 activity reflecting the changes in ppGpp (Fig. S4). Several factors can contribute to this

increase in translation rate: the shorter than average length of ribosomal genes (Fig. S12),

whose translation is not affected as strongly by translation inhibition compared to longer genes;

the high affinity of RNAP for ribosomal promoters, insuring transcription despite a decrease in

the amount of free RNAP (Fig. S11); the decrease in ppGpp in the growth media lacking

casamino acids (Fig. S4), resulting in an increase in both ribosomal promoter activity (Fig. 2)[5,43]

and in the fraction of active ribosomes due to the effect of ppGpp on ribosome assembly[7,8,19],

and the increase in translation elongation rate[19]. The higher increase in translation rate in the

growth media without cAA is consistent with a larger fraction of inactive ribosomes stored in

the cells growing at these slower growth rates that can be quickly reactivated by a decrease in

ppGpp concentration[19,44–46]. Furthermore, gene length could also have an effect on translation

via increased probability of ribosome recycling on shorter mRNAs, increasing the fraction of

ribosomes translating ribosomal genes[47].

The similarity in the pattern of the change in GFP expression from the rrnBP1 and P5 promoters

despite their different regulation by ppGpp indicates that upon translation limitation the main

effect of ppGpp on gene expression from high affinity promoters is via the translation rate

rather than the transcription rate (Fig. 1 and Fig. S16). The effect of decreasing ppGpp on

transcription from the P1 promoter is just strong enough to compensate for the decreased

availability of RNAP (Fig. S16). In the richer growth media, where ppGpp levels are low and the

ribosomes are working at near maximum capacity, a specific induction of the ribosomal

promoter is practically absent and the increase in translation rate is lower, consistent with a

smaller change in ppGpp levels. The decrease in transcription capacity however is maintained.

In these growth conditions, in rich media when ppGpp levels are low, a decrease in transcription

rate when translation is compromised might play a crucial role for the cell's survival, as the

capacity of the cell to increase its translation capacity to respond to the presence of the inhibitor

is limited by the small fraction of inactive ribosomes[19,48].

The ppGpp independent decrease in transcription capacity can be explained by the fact that the

RNA polymerase core contains two of the longest proteins in *E. coli*, the β and β' subunits (Fig.

S12), and by the increased probability that a ribosome will stall before reaching the end of a

long mRNA than a short mRNA, particularly in the presence of inhibitors such as

chloramphenicol, as shown by the difference between the expression of the *lacZ* and *gfp* genes

(Fig. 3). These results can shed light on a recent study by Dai et al[19] where it was proposed that

in the presence of sublethal concentrations of Cm, despite an increase in the translation

elongation rate due to a higher concentration of ternary complexes, the reduction in the total

protein production rate results from a decrease in the active ribosome fraction, or the fraction

of ribosomes that can reach the end of a mRNA in the presence of the inhibitor[19]. Here we

identify RNAP as one of the genes that is likely to be most affected by the decrease in ribosome

processivity, while shorter genes are affected to a lesser extent.

The extreme length of the β and β' subunits is conserved throughout bacteria[49]. In the case of

Helicobacteraceae and *Wolbachia* the two genes are even fused together[49,50]. The gene length

of β and β' subunits can vary in different strains since they are composed of independent

structural modules separated by spacers of differing length[49]. In *E. coli* the spacer sequences,

that account for more than 25% of the total sequence, can be deleted without causing a

significant decrease in activity. In archea and chloroplasts some of the conserved protein

modules are found in separate genes and in *E. coli* they can be split from each other to produce

an active enzyme[51]. The reason why the RNAP and ribosomal proteins find themselves at

opposite ends of the spectrum of gene lengths in bacteria is likely linked to the assembly

process, structural flexibility and stability of the final multi-protein complex[52,53]; however, these

results suggest that it could also play an important role for the cell's survival in the presence of

antibiotics that decrease translation processivity.

It is possible to obtain an estimate the change in ppGpp from the ratio of the Gpr of the P1 and

P5 promoters since they both have a similar affinity for RNA polymerase while P1 (*rrnBP1)* is

also negatively regulated by ppGpp (Fig. S4). ppGpp decreases at the lower Cm concentrations

and then remains constant or increases again as Cm is increased, depending on the growth

medium. An increase in ppGpp coupled to an activation of the RpoS regulon has been implicated

in a general hormetic stress response to sublethal levels of antibiotics[1] and in the emergence of

tolerance and persister cells[3], a subset of the population that resists antibiotic inhibition due to

decreased growth[54,55]. Here we measured gene expression when the cells are still in exponential

phase, before the amounts of amino acids become limiting and before the adaptation to early

stationary phase. However, the repartition of the RNAP core among the sigma factors

associated with stress response[12,56] could also contribute to a further decrease the amount of

free sigma 70 RNAP as the cells enter stationary phase or in the presence of other antibiotics

that are not specifically targeting translation[1]. The results presented here provide a new cellular

mechanism by which bacterial cells can decrease their growth rate in response to antibiotic

stress[1,57,58] and can help explain the observation from a previous report that the induction of

persistence and β-lactam tolerance by chloramphenicol is also independent of the presence of

the RelA enzyme[59]. In conclusion, the results presented here provide evidence that in the

presence of sublethal concentrations of ribosome inhibitors, it is not translation that becomes

limiting for the cell's growth rate, or the ppGpp dependent repartition of RNAP between

ribosomal and non-ribosomal promoters, but it is the decrease in total transcription capacity.

Figure 4. Summary. Orange arrows show translation effects and blue arrows show transcription effects. **A.** In the absence of translation limitation ribosomes translate both ribosomal and non-ribosomal genes with similar rates. The amount of RNAP available is regulated in part by the changes in ppGpp and the ensuing transcription rate of ribosomal promoters. **B.** The decrease in ribosome processivity by inhibitors such as chloramphenicol results in translation of longer genes being prematurely terminated more frequently than translation of shorter genes. RNAP subunits β and β' are among the longest genes in *E. coli*, while ribosomal proteins are among the shortest. The decrease in free RNAP can be measured by a decreasing ratio of high affinity to low affinity promoter genes expression rate. The decrease in ribosome processivity increases the probability of mRNA degradation, thus penalizing the expression of genes at the end of operons. In nutrient poor media, inhibition of ribosome activity by chloramphenicol increases the pool of amino acids and decreases the levels of ppGpp, increasing both ribosome production and ribosome activity. Ab: antibiotic.

Materials and Methods

Strains, promoters and reporters

GFP and β-galactosidase were used as the reporter proteins to measure the rate of gene expression from a specific promoter. The GFP gene used is *gfpmut2* coding for a fast-folding GFP[60]. The β-galactosidase gene is a 5'-end-modified *lacZ* from the pCMVbeta plasmid[61]. Comparison with the wild type *lacZ* gene shows that the additional 23 amino acids do not change the results obtained with this version of the reporter gene (data not shown). The promoters include two constitutive promoters (P5, obtained from T5 phage and PLtet, i.e. $P_{LtetO\text{-}1}$[31] and a shortened version of a rRNA promoter (*rrnB*P1 without the upstream Fis and H-NS sites) (Table S1). The constructs of P1-*gfpmut2*, P5-*gfpmut2*, PLtet-*gfpmut2* and P5-*lacZ* with a divergent kanamycin resistance gene were inserted in the chromosome of the BW25113 *Escherichia coli* strain. P1-*gfpmut2* and P5-*gfpmut2* (for Fig. 1) were inserted at position 258235 between the convergent *crl* and *phoE* genes, PLtet-*gfpmut2* was at position 356850 between *cynR* and *codA*, and P5-*lacZ* and P5-*gfpmut2* (for Fig. 3) were at position 1395689 between *uspE* and *ynaJ*. Genome position did not have an effect on the change in reporter gene expression as a function of Cm. The double fluorescent protein system (RFP-GFP constructs) has been described previously[62]. The ribosome binding sites (RBS) i.e. Shine-Dalgarno sequences, used in above constructs are all similar to the consensus UAAGGAGGU[63]. The RBS for GFP (*gfpmut2*) and β-gal (*lacZ*) is GAAGGAGAU, for RFP (*mCherry*) it is AGAGGAGAA.

Bacterial growth and fluorescence measurements.

Bacterial growth was carried out in M9 minimal growth medium supplemented with 0.5% glycerol (gly), 0.5% glucose (glu), 0.5% glycerol+0.2% casamino acids (cAA+gly) and 0.5% glucose+0.2% casamino acids (cAA+glu). The pre-culture was obtained from the inoculation of one bacterial colony in LB growth medium. After overnight growth, the seed culture was washed

once with PBS and diluted 200 times with the corresponding growth medium containing a

specific concentration of chloramphenicol (Cm). This culture was diluted again 200 times once

it reached exponential phase. The cultures were grown in flasks, shaking at 37°C and 170 rpm,

and optical density and fluorescence were measured with a plate reader (Tecan, Infinite 200Pro)

every 30 - 50 min. Alternatively, the cultures were grown in a 96 well plate, with 150 μL of

bacterial culture per well covered by 70μL mineral oil. The culture plate was kept at 37°C in the

plate reader, shaking and measuring fluorescence and OD_{600} every 5 minutes. The auto-

fluorescence measured from the wild type strain Bw25113 was subtracted from the

fluorescence of the fluorescent strains at the same OD (dependent on the medium). The

experimental procedure of the β-galactosidase assay followed the protocol of Zhang et al.[64]

except that the bacterial strains were cultivated in flasks instead of 48-well plates[18,65,66]. The

measurement of RFP-GFP constructs followed the protocol described previously[62].

Analysis of GFP reporter expression data.

Experimental data obtained from the plate reader were analyzed with Matlab to obtain growth

rate, protein concentration and protein expression rate. The pipeline is shown in Fig. S2. The

window in the growth curve corresponding to the exponential growth phase was defined as a

linear range between an upper and a lower threshold in the growth curve plot of $\log(OD_{600})$

versus time (the thresholds determined manually or from an automated method[67] gave similar

results). Growth rate was derived from the slope of $\log(OD_{600})$ versus time in exponential phase

(Fig. S2B and C). GFP concentration was derived as the slope of the plot of GFP versus OD_{600} in

the exponential growth phase (Fig. S2B). β-galactosidase concentration in Miller Units was

obtained by the following formula (Fig. S2C)

$$\beta - \text{gal activity} \triangleq 1000 \cdot A \cdot \frac{1}{0.01} \cdot 20 \cdot \frac{1}{OD_{600}} = 2 \times 10^6 \times \frac{A}{OD_{600}} \quad (2)$$

393   where A comes from the fit of OD_{600} as a function of time with the formula $A(1 - e^{-\gamma t})/\gamma$

394   and γ is a decay factor from taking into account that the reaction product o-nitrophenol is

395   volatile[64]. The rate of protein expression is defined as the product of protein concentration and

396   growth rate (μ).

References

1. Mathieu, A. *et al.* Discovery and Function of a General Core Hormetic Stress Response in E. coli Induced by Sublethal Concentrations of Antibiotics. *Cell Reports* **17**, 46–57 (2016).

2. Kohanski, M. A., Dwyer, D. J., Hayete, B., Lawrence, C. A. & Collins, J. J. A Common Mechanism of Cellular Death Induced by Bactericidal Antibiotics. *Cell* **130**, 797–810 (2007).

3. Levin-Reisman, I. *et al.* Antibiotic tolerance facilitates the evolution of resistance. *Science (New York, N.Y.)* **355**, 826–830 (2017).

4. Andersson, D. I. & Hughes, D. Microbiological effects of sublethal levels of antibiotics. *Nature Reviews. Microbiology* **12**, 465–478 (2014).

5. Hauryliuk, V., Atkinson, G. C., Murakami, K. S., Tenson, T. & Gerdes, K. Recent functional insights into the role of (p)ppGpp in bacterial physiology. *Nat Rev Micro* **13**, 298–309 (2015).

6. Murray, H. D., Schneider, D. A. & Gourse, R. L. Control of rRNA Expression by Small Molecules Is Dynamic and Nonredundant. *Molecular Cell* **12**, 125–134 (2003).

7. Nunes-Alves, C. Bacterial physiology: (p)ppGpp target ribosome assembly. *Nat Rev Micro* **14**, 266–266 (2016).

8. Zhang, Y., Zborníková, E., Rejman, D. & Gerdes, K. Novel (p)ppGpp Binding and Metabolizing Proteins of Escherichia coli. *MBio* **9**, (2018).

9. Haugen, S. P., Ross, W. & Gourse, R. L. Advances in bacterial promoter recognition and its control by factors that do not bind DNA. *Nat. Rev. Microbiol.* **6**, 507–519 (2008).

10. Gummesson, B. *et al.* Increased RNA polymerase availability directs resources towards growth at the expense of maintenance. *The EMBO Journal* **28**, 2209–2219 (2009).

11. Klumpp, S., Zhang, Z. & Hwa, T. Growth rate-dependent global effects on gene expression in bacteria. *Cell* **139**, 1366–75 (2009).

12. Mauri, M. & Klumpp, S. A model for sigma factor competition in bacterial cells. *PLoS Comput. Biol.* **10**, e1003845 (2014).

13. Nyström, T. Growth versus maintenance: a trade-off dictated by RNA polymerase availability and

sigma factor competition? *Mol. Microbiol.* **54**, 855–862 (2004).

14. Stracy, M. *et al.* Live-cell superresolution microscopy reveals the organization of RNA polymerase

in the bacterial nucleoid. *Proc. Natl. Acad. Sci. U.S.A.* **112**, E4390-4399 (2015).

15. Paul, B. J., Ross, W., Gaal, T. & Gourse, R. L. rRNA transcription in Escherichia coli. *Annu. Rev.*

Genet. **38**, 749–770 (2004).

16. Zhang, X., Liang, S.-T. & Bremer, H. Feedback control of ribosome synthesis in Escherichia coli is

dependent on eight critical amino acids. *Biochimie* **88**, 1145–1155 (2006).

17. Potrykus, K., Murphy, H., Philippe, N. & Cashel, M. ppGpp is the major source of growth rate

control in E. coli. *Environmental Microbiology* **13**, 563–575 (2011).

18. Scott, M., Gunderson, C. W., Mateescu, E. M., Zhang, Z. & Hwa, T. Interdependence of cell growth

and gene expression: origins and consequences. *Science* **330**, 1099–1102 (2010).

19. Dai, X. *et al.* Reduction of translating ribosomes enables Escherichia coli to maintain elongation

rates during slow growth. *Nature Microbiology* **2**, 16231 (2016).

20. Bremer, H. & Dennis, P. Modulation of Chemical Composition and Other Parameters of the Cell by

Growth Rate. in *Escherichia coli and Salmonella typhimurium: cellular and Molecular Biology* (eds.

Neidhardt, F. C. et al.) 1553–1569 (American Society for Microbiology, 1996).

21. Travers, A. & Muskhelishvili, G. DNA supercoiling - a global transcriptional regulator for

enterobacterial growth? *Nat. Rev. Microbiol.* **3**, 157–169 (2005).

22. Brambilla, E. & Sclavi, B. Gene regulation by H-NS as a function of growth conditions depends on

chromosomal position in Escherichia coli. *G3 (Bethesda)* **5**, 605–614 (2015).

23. Dennis, P. P., Ehrenberg, M. & Bremer, H. Control of rRNA synthesis in Escherichia coli: a systems

biology approach. *Microbiol. Mol. Biol. Rev.* **68**, 639–668 (2004).

24. Afflerbach, H., Schröder, O. & Wagner, R. Effects of the Escherichia coli DNA-binding protein H-NS

on rRNA synthesis in vivo. *Molecular Microbiology* **28**, 641–653 (1998).

25. Gaal, T. & Gourse, R. L. Guanosine 3'-diphosphate 5'-diphosphate is not required for growth rate-

dependent control of rRNA synthesis in Escherichia coli. *Proceedings of the National Academy of*

Sciences **87**, 5533–5537 (1990).

26. Hernandez, V. J. & Bremer, H. Characterization of RNA and DNA synthesis in Escherichia coli strains

devoid of ppGpp. *J. Biol. Chem.* **268**, 10851–10862 (1993).

27. Bartlett, M. S., Gaal, T., Ross, W. & Gourse, R. L. Regulation of rRNA transcription is remarkably

robust: FIS compensates for altered nucleoside triphosphate sensing by mutant RNA polymerases

at Escherichia coli rrn P1 promoters. *J. Bacteriol.* **182**, 1969–1977 (2000).

28. Metzl-Raz, E. *et al.* Principles of cellular resource allocation revealed by condition-dependent

proteome profiling. *Elife* **6**, (2017).

29. Hui, S. *et al.* Quantitative proteomic analysis reveals a simple strategy of global resource allocation

in bacteria. *Molecular systems biology* **11**, 784 (2015).

30. Traxler, M. F. *et al.* The global, ppGpp-mediated stringent response to amino acid starvation in

Escherichia coli. *Molecular microbiology* **68**, 1128–1148 (2008).

31. Lutz, R. & Bujard, H. Independent and tight regulation of transcriptional units in Escherichia coli via

the LacR/O, the TetR/O and AraC/I1-I2 regulatory elements. *Nucleic acids research* **25**, 1203–1210

(1997).

32. Traxler, M. F. *et al.* Discretely calibrated regulatory loops controlled by ppGpp partition gene

induction across the 'feast to famine' gradient in Escherichia coli. *Mol. Microbiol.* **79**, 830–845

(2011).

33. Sendy, B., Lee, D. J., Busby, S. J. W. & Bryant, J. A. RNA polymerase supply and flux through the lac

operon in Escherichia coli. *Philos. Trans. R. Soc. Lond., B, Biol. Sci.* **371**, (2016).

34. Klumpp, S. & Hwa, T. Growth-rate-dependent partitioning of RNA polymerases in bacteria. *Proc*

Natl Acad Sci U S A **105**, 20245–50 (2008).

35. Berg, O. G. & von Hippel, P. H. Selection of DNA binding sites by regulatory proteins. Statistical-

mechanical theory and application to operators and promoters. *J. Mol. Biol.* **193**, 723–750 (1987).

36. Tsung, K., Inouye, S. & Inouye, M. Factors affecting the efficiency of protein synthesis in

Escherichia coli. Production of a polypeptide of more than 6000 amino acid residues. *Journal of*

Biological Chemistry **264**, 4428–4433 (1989).

37. Harvey, R. J. & Koch, A. L. How partially inhibitory concentrations of chloramphenicol affect the

growth of Escherichia coli. *Antimicrob. Agents Chemother.* **18**, 323–337 (1980).

38. Scott, M. & Hwa, T. Bacterial growth laws and their applications. *Curr. Opin. Biotechnol.* **22**, 559–

565 (2011).

39. Davies, J., Spiegelman, G. B. & Yim, G. The world of subinhibitory antibiotic concentrations. *Current*

Opinion in Microbiology **9**, 445–453 (2006).

40. Deris, J. B. *et al.* The Innate Growth Bistability and Fitness Landscapes of Antibiotic-Resistant

Bacteria. *Science* **342**, 1237435 (2013).

41. de Jong, H., Geiselmann, J. & Ropers, D. Resource Reallocation in Bacteria by Reengineering the

Gene Expression Machinery. *Trends Microbiol.* **25**, 480–493 (2017).

42. Hausser, J., Mayo, A., Keren, L. & Alon, U. Central dogma rates and the trade-off between precision

and economy in gene expression. *Nature Communications* **10**, 68 (2019).

43. Milon, P. *et al.* The nucleotide-binding site of bacterial translation initiation factor 2 (IF2) as a

metabolic sensor. *Proceedings of the National Academy of Sciences* **103**, 13962–13967 (2006).

44. Bakshi, S., Choi, H. & Weisshaar, J. C. The spatial biology of transcription and translation in rapidly

growing Escherichia coli. *Front Microbiol* **6**, 636 (2015).

45. Gausing, K. Regulation of ribosome synthesis in E. coli. *Trends in Biochemical Sciences* **7**, 65–67

(1982).

46. McGinnis, J. L. *et al.* In-cell SHAPE reveals that free 30S ribosome subunits are in the inactive state.

PNAS **112**, 2425–2430 (2015).

47. Fernandes, L. D., Moura, A. P. S. de & Ciandrini, L. Gene length as a regulator for ribosome

recruitment and protein synthesis: theoretical insights. *Scientific Reports* **7**, 17409 (2017).

48. Greulich, P., Scott, M., Evans, M. R. & Allen, R. J. Growth-dependent bacterial susceptibility to

ribosome-targeting antibiotics. *Mol Syst Biol* **11**, n/a-n/a (2015).

49. Lane, W. J. & Darst, S. A. Molecular Evolution of Multi-subunit RNA Polymerases: Sequence

Analysis. *J Mol Biol* **395**, 671 (2010).

50. Zakharova, N. *et al.* Fused and Overlapping rpoB and rpoC Genes in Helicobacters,

Campylobacters, and Related Bacteria. *Journal of Bacteriology* **181**, 3857 (1999).

51. Severinov, K. *et al.* Structural modules of the large subunits of RNA polymerase. Introducing

archaebacterial and chloroplast split sites in the beta and beta' subunits of Escherichia coli RNA

polymerase. *Journal of Biological Chemistry* **271**, 27969–74 (1996).

52. Davis, J. H. & Williamson, J. R. Structure and dynamics of bacterial ribosome biogenesis. *Phil. Trans.*

R. Soc. B **372**, 20160181 (2017).

53. Murakami, K. S. & Darst, S. A. Bacterial RNA polymerases: the wholo story. *Curr. Opin. Struct. Biol.*

13, 31–39 (2003).

54. Balaban, N. Q., Merrin, J., Chait, R., Kowalik, L. & Leibler, S. Bacterial persistence as a phenotypic

switch. *Science* **305**, 1622–1625 (2004).

55. Lewis, K. Persister cells. *Annu Rev Microbiol* **64**, 357–72 (2010).

56. Gruber, T. & Gross, C. Multiple sigma subunits and the partitioning of bacterial transcription space.

Annu Rev Microbiol **57**, 441–66 (2003).

57. Germain, E., Roghanian, M., Gerdes, K. & Maisonneuve, E. Stochastic induction of persister cells by

HipA through (p)ppGpp-mediated activation of mRNA endonucleases. *Proc Natl Acad Sci U S A* **112**,

5171–5176 (2015).

58. Balaban, N. Persistence: mechanisms for triggering and enhancing phenotypic variability. *Current*

opinion in genetics & development **21**, 768–75 (2011).

59. Kudrin, P. *et al.* Subinhibitory Concentrations of Bacteriostatic Antibiotics Induce relA-Dependent

and relA-Independent Tolerance to β-Lactams. *Antimicrobial Agents and Chemotherapy* **61**,

e02173-16 (2017).

60. Cormack, B. P., Valdivia, R. H. & Falkow, S. FACS-optimized mutants of the green fluorescent

protein (GFP). *Gene* **173**, 33–38 (1996).

61. MacGregor, G. R. & Caskey, C. T. Construction of plasmids that express E. coli beta-galactosidase in

mammalian cells. *Nucleic Acids Res* **17**, 2365 (1989).

62. Li, R., Zhang, Q., Li, J. & Shi, H. Effects of cooperation between translating ribosome and RNA

polymerase on termination efficiency of the Rho-independent terminator. *Nucleic acids research*

gkv1285 (2015).

63. Shine, J. & Dalgarno, L. The 3′-Terminal Sequence of Escherichia coli 16S Ribosomal RNA:

Complementarity to Nonsense Triplets and Ribosome Binding Sites. *Proc Natl Acad Sci U S A* **71**,

1342–1346 (1974).

64. Zhang, Q., Li, R., Li, J. & Shi, H. Optimal Allocation of Bacterial Protein Resources under Nonlethal

Protein Maturation Stress. *Biophys. J.* **115**, 896–910 (2018).

65. Zhang, X. & Bremer, H. Control of the Escherichia coli rrnB P1 promoter strength by ppGpp. *Journal*

of Biological Chemistry **270**, 11181–11189 (1995).

66. Shetty, R. P. Applying engineering principles to the design and construction of transcriptional

devices. (Massachusetts Institute of Technology, 2008).

67. Rudge, T. J. *et al.* Characterization of Intrinsic Properties of Promoters. *ACS Synth Biol* **5**, 89–98

(2016).